

ON BOUNDARY – VALUE PROBLEMS FOR THE LAPLACIAN IN BOUNDED AND IN UNBOUNDED DOMAINS WITH PERFORATED BOUNDARIES

Gregory A.Chechkin[‡], Rustem R.Gadyl'shin^{‡#}

[‡] Department of Differential Equations
Faculty of Mechanics and Mathematics
Moscow State University
Moscow 119899, Russia
checkkin@mech.math.msu.su

[‡] Institute of Mathematics with Computing Center
Russian Academy of Sciences
Ufa 450077, Russia

[#] Department of Mathematical Analysis
Faculty of Physics and Mathematics
Bashkir State Pedagogical University
Ufa 450000, Russia
gadylshin@bspu.ru

CORRESPONDENCE ADDRESS:

Department of Differential Equations,
Faculty of Mechanics and Mathematics
Moscow State University, Moscow 119899, Russia.
checkkin@mech.math.msu.su
Phone: +7-(095)-113-4535 Fax: +7-(095)-939-2090

running head: ON BOUNDARY – VALUE PROBLEMS ...

The research of the first author was partially supported by Russian Foundation for Basic Research (RFBR, project No 02-01-00868). The second author was partially supported by Russian Foundation for Basic Research (RFBR, project No 02-01-00693) and Scientific programm “Universities of Russia”

Abstract.

In the paper we consider boundary – value problems with rapidly alternating type of boundary conditions, including problems in domains with perforated boundaries. We present the classification of homogenized (limit) problems depending on the ratio of small parameters, which characterize the diameter of parts of the boundary with different types of boundary conditions or on the ratio of small parameters, which characterize the diameter and the distance between holes. Also we studied the analogue of the Helmholtz resonator for domains with perforated boundary.

2000. *Mathematical Subject Classification.* 35B25, 35B27, 35J25

Keywords. homogenization, rapidly alternating boundary conditions, perforated boundary, spectral problems

0 Introduction

We deal with boundary – value problems with rapidly alternating type of boundary conditions and problems in domains with perforated boundaries. Problems of this kind attracted the attention of mathematicians from the mid-1960-s (see, for instance, [1] – [32]). Such problems appear in physics and engineering sciences, when one studies, for example, the scattering of acoustic waves on the small periodic obstacles, the behavior of partially fastened membranes and many others. The engineering applications of such problems could be also found in construction of atomic power stations in space antennas etc. One can study the problem of permeation of fuel through the walls of plastic tank. In order to reduce permeation of fuel, the inner boundary of the container is coated with thin barrier layer of fluorine by a blow molding process. The resulting thin layer, however, typically has flaws: it leaves many small patches uncovered. This model is described in more details in [33] and [34].

In this work we consider boundary – value problem in a 3D domain for the Laplacian. We assume that the boundary of the domain consists of two parts. One of them has purely periodic microstructure. It could be rapidly alternating spots or periodically situated holes. In the first case we have bounded domain with micro inhomogeneous structure of the boundary and in the second one we have two domains connected through the holes. In the second case we study problems in both bounded domains or in one bounded and one unbounded domains. We give a complete classification of homogenized problems in their dependence on the ratio of the small parameters, characterizing the frequency of the periodical change of the boundary conditions in the first case or on the ratio of the small parameters characterizing the diameter and the distance between holes in the second case.

It should be noted that in the first case on the base of other methods the convergence of solutions to such problems was proved in [12] for more general situation. Also nonperiodic boundary structure was considered in [19] and [13]. On the other hand, the direct combination of the approaches from [11] (see, also [10]) and [35] (see, also [36]) gives an opportunity to obtain the estimate for the rate of convergence for solutions in the periodical situation (see, also [17]). In this paper for the periodical boundary microstructure we demonstrate the much shorter proof (than in [12] and in [11] and [17]) of the convergence theorem. We use the homogenization methods [37] – [41] and the method of matching asymptotic expansions [42] – [44]. More precisely, we arrange the approaches of [11] and [35] in the most rational form. This combination allows to get clear expressible formulae in a short way.

Also we study problems in domains with perforated boundaries. Similar problems were considered in [1], [45] – [49]. In present paper we develop the rational approach (mentioned above) for problems in domains with periodically perforated boundaries. We find the ratio of the small parameters characterizing the diameter and the distance between holes, that implies in the limit the decomposition of the original problem to two independent problems. In the case of unbounded external domain we show that the decomposition involves the appearance of poles (scattering frequencies) with small imaginary part of the analytical continuation of solutions to the original problem. It is wellknown that such poles for the Helmholtz resonator do exist (see, for instance, [50] – [53]). Namely these poles induce the resonance in the Helmholtz resonator [50], [54] – [57]. Remind that the classical Helmholtz resonator could be described by the boundary – value problem for the Helmholtz equation in an unbounded domain outside the surface with a small aperture [58], [59]. The model 2-dimensional analogue of the Helmholtz resonator in periodically perforated domain was considered in [21],[30], [31]. The author discovered the resonances for this analogue of the Helmholtz resonator. In this paper we consider 3-dimensional analogue of the Helmholtz resonator in homogenization theory. We proved the existence of the scattering frequencies with small imaginary part.

In the first section we introduce notation, describe the domains, set the problems and formulate six basic theorems and one auxiliary theorem which is of independent interest. The last theorem is proved in the Section 2.

Problems with rapidly alternating boundary conditions is considered in Sections 3 and 4.

Sections 5 – 8 are devoted to investigations of problems in domains with perforated boundaries. In Sections 5, 6 we consider the case when connected domains are bounded and in Sections 7, 8 we study the case when one of the domains is unbounded, a 3D analogue of the Helmholtz resonator in homogenization.

1 Statements

Let Ω be a bounded domain in \mathbb{R}^3 with a C^∞ -boundary Γ . We suppose that Ω lies in the half-space $x_3 < 0$, $\Gamma_1 = \text{int}(\{x : x_3 = 0\} \cap \Gamma)$, $\text{mes}_2\Gamma_1 \neq 0$. Denote by ω a two-dimensional bounded domain with a smooth boundary on the plane $x_3 = 0$. We suppose that $0 < \varepsilon, \delta \ll 1$ are small parameters. Introduce the following notation: $\omega_\varepsilon = \{x : x\varepsilon^{-1} \in \omega\}$, $\Pi_\varepsilon = \{x : x = (2n, 2m, 0) + x', x' \in \omega_\varepsilon, n, m \in \mathbb{Z}\}$, $\Pi_\varepsilon^\delta = \{x : \delta^{-1}x \in \Pi_\varepsilon\}$, $\Gamma_2 = \Gamma \setminus \overline{\Gamma_1}$, $\Gamma_{\varepsilon,\delta}^D = \Gamma_1 \cap \Pi_\varepsilon^\delta$, and $\Gamma_{\varepsilon,\delta}^S = \Gamma_1 \setminus \overline{\Gamma_{\varepsilon,\delta}^D}$ (see Fig. 1.).




Figure 1:

We consider the case when $\delta = \delta(\varepsilon)$ depends on ε and

$$\lim_{\varepsilon \rightarrow 0} \frac{\varepsilon}{\delta(\varepsilon)} = p, \quad p \in [0, \infty].$$

Our goal of the first part of the work is to prove the following two auxiliary statement:

Theorem 1.1. *Let $f \in L_2(\Omega)$, $q \geq 0$, $p = \infty$. Then the solution of the boundary value problem*

$$\begin{cases} \Delta u_{\varepsilon,\delta} = f, & x \in \Omega, \\ u_{\varepsilon,\delta} = 0, & x \in \Gamma_{\varepsilon,\delta}^D \cup \Gamma_2, \\ \frac{\partial u_{\varepsilon,\delta}}{\partial x_3} + qu_{\varepsilon,\delta} = 0, & x \in \Gamma_{\varepsilon,\delta}^S \end{cases} \quad (1.1)$$

converges to the solution of the boundary - value problem

$$\begin{cases} \Delta u_0 = f, & x \in \Omega, \\ u_0 = 0, & x \in \Gamma \end{cases} \quad (1.2)$$

in $H^1(\Omega)$.

Theorem 1.2. *Let $f \in L_2(\Omega)$, $q \geq 0$, $p < \infty$, $Q = q + c_\omega p$. Then the solution of (1.1) converges to the solution of the boundary - value problem*

$$\begin{cases} \Delta u_0 = f, & x \in \Omega, \\ u_0 = 0, & x \in \Gamma_2, \\ \frac{\partial u_0}{\partial x_3} + Qu_0 = 0, & x \in \Gamma_1. \end{cases} \quad (1.3)$$

in $H^1(\Omega)$, if $p = 0$, and weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$, if $p > 0$.

Hereafter, c_ω is the capacity of the plate ω ([60], [61]). This wellknown constant is positive and, for instance, if ω is the unit disk, then $c_\omega = 2\pi^{-1}$ (see, [60]).

Later on we consider problems in domains with perforated boundary. In Sections 5 and 6 we study problems in bounded domains.

Let $\tilde{\Omega}$ be a bounded domain in \mathbb{R}^3 with a C^∞ -boundary $\tilde{\Gamma}$ such that $\bar{\Omega} \subset \tilde{\Omega}$ (see Fig. 2.).




Figure 2:

Theorem 1.3. Let $p = \infty$, $F \in L_2(\tilde{\Omega})$, f and \tilde{f} be the restrictions of F in Ω and in $\tilde{\Omega} \setminus \bar{\Omega}$, respectively. Then the solution of the boundary – value problem

$$\begin{cases} \Delta u_{\varepsilon,\delta} = F, & x \in \tilde{\Omega} \setminus \overline{\Gamma_{\varepsilon,\delta}^D \cup \Gamma_2}, \\ u_{\varepsilon,\delta} = 0, & x \in \Gamma_{\varepsilon,\delta}^D \cup \Gamma_2 \cup \tilde{\Gamma}, \end{cases} \quad (1.4)$$

converges strongly in $H^1(\tilde{\Omega})$ as $\varepsilon \rightarrow 0$ to the function

$$u(x) = \begin{cases} u_0(x), & x \in \Omega, \\ \tilde{u}_0(x), & x \in \tilde{\Omega} \setminus \bar{\Omega}, \end{cases}$$

where $u_0(x)$ is a solution of Problem (1.2) and $\tilde{u}_0(x)$ satisfies the boundary – value problem

$$\begin{cases} \Delta \tilde{u}_0 = \tilde{f}, & x \in \tilde{\Omega} \setminus \bar{\Omega}, \\ \tilde{u}_0 = 0, & x \in \Gamma \cup \tilde{\Gamma}. \end{cases} \quad (1.5)$$

Theorem 1.4. Let $p = 0$, $F \in L_2(\tilde{\Omega})$, f and \tilde{f} be the restrictions of F in Ω and in $\tilde{\Omega} \setminus \bar{\Omega}$, respectively. Then the solution of the boundary – value problem

$$\begin{cases} \Delta u_{\varepsilon,\delta} = F, & x \in \tilde{\Omega} \setminus \overline{(\Gamma_2 \cup \Gamma_{\varepsilon,\delta}^S)}, \\ u_{\varepsilon,\delta} = 0, & x \in \Gamma_2 \cup \tilde{\Gamma}, \\ \frac{\partial u_{\varepsilon,\delta}}{\partial x_3} = 0, & x \in \Gamma_{\varepsilon,\delta}^S, \end{cases} \quad (1.6)$$

converges strongly in $H^1(\tilde{\Omega} \setminus \bar{\Gamma}_1)$ as $\varepsilon \rightarrow 0$ to the function

$$u(x) = \begin{cases} u_0(x), & x \in \Omega, \\ \tilde{u}_0(x), & x \in \tilde{\Omega} \setminus \bar{\Omega}, \end{cases}$$

where $u_0(x)$ is a solution of the boundary – value problem

$$\begin{cases} \Delta u_0 = f, & x \in \Omega, \\ u_0 = 0, & x \in \Gamma_2, \\ \frac{\partial u_0}{\partial x_3} = 0, & x \in \Gamma_1, \end{cases} \quad (1.7)$$

and $\tilde{u}_0(x)$ is a solution of the boundary – value problem

$$\begin{cases} \Delta \tilde{u}_0 = \tilde{f}, & x \in \tilde{\Omega} \setminus \bar{\Omega}, \\ \tilde{u}_0 = 0, & x \in \Gamma_2 \cup \tilde{\Gamma}, \\ \frac{\partial \tilde{u}_0}{\partial x_3} = 0, & x \in \Gamma_1. \end{cases} \quad (1.8)$$

Denote by σ the square $(-1, 1) \times (-1, 1)$ in the plane $x_3 = 0$ and let $\Sigma = \sigma \times (0, -\infty)$. Denote by $C^\infty(\overline{\Sigma}, \omega_\varepsilon)$ the set of C^∞ -functions vanishing in a neighborhood of ω_ε and having the finite Dirichlet integral. We define the space $\tilde{H}^1(\Sigma; \omega_\varepsilon)$ as the closure of $C^\infty(\overline{\Sigma})$ by the norm

$$\|v\|_1 = \left(\int_{\Sigma} |\nabla v|^2 dx + \int_{\sigma} v^2 ds \right)^{1/2}.$$

The proofs of Theorems 1.1 and 1.3 are based on the following statement.

Theorem 1.5. *Asymptotics of*

$$\lambda_\varepsilon = \inf_{v \in \tilde{H}^1(\Sigma; \omega_\varepsilon) \setminus \{0\}} \frac{\int_{\Sigma} |\nabla v|^2 dx}{\int_{\sigma} v^2 ds} \quad (1.9)$$

reads as follows:

$$\lambda_\varepsilon = \varepsilon \frac{\pi c_\omega}{2} + o(\varepsilon).$$

The last part of the paper is devoted to the investigation of problems in unbounded domains. Assume that F is a function from $L_2(\mathbb{R}^3)$ with bounded support. We consider the following boundary – value problems:

$$\begin{cases} (\Delta + k^2) u_{\varepsilon, \delta} = F, & x \in \mathbb{R}^3 \setminus \overline{\Gamma_{\varepsilon, \delta}^D \cup \Gamma_2}, \\ u_{\varepsilon, \delta} = 0, & x \in \Gamma_{\varepsilon, \delta}^D \cup \Gamma_2, \end{cases} \quad (1.10)$$

$$\begin{cases} (\Delta + k^2) u_{\varepsilon, \delta} = F, & x \in \mathbb{R}^3 \setminus \overline{(\Gamma_2 \cup \Gamma_{\varepsilon, \delta}^S)}, \\ u_{\varepsilon, \delta} = 0, & x \in \Gamma_2, \quad \frac{\partial u_{\varepsilon, \delta}}{\partial x_3} = 0, \quad x \in \Gamma_{\varepsilon, \delta}^S, \end{cases} \quad (1.11)$$

with the radiation condition

$$u_{\varepsilon, \delta} = O(r^{-1}), \quad \frac{\partial u_{\varepsilon, \delta}}{\partial r} - ik u_{\varepsilon, \delta} = o(r^{-1}), \quad r \rightarrow \infty. \quad (1.12)$$

for k such that $\text{Im } k \geq 0$. Here and throughout $r = |x|$.

Theorem 1.6. *Let $p = \infty$. Suppose also that f and \tilde{f} are the restrictions of F in Ω and in $\mathbb{R}^3 \setminus \overline{\Omega}$, respectively. Then the solution to Problem (1.10), (1.12) converges strongly in $H_{\text{loc}}^1(\mathbb{R}^3)$ as $\varepsilon \rightarrow 0$ to the function*

$$u(x) = \begin{cases} u_0(x), & x \in \Omega, \\ \tilde{u}_0(x), & x \in \tilde{\Omega} \setminus \overline{\Omega}, \end{cases}$$

where $u_0(x)$ is a solution of the boundary – value problem

$$\begin{cases} -\Delta u_0 = k^2 u_0 - f, & x \in \Omega, \\ u_0 = 0, & x \in \Gamma, \end{cases} \quad (1.13)$$

and $\tilde{u}_0(x)$ is a solution of the boundary – value problem

$$\begin{cases} (\Delta + k^2) \tilde{u}_0 = \tilde{f}, & x \in \mathbb{R}^3 \setminus \bar{\Omega}, \\ \tilde{u}_0 = 0, & x \in \Gamma, \end{cases} \quad (1.14)$$

with the radiation condition

$$\tilde{u}_0 = O(r^{-1}), \quad \frac{\partial \tilde{u}_0}{\partial r} - ik\tilde{u}_0 = o(r^{-1}), \quad r \rightarrow \infty. \quad (1.15)$$

Here it is assumed that k^2 is not an eigenvalue of Problem (1.13).

If $k^2 = k_0^2$ is an eigenvalue to Problem (1.13), then there is a pole τ_ε of the analytic continuation of the solution of (1.10), (1.12) in the half plane $\text{Im } k < 0$, converging to k_0 as $\varepsilon \rightarrow 0$.

Theorem 1.7. Let $p = 0$, f and \tilde{f} be the restrictions of F in Ω and in $\mathbb{R}^3 \setminus \bar{\Omega}$, respectively. Then the solution to Problem (1.11), (1.12) converges strongly in $H_{\text{loc}}^1(\mathbb{R}^3 \setminus \bar{\Gamma}_1)$ as $\varepsilon \rightarrow 0$ to the function

$$u(x) = \begin{cases} u_0(x), & x \in \Omega, \\ \tilde{u}_0(x), & x \in \tilde{\Omega} \setminus \bar{\Omega}, \end{cases}$$

where $u_0(x)$ is a solution of the boundary – value problem

$$\begin{cases} -\Delta u_0 = k^2 u_0 - f, & x \in \Omega, \\ u_0 = 0, & x \in \Gamma_2, \\ \frac{\partial u_0}{\partial x_3} = 0, & x \in \Gamma_1 \end{cases} \quad (1.16)$$

and $\tilde{u}_0(x)$ is a solution of the boundary – value problem

$$\begin{cases} (\Delta + k^2) \tilde{u}_0 = \tilde{f}, & x \in \mathbb{R}^3 \setminus \bar{\Omega}, \\ \tilde{u}_0 = 0, & x \in \Gamma_2, \quad \frac{\partial \tilde{u}_0}{\partial x_3} = 0, & x \in \Gamma_1 \end{cases} \quad (1.17)$$

with the radiation condition (1.15).

Here it is assumed that k^2 is not an eigenvalue of Problem (1.16).

If $k^2 = k_0^2$ is an eigenvalue of the boundary – value problem (1.16), then there is a pole τ_ε of the analytic continuation of the solution of (1.11), (1.12) in the half plane $\text{Im } k < 0$, converging to k_0 as $\varepsilon \rightarrow 0$.

The notion of an analytic continuation is classical and we shall give all the necessary definitions in section 7.

2 Proof of Theorem 1.5

Suppose that $G \in C^\infty(\overline{\Sigma} \setminus \{0\})$ is a bounded σ -periodic solution of the boundary – value problem

$$\Delta G = 0, \quad x \in \Sigma, \quad \frac{\partial G}{\partial x_3} = -\frac{1}{4}, \quad x \in \sigma \setminus \{0\}, \quad (2.1)$$

with the asymptotics $G(x) \sim (2\pi r)^{-1}$ as $r = |x| \rightarrow 0$. The existence theorem of this Green function is wellknown. In more detail the asymptotics of G reads as follows (see, for instance, [62]):

$$\begin{aligned} G(x) &= C_\Sigma + O\left(\exp\left\{\frac{\pi x_3}{2}\right\}\right), \quad x_3 \rightarrow -\infty, \\ G(x) &= \frac{1}{2\pi r} - \frac{1}{4}x_3 + O(r^2), \quad r \rightarrow 0, \end{aligned} \quad (2.2)$$

where C_Σ is a fixed constant. It is easy to show that the right-hand side of the asymptotics (2.2) are a differentiable function. In view of the evenness the function G satisfies the homogeneous Neumann boundary condition in $\partial\Sigma \setminus \overline{\sigma}$.

It is known (see, for instance [63]) that there exists a harmonic in $\mathbb{R}_-^3 = \{x : x_3 < 0\}$ function $X_0 \in H_{loc}^1(\mathbb{R}_-^3) \cap C^\infty(\overline{\mathbb{R}_-^3} \setminus \partial\omega)$ which vanishes at infinity and satisfies the boundary conditions $X_0 = 1$ in ω and $\partial X_0 / \partial x_3 = 0$ on $\gamma = \{x : x_3 = 0, (x_1, x_2) \notin \overline{\omega}\}$. In addition the function X_0 has the differentiable asymptotics

$$X_0(x) = c_\omega r^{-1} + \sum_{i=1}^2 c_i \frac{\partial}{\partial x_i} r^{-1} + O(r^{-3}) \quad \text{as } r \rightarrow \infty. \quad (2.3)$$

It should be noted, that the capacity of ω is defined namely as the coefficient of r^{-1} in (2.3).

Denote by $\chi(t)$ a smooth cut-off function equals to one for $t < 1/3$ and equals to zero for $t > 2/3$ and suppose that

$$\widetilde{W}^\varepsilon(x, \beta) = 1 - \varepsilon 2\pi c_\omega (1 - \chi(r\varepsilon^{-\beta})) G(x) - \chi(r\varepsilon^{-\beta}) X_0(x\varepsilon^{-1}),$$

where $\beta > 0$. Taking into account (2.1)–(2.3), one can obtain the following statement.

Lemma 2.1. *The function $\widetilde{W}^\varepsilon \in \widetilde{H}^1(\Sigma; \omega_\varepsilon) \cap C^\infty(\overline{\Sigma} \setminus \partial\omega_\varepsilon)$ satisfies the estimates*

$$\|\widetilde{W}^\varepsilon - 1\|_{L_2(\sigma)} = o(1), \quad \varepsilon \rightarrow 0, \quad |\nabla \widetilde{W}^\varepsilon| = O\left(\exp\left\{\frac{\pi x_3}{2}\right\}\right), \quad x_3 \rightarrow -\infty$$

and it is a solution to the boundary - value problem

$$\begin{cases} \Delta \widetilde{W}^\varepsilon = \widetilde{F}_\varepsilon, & x \in \Sigma, \\ \frac{\partial \widetilde{W}^\varepsilon}{\partial x_3} = \varepsilon \frac{\pi}{2} c_\omega \widetilde{W}^\varepsilon + \widetilde{h}_\varepsilon, & x \in \sigma \setminus \overline{\omega_\varepsilon}, \\ \widetilde{W}^\varepsilon = 0, & x \in \omega_\varepsilon, \\ \frac{\partial \widetilde{W}^\varepsilon}{\partial \nu} = 0, & x \in \partial \Sigma \setminus \overline{\sigma}, \end{cases}$$

where $\text{supp} \widetilde{F}_\varepsilon$ lies above the plane $x_3 = -\frac{2}{3}\varepsilon^\beta$.

If $\beta < 2/5$, then

$$\|\widetilde{F}_\varepsilon\|_{L_2(\Sigma)} + \|\widetilde{h}_\varepsilon\|_{L_2(\sigma)} = o(\varepsilon), \quad \varepsilon \rightarrow 0.$$

By a standard way (see, for instance, [11]) it is easy to show that infimum in (1.9) is attained at some harmonic function W^ε and

$$\int_{\Sigma} (\nabla W^\varepsilon, \nabla v) dx = \lambda_\varepsilon \int_{\sigma} W^\varepsilon v ds \quad (2.4)$$

for any $v \in \widetilde{H}^1(\Sigma; \omega_\varepsilon)$, where λ_ε is defined in (1.9).

Due to Lemma 2.1 we have

$$\int_{\Sigma} \widetilde{F}_\varepsilon W^\varepsilon dx - \int_{\sigma} \widetilde{h}_\varepsilon W^\varepsilon ds = - \int_{\Sigma} (\nabla \widetilde{W}^\varepsilon, \nabla W^\varepsilon) dx + \varepsilon \frac{\pi c_\omega}{2} \int_{\sigma} \widetilde{W}^\varepsilon W^\varepsilon ds. \quad (2.5)$$

From (2.4) and (2.5) we deduce the expression

$$\int_{\Sigma} \widetilde{F}_\varepsilon W^\varepsilon dx - \int_{\sigma} \widetilde{h}_\varepsilon W^\varepsilon ds = \left(\varepsilon \frac{\pi c_\omega}{2} - \lambda_\varepsilon \right) \int_{\sigma} \widetilde{W}^\varepsilon W^\varepsilon ds. \quad (2.6)$$

Denote by τ the maximum $\max_{x' \in \omega} \{|x'|\}$. Obviously, the function

$$w^\varepsilon(x) = 1 - \chi(3\tau\varepsilon^{-1}r)$$

belongs to $\widetilde{H}^1(\Sigma; \omega_\varepsilon)$ and $\|\nabla w^\varepsilon\|_{L_2(\Sigma)} + \|w^\varepsilon - 1\|_{L_2(\sigma)} \rightarrow 0$ as $\varepsilon \rightarrow 0$. Hence,

$$\lambda_\varepsilon \rightarrow 0, \quad \varepsilon \rightarrow 0. \quad (2.7)$$

We suppose that W^ε is normalized, i.e. $\int_{\sigma} (W^\varepsilon)^2 ds = 1$. It should be noted that (2.7) and (1.9) lead to $\|\nabla W^\varepsilon\|_{L_2(\Sigma)} \rightarrow 0$. Hence, for any fixed $R < 0$,

there exists a representation $W^\varepsilon(x) = w_R^\varepsilon(x) + c_R(\varepsilon)$, where $\|w_R^\varepsilon\|_{H^1(\Sigma_R)} \rightarrow 0$ as $\varepsilon \rightarrow 0$ and $\Sigma_R = \Sigma \cup \{x : x_3 > R\}$. Therefore, $\|w_R^\varepsilon\|_{L_2(\sigma)} \rightarrow 0$ as $\varepsilon \rightarrow 0$ and then $c_R(\varepsilon) \rightarrow 1$. The latter convergences imply that

$$\|W^\varepsilon\|_{L_2(\Sigma_R)} \leq C_R, \quad \|W^\varepsilon - 1\|_{L_2(\sigma)} \rightarrow 0, \quad \varepsilon \rightarrow 0, \quad (2.8)$$

where the constant C_R is independent of ε . Now (2.6), (2.8) and Lemma 2.1 (with $\beta < 2/5$) prove Theorem 1.5.

3 Proof of Theorem 1.1

We consider Problem (1.1)–(1.8) in a weak sense (see, for instance, [64]).

Definition 3.1. Denote by $C^\infty(Q; S)$ the set of functions from $C^\infty(\overline{Q})$ vanishing in a neighborhood of $S \subset \overline{Q}$.

Define the space $H^1(Q; S)$ as the closure of the set of functions from $C^\infty(Q; S)$ by the norm of the Sobolev space $H^1(Q)$.

Definition 3.2. A function $u_{\varepsilon, \delta} \in H^1(\Omega; \Gamma_{\varepsilon, \delta}^D \cup \Gamma_2)$ is called a solution of (1.1), if there holds the integral identity

$$\int_{\Omega} \nabla u_{\varepsilon, \delta} \nabla v \, dx + q \int_{\Gamma_{\varepsilon, \delta}^S} u_{\varepsilon, \delta} v \, ds = - \int_{\Omega} f v \, dx \quad (3.1)$$

for any $v \in H^1(\Omega; \Gamma_{\varepsilon, \delta}^D \cup \Gamma_2)$.

In a similar way we define solutions of the boundary – value problem (1.2).

Definition 3.3. The function $u_0 \in H^1(\Omega; \Gamma)$ is a solution of (1.2), if it satisfies the identity

$$\int_{\Omega} \nabla u_0 \nabla v \, dx = - \int_{\Omega} f v \, dx, \quad (3.2)$$

for any $v \in H^1(\Omega; \Gamma)$.

Since the usual H^1 -norm and the norm $\|u\|'_{H^1(\Omega)} = \|\nabla u\|_{L_2(\Omega)}$ are equivalent in $H^1(\Omega; \Gamma_2)$ and $H^1(\Omega; \Gamma_{\varepsilon, \delta}^D \cup \Gamma_2) \subset H^1(\Omega; \Gamma_2)$, we deduce from (3.1) the uniform estimate

$$\|u_{\varepsilon, \delta}\|_{H^1(\Omega)} \leq C \|f\|_{L_2(\Omega)}. \quad (3.3)$$

Lemma 3.1. *For any $v \in H^1(\Omega; \Gamma_{\varepsilon, \delta}^D \cup \Gamma_2)$ the following estimate holds:*

$$\|v\|_{L_2(\Gamma)} \leq C \left(\frac{\delta}{\varepsilon} \right)^{1/2} \|v\|_{H^1(\Omega)}. \quad (3.4)$$

Proof. Consider the extension of v by zero into $\mathbb{R}_-^3 \setminus \Omega$. Obviously, this extension belongs to $H^1(\mathbb{R}_-^3)$. Assume that $\sigma_\delta^{(m,n)} = \{x : x = \delta(2m, 2n, 0) + x', x' \delta^{-1} \in \sigma\}$, $\Sigma_\delta^{(m,n)} = \sigma_\delta^{(m,n)} \times (-\infty, 0)$. Denote by Σ_δ^∂ the union of $\Sigma_\delta^{(m,n)}$ such that $\sigma_\delta^{(m,n)} \cap \partial\Gamma_1 \neq \emptyset$, by Σ_δ^{in} the union of $\Sigma_\delta^{(m,n)}$ such that $\sigma_\delta^{(m,n)} \subset \Gamma_1$, and by Σ_δ^{ex} the union of $\Sigma_\delta^{(m,n)}$ such that $\sigma_\delta^{(m,n)} \cap \Gamma_1 = \emptyset$. Then

$$\|v\|_{L_2(\sigma_\delta^{(m,n)})} \leq C \left(\frac{\delta}{\varepsilon} \right)^{1/2} \|\nabla v\|_{L_2(\Sigma_\delta^{(m,n)})}, \quad \Sigma_\delta^{(m,n)} \subset \Sigma_\delta^{ex} \quad (3.5)$$

and, due to (1.9) and Theorem 1.5 the following estimate takes place:

$$\|v\|_{L_2(\sigma_\delta^{(m,n)})} \leq C \left(\frac{\delta}{\varepsilon} \right)^{1/2} \|\nabla v\|_{L_2(\Sigma_\delta^{(m,n)})}, \quad \Sigma_\delta^{(m,n)} \subset \Sigma_\delta^{in}. \quad (3.6)$$

Now, suppose that $\Sigma_\delta^{(m,n)} \subset \Sigma_\delta^\partial$, $\omega_\varepsilon^\delta = \{x : x = \delta(2m, 2n, 0) + x', x' \delta^{-1} \in \omega\}$, and $\tilde{\omega}_\varepsilon^\delta$ are the subset of $\sigma_\delta^{(m,n)}$, where the function v equals to zero. Since, $\omega_\varepsilon^\delta \subset \tilde{\omega}_\varepsilon^\delta$, then (1.9) and Theorem 1.5 imply that

$$\|v\|_{L_2(\sigma_\delta^{(m,n)})} \leq C \left(\frac{\delta}{\varepsilon} \right)^{1/2} \|\nabla v\|_{L_2(\Sigma_\delta^{(m,n)})}, \quad \Sigma_\delta^{(m,n)} \subset \Sigma_\delta^\partial. \quad (3.7)$$

Estimates (3.5)–(3.7) give estimate (3.4). Lemma is proved.

Let $p = \infty$, suppose also that v is a function from $C_0^\infty(\Omega)$ and $\{\varepsilon_n\}$ is a sequence which tends to zero as $n \rightarrow \infty$. Due to the embedding theorems, weak compactness of the bounded set in $H^1(\Omega)$ and estimates (3.3), (3.4) there exists a subsequence of this sequence such that $u_{\varepsilon, \delta} \rightarrow u_0 \in H^1(\Omega; \Gamma)$ at this subsequence weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$. Passing to the limit in (3.1) on this subsequence we obtain (3.2). Hence, u_0 is the solution of (1.2). On the other hand, due to the arbitrariness of choosing of the sequence of $\{\varepsilon_n\}$, we have that $u_{\varepsilon, \delta} \rightarrow u_0$ as $\varepsilon \rightarrow 0$ (weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$). Taking in account the latter convergences and coming to the limit in (3.1) with $v = u_{\varepsilon, \delta}$, we obtain convergence

$$\|u_{\varepsilon, \delta}\|_{H^1(\Omega)} \rightarrow \|u_0\|_{H^1(\Omega)}.$$

Hence, $u_{\varepsilon, \delta} \rightarrow u_0$ as $\varepsilon \rightarrow 0$ strongly in $H^1(\Omega)$. Theorem is proved.

4 Proof of Theorem 1.2

Let us introduce the notation

$$W_\varepsilon(x; \beta) = \widetilde{W}^\varepsilon(x; \beta) - \varepsilon^2 2\pi (1 - \chi(r\varepsilon^{-\beta})) \sum_{i=1}^2 c_i \frac{\partial}{\partial x_i} G(x) + \varepsilon \frac{\pi c_\omega}{2} \chi(r\varepsilon^{-\beta}) x_3.$$

After this correction of the function $\widetilde{W}^\varepsilon(x; \beta)$ we obtain the following statement.

Lemma 4.1. *The σ -periodic function $W_\varepsilon(x; \beta) \in H_{loc}^1(\Sigma) \cap C^\infty(\overline{\Sigma} \setminus \partial\omega_\varepsilon)$ satisfies the estimates*

$$\begin{aligned} \|W_\varepsilon - 1 + \varepsilon 2\pi c_\omega C_\Sigma\|_{L_2(\Sigma)} &= o(1), \quad \varepsilon \rightarrow 0, \\ |\nabla W_\varepsilon| + |W_\varepsilon - 1 + \varepsilon 2\pi c_\omega C_\Sigma| &= O\left(\exp\left\{\frac{\pi x_3}{2}\right\}\right), \quad x_3 \rightarrow -\infty, \quad \varepsilon \rightarrow 0 \end{aligned}$$

and W_ε is a solution to the boundary - value problem

$$\begin{cases} \Delta W_\varepsilon = F_\varepsilon, & x \in \Sigma, \\ \frac{\partial W_\varepsilon}{\partial x_3} = \varepsilon \frac{\pi}{2} c_\omega W_\varepsilon + h_\varepsilon, & x \in \sigma \setminus \overline{\omega_\varepsilon}, \\ W_\varepsilon = 0, & x \in \omega_\varepsilon, \end{cases}$$

where $\text{supp} F_\varepsilon$ lies above the plane $x_3 = -\frac{2}{3}\varepsilon^\beta$.

If $\frac{1}{5} < \beta < \frac{3}{7}$, then

$$\|F_\varepsilon\|_{L_2(\Sigma)} = o(\varepsilon^{3/2}), \quad \|h_\varepsilon\|_{L_2(\sigma)} = o(\varepsilon).$$

Remark 4.1. We use our correction only to improve estimates of the right-hand side of the equation. This estimate will be used in the proof of Theorem 1.2.

Denote by $W_\varepsilon^\delta(x)$ the expression

$$1 + \chi(-x_3 \delta^{-1/2}) \left(\frac{1}{1 - \varepsilon 2\pi c_\omega C_\Sigma} W_\varepsilon \left(\frac{x}{\delta}; \frac{1}{4} \right) - 1 \right).$$

Lemma 4.1 implies the following result, which is a key to prove Theorem 1.2.

Lemma 4.2. *If $\varepsilon \rightarrow 0$ and $\delta \rightarrow 0$, then*

$$\begin{aligned} \|\Delta W_\varepsilon^\delta\|_{L_2(\Omega)} &= o\left(\left(\frac{\varepsilon}{\delta}\right)^{3/2}\right) + o(1), & \|W_\varepsilon^\delta - 1\|_{L_2(\Omega)} &= o(1), \\ \left\| \frac{\partial W_\varepsilon^\delta}{\partial x_3} - \frac{\pi c_\omega \varepsilon}{2\delta} \right\|_{L_2(\Gamma_1)} &= o\left(\frac{\varepsilon}{\delta}\right). \end{aligned}$$

Note, we consider Problem (1.3) in a distributional sense. Therefore, a function $u_{\varepsilon,\delta} \in H^1(\Omega; \Gamma_2)$ is called a solution of (1.3), if there holds the integral identity

$$\int_{\Omega} \nabla u_0 \nabla v \, dx + Q \int_{\Gamma_1} u_0 v \, ds = - \int_{\Omega} f v \, dx, \quad (4.1)$$

for any $v \in H^1(\Omega; \Gamma_2)$, respectively.

Let v be a function from $C^\infty(\Omega; \Gamma_2)$. Then (3.1) implies that

$$\int_{\Omega} (\nabla u_{\varepsilon,\delta}, \nabla (W_\varepsilon^\delta v)) \, dx + q \int_{\Gamma_{\varepsilon,\delta}^S} u_{\varepsilon,\delta} v W_\varepsilon^\delta \, ds = - \int_{\Omega} f v W_\varepsilon^\delta \, dx. \quad (4.2)$$

Keeping in mind the definition of $H^1(\Omega; S)$, the Green formula and (4.2) we deduce

$$\begin{aligned} & - \int_{\Omega} \Delta v W_\varepsilon^\delta u_{\varepsilon,\delta} \, dx - \int_{\Omega} v \Delta W_\varepsilon^\delta u_{\varepsilon,\delta} \, dx + I_{\varepsilon,\delta} + \\ & + \int_{\Gamma_{\varepsilon,\delta}^S} \left(q v W_\varepsilon^\delta + \frac{\partial v}{\partial x_3} W_\varepsilon^\delta + \frac{\partial W_\varepsilon^\delta}{\partial x_3} v \right) u_{\varepsilon,\delta} \, ds = - \int_{\Omega} f v W_\varepsilon^\delta \, dx, \end{aligned} \quad (4.3)$$

where

$$I_{\varepsilon,\delta} = -2 \int_{\Omega} (\nabla v, \nabla W_\varepsilon^\delta) u_{\varepsilon,\delta} \, dx. \quad (4.4)$$

Integrating by parts (4.4), we obtain that

$$I_{\varepsilon,\delta} = 2 \int_{\Omega} ((\nabla v, \nabla u_{\varepsilon,\delta}) + u_{\varepsilon,\delta} \Delta v) W_\varepsilon^\delta \, dx - 2 \int_{\Gamma_1} (W_\varepsilon^\delta - 1) u_{\varepsilon,\delta} \frac{\partial v}{\partial x_3} \, ds. \quad (4.5)$$

Let $\{\varepsilon_n\}$ be a sequence which tends to zero as $n \rightarrow \infty$. Due to the embedding theorems, weak compactness of the bounded set in $H^1(\Omega)$ and estimate (3.3), there exists a subsequence $\{\varepsilon'_n\}$, such that $u_{\varepsilon,\delta} \rightarrow u_0 \in H^1(\Omega; \Gamma_2)$ weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$ as $\varepsilon = \varepsilon'_n \rightarrow 0$. Using Lemma 4.2 and passing to the limit in (4.3), and (4.5) as $\varepsilon'_n \rightarrow 0$, we obtain that

$$- \int_{\Omega} \Delta v u_0 \, dx + \int_{\Gamma_1} u_0 \frac{\partial v}{\partial x_3} \, ds + Q \int_{\Gamma_1} u_0 v \, ds = - \int_{\Omega} f v \, dx. \quad (4.6)$$

Using the Green formula and (4.6), we obtain (4.1). By means of the definition of $H^1(\Omega; S)$ the function u_0 is a solution of (1.3). On the other hand, due to the arbitrariness of choosing of the sequence $\{\varepsilon_n\}$, we obtain that $u_{\varepsilon,\delta} \rightarrow u_0$ as $\varepsilon \rightarrow 0$ (weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$). For $p = 0$, the strong convergence of $u_{\varepsilon,\delta} \rightarrow u_0$ in $H^1(\Omega)$ proves similarly as in the proof of Theorem 1.2. Theorem is proved.

5 Proof of Theorem 1.3

Definition 5.1. *The function $\tilde{u}_0 \in H^1(\tilde{\Omega} \setminus \tilde{\Omega}; \Gamma \cup \tilde{\Gamma})$ is a solution of (1.5), if it satisfies the identity*

$$\int_{\tilde{\Omega} \setminus \tilde{\Omega}} \nabla \tilde{u}_0 \nabla v \, dx = - \int_{\tilde{\Omega} \setminus \tilde{\Omega}} \tilde{f} v \, dx, \quad (5.1)$$

for any $v \in H^1(\tilde{\Omega} \setminus \tilde{\Omega}; \Gamma \cup \tilde{\Gamma})$.

Note, that for boundary – value problem (1.4) (for boundary – value problem (1.6)), the surface $\Gamma_{\varepsilon,\delta}^D \cup \Gamma_2$ (the surface $\Gamma_2 \cup \Gamma_{\varepsilon,\delta}^S$) is considered as two-sides. Keeping in mind this remark, a function $u_{\varepsilon,\delta} \in H^1(\tilde{\Omega} \setminus \overline{\Gamma_{\varepsilon,\delta}^D \cup \Gamma_2}; \tilde{\Gamma} \cup \Gamma_{\varepsilon,\delta}^D \cup \Gamma_2)$, or $u_{\varepsilon,\delta} \in H^1(\tilde{\Omega} \setminus \overline{(\Gamma_2 \cup \Gamma_{\varepsilon,\delta}^S)}; \tilde{\Gamma} \cup \Gamma_2)$ is called a solution of (1.4), or (1.6), if there holds the integral identity

$$\int_{\Omega} \nabla u_{\varepsilon,\delta} \nabla v \, dx + \int_{\tilde{\Omega} \setminus \tilde{\Omega}} \nabla u_{\varepsilon,\delta} \nabla v \, dx = - \int_{\tilde{\Omega}} F v \, dx \quad (5.2)$$

for any $v \in H^1(\tilde{\Omega} \setminus \overline{\Gamma_{\varepsilon,\delta}^D \cup \Gamma_2}; \tilde{\Gamma} \cup \Gamma_{\varepsilon,\delta}^D \cup \Gamma_2)$, or $v \in H^1(\tilde{\Omega} \setminus \overline{(\Gamma_2 \cup \Gamma_{\varepsilon,\delta}^S)}; \tilde{\Gamma} \cup \Gamma_2)$, respectively.

Using the integral identity (5.2) of Problem (1.4) and keeping in mind the definition of the space $H^1(\tilde{\Omega}; \tilde{\Gamma} \cup \Gamma_{\varepsilon,\delta}^D) \cup \Gamma_2$, we conclude that the uniform estimates

$$\|u_{\varepsilon,\delta}\|_{H^1(\Omega)} \leq C \|F\|_{L_2(\tilde{\Omega})} \quad (5.3)$$

and

$$\|u_{\varepsilon,\delta}\|_{H^1(\tilde{\Omega} \setminus \tilde{\Omega})} \leq C \|F\|_{L_2(\tilde{\Omega})} \quad (5.4)$$

hold true.

Lemma 5.1. *For any $v \in H^1(\tilde{\Omega}; \tilde{\Gamma} \cup \Gamma_{\varepsilon, \delta}^D \cup \Gamma_2)$ the following estimates:*

$$\|v\|_{L_2(\Gamma)} \leq C \left(\frac{\delta}{\varepsilon}\right)^{1/2} \|v\|_{H^1(\Omega)} \quad (5.5)$$

and

$$\|v\|_{L_2(\Gamma)} \leq C \left(\frac{\delta}{\varepsilon}\right)^{1/2} \|v\|_{H^1(\tilde{\Omega} \setminus \Omega)} \quad (5.6)$$

are valid.

The proof of this Lemma is similar to the proof of Lemma 3.1.

Let $p = \infty$, suppose also that v is a function from $C_0^\infty(\Omega)$ and $\{\varepsilon_n\}$ is a sequence which tends to zero as $n \rightarrow \infty$. Due to the embedding theorems, weak compactness of the bounded set in $H^1(\Omega)$ and estimates (5.3), (5.5) there exists a subsequence of this sequence such that $u_{\varepsilon, \delta} \rightarrow u_0 \in H^1(\Omega; \Gamma)$ at this subsequence weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$. Multiplying the equation (5.2) of Problem (1.4) by a test-function from $H^1(\tilde{\Omega}; \tilde{\Gamma} \cup \Gamma_{\varepsilon, \delta}^D \cup \Gamma_2)$, integrating over Ω and passing to the limit on this subsequence we obtain the integral identity (3.2) of Problem (1.2). Hence, u_0 is a solution of (1.2). On the other hand, due to the arbitrariness of choosing of the sequence of $\{\varepsilon_n\}$, we have that $u_{\varepsilon, \delta} \rightarrow u_0$ as $\varepsilon \rightarrow 0$ weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$. The strong convergence of $u_{\varepsilon, \delta} \rightarrow u_0$ in $H^1(\Omega)$ proves similarly as in the proof of Theorem 1.2.

Acting in the same way, we obtain the convergence of $u_{\varepsilon, \delta}$ in the domain $\tilde{\Omega} \setminus \Omega$. Theorem is proved.

Remark 5.1. It is easy to see that the statement of Theorem holds true if we consider instead of the fixed function F the oscillating function $F_{\varepsilon, \delta}$ such that $F_{\varepsilon, \delta} \rightharpoonup F$ weakly in $L_2(\tilde{\Omega})$.

6 Proof of Theorem 1.4

Denote by $\widehat{W}^\varepsilon(x)$ the even continuation of the function

$$1 - \chi(r\varepsilon^{-1/2})X_0(x\varepsilon^{-1})$$

defined in Σ , with respect to x_3 . We conserve the same notation for the σ -periodic translation of the function $\widehat{W}^\varepsilon(x)$ on the plane $x_3 = 0$. Taking into account the definition of X_0 one can obtain the following statement.

Lemma 6.1. *Let $p = 0$, $\widehat{W}^{\varepsilon, \delta}(x) = \widehat{W}^\varepsilon\left(\frac{x}{\delta}\right)$. Then $\widehat{W}^{\varepsilon, \delta} \in H^1(\tilde{\Omega}; \Pi_\varepsilon^\delta)$,*

$$\|\widehat{W}^{\varepsilon, \delta} - 1\|_{H^1(\tilde{\Omega})} = o(1), \quad \varepsilon \rightarrow 0.$$

Let v be a function from $C^\infty(\Omega; \Gamma_2)$, \tilde{v} be a function from $C^\infty(\tilde{\Omega} \setminus \bar{\Omega}; \tilde{\Gamma} \cup \Gamma_2)$. Then, due to lemma 6.1

$$\|v - \widehat{W}^{\varepsilon, \delta} v\|_{H^1(\Omega)} + \|\tilde{v} - \widehat{W}^{\varepsilon, \delta} \tilde{v}\|_{H^1(\tilde{\Omega} \setminus \bar{\Omega})} \xrightarrow{\varepsilon \rightarrow 0} 0, \quad (6.1)$$

the continuation of $\widehat{W}^{\varepsilon, \delta} v$ into $\tilde{\Omega} \setminus \bar{\Omega}$ by zero belongs $H^1(\tilde{\Omega} \setminus \overline{\Gamma_2 \cup \Gamma_{\varepsilon, \delta}^S}; \tilde{\Gamma} \cup \Gamma_2)$, and the continuation of $\widehat{W}^{\varepsilon, \delta} \tilde{v}$ into Ω by zero belongs $H^1(\tilde{\Omega} \setminus \overline{\Gamma_2 \cup \Gamma_{\varepsilon, \delta}^S}; \tilde{\Gamma} \cup \Gamma_2)$, too. So, (5.2) can be written as

$$\int_{\Omega} \left(\nabla u_{\varepsilon, \delta}, \nabla \left(\widehat{W}^{\varepsilon, \delta} v \right) \right) dx = - \int_{\Omega} f v \widehat{W}^{\varepsilon, \delta} dx \quad (6.2)$$

or

$$\int_{\tilde{\Omega} \setminus \bar{\Omega}} \left(\nabla u_{\varepsilon, \delta}, \nabla \left(\widehat{W}^{\varepsilon, \delta} \tilde{v} \right) \right) dx = - \int_{\tilde{\Omega} \setminus \bar{\Omega}} \tilde{f} \tilde{v} \widehat{W}^{\varepsilon, \delta} dx, \quad (6.3)$$

where f and \tilde{f} are the restrictions of F on Ω and $\tilde{\Omega} \setminus \bar{\Omega}$, respectively.

From the integral identity (5.2) we could deduce the uniform boundedness of the function $u_{\varepsilon, \delta}$ in $H^1(\tilde{\Omega})$. Let $\{\varepsilon_n\}$ be a sequence which tends to zero as $n \rightarrow \infty$. Due to the embedding theorems, weak compactness of the bounded set of functions in $H^1(\tilde{\Omega})$, there exists a subsequence $\{\varepsilon'_n\}$, such that $u_{\varepsilon, \delta} \rightarrow u_*$ weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$ as $\varepsilon'_n \rightarrow 0$, $u_* \in H^1(\Omega; \bar{\Gamma}_2)$, and $u_{\varepsilon, \delta} \rightarrow \tilde{u}_*$ weakly in $H^1(\tilde{\Omega} \setminus \bar{\Omega})$ and strongly in $L_2(\tilde{\Omega} \setminus \bar{\Omega})$ as $\varepsilon'_n \rightarrow 0$, $\tilde{u}_* \in H^1(\tilde{\Omega} \setminus \bar{\Omega}; \bar{\Gamma}_2)$.

Keeping in mind (6.1) and passing to the limit in (6.2), and (6.3) as $\varepsilon'_n \rightarrow 0$, we obtain

$$\int_{\Omega} (\nabla u_*, \nabla v) dx = - \int_{\Omega} f v, dx \quad (6.4)$$

and

$$\int_{\tilde{\Omega} \setminus \bar{\Omega}} (\nabla u_*, \nabla \tilde{v}) dx = - \int_{\tilde{\Omega} \setminus \bar{\Omega}} \tilde{f} \tilde{v} dx \quad (6.5)$$

respectively. Due to the uniqueness of solutions to Problems (1.7) and (1.8) we conclude that $u_* \equiv u_0$ and $\tilde{u}_* \equiv \tilde{u}_0$.

On the other hand, due to the arbitrariness of choosing of the sequence $\{\varepsilon_n\}$, we obtain that $u_{\varepsilon, \delta} \rightarrow u_0$ as $\varepsilon \rightarrow 0$ (weakly in $H^1(\Omega)$ and strongly in $L_2(\Omega)$) and $u_{\varepsilon, \delta} \rightarrow \tilde{u}_0$ weakly in $H^1(\tilde{\Omega} \setminus \bar{\Omega})$ and strongly in $L_2(\tilde{\Omega} \setminus \bar{\Omega})$ as $\varepsilon \rightarrow 0$.

The proof of the strong convergence of $u_{\varepsilon, \delta} \rightarrow u_0$ in $H^1(\Omega)$ and $u_{\varepsilon, \delta} \rightarrow \tilde{u}_0$ in $H^1(\tilde{\Omega} \setminus \bar{\Omega})$ is similar to that one in Theorem 1.2. Theorem is proved.

Remark 6.1. It is easy to see that the statement of Theorem holds true if we consider instead of the fixed function F the oscillating function $F_{\varepsilon,\delta}$ such that $F_{\varepsilon,\delta} \rightharpoonup F$ weakly in $L_2(\tilde{\Omega})$.

7 Construction of analytic continuations of solutions

In this section we give the construction of the solutions of the problems (1.10)–(1.17) and their analytic continuations. This construction reproduces enough standard construction [38] given in [65], for the Helmholtz resonator and in [21]–[30] for its two-dimensional analogue in homogenization.

If X is a notation for some Banach space (for instance, $X = L_2$), then $X_{loc}(D) \stackrel{def}{=} \{u : u \in X(D \cap S(R)) \quad \forall R\}$, where $S(R)$ is the open ball of radius R centered at the origin. We say that a sequence converges in $X_{loc}(D)$, if it converges in $X(D \cap S(R))$ for all R . Let $\mathcal{B}(X, Y)$ be the Banach space of bounded linear operators mapping the Banach space X into the Banach space Y , $\mathcal{B}(X) \stackrel{def}{=} \mathcal{B}(X, X)$, $\mathcal{B}(Y, X_{loc}(D))$ be the set of maps $\mathcal{A} : Y \rightarrow X_{loc}(D)$ such that $\mathcal{A} \in \mathcal{B}(Y, X(D \cap S(R)))$ for all R . We indicate by $\mathcal{B}^h(X, Y)$ (by $\mathcal{B}^m(X, Y)$) the set of holomorphic (meromorphic) operator-valued functions whose values belong to $\mathcal{B}(X, Y)$; $\mathcal{B}^{h(m)}(X, X) \stackrel{def}{=} \mathcal{B}^{h(m)}(X)$, $\mathcal{B}^{h(m)}(X, Y_{loc}(D)) \stackrel{def}{=} \{\mathcal{A} : \mathcal{A} \in \mathcal{B}^{h(m)}(X, Y(D \cap S(R))) \quad \forall R\}$.

Let L be any number such that $\tilde{\Omega} \subset S(L/3)$. So, we can put $\tilde{\Omega} = S(L)$. Let's consider two families of boundary – value problems in bounded domains:

$$\begin{cases} \Delta u_{\varepsilon,\delta} = \Delta w, & x \in \tilde{\Omega} \setminus \overline{\Gamma_{\varepsilon,\delta}^D \cup \Gamma_2}, \\ u_{\varepsilon,\delta} = 0, & x \in \Gamma_{\varepsilon,\delta}^D \cup \Gamma_2, \quad u_{\varepsilon,\delta} = w, \quad x \in \tilde{\Gamma}, \end{cases} \quad (7.1)$$

and

$$\begin{cases} \Delta u_{\varepsilon,\delta} = \Delta w, & x \in \tilde{\Omega} \setminus \overline{(\Gamma_2 \cup \Gamma_{\varepsilon,\delta}^S)}, \\ u_{\varepsilon,\delta} = 0, & x \in \Gamma_2, \quad u_{\varepsilon,\delta} = w, \quad x \in \tilde{\Gamma}, \\ \frac{\partial u_{\varepsilon,\delta}}{\partial x_3} = 0, & x \in \Gamma_{\varepsilon,\delta}^S, \end{cases} \quad (7.2)$$

where $w \in H^2(\tilde{\Omega})$. Denote by $\sigma_\varepsilon^{(1)}$ an operator whose value on $w \in H^2(\tilde{\Omega})$ is the solution $u_{\varepsilon,\delta(\varepsilon)} \in H^1(\tilde{\Omega} \setminus \overline{\Gamma_{\varepsilon,\delta}^D \cup \Gamma_2})$ of the problem (7.1) and denote by $\sigma_\varepsilon^{(2)}$ an operator whose value on $w \in H^2(\tilde{\Omega})$ is the solution $u_{\varepsilon,\delta(\varepsilon)} \in H^1(\tilde{\Omega} \setminus \overline{(\Gamma_2 \cup \Gamma_{\varepsilon,\delta}^S)})$ of the problem (7.2). Similarly, denote by $\sigma_0^{(1)}$ an operator

whose value on $w \in H^2(\tilde{\Omega})$ is the pair of the solution $u_0 \in H^1(\Omega)$ and $\tilde{u}_0 \in H^1(\tilde{\Omega} \setminus \bar{\Omega})$ of the problems

$$\begin{cases} \Delta u_0 = \Delta w, & x \in \Omega, \\ u_0 = 0, & x \in \Gamma, \end{cases} \quad (7.3)$$

and

$$\begin{cases} \Delta \tilde{u}_0 = \Delta w, & x \in \tilde{\Omega} \setminus \bar{\Omega}, \\ \tilde{u}_0 = 0, & x \in \Gamma, \quad \tilde{u}_0 = w, \quad x \in \tilde{\Gamma}, \end{cases} \quad (7.4)$$

respectively, and denote by $\sigma_0^{(2)}$ an operator whose value on $w \in H^2(\tilde{\Omega})$ is the pair of the solution $u_0 \in H^1(\Omega)$ and $\tilde{u}_0 \in H^1(\tilde{\Omega} \setminus \bar{\Omega})$ of the problems

$$\begin{cases} \Delta u_0 = \Delta w, & x \in \Omega, \\ u_0 = 0, & x \in \Gamma_2, \\ \frac{\partial u_0}{\partial x_3} = 0, & x \in \Gamma_1, \end{cases} \quad (7.5)$$

$$\begin{cases} \Delta \tilde{u}_0 = \Delta w, & x \in \tilde{\Omega} \setminus \bar{\Omega}, \\ \tilde{u}_0 = 0, & x \in \Gamma_2, \quad \tilde{u}_0 = w, \quad x \in \tilde{\Gamma}, \\ \frac{\partial \tilde{u}_0}{\partial x_3} = 0, & x \in \Gamma_1, \end{cases} \quad (7.6)$$

respectively. Set

$$(\Delta + k^2)^{-1} g \stackrel{def}{=} -\frac{1}{4\pi} \int_{S(L)} \frac{e^{ik|x-y|}}{|x-y|} g(y) dy, \quad x \in \mathbb{R}^3,$$

Hereinafter we use the same notation for a function from $L_2(S(L))$ and its continuation by zero outside $S(L)$ meaning the latter as a function from $L_2(\mathbb{R}^3)$. Next, let p_L be the operator of restriction to $S(L)$,

$$\begin{aligned} A_\mu^{(j)}(k) &\stackrel{def}{=} \left(1 + \chi \left(\frac{r}{L} \right) (\sigma_\mu^{(j)} p_L - 1) \right) (\Delta + k^2)^{-1}, \\ T_\mu^{(j)}(k) g &\stackrel{def}{=} \left((\Delta + k^2) \left(\chi \left(\frac{r}{L} \right) \right) \left((1 - \sigma_\mu^{(j)} p_L) (\Delta + k^2)^{-1} \right) \right. \\ &\quad \left. + 2 \sum_{i=1}^2 \frac{\partial}{\partial x_i} \left(\chi \left(\frac{r}{L} \right) \right) \frac{\partial}{\partial x_i} \left((1 - \sigma_\mu^{(j)} p_L) (\Delta + k^2)^{-1} \right) \right) g, \end{aligned}$$

From the definitions of $T_\mu^{(j)}(k)$ it follows that for $g \in L_2(S(L))$, the function $T_\mu^{(j)}(k)g \in L_2(\mathbb{R}^3)$ and $\text{supp } T_\mu^{(j)}(k)g \subset \overline{S(L)}$. For this reason, the maps $T_\mu^{(j)}(k)$ and $B_\mu^{(j)}(k) = I - T_\mu^{(j)}(k)$ can be considered as operators from $L_2(S(L))$ into $L_2(S(L))$. Under this interpretation, from the definitions of $A_\mu^{(j)}(k)$ and $T_\mu^{(j)}(k)$ the following statements hold.

Lemma 7.1. For $k \in \mathbb{C}$,

a) mappings $A_\mu^{(1)}(k) \in \mathcal{B}^h(L_2(S(L)), H_{loc}^1(\mathbb{R}^3))$,

$$A_\varepsilon^{(2)}(k) \in \mathcal{B}^h\left(L_2(S(L)), H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Gamma_{\varepsilon, \delta(\varepsilon)}})\right),$$

$$A_0^{(2)}(k) \in \mathcal{B}^h(L_2(S(L)), H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Gamma_1})),$$

$$T_\mu^{(j)}(k) \in \mathcal{B}^h(L_2(S(L))),$$

and, for any fixed k and μ , $T_\mu^{(j)}(k)$ is a compact operator in $L_2(S(L))$;

b) the function $u_{\varepsilon, \delta(\varepsilon)} = A_\varepsilon^{(1)}(k)g$ satisfies (1.10) for $F = (I - T_\varepsilon^{(1)}(k))g$, where I is the identity mapping, the function $u_{\varepsilon, \delta(\varepsilon)} = A_\varepsilon^{(2)}(k)g$ satisfies (1.11) for $F = (I - T_\varepsilon^{(2)}(k))g$, the restriction u_0 of $A_0^{(1)}(k)g$ to Ω satisfies (1.13), where f is the restriction of $(I - T_0^{(1)}(k))g$ to Ω , the restriction \tilde{u}_0 of $A_0^{(1)}(k)g$ to $\mathbb{R}^3 \setminus \overline{\Omega}$ satisfies (1.14), where \tilde{f} is the restriction of $(I - T_0^{(1)}(k))g$ to $S(L) \setminus \overline{\Omega}$, the restriction u_0 of $A_0^{(2)}(k)g$ to Ω satisfies (1.16), where f is the restriction of $(I - T_0^{(2)}(k))g$ to Ω , the restriction \tilde{u}_0 of $A_0^{(2)}(k)g$ to $\mathbb{R}^3 \setminus \overline{\Omega}$ satisfies (1.17), where \tilde{f} is the restriction of $(I - T_0^{(2)}(k))g$ to $S(L) \setminus \overline{\Omega}$, and, for $\text{Im } k \geq 0$, the functions $u_{\varepsilon, \delta(\varepsilon)}$ and \tilde{u}_0 also satisfies the radiation conditions (1.12), (1.15).

The square root of the eigenvalue is called the eigenfrequency of the boundary – value problem. Denote by $\Sigma^{(1)}$ and $\Sigma^{(2)}$ the sets of eigenfrequencies of boundary – value problems (1.13) and (1.16), respectively.

Proposition 7.1. If $\text{Im } k \geq 0$ then the solution of the perturbed problem (1.10) ((1.11)), (1.12) and the limit external problem (1.14) ((1.17)), (1.15) are unique. If $k \notin \Sigma^{(1)}$ ($k \notin \Sigma^{(2)}$) then the solution of the limit internal problem (1.13) ((1.16)) is unique.

The proof of this statement is wellknown (for the three-dimensional external Neumann problem outside nonclosed surfaces see, for instance in [65]).

Lemma 7.2. If $g \neq 0$ and $g \in L_2(S(L))$, then $A_\mu^{(j)}(k)g \neq 0$.

The proof of this statement is completely identical to the proof of Lemma 2.2 in [65] and Lemma 3.3 in [30].

From Proposition 7.1, Lemma 7.2 and the compactness of the operator $T_\mu^{(j)}(k)$ we deduce the following Lemma.

Lemma 7.3. *If $\text{Im } k > 0$ or $k > 0$, and $k \notin \Sigma^{(j)}$, then there exists the operator $\left(B_\mu^{(j)}\right)^{-1}(k) \in \mathcal{B}(L_2(S(L)))$.*

Later we will use the following statement from [38]

Proposition 7.2. *Suppose that D is a connected domain of the complex plane, $T(k)$ ($k \in D$) is a holomorphic family of compact operators in a Banach space \mathcal{X} and there exists a point $k_* \in D$, such that $(I - T(k_*))^{-1} \in \mathcal{B}(\mathcal{X})$. Then $(I - T(k))^{-1}$ is a meromorphic function in D with values in $\mathcal{B}(\mathcal{X})$.*

From Proposition 7.2 and Lemma 7.3 the following Lemma follows:

Lemma 7.4. $\left(B_\mu^{(j)}\right)^{-1}(k) \in \mathcal{B}^m(L_2(S(L)))$ in \mathbb{C} .

Denote $\mathcal{A}_\mu^{(j)}(k) \stackrel{\text{def}}{=} A_\mu^{(j)}(k) \left(B_\mu^{(j)}\right)^{-1}(k)$

Theorem 7.1. *For $k \in \mathbb{C}$,*

a) *mappings $\mathcal{A}_\mu^{(1)}(k) \in \mathcal{B}^m(L_2(S(L)), H_{loc}^1(\mathbb{R}^3))$,*

$$\mathcal{A}_\varepsilon^{(2)}(k) \in \mathcal{B}^m\left(L_2(S(L)), H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Gamma_{\varepsilon, \delta(\varepsilon)}^S})\right),$$

$$\mathcal{A}_0^{(2)}(k) \in \mathcal{B}^m\left(L_2(S(L)), H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Gamma_1})\right);$$

b) *the function $u_{\varepsilon, \delta(\varepsilon)} = \mathcal{A}_\varepsilon^{(1)}(k)F$ satisfies (1.10), the function $u_{\varepsilon, \delta(\varepsilon)} = \mathcal{A}_\varepsilon^{(2)}(k)F$ satisfies (1.11), the restriction u_0 (\tilde{u}_0) of $\mathcal{A}_0^{(1)}(k)F$ to Ω (to $\mathbb{R}^3 \setminus \overline{\Omega}$) satisfies (1.13) ((1.14)), where f (\tilde{f}) is the restriction of F to Ω (to $S(L) \setminus \overline{\Omega}$), the restriction u_0 (\tilde{u}_0) of $\mathcal{A}_0^{(2)}(k)F$ to Ω (to $\mathbb{R}^3 \setminus \overline{\Omega}$) satisfies (1.16) ((1.17)), where f (\tilde{f}) is the restriction of F to Ω (to $S(L) \setminus \overline{\Omega}$), and, for $\text{Im } k \geq 0$, the functions $u_{\varepsilon, \delta(\varepsilon)}$ and \tilde{u}_0 also satisfies the radiation conditions (1.12), (1.15);*

c) *if $\text{supp } F \subset S(T)$, then the function $\mathcal{A}_\varepsilon^{(j)}(k)F$ does not depend on $L \geq T$; $j = 1, 2$;*

d) *the set of poles of the operators $\mathcal{A}_\varepsilon^{(j)}(k)$ and $\left(B_\mu^{(j)}(k)\right)^{-1}$ coincide for fixed j ; $j = 1, 2$;*

e) *the set of poles of the operator $\mathcal{A}_\varepsilon^{(j)}(k)$ does not depend on L ; $j = 1, 2$.*

Proof. The statements a) and b) follow from Lemmas 7.1, 7.4. The statement c) follows from Proposition 7.1 and the uniqueness of the analytic continuation. The statement d) follows from Lemma 7.2 and the definition of the operators $\mathcal{A}_\varepsilon^{(j)}(k)$; $j = 1, 2$. Let us prove the statement e) for $j = 1$.

Denote by $\mathcal{A}_{\varepsilon,t}^{(1)}$ the operator $\mathcal{A}_{\varepsilon}^{(1)}$ defined for $L = t$. Suppose $a > b$. It is obvious that the set of poles of $\mathcal{A}_{\varepsilon,b}^{(1)}$ is a subset of the set of poles of $\mathcal{A}_{\varepsilon,a}^{(1)}$. Now we show the inverse inclusion. Suppose that $\text{supp}F \subset S(a)$ and assume that

$$W = (1 - \chi(rb^{-1}))(\Delta + k^2)^{-1}F.$$

Since the support $\text{supp}(F - (\Delta + k^2)W) \subset \overline{S(b)}$, then due to b) the solution of the perturbed problem (1.10), (1.12) can be defined by one of the following formulae

$$u_{\varepsilon,\delta} = \mathcal{A}_{\varepsilon,a}^{(1)}(k)F, \quad u_{\varepsilon,\delta} = \mathcal{A}_{\varepsilon,a}^{(1)}(k)(F - (\Delta + k^2)W) + W.$$

Since W is holomorphic the set of poles of $\mathcal{A}_{\varepsilon,a}^{(1)}$ is a subset of the set of poles of $\mathcal{A}_{\varepsilon,b}^{(1)}$. The case $j = 2$ can be proved in analogous way. Theorem is proved.

8 Proof of Theorems 1.6 and 1.7

By the definition of operators $\sigma_{\delta}^{(j)}$, Statements of Theorems 1.3 , 1.4 and Remarks 5.1 and 6.1 we deduce the following Lemma:

Lemma 8.1. *If $p = \infty$, then*

$$\|\sigma_{\varepsilon}^{(1)} - \sigma_0^{(1)}\|_{\mathcal{B}(W_2^2(S(L)), W_2^1(S(L)))} \xrightarrow{\varepsilon \rightarrow 0} 0.$$

If $p = 0$, then

$$\|\sigma_{\varepsilon}^{(2)} - \sigma_0^{(2)}\|_{\mathcal{B}(W_2^2(S(L)), W_2^1(S(L) \setminus \overline{\Omega} \cup \Omega))} \xrightarrow{\varepsilon \rightarrow 0} 0.$$

Definition 8.1. *Suppose that \mathcal{D}_{μ} is a family of operators acting from the Banach space \mathcal{X} in $L_{2,loc}(\mathbb{R}^3)$ (in $H_{loc}^1(\mathbb{R}^3)$, $H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Omega})$).*

We say that $\mathcal{D}_{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} \mathcal{D}_0$ strongly (weakly, by the norm) in $\mathcal{B}(\mathcal{X}, L_{2,loc}(\mathbb{R}^3))$ (in $\mathcal{B}(\mathcal{X}, H_{loc}^1(\mathbb{R}^3))$, $\mathcal{B}(\mathcal{X}, H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Omega}))$), if $\mathcal{D}_{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} \mathcal{D}_0$ strongly (weakly, by the norm) in $\mathcal{B}(\mathcal{X}, L_2(S(R)))$ (in $\mathcal{B}(\mathcal{X}, H^1(S(R)))$, $\mathcal{B}(\mathcal{X}, H^1(S(R) \setminus \overline{\Omega}))$) for any $R > 0$.

The next Lemma follows from the definitions of operators $A_{\mu}^{(j)}(k)$, $T_{\mu}^{(j)}(k)$, and Lemma 8.1:

Lemma 8.2. *Assume that K is an arbitrary compact set in \mathbb{C} . Then*

- a) if $p = \infty$, then $A_{\varepsilon}^{(1)}(k) \xrightarrow{\varepsilon \rightarrow 0} A_0^{(1)}(k)$ by the norm $\mathcal{B}(L_2(S(L)), H_2^1(\mathbb{R}^3))$ and $T_{\varepsilon}^{(1)}(k) \xrightarrow{\varepsilon \rightarrow 0} T_0^{(1)}(k)$ by the norm $\mathcal{B}(L_2(S(L)))$ uniformly on $k \in K$;*
- b) if $p = 0$, then $A_{\varepsilon}^{(2)}(k) \xrightarrow{\varepsilon \rightarrow 0} A_0^{(2)}(k)$ by the norm $\mathcal{B}(L_2(S(L)), H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Omega} \cup \Omega))$ and $T_{\varepsilon}^{(2)}(k) \xrightarrow{\varepsilon \rightarrow 0} T_0^{(2)}(k)$ by the norm $\mathcal{B}(L_2(S(L)))$ uniformly on $k \in K$.*

Later on we shall use the following statement from [38]:

Proposition 8.1. *Suppose that D is a connected domain in the complex plane, $T(k, \mu)$ is a family of compact operators in Banach space \mathcal{X} , defined for $k \in D$ and $\mu \in [0, \mu_0]$, such that it is holomorphic on k for each μ and continuous on $D \times [0, \mu_0]$ by the norm $\mathcal{B}(\mathcal{X})$. Furthermore assume that there exists the point $k_0 \in D$, such that $(I - T(k_0, \mu))^{-1} \in \mathcal{B}(\mathcal{X})$ for any $\mu \in (0, \mu_0)$. Then*

- a) $(I - T(k, \mu))^{-1}$ (for any μ) is a meromorphic function in D with values in $\mathcal{B}(\mathcal{X})$;
- b) if k_* is not a pole $(I - T(k, \mu_*))^{-1}$, then the operator-values function $(I - T(k, \mu))^{-1}$ is continuous by the norm in a neighborhood of (k_*, μ_*) ;
- c) the poles $(I - T(k, \mu))^{-1}$ depend on μ continuously.

Denote by $\Sigma_\mu^{(j)}$ the set of the poles of the operator $\mathcal{A}_\mu^{(j)}$. By Lemmas 7.4 and 8.2 the family $T_\mu^{(j)}(k)$ satisfies the conditions of Proposition 8.1. Then the following Lemma holds true:

Lemma 8.3. a) *Assume that $p = \infty$. If K is an arbitrary compact set in \mathbb{C} , such that $K \cap \Sigma_0^{(1)} = \emptyset$, then $(B_\varepsilon^{(1)}(k))^{-1} \xrightarrow{\varepsilon \rightarrow 0} (B_0^{(1)}(k))^{-1}$ by the norm $\mathcal{B}(L_2(S(L)))$ uniformly on $k \in K$. If $\tau_0 \in \Sigma_0^{(1)}$, then there exists the pole $\tau_\varepsilon \in \Sigma_\varepsilon^{(1)}$, converging to τ_0 as $\varepsilon \rightarrow 0$.*

b) *Assume that $p = 0$. If K is an arbitrary compact set in \mathbb{C} , such that $K \cap \Sigma_0^{(2)} = \emptyset$, then $(B_\varepsilon^{(2)}(k))^{-1} \xrightarrow{\varepsilon \rightarrow 0} (B_0^{(2)}(k))^{-1}$ by the norm $\mathcal{B}(L_2(S(L)))$ uniformly on $k \in K$. If $\tau_0 \in \Sigma_0^{(2)}$, then there exists the pole $\tau_\varepsilon \in \Sigma_\varepsilon^{(2)}$, converging to τ_0 as $\varepsilon \rightarrow 0$.*

Finally from Lemmas 8.2 and 8.3 we deduce.

Theorem 8.1. a) *Suppose that $p = \infty$. If K is an arbitrary compact set in \mathbb{C} , such that $K \cap \Sigma_0^{(1)} = \emptyset$, then $\mathcal{A}_\varepsilon^{(1)}(k) \xrightarrow{\varepsilon \rightarrow 0} \mathcal{A}_0^{(1)}(k)$ by the norm of the space $\mathcal{B}(L_2(S(L)), H_{loc}^1(\mathbb{R}^3))$ uniformly on $k \in K$. If $\tau_0 \in \Sigma_0^{(1)}$, then there exists a pole $\tau_\varepsilon \in \Sigma_\varepsilon^{(1)}$, converging to τ_0 as $\varepsilon \rightarrow 0$.*

b) *Suppose that $p = 0$. If K is an arbitrary compact set in \mathbb{C} , such that $K \cap \Sigma_0^{(2)} = \emptyset$, then $\mathcal{A}_\varepsilon^{(2)}(k) \xrightarrow{\varepsilon \rightarrow 0} \mathcal{A}_0^{(2)}(k)$ by the norm $\mathcal{B}(L_2(S(L)), H_{loc}^1(\mathbb{R}^3 \setminus \overline{\Gamma_1}))$ uniformly on $k \in K$. If $\tau_0 \in \Sigma_0^{(2)}$, then there exists a pole $\tau_\varepsilon \in \Sigma_\varepsilon^{(2)}$, converging to τ_0 as $\varepsilon \rightarrow 0$.*

Theorems 1.6 and 1.7 are the implication of Theorems 8.1 and 7.1.

Acknowledgments.

The paper was completed in Narvik University College (HiN) in the Northern Norway during the cold January 2003. The authors express thanks to Narvik University College for the wonderful conditions to work and for the support.

References

- [1] V.A. Marchenko, E.Ya.Khruslov, "Boundary – Value Problems in Domains with Fine–Grained Boundaries", Naukova Dumka, Kyiv, 1974. (in Russian).
- [2] A.Damlamian and Li Ta-Tsien (Li Daqian), Homogénéisation sur le bord pour des problèmes elliptiques, *C. R. Acad. Sci. Paris, Série I* **229** (17, 1984), 859–862.
- [3] A.Damlamian and Li Ta-Tsien (Li Daqian), Boundary Homogeniation for Elliptic Problems, *J. Math. Pure et Appl.* **66** (1987), 351–361.
- [4] M.Lobo (Lobo–Mildago) and M.E.Pérez, Comportement asymptotique d'un corps élastique dont une surface présente de petites zones de collage *C. R. Acad. Sci. Paris, Série II* **304** (1987), 195–198.
- [5] M.Lobo (Lobo–Mildago) and M.E.Pérez, Asymptotic Behavior of an Elastic Body with a Surface Having Small Stuck Regions, *RAIRO Model. Math. & Analyze Numer.* **22** (4, 1988), 609–624.
- [6] G.A.Chechkin, On Boundary – Value Problems for a Second Order Elliptic Equation with Oscillating Boundary Conditions, *in* "Nonclassical Partial Differential Equations", Institute for Mathematics of the Siberian Division of the Academy of Sciences of the Soviet Union Press, ed. V.N.Vragov p. 95–104. 1988. Novosibirsk (in Russian). (Reported in Referent. Math., 1989, 12B442, p.62)
- [7] G.A.Chechkin, Spectral properties of an elliptic problem with rapidly oscillating boundary conditions, *in* "Boundary value problems for non-classical equations in mathematical physics", Institute for Mathematics of the Siberian Division of the Academy of Sciences of the Soviet Union Press, ed. V.N.Vragov p. 197–200, 1989. Novosibirsk (in Russian)
- [8] A.Brillard, M.Lobo and M.E.Pérez, Homogénéisation des frontières par epi-convergence en élasticité lineaire, *RAIRO Model. Math. & Analyfze Numer.* **24** (1, 1990), 5–26.

- [9] A.Brillard, M.Lobo and M.E.Pérez, Un probleme d'homogénéisation de frontière en élasticité lineaire pour un corps cylindrique, *C. R. Acad. Sci. Paris, Série II* **311** (1990), 15–20.
- [10] G.A.Chechkin, "Elliptic Boundary – Value Problems with Rapidly Alternating Boundary Conditions" Ph.D Thesys. Moscow State University (in Russian), 1992.
- [11] G.A.Chechkin, Average of Boundary – Value Problems with Singular Perturbation in Boundary Condition, *Matem. Sbornik* **184** (6, 1993), 99–150. English transl. in *Russian Acad. Sci. Sb. Math.* **79** (1, 1994), 191–222.
- [12] A.Friedman, C.Huang and J.Yong, Effective Permeability of the Boundary of a Domain, *Commun. in Partial Differential Equations* **20** (1995), 59–102.
- [13] O.A.Oleinik and G.A.Chechkin, On Asymptotics of Solutions and Eigenvalues of the Boundary – Value Problems with Rapidly Alternating Boundary Conditions for the System of Elasticity *Rendiconti Lincei: Matematica e Applicazioni. Serie IX* **7** (1, 1996), 5–15.
- [14] R.R.Gadyl'shin, Asymptotics of the minimum eigenvalue for a circle with fast oscillating boundary conditions, *C. R. Acad. Sci., Paris, Sér. I.* **323** (3, 1996), 319–323.
- [15] R.R.Gadyl'shin, On the eigenvalue asymptotics for periodically clamped membranes, *Algebra Anal.* **10** (1,1998), 3–19. English transl. *St. Petersburg. Math. J.* **10** (1, 1999), 1–14.
- [16] R.R.Gadyl'shin, Boundary – value problem for the Laplacian with rapidly oscillating boundary conditions, *Dokl. Akad. Nauk, Ross. Akad. Nauk.* **362** (4, 1998), 456–459. English transl. *Dokl. Math.* **58** (2, 1998), 293–296.
- [17] G.A.Chechkin and R.R.Gadyl'shin, Boundary – Value Problem for the Laplacian with Rapidly Oscillating Type of the Boundary Conditions in Many-Dimensional Domain, *Sibirskii Matematicheskii Zhurnal* **40** (2, 1999), 271–287. English transl. in *Siberian Mathematical Journal* **40** (2, 1999), 229–244.
- [18] D.I.Borisov, R.R.Gadyl'shin, On the spectrum of the Laplacian with frequently alternating boundary conditions, *Teor. Mat. Fiz.* **118** (3, 1999), 347–353. English transl. *Theor. Math. Phys.* **118** (3, 1999), 272–277.

- [19] A.Yu.Beliaev and G.A.Chechkin, Averaging Operators with Boundary Conditions of Fine-scaled Structure, *Math. Zametki* **65** (4, 1999), 496–510. English transl. in *Math. Notes* **65** (4, 1999), 418–429.
- [20] R.R.Gadyl'shin, Asymptotics of the eigenvalues of a boundary – value problem with rapidly oscillating boundary conditions, *Differ. Uravn.* **35** (4, 1999), 540–551. English transl. *Differ. Equations.* **35** (4, 1999), 540–551.
- [21] R.R.Gadyl'shin, On an analog of the Helmholtz resonator in the averaging theory, *C. R. Acad. Sci. Paris. Sér. I. Math.* **329** (12, 1999), 1121–1126.
- [22] G.A.Chechkin, E.I.Doronina, On the Averaging of Solutions of a Second - Order Elliptic Equation with Nonperiodic Rapidly Changing Boundary Conditions, *Moscow Univ. Math. Bull.* **56** (1, 2001), 14–19 (Translated from *Vestnik Moskovskogo Universiteta. Ser 1. Mathematics. Mechanics.* **1** (2001), 14–19.)
- [23] G.A.Chechkin, E.I. Doronina, On the Asymptotics of the Spectrum of a Boundary – Value Problems with Nonperiodic Rapidly Alternating Boundary Conditions, in "Functional Differential Equations" (Eds. E.Mitidieri, S.Pohozaev, A.Skubachevskii), Vol. 8, No.1-2, P. 111-122, Marcel Dekker, New York, 2001.
- [24] D.I.Borisov, Two-parameter asymptotics in a boundary – value problem for the Laplacian, *Matematicheskie zametki* **70** (4, 2001), 520–534; translated in *Mathematical Notes* **70** (3/4, 2001), 471–485.
- [25] D.I.Borisov, The asymptotics for the eigenelements of the Laplacian in a cylinder with frequently alternating boundary conditions, *C. R. Acad. Sci. Paris, Série IIb.* **329** (10, 2001), 717–721.
- [26] R.R.Gadyl'shin, Homogenization and asymptotics for a membrane with closely spaced clamping points, *Zh. Vychisl. Mat. Mat. Fiz.* **41** (12, 2001), 1857–1869 English transl. *Comput. Math. Math. Phys.* **41** (12, 2001), 1765–1776.
- [27] D.I.Borisov, On a Laplacian with frequently nonperiodically alternating boundary conditions, *Doklady Akademii Nauk* **383** (4, 2002), 443–445; translated in *Dokl. Math.* **65** (2, 2002), 224–226.

- [28] D.I.Borisov, Boundary – value problem in a cylinder with frequently changing type of boundary condition, *Matematicheskii Sbornik* **193** (7, 2002), 37–68; translated in *Sb. Math.* **193** (7, 2002), 977–1008.
- [29] D.I.Borisov, On a singularly perturbed boundary – value problem for Laplacian in a cylinder, *Differentsial'nye Uravneniya.* **38** (8, 2002), 1071–1078; translated in *Differ. Equations* **38** (8, 2002), 1140–1148.
- [30] R.R.Gadyl'shin, On analogs of Helmholtz resonator in averaging theory, *Mat. Sb.* **193** (11, 2002), 43–70. English transl. *Sb. Math.* **193** (11, 2002).
- [31] R.R.Gadyl'shin, On a model analogue of the Helmholtz resonator in homogenization, *Trudy Matem. Inst. Steklova* **236** (2002), 79–86. English transl. *Proc. Steklov Inst. Math.* **236** (2002), 70–77.
- [32] D.I.Borisov, On a model boundary – value problem for Laplacian with frequently alternating type of boundary condition, *Asymptotic Analysis* **00** (2003) (to appear).
- [33] G.Rosi and M.Nulman, Effect of Local Flows in Polymetric Permeation Reducing Barriers, *J. Appl. Phys.* **74** (1993), 5471–5475.
- [34] A.Friedman, "Mathematics in Industrial Problem. Part 6", IMA Volume in Mathematics and Its Applications, Vol. 57, Springer-Verlag, New York, 1993.
- [35] R.R.Gadyl'shin (R.R.Gadyl'shin), Asymptotic Properties of an Eigenvalue of a Problem for a Singularly Perturbed Self-Adjoint Elliptic Equation with a Small Parameter in the Boundary Condition, *Differentsial'nye Uravneniya* **22** (1986), 640–652. English transl. in *Differential Equations* **22** (1986), 474–483.
- [36] R.R.Gadyl'shin, Perturbation of the Laplacian spectrum when there is a change in the type of boundary condition on small part of the boundary, *Zh. Vychisl. Mat. Mat. Fiz.* **36** (7, 1996), 77–88 English transl. *Comput. Math. Math. Phys.* **36** (7, 1996), 889–898.
- [37] A.Bensoussan, and J.-L.Lions, and G.Papanicolau, "Asymptotic analysis for periodic structures", North-Holland, Amsterdam, 1978.
- [38] É.Sanchez-Palencia, "Homogenization Techniques for Composit Media", Springer-Verlag, Berlin – New York, 1987.
- [39] O.A.Oleinik, A.S.Shamaev and G.A.Yosifian, "Mathematical Problems in Elasticity and Homogenization", North-Holland, Amsterdam, 1992.

- [40] G.Dal Maso, "Introduction to Γ – Convergence", Birkhäuser, Boston, 1993.
- [41] V.V.Jikov, S.M.Kozlov and O.A.Oleinik, "Homogenization of Differential Operators and Integral Functionals", Springer–Verlag, Berlin — New York, 1994.
- [42] M.D.Van Dyke, "Perturbation Method in Fluid Mechanics", Academic Press, New York, 1964.
- [43] A.H.Nayfeh, "Perturbation Methods", John Wiley, New York, 1973.
- [44] A.M.Il'in, "Matching of Asymptotic Expansions of Solutions of Boundary – Value Problems", AMS, Providence, 1992.
- [45] É.Sanchez-Palencia, Boundary Value Problems in Domains Containing Perforated Walls, *in* "Nonlinear Partial Differential Equations and Their Applications", College de France Seminar III, editors: H.Brezis and J.L.Lions, Research Notes in Mathematics, Vol. 70, pp. 309–325, Pittman, London, 1982.
- [46] D.Cioranescu and F.Murat, Un terme etrange venu d'ailleurs, *in* Non-linear Partial Differential Equations and Their Applications. Res. Notes in Math., Vol. 60, pp. 98–138. Pittman, London, 1982.
- [47] F.Murat, The Neumann sieve, *in* "Nonlinear Problems", editors: A.Marino, L. Modica, S.Spagnolo and M.Degiovanni, Research Notes in Mathematics, Vol. 127, pp. 24–32, Pittman, London, 1985.
- [48] G.Allaire, Homogenization of the Navier–Stokes Equations in Open Sets Perforated with Tiny Holes I: Abstract Framework, a Volume Distribution of Holes, *Arch. Rat. Mech. Anal.* **113** (1991), 209–259.
- [49] G.Allaire, Homogenization of the Navier–Stokes Equations in Open Sets Perforated with Tiny Holes II: Noncritical Sizes of the Holes for a Volume Distribution and Surface Distribution of Holes, *Arch. Rat. Mech. Anal.* **113** (1991), 261–298.
- [50] A.A.Arsen'ev, On the singularities of the analytic continuation and the resonance properties of a solution of the dispersion problem for the Helmholtz equation, *Sov. Math., Dokl.* **12** (1971), 476–477; translation from *Dokl. Akad. Nauk SSSR* **197** (1971), 511–512.
- [51] J.T.Beale, Scattering frequencies of resonators, *Comm. Pure and Applied Math.* **26** (1973), 549–564.

- [52] P.D.Hislop, A.Martinez, Scattering resonances of Helmholtz resonator, *Indiana Univ. Math. J.* **40** (1991), 767–788.
- [53] R.R.Gadyl'shin, Surface potentials and the method of matching asymptotic expansions in the problem of the Helmholtz resonator, *Algebra Anal.* **4** (2, 1992), 88–115. English transl. *St. Petersburg. Math. J.* **4** (1, 1993), 273–296.
- [54] R.R.Gadyl'shin, On scattering frequencies of acoustic resonator, *C. R. Acad. Sci., Paris, Sér. I* **316** (9, 1993), 959–963.
- [55] R.R.Gadyl'shin, On acoustic Helmholtz resonator and on its electromagnetic analogue, *J. Math. Phys.* **35** (1994), 3464–3481.
- [56] R.R.Gadyl'shin, Asymptotics of scattering frequencies with small imaginary parts for an acoustic resonator, *RAIRO Modelisation Math. & Anal. Numer.* **28** (6, 1994), 761–780.
- [57] R.R.Gadyl'shin, Existence and asymptotics of poles with small imaginary part for the Helmholtz resonator, *Usp. Mat. Nauk* **52** (1, 1997), 3–76. English transl. *Russ. Math. Surv.* **52** (1, 1997), 1–72.
- [58] Lord Rayleigh, The theory of Helmholtz resonator, *Proc. Roy. Soc. London A.* **92** (1916), 265–275.
- [59] J.W.Miles, Scattering by a spherical cap, *J. Acoust. Soc. Amer.* **50** (2, 1971), 892–903.
- [60] N.S.Landkof, "Foundations of Modern Potential Theory", Springer-Verlag, Berlin–New York, 1972.
- [61] G.Pólya and G.Szegö, "Isoperimetric Inequalities in Mathematical Physics", Princeton Univ. Press, Princeton, 1951.
- [62] A.G.Belyaev, G.A.Chechkin and R.R.Gadyl'shin, Effective Membrane Permeability: Estimates and Low Concentration Asymptotics, *SIAM J. Appl. Math.* **60** (1, 2000), 84–108
- [63] G.I.Èskin, "Boundary – Value Problems for Elliptic Pseudodifferential Equations", AMS, Providence, 1981.
- [64] O.A.Ladyzhenskaya, "The boundary – value problems of mathematical physics", Applied Mathematical Sciences, Vol. 49. Springer-Verlag, Berlin–New York, 1985, 322 pp.

- [65] R.R.Gadyl'shin, Resonator systems, *Izv. Ross. Akad. Nauk, Ser. Mat.* **64** (3, 2000), 51–96. English transl. *Izv. Math.* **64** (3, 2000), 487–529.