Topological Phenomena in the Real Periodic Sine-Gordon Theory

P.G.Grinevich¹

Landau Institute for Theoretical Physics of the Russian Academy of Sciences, Moscow 117940 Kosygin Street 2, e-mail pgg@landau.ac.ru

S.P.Novikov²

IPST, University of Maryland-College Park, MD 20742-2431 and Landau Institute for Theoretical Physics, Moscow, e-mail novikov@ipst.umd.edu

1 Introduction. Algebro-Geometrical Solutions and the Reality Reductions. The topological charge.

The algebro-geometrical (or finite-gap) solutions play a very important role in the modern theory of soliton equations with periodic and quasiperiodic boundary conditions. The main property characterizing these solutions is the following: the wave function of the auxiliary linear operator is meromorphic in spectral parameter on the finite part of an algebraic Riemann surface. It allows to apply the powerful tools of classical algebraic geometry, and to write explicit representations for solutions and wave functions in terms of Riemann Θ -functions of several variables. Usually the algebro-geometrical solutions are dense in the space of the periodic ones, and if the nonlinearity tends to zero, they degenerate to finite Fourier sums. Therefore their role in the soliton theory is rather similar to the role of finite Fourier series for the linear PDE's.

The algebro-geometrical solutions of solition equations were first introduced in the paper [1] in 1974 dedicated to the integration of the Korteveg-

¹The author was partially supported by the RFBR Grant No 01-01-00874-A. Numerical simulations were partially performed on a computer, donated to P.G.Grinevich by the Humboldt Foundation.

²The author was partially supported by NSF Grant DMS 0072700

de Vries (KdV) equation with the periodic boundary conditions. Real nonsingular periodic algebro-geometrical KdV solutions can be characterised by the following property: the number of gaps (forbidden bands or zones) in the spectrum of the auxiliary Schrödinger operator acting in the Hilbert Space $L_2(R)$ is finite. This characterization is not valid for other systems (for example the spectrum of non-selfadjoint auxiliary operator for the self-focusing Nonlinear Schrödinger equation (NLS) has no gaps for any regular potential), but neverthereless the algebro-geometrical solutions are called finite-gap.

The complete theory of algebro-geometrical KdV solutions involving the spectral theory of periodic (quasiperiodic) finite-gap Schrödinger operators on the line and its unification with algebraic geometry and analysis on the Riemann surfaces, time dynamics and algebro-geometric hamiltonian aspects was constructed in the works [2, 3, 4, 5, 6, 7] (see in the survey article [8]) and in the works [9, 10]. Complex finite-gap solutions of the NLS and Sine-Gordon (SG) equations were constructed in [11, 12]. Finite-gap solutions of the 2+1 Kadomtsev-Petviashvili (KP) equation were constructed in [13] on a basis of a purely algebraic formulation of the finite-gap approach developed in this paper. The algebro-geometric spectral theory of the stationary periodic 2D Schrödinger operator was developed in the works [14, 15]. In these papers it was shown, that the proper analog of the finite-gap constraint in the 2dimensional case is the following: the restriction of the Bloch variety to one energy level is algebraic, and the spectral data is "collected" at this energy level only. Such potentials can be treated as "integrable" at one energy. It means, that the wave functions for this energy and arbitrary complex quaismomenta can be written explicitly in terms of Riemann Θ -functions, but the information about the spectral properties at other energies is very restricted.

The algebro-geometrical approach allows to obtain explicit compact formulas for solutions. But the Θ -functions of several variables are very complicated from the analytic point of view; therefore the existence of such formulas does not lead sometimes to the simple solution of the problem of selecting physically or geometrically meaningful solutions among generic complex ones. For example, it very easy to describe reality conditions for the KdV, defocusing NLS or Sinh-Gordon equation

$$\frac{\partial^2 u}{\partial \xi \partial \eta} = 4 \sinh u(\xi, \eta).$$

(in all these cases the auxiliary linear operators are self-adjoint). However,

the reality reductions for the self-focusing NLS or the Sine-Gordon equation found in [16] are rather nontrivial. It was shown in [15] that analogous reductions are responsible for the absence of magnetic field it the theory of the 2D Schrödinger operator. A good review of reality reductions can be found if [17].

The Sine-Gordon equation

$$u_{tt} - u_{xx} + \sin u(x, t) = 0 \tag{1}$$

derived in the XIX century geometry in the light-cone representation

$$u_{\xi\eta} = 4\sin u, \quad u = u(\xi, \eta), \tag{2}$$

where

$$x = 2(\xi + \eta), \quad t = 2(\xi - \eta), \quad \partial_{\xi} = 2\partial_x + 2\partial_t, \quad \partial_{\eta} = 2\partial_x - 2\partial_t.$$
(3)

is one the most important systems of the soliton theory. One it's distinguished properties is the folowing. Usually if we fix the spectral curve (or equivalently all local conservation laws), then the level set is a connected Jacoby torus. It is true for KdV, defocusing and self-focusing NLS, the both real reductions of the KP equation. But in the Sine-Gordon case the level set is a union of 2^m real Jacoby tori, where m denotes the number of pairs of negative branch points. As a corollary, the result of averaging a generic functional depends not only on the spectral curve, but also on the component. This property is also essential if we calculate the action-angle variables (this problem was studied in [18, 19]).

The averaged densities of conservation laws arose naturally as hamiltonians of the Whitham equations. These equations describe the evolution of parametrs of the asymptotic solutions in the form of a slowly varying Nphase wavetrains. The Whitham equations for the Sine-Gordon were derived in the paper [20], but the connectivity components of the level sets were not discussed. Let us point out, that the Whitham equations are hyperbolic only in the "stable" case, when all branch point are negative and the number of components is always greater than 1.

It is possible to extract from the calculations of [20], that for all classical local conservation laws except the topological charge the result of averaging is the same for all components, but this property was never pointed out explicitly. In Section 4 we give a simple explanation of it. **Remark.** The number of real tori was calculated in [16], their characterization in terms of the complex Jacoby torus was obtained in [21], see also [22]. An elementary description of these tori was obtained by the authors in [23, 24], see Section 2 below.

Another interesting property of the Sine-Gordon equation is the following. In the theory of constant negative curvature surfaces $u(\xi, \eta)$ denotes the angle between asymptotic lines. In the theory of Josephson jucntions u has the meaning of a phase. In both situations it is defined up to an additive constant $c = 2\pi n$. Therefore it is natural to call a solution u(x,t) space-periodic with the period T if $\exp\{iu(x+T)\} = \exp\{iu(x)\}$. The integer n such, that $u(x+T) = u(x) + 2\pi n$ is called **Topological charge**, the real number $\bar{n} = n/T$ denotes the **Density of topological charge**. These quantities are the "most stable" conservation laws surviving all real periodic nonintegrable perturbations. The density of topological charge can be naturally extended to all nonsingular quasiperiodic solutions using the formula

$$\bar{n} = \lim_{T \to \infty} [u(x+T) - u(x)]/2\pi T$$

One of the most natural problems is the following: How to calculate the topological charge of real periodic SG solutions in terms of the Spectral Data: the Riemann surface and the divisor? It turned out, that nobody succeeded to extract the answer from the explicit description of the real components described in terms of the Jacoby torus and Θ -functional formulas, found in the works [21, 22]. An attempt has been made almost 20 years ago in the work [18] to solve this problem. For this purpose the so-called "Algebro-Topological" approach was developed in [18], and an interesting formula has been proposed. However, as it was pointed out in [19], the idea of the proof presented in [18] works only if the spectral curve is sufficiently close to a degenerate one. A complete solution using a new development of the main idea of the work [18] was obtained only recently by the authors in the papers [23] (for the so-called stable curves) and [24] for generic ones. Our proof is based on a new effective description of the real components in terms of divisors. We present a summary of these results in Section 3.

Remark. Formally we can define topological charge for any non-singular complex solution. But if the solution has poles, this quantity became ill-defined, and no natural non-singularity condition in terms of the spectral data for complex potentials is known. It explains why only real solutions are discussed.

2 Finite-gap Sine-Gordon solutions.

The "soliton-type" solutions of Sine-Gordon were found already in the XIX century using substitution discovered by Bianchi and S.Lie. Now this method is called "Backlund transformation".

In the early 1970's G.Lamb found out (see [25]), that the Sine-Gordon equation is analogous to KdV in the following sence: the so-called "inverse scattering method" can be applied to it. The modern approach was started in the work [26]. The critical point of this approach is the following zerocurvature representation for SG

$$\Psi_x = \frac{1}{4}(U+V)\Psi, \quad \Psi_t = \frac{1}{4}(U-V)\Psi, \quad (4)$$

where

$$U = U(\lambda, x, t) = \begin{bmatrix} i(u_x + u_t) & 1\\ -\lambda & -i(u_x + u_t) \end{bmatrix},$$
(5)

$$V = V(\lambda, x, t) = \begin{bmatrix} 0 & -\frac{1}{\lambda}e^{iu} \\ e^{-iu} & 0. \end{bmatrix}.$$
 (6)

To construct generic complex finite-gap SG solutions assume, that we have the following spectral data:

- 1. A nonsingular hyperelliptic Riemann surface $\Gamma \ [\mu^2 = R(\lambda)]$, where $R(\lambda) = \prod_{k=0}^{2g} (\lambda E_k)$, such that $E_0 = 0$ and $E_i \neq E_j, i, j = 0, 1, \ldots, 2g$. It has exactly 2g + 2 branch points $E_0 = 0, E_1, \ldots, E_{2g}, \infty$; the genus of Γ is equal to g. A point $\gamma \in \Gamma$ is by definition a pair of complex numbers $\gamma = (\lambda, \mu)$ such, that $\mu^2 = R(\lambda)$.
- 2. A divisor D of degree g, i.e. set (or formal sum) of g points $D = \gamma_1 + \ldots + \gamma_g$.

For generic data Γ , D there exists an unique two-component "Baker-Akhiezer" vector-function $\Psi(\gamma, x, t)$ such that

1. For fixed (x,t) the function $\Psi(\gamma, x, t)$ is meromorphic in the variable $\gamma \in \Gamma$ outside the points $0, \infty$ and has at most 1-st order poles at the divisor points $\gamma_k, k = 1, \ldots, g$.

2. $\Psi(\gamma, x, t)$ has essential singularities at the points 0, ∞ with the following asymptotics:

$$\Psi(\gamma, x, t) = \begin{pmatrix} 1 + o(1) \\ i\sqrt{\lambda} + O(1) \end{pmatrix} e^{i\sqrt{\lambda}\frac{x+t}{4}} \quad \text{as} \quad \lambda \to \infty,$$
(7)

$$\Psi(\gamma, x, t) = \begin{pmatrix} \phi_1(x, t) + o(1) \\ i\sqrt{\lambda}\phi_2(x, t) + O(\lambda) \end{pmatrix} e^{-\frac{i}{\sqrt{\lambda}}\frac{x-t}{4}} \text{ as } \lambda \to 0, \quad (8)$$

where $\phi_1(x,t)$, $\phi_2(x,t)$ are some functions of the variables x, t.

Denote the divisor of zeroes of the first component ψ_1 by $D(x,t) = \sum_j \gamma_j(x,t)$. The vector-function $\Psi(\gamma, x, t)$ satisfies to the zero-curvature equations (4) with potential given by the formula

$$u(x,t) = i \ln \frac{\phi_2(x,t)}{\phi_1(x,t)},$$
(9)

and the function u(x,t) solves the SG equation (2). In terms of the divisor of zeroes D(x,t) the potential u(x,t) can be easily written explicitly:

$$e^{iu(x,t)} = \frac{\prod_{j=1}^{g} (-\lambda_j(x,t))}{\sqrt{\prod_{j=1}^{2g} E_j}}.$$
 (10)

The x and t dynamics of the divisor D(x,t) can be described it terms of Dubrovin equations (see [24]). From (10) it follows, that the potential u(x,t)is singular at the point (x_0, t_0) if and only if one of the points $\gamma_k(x_0, t_0)$ coincides with 0 or ∞ . Collisions of 2 or more divisor points result in singularities in the solutions of Dubrovin equations, but all symmetric combinations of the divisor points including the potential u(x,t) remain non-singular.

Remark. The first component ψ_1 of the Baker-Akhiezer vector-function Ψ is a partial case of the general scalar two-point Baker-Akhiezer functions satisfying to the second order linear Schrödinger equation $L_1\psi_1 = 0$ invented in the work [14]. Here we have

$$L_1 = \frac{\partial^2}{\partial \xi \partial \eta} + A(\xi, \eta) \frac{\partial}{\partial \eta} + W(\xi, \eta),$$

 $-A = \partial_{\xi} \log \phi_1$. In this special case our surface Γ is hyperelliptic, the selected points coincide with $0, \infty$ and corresponding local parameters are chosen as $\sqrt{\lambda}, 1/\sqrt{\lambda}$. The second function ψ_2 satisfies to similar equation $L_2\psi_2 = 0$ where the operator L_2 is obtained from L_1 by the so-called Laplace Transformations and vice versa. It is known that cyclic Laplace Chains of the length 2 lead to the SG equation in the general complex case (see the work [27]).

Lemma 1 Assume, that the spectral data (Γ, D) , $D = \gamma_1 + \ldots, \gamma_g$ satisfy the following reality constraints:

1. The set of branch points of the Riemann surface Γ contains real points or complex conjugate pairs of points only, and all real branch points are nonpositive. Without loss of generality we may assume, that the first 2m + 1 of them $E_0 = 0, E_1, E_2, \ldots, E_{2m}$ are real and $0 > E_1 > E_2 >$ $\ldots > E_{2m}$, and for k > m we have $E_{2k} = \overline{E_{2k-1}}$, Im $E_{2k} \neq 0$. Then the spectral curve Γ is real and admits a $\mathbb{Z}_2 \times \mathbb{Z}_2$ -group of involutions generated by the standard holomorphic involution

$$\sigma: \Gamma \to \Gamma: \quad \sigma(\lambda, \mu) = (\lambda, -\mu), \tag{11}$$

transposing the sheets, and by the antiholomorphic involution

$$\tau: \Gamma \to \Gamma: \quad \tau(\lambda, \mu) = (\bar{\lambda}, \bar{\mu}),$$
(12)

(complex conjugation).

2. There exists a meromophic differential Ω with two simple poles at the points $0, \infty$ such, that the zeroes of Ω are located at the points $D + \tau D$. We shall call such divisors **admissible**. Without loss of generality we can normalize Ω to be real: $\tau \Omega = \overline{\Omega}$.

Then the potential u(x,t) is real and non-singular.

Remark The constraints on the spectral curve we found in [12]. The characterisation of admissible divisors and the proof of non-singularity were obtained in the work [16].

It is easy to check (see, for example [24])

Lemma 2 If D is admissible divisor, then the divisor of zeroes $D(x,t) = \gamma_1(x,t) + \ldots + \gamma_q(x,t)$ is admissible for all x, t

Assume, that we have a real meromorphic differential Ω with exactly 2 simple poles at the points $0, \infty$. We can associate an admissible divisor D to it if and only if 2g zeroes of Ω can be can be presented as an union of 2 subsets D and D_1 such, that $D_1 = \tau D$. It is possible if and only if all real roots of Ω have even multiplicity.

Definition 1 Assume that the spectral curve Γ satisfy the reality condinitions formulated above. A real meromorphic differential Ω on Γ with exactly 2 simple poles at the points $0, \infty$ is called **admissible** if the multiplicity of all real roots is even.

Without loss of generality we may assume that the residued of Ω at the points 0 and ∞ are equal to +1 and -1 respectively. Then we have

$$\Omega = \left(1 - \frac{\lambda P_{g-1}(\lambda)}{R(\lambda)^{1/2}}\right) \frac{d\lambda}{2\lambda},\tag{13}$$

where $P_{g-1}(\lambda)$ is a polynomial of degree at most g-1 with real coefficients. Let us call the polynomial $P_{g-1}(\lambda)$ admissible if Ω is an admissible differential.

Denote the map associating an admissible polynomial with an admissible divisor by $\Pi: D \to P_{g-l}(\lambda)$. The inverse map Π^{-1} is multivalued. If $P_{g-1}(\lambda)$ is an admissible polynomial, then the equation

$$\begin{cases} \mu = \lambda P_{g-1}(\lambda) \\ R(\lambda) = \mu^2, \end{cases}$$
(14)

has 2g + 1 roots. One of them is the point (0, 0), other 2g roots form g pairs. To define an admissible divisor we have to choose one point from each pair: therefore we have at most 2^g possibilities depending on the number of real points. In the generic case we have no real roots, so the number is equal to 2^g in this case.

The description of connected components suggested by the authors in [23, 24] is based on the following simple observation:

Consider a pair of functions of the real variable λ :

$$f_{\pm}(\lambda) = \pm \frac{\sqrt{R(\lambda)}}{\lambda}.$$
 (15)

We assume, that the functions $f_{\pm}(\lambda)$ are defined only for such $\lambda \neq 0$ that $R(\lambda) \geq 0$, i.e. these functions are real-valued. Let us draw the graphs of these functions $y = f_{+}(\lambda)$, $y = f_{-}(\lambda)$, and fill in the following domains:

$$\lambda < 0, y^2 < \frac{R(\lambda)}{\lambda^2}, \lambda > 0, y^2 > \frac{R(\lambda)}{\lambda^2}.$$
(16)



Fig 1.

Lemma 3 The polynomial $P_{g-1}(\lambda)$ is admissible if and only if the graph of $P_{g-1}(\lambda)$ does not cross the black open domains (but it can touch their boundaries).

Now we can associate with each admissible polynomial a **Topological Type**, i.e. a collection of m numbers $s_k = \pm 1, k = 1, \ldots, m$ defined by the following rule:

 $s_k = 1$ if $P_{g-1}(\lambda) \ge f_+(\lambda)$ as $E_{2k} \le \lambda \le E_{2k-1}$, $s_k = -1$ if $P_{g-1}(\lambda) \le f_-(\lambda)$ as $E_{2k} \le \lambda \le E_{2k-1}$. Using simple algebraic estimates (see [24]) it is easy to check:

Lemma 4 Each topological type is presented by the convex subset in the space of polynomials P_{g-1} . Each of these 2^m connected components is non-empty and depends continuously on the branch points E_1, \ldots, E_{2g} . The x and t dynamics of admissible divisors may be rather non-trivial (a number of numeric experiments are discussed in [22]). In contrast with the self-ajoint case (KdV, defocusing NLS, sinh-Gordon) the admissible position of a given divisor point depends on the position of other ones (the admissibility constraints are non-local). The trajectory of a single divisor point may be not well-defined because divisor points may collide and a small variation of parameters near the collision point results in a bifurcation of trajectories. If the potential u(x,t) is periodic with the period T, the corresponding divisor is generically periodic only modulo permutations. But the following properties can be easily proved using the characterization formulated above.

Lemma 5 Projections of the points of admissible divisors to the λ -plane could not lie in the segments $[E_1, E_0]$, $[E_3, E_2], \ldots, (-\infty, E_{2m}]$. Moreover, it is possible to choose an open neighborhood of these segments such, that the projections of these points could not lie in it.

Lemma 6 Assume, that the projection of a point $\gamma_s = (\lambda_s, \mu_s)$ of an admissible divisor to the λ -plane lie in the segments $[E_{2k}, E_{2k-1}]$; then $\mu_s < 0$ if $s_k = 1$ and $\mu_s > 0$ if $s_k = -1$.

3 The topological charge of real solutions

It is well-known (see, for example [21, 22]) that after the Abel transform the *x*-dynamics of Sine-Gordon corresponds to the motion along the straight line

$$\vec{X} = \vec{X}_0 + \vec{U}x,\tag{17}$$

in the Jacoby torus, where \vec{U} denotes the noramalized vector of *b*-periods of the quasimomentum differential dp.

$$U^k = -\frac{1}{2\pi} \oint_{b_k} dp.$$
(18)

Let us recall, that by definition dp is a meromorphic differential with zero a-periods and two second order poles at 0, ∞ such, that

$$dp = \begin{cases} \left(\frac{1}{8\sqrt{\lambda}} + o\left(\frac{1}{\lambda}\right)\right) d\lambda \text{ as } \lambda \to \infty, \\ \left(\frac{1}{8\lambda\sqrt{\lambda}} + o\left(\frac{1}{\lambda}\right)\right) d\lambda \text{ as } \lambda \to 0. \end{cases}$$
(19)

The real part of the Jacoby torus is isomorphic to the factor $\mathbb{R}^n/\mathbb{Z}^n$, therefore we can apply the following simple analytic lemma:

Lemma 7 Let $u(\vec{X}), X \in \mathbb{R}^n$ be a smooth function in \mathbb{R}^n such, that $\exp(iu(\vec{X}))$ is single-valued on the torus $\mathbb{R}^n/\mathbb{Z}^n$, i.e. $\exp(iu(\vec{X} + \vec{N})) = \exp(iu(\vec{X}))$ for any integer vector \vec{N} . Denote by u(x) restriction of $u(\vec{X})$ to the straight line $\vec{X} = \vec{X}_0 + x \cdot \vec{U}$. Then the density of topological charge $\bar{n} = \lim_{T \to \infty} [u(x+T) - iu(x+T)]$

 $u(x)]/2\pi T$ is well-defined; it does not depend on the point \vec{X}_0 and can be expressed by the following formula:

$$\bar{n} = \sum_{k=1}^{n} n_k U^k, \tag{20}$$

where $\vec{U} = (U^1, U^2, \dots, U^n)$, and n_k are topological charges along the basic cycles \mathcal{A}_k , $k = 1, \dots, n$:

$$u(X^{1}, X^{2}, \dots, X^{k} + 1, \dots, X^{n}) - u(X^{1}, X^{2}, \dots, X^{k}, \dots, X^{n}) = 2\pi n_{k}.$$
(21)

The main step is the calculation of the the charges along the basic cycles \mathcal{A}_k . To obtain a convenient answer the choice of canonical basis is very important. For **stable** surfaces (surfaces without complex branch points) the proper system of *a*-cycles was constructed in [18].



Fig 2.

We work with the following system of cuts $[E_1, E_0]$, $[E_3, E_2]$,..., $(-\infty, E_{2g}]$. Let us denote the sheet containing the line $\lambda \in \mathbb{R}$, $\lambda > 0$ $\mu > 0$ by G_+ and the second sheet by G_- (see Fig 2).

For the generic real curves with 2m negative branch points E_1, \ldots, E_{2m} and g - m complex adjoint pairs $E_{2j-1}, E_{2j} = \overline{E}_{2j-1}, j = m + 1, \ldots, g$ we choose first m a-cycles exactly as in the stable case. The last g - m a-cycles should be chosen as coverings on Γ over the path on the λ -plane connecting the points E_{2j-1} and E_{2j} for j > m. All these path should not meet each other. All of them should cross positive part of the line $\lambda > 0$ in one point κ_j (i.e. in two points of Riemann surface) such that $\kappa_{m+1} < \kappa_{m+2} < \ldots < \kappa_g$. This basis depends on the order of the complex conjugate pairs only (see Fig 3).



We assume also that the cycles b_1, b_2, \ldots, b_m are the ovals lying over the segments $[E_2, E_1], [E_4, E_3], [E_{2m}, E_{2m-1}]$ respectively.

For the complex conjugate pairs of branch points E_{2j} , E_{2j-1} , j > m, we make the cuts connecting each of these points with $i\infty$ or $-i\infty$ such that the cuts do not intersect the real line. As before we denote the sheet containing the line $\lambda \in \mathbb{R}$, $\lambda > 0$ $\mu > 0$ by G_+ and the second sheet by G_- . The orientation of the cycles b_l , $l = 1, \ldots, m$ coincides with the standard orientation of the negative semi-line \mathbb{R}_{-} at the sheet G_{+} ; they are opposite at the sheet G_{-} .

Consider a basic cycle \mathcal{A}_k on the real component of Jacoby torus, represented by the closed curve B_k on this torus. The image of this cycle in the surface Γ (i.e. image of it under the inverse Abel map) is a closed oriented curve C_k (may be consisting from several connected components) homological to the cycle $a_k \in H_1(\Gamma, \mathbb{Z})$. From Lemma 5 it follows, that this curve does not touch the closed segments on the real line $[-\infty, E_{2m}], \ldots, [E_3, E_2], [E_1, 0]$, and if it crosses the negative semi-line at the inteval $[E_{2l}, E_{2l-1}]$, then it happens in the sheet G_+ if $(-1)^{l-1}s_l > 0$ and in the sheet G_- if $(-1)^{l-1}s_l < 0$. From Formula 10 it follows, that the topological charge along the cycle \mathcal{A}_k is given by

$$n_k = \hat{C}_k \circ \mathbb{R}_-,\tag{22}$$

where \tilde{C}_k is the projection of C_k to the λ -plane, and \circ denotes the intersection index of curves on the λ -plane. Taking into account that any intercetion with the negative semi-line is the intersection with one of the *b*-cycles we obtain:

Lemma 8 The topological charge along the cycle \mathcal{A}_k on the real torus in the Jacobian variety is equal to the intersection index of the corresponding curve C_k with the **Counting Cycle of the real component** on the Riemann surface Γ

$$n_k = C_k \circ \left(\sum_{l=1}^m (-1)^{l-1} s_l b_l \right).$$
 (23)

Let us mention that the cycle C_k is homotopic to a_k on Γ . Therefore $C_k \circ b_l = \delta_{kl}$ and

$$\begin{cases} n_k = (-1)^{k-1} s_k & \text{for } k \le m \\ n_k = 0 & \text{for } k > m. \end{cases}$$
(24)

Combining (20) with (24), we obtain our main result:

Theorem 1 For any real solution the density of topological charge along the variable x is given by the formula

$$\bar{n} = \frac{1}{2} \sum_{k=1}^{m} (-1)^{k-1} s_k U^k, \qquad (25)$$

where the components U^k of the vector \vec{U} are defined by (18).

4 Averaging of the local conservation laws and topological types

For a generic functional

$$F[u] = \int f(e^{iu}, e^{-iu}, u_x, u_{xx}, \dots, u_t, u_{tx}, u_{txx}, \dots) dx$$
(26)

the result of averaging over a real componet of the Jacoby torus depends on the topolgical type. For example by avearging of u_x we get the topological charge. But if we average the Sine-Gordon Hamiltonian

$$H[u] = \int \left[\frac{u_t^2}{2} + \frac{u_x^2}{2} + (1 - \cos(u))\right] dx$$
(27)

the result is the same for all components. Therefore it is natural to formulate the following:

Problem. How to characterize the functional such, that the averaging does not depend on the topological type?

A complete classification seems to be a complicated problem. But it is rather easy to formulate a natural sufficient condition.

If $f(e^{iu}, e^{-iu}, u_x, u_{xx}, \dots, u_t, u_{tx}, u_{txx}, \dots)$ is a polynomial of its arguments then it can be written as a symmetric rational function of the divisor coordinates $f = \tilde{f}(\lambda_1(x, t), \dots, \lambda_g(x, t), \mu_1(x, t), \dots, \mu_g(x, t)).$

Lemma 9 Consider the following form on the $\underbrace{\Gamma \times \Gamma \times \ldots \Gamma}_{g \text{ times}}$

$$W_f = \tilde{f}(\lambda_1, \dots, \lambda_g, \mu_1, \dots, \mu_g) \prod_{k < l} (\lambda_k - \lambda_l) \prod_{k=1}^g \frac{d\lambda_k}{\mu_k}$$
(28)

If we fix all divisor points except γ_1 we obtain a meromorphic differntial on Γ with poles only at 0 and ∞ . Assume that the residues of this differential are equal to 0 identically in $\gamma_2, \ldots \gamma_g$. Then for a given spectral curve the averaging does not depend on the topological type

Proof. Following [20] we can calculate the averaging of F[u] using the formula:

$$\overline{F[u]} = C \oint_{a_1} \dots \oint_{a_g} W_f$$
(29)

where C is the normalization constant, and **all cycles** a_k have the proper topological type. The last condition was not discussed in [20]. The deformations of the integration cycles do not affect the integral if we do not cross 0. If the condition of Lemma 9 is fulfilled, then the we can move the integration path through 0 and "reach" any topological type without changing the integral.

As a corollary we immediately get the following statement:

Theorem 2 Assume that the functional F[u] is invariant with respect to the following symmetries: $\sigma_k : (\lambda_k, \mu_k) \to (\lambda_k, -\mu_k), \ \sigma_k : (\lambda_l, \mu_l) \to (\lambda_l, \mu_l)$ for $l \neq k$. Then the result of averaging does not depend on the topological type.

Important example. Consider the densities of "higher SG hamiltonians" defined as odd expansion coefficients of the function

$$\frac{\Psi_x(\lambda, x, t)}{\Psi(\lambda, x, t)} \tag{30}$$

at the points 0 and ∞ . Direct calculation shows, that

$$\frac{\Psi_x}{\Psi} = \frac{1}{4} \left[\frac{\Psi_\xi}{\Psi} + \frac{\Psi_\eta}{\Psi} \right] \tag{31}$$

where

$$\frac{\Psi_{\xi}}{\Psi} = i \frac{\mu + Q^{\xi}(\lambda)}{(\lambda - \lambda_1(x, t)) \dots (\lambda - \lambda_g(x, t))},$$
(32)

$$\frac{\Psi_{\eta}}{\Psi} = i \frac{[\mu + \lambda Q^{\eta}(\lambda)][-\lambda_1(x,t), \dots (-\lambda_g(x,t))]}{\lambda(\lambda - \lambda_1(x,t)) \dots (\lambda - \lambda_g(x,t))\sqrt{E_1 \dots E_{2g}}},$$
(33)

 $Q^{\xi}(\lambda), Q^{\eta}(\lambda)$ are polynomials of degree g-1 determined by the following conditions:

$$Q^{\xi}(\lambda_k(x,t)) = \mu_k(x,t), \quad Q^{\eta}(\lambda_k(x,t)) = \frac{\mu_k(x,t)}{\lambda_k(x,t)}.$$
(34)

We see that the odd part of $\Psi_x \Psi^1$ depends on $\lambda_k(x,t)$ but not $\mu_k(x,t)$. Therefore the averaging of the "higher SG hamiltonians" does not depend on the topological type.

References

- Novikov, S.P. The periodic problem for the Korteweg-de Vries equation.
 I. Funkts. Anal. i ego Pril. 8 (1974), no. 3, 54–66.; English translation: Funct. Anal. Appl. 8 (1974), 236–246.
- [2] Dubrovin, B.A.; Novikov, S.P. Periodic and conditionally periodic analogs of the many-soliton solutions of the Korteweg-de Vries equation. *Zh. Eksper. Teoret. Fiz.* **67** (1974), no. 6, 2131–2144; English translation: *Sov. Phys-JETP* **40** (1974), 1058–1063.
- [3] Dubrovin, B.A.; Novikov, S.P. A periodicity problem for the Kortewegde Vries and Sturm-Liouville equations. Their connections with algebraic geometry. *Dokl. Akad. Nauk SSSR* **219** (1974), no. 3, 531–534; English translation: *Sov. Math. Dokl.* **15** (1974), 1597–1601.
- [4] Dubrovin, B.A. Periodic problem for the Kortweg-de Vries equation in the class of finite-band potentials. *Funkts. Anal. i ego Pril.* 9 (1975), no. 3, 41–51; English translation: *Funct. Anal. Appl.* 9 (1975), 215–223.
- [5] Its, A.R.; Matveev, V.B. Hill operators with finitely many gaps. Funkts. Anal. i ego Pril. 9 (1975), no. 1, 69–70; English translation: Funct. Anal. Appl. 9 (1975), 65–66.
- [6] Lax, P.D. Periodic solutions of the KdV equation. Comm. Pure Appl. Math. 28 1975, 141–188.
- [7] McKean, H.P.; van Moerbeke, P. The spectrum of Hill's equation. Invent. Math. 30 (1975), no. 3, 217–274.
- [8] Dubrovin, B.A.; Matveev, V.B.; Novikov, S.P. Non-linear equations of Korteweg-de Vries type, finite-zone linear operators and Abelian varieties. Uspekhi Mat. Nauk **31** (1976), no. 1, 55–136; English translation: Russ. Math. Surveys **31** (1976), no. 1, 59–146.
- [9] Flaschka, H.; McLaughlin, D.W. Canonically conjugate variables for the Korteweg-de Vries equation and the Toda lattice with periodic boundary conditions. *Progr. Theoret. Phys.* 55 (1976), no. 2, 438–456.
- [10] Novikov, S.P.; Veselov, A.P. On Poisson brackets compatible with algebraic geometry and Korteweg-de Vries dynamics on the set of finite-zone

potentials. Dokl. Akad. Nauk SSSR **266** (1982), no. 3, 533–537; English translation: Sov. Math. Dokl. **26** (1982), 357–362.

- [11] Kozel, V.A.; Kotlyarov, V.P. Almost periodic solutions of the equation $u_{tt} u_{xx} + \sin u = 0$. (Russian) *Dokl. Akad. Nauk Ukrain. SSR* Ser. A, (1976), no. 10, 878–881.
- [12] Its, A.R.; Kotlyarov, V.P. Explicit formulas for solutions of the Nonlinear Schrödinger equation. (Russian) Dokl. Akad. Nauk Ukrain. SSR Ser. A, (1976), no. 11, 965–968;
- [13] Krichever, I.M. An algebraic-geometric construction of the Zakharov-Shabat equations and their periodic solutions. *Dokl. Akad. Nauk SSSR* 227 (1976), no. 2, 291–294; English translation: *Sov. Math. Dokl.* 17 (1976), 394–397.
- [14] Dubrovin, B.A.; Krichever, I.M.; Novikov, S.P. The Schrödinger equation in a periodic field and Riemann surfaces. *Dokl. Akad. Nauk SSSR* 229 (1976), no. 1, 15–18; English translation: *Sov. Math. Dokl.* 17 (1976), 947–951.
- [15] Novikov, S.P.; Veselov, A.P. Finite-zone, two-dimensional Schrödinger operators. Potential operators. *Dokl. Akad. Nauk SSSR* **279** (1984), no. 4, 784–788. English translation: *Sov. Math. Dokl.* **30** (1984), 705– 708; Finite-zone, two-dimensional potential Schrödinger operators. Explicit formulas and evolution equations. *Dokl. Akad. Nauk SSSR* **279** (1984), no. 1, 20–24; English translation: Sov. Math. Dokl., **30** (1984), 588–591.
- [16] Cherednik, I.V. Reality conditions in "finite-zone" integration. Dokl. Akad. Nauk SSSR 252 (1980), no. 5, 1104–1108; English translation: Sov. Phys. Dokl. 25 (1980), 450–452.
- [17] Natanzon, S.M. Moduli of real algebraic surfaces, and their superanalogues. Differentials, spinors and Jacobians of real curves. Uspekhi Mat. Nauk 54 (1999), no. 6, 3–60; English translation: Russ. Math. Surveys 54 (1999), no. 6, 1091–1147.
- [18] Dubrovin, B.A.; Novikov, S.P. Algebro-geometrical Poisson brackets for real finite-zone solutions of the Sine-Gordon equation and the nonlinear

Schrödinger equation. Dokl. Akad. Nauk SSSR **267** (1982), no. 6, 1295–1300; English translation: Sov. Math. Dokl. **26** (1982), no. 3, 760–765.

- [19] Novikov, S.P. Algebrotopological approach to the reality problems. Real action variables in the theory of finite-zone solutions of the Sine-Gordon equation. Zap. Nauchn. Sem. LOMI: Differential geometry, Lie groups and mechanics. VI. 133 (1984), 177–196; English Translation: J. Soviet Math. 31(1985), no. 6, 3373–3387
- [20] Ercolani, N.; Forest M.G.; McLaughlin D.W.; Montgomery R. Hamiltonian structure for th emodulation equations of a Sine-Gordon wavetrain. *Duke Mathematical Journal* 55 (1987), no 4, 949–983.
- [21] Dubrovin, B.A.; Natanzon, S.M. Real two-zone solutions of the sine-Gordon equation. *Funkts. Anal. i ego Pril.* **16** (1982), no. 1, 27–43; English translation: *Funct. Anal. Appl.* **16** (1982), 21–33.
- [22] Ercolani, N.M.; Forest, M.G. The geometry of real Sine-Gordon wavetrains. Comm. Math. Phys. 99 (1985), no. 1, 1–49.
- [23] Grinevich, P.G.; Novikov, S.P. Real finite-gap Sine-Gordon solutions: a formula for topological charge. (in Russian) Uspekhi. Mat. Nauk 56 (2001), no. 5, 181–182.; English translation: Russ. Math. Surveys 56 (2001), no. 5, 980–981.
- [24] Grinevich, P.G.; Novikov, S.P Topological Charge of the real periodic finite-gap Sine-Gordon solutions. To appear in *Commun. Pure. Appl. Math.*.
- [25] Lamb, G.L. Analytical description of ultrashort optical pulse propagation in a resonant medium. *Rev. Mod. Phys.* 43 (1971), 99–124.
- [26] Ablowitz, M.J.; Kaup, D.J.; Newell, A.S.; Segur, H. Method for solving the Sine-Gordon equation. *Phys. Rev. Lett.* **30** (1973), no. 25. 1262–1264.
- [27] Novikov, S.P.; Veselov, A.P. Exactly solvable two-dimensional Schrödinger operators and Laplace transformations. In: Solitons, geometry, and topology: on the crossroad. *Amer. Math. Soc. Transl.* Ser. 2, **179** (1997), 109–132.