

Quotient Equations and Integrals of Motion for Scalar Field

S.S. Moskaliuk

Bogolyubov Institute for Theoretical Physics
Metrolohichna Str., 14-b, Kyiv-143, Ukraine, UA-03143
e-mail: mss@bitp.kiev.ua

Abstract

In this article a group-theoretical aspect of the method of dimensional reduction is presented. Then, on the base of symmetry analysis for an anisotropic space geometrical description of dimensional reduction of equation for scalar field is given. Formula for calculating components of the energy-momentum tensor from the variables of the field factor-equations is derived.

Much well-known researches [1,2] are devoted to quantization of scalar field in various anisotropic models. Here we shall briefly outline the results of this research as applied to our task, following [2].

The covariant equation for massive scalar conform-invariant field in metric

$$ds^2 = dt^2 - \sum_{i=1}^3 A_i^2(t)(dx^i)^2 \quad (1)$$

is of the form

$$\nabla_\mu \nabla^\mu \varphi + (m^2 + R/6)\varphi = 0,$$

where ∇_μ is covariant derivative and R is scalar curvature.

The solution of this equation can be written as $\varphi_k(x) = (16\pi)^{-3/2}a(\eta)^{-1} \times \tilde{g}_k(\eta) \exp(-i\mathbf{k}\mathbf{x})$ [3,4]. The temporal function $\tilde{g}_k(\eta)$ satisfies the equation of oscillator type

$$\ddot{\tilde{g}}_k + [\mu^2(\eta)K_0^2(\eta) + Q(\eta)]\tilde{g}_k = 0. \quad (2)$$

Here and after in this article we shall use the following notation:

$$\begin{aligned} k &= (K_0, \mathbf{k}), \\ \mathbf{k} &= (k_1, k_2, k_3) = k(\sin(\theta) \cos(\varphi), \sin(\theta) \sin(\varphi), \cos(\theta)), \\ K_0^2(\eta, \mathbf{k}) &= k^2 + m^2 g^2(\eta, \theta, \varphi), \\ g &= a(\eta)/\mu(\eta, \theta, \varphi), \quad C_i = \dot{\alpha}_i/\alpha_i, \\ \mu^2(\eta, \theta, \varphi) &= \frac{\sin^2(\theta) \cos^2(\varphi)}{\alpha_1^2(\eta)} + \frac{\sin^2(\theta) \sin^2(\varphi)}{\alpha_2^2(\eta)} + \frac{\cos^2 \varphi}{\alpha_3^2(\eta)}, \\ Q &= [(C_1 - C_2)^2 + (C_2 - C_3)^2 + (C_1 - C_3)^2]. \end{aligned} \quad (3)$$

Following Lagrange method, we pursue the solutions of the equation (2) in the form

$$\begin{aligned} \tilde{g}_k(\eta) &= (\mu K_0)^{-1} [\alpha_k^* e_+ + \beta_k e_-], \\ \dot{\tilde{g}}_k(\eta) &= -i\mu K_0 [\alpha_k^* e_+ - \beta_k e_-], \\ e_\pm &= \exp \left(\pm i \int_{\eta_0} K_0(\eta') d\eta' \right), \end{aligned}$$

rendering instead of one differential equation of the second order for $\tilde{g}_k(\eta)$ a set of two linear equations of the first order for complex-valued functions $\alpha_k(\eta)$ and $\beta_k(\eta)$, related by additional condition $|\alpha_k(\eta)|^2 - |\beta_k(\eta)|^2 = 1$:

$$\begin{aligned}\dot{\alpha}_k &= \left(\frac{\mathcal{W}}{2} - i\frac{\widetilde{\mathcal{W}}}{2}\right) \beta_k^* e_+^2 - i\frac{\widetilde{\mathcal{W}}}{2}\alpha_k, \\ \dot{\beta}_k^* &= \left(\frac{\mathcal{W}}{2} + i\frac{\widetilde{\mathcal{W}}}{2}\right) \alpha_k e_-^2 + i\frac{\widetilde{\mathcal{W}}}{2}\beta_k^*,\end{aligned}\quad (4)$$

where $\mathcal{W} = \dot{\mu}/\mu + \dot{K}_0/K_0$; $\widetilde{\mathcal{W}} = Q/(\mu K_0)$.

In actual practice it is more convenient to change from two complex-valued functions $\alpha_k(\eta)$ and $\beta_k(\eta)$ to three real-valued functions

$$\begin{aligned}S_k(\eta) &= |\beta_k|^2, \\ U_k(\eta) &= 2\text{Re}(\alpha_k \beta_k^* e_-^2), \\ V_k(\eta) &= 2\text{Im}(\alpha_k \beta_k^* e_-^2),\end{aligned}\quad (5)$$

for which a set of three linear equations can be derived [5]

$$\begin{aligned}\dot{S}_k &= \frac{\mathcal{W}}{2}U_k + \frac{\widetilde{\mathcal{W}}}{2}V_k, \\ \dot{U}_k &= \mathcal{W}(2S_k + 1) - (\widetilde{\mathcal{W}} + 2K_0)V_k, \\ \dot{V}_k &= \widetilde{\mathcal{W}}(2S_k + 1) - (\widetilde{\mathcal{W}} + 2K_0)U_k\end{aligned}\quad (6)$$

with initial conditions $S_k = U_k = V_k = 0$ at $\eta = \eta_0$.

As it can be seen, the structure of the equations (6), is similar to that of the set of equations, derived for vector massless files and massive spinor field [6,7].

The introduced functions and notation enable us to write vacuum averages for normally ordered operator of energy-momentum tensor $T_{\mu\nu} = \langle 0_{in}|N_\eta T_{\mu\nu}(\eta, \mathbf{x})|0_{in}\rangle$ for scalar field [2]:

$$T_0^0(\eta) = \frac{1}{(2\pi)^3 a^4(\eta)} \int d^3\mathbf{k} \mu(\eta, \theta, \varphi) K_0(\eta, \mathbf{k}) (S_k(\eta) -$$

$$\begin{aligned}
& - \frac{Q(\eta)}{2\mu^2(\eta, \theta, \varphi)K_0^2(\eta, k)} \left[S_k(\eta) + \frac{1}{2}U_k(\eta) \right] \Bigg), \\
T_i^i(\eta) &= \frac{1}{(2\pi)^3 a^4(\eta)} \int d^3\mathbf{k} \mu(\eta, \theta, \varphi) K_0(\eta, \mathbf{k}) \times \\
& \times \left(\frac{k_i^2}{\alpha_i^2 \mu^2 K_0^2} \left[S_k(\eta) + \frac{1}{2}U_k(\eta) \right] - \frac{1}{6}U_k(\eta) + \right. \\
& \left. + \frac{1}{6\mu^2 K_0^2} (C_i - Q) \left[S_k(\eta) + \frac{1}{2}U_k(\eta) \right] - \frac{C_i}{18\mu K_0} V_k \right), \\
T_\mu^\mu(\eta) &= \frac{1}{(2\pi)^3 a^4(\eta)} \int d^3\mathbf{k} \mu(\eta, \theta, \varphi) K_0(\eta, \mathbf{k}) \times \\
& \times \left(\frac{m^2 g^2(\eta, \theta, \varphi)}{K_0^2(\eta, \mathbf{k})} \left[S_k(\eta) + \frac{1}{2}U_k(\eta) \right] \right). \tag{7}
\end{aligned}$$

In such a manner, solving the set of equations (6) and examining the properties of functions S_k , U_k and V_k , we get the information on energy-momentum tensor for scalar field.

The author is grateful to Yu.A. Danilov for helpful discussions.

References

- [1] Grib A.A., Mamaev S.G., Mostopanenko V.M. *Vacuum quantum effects in the strong fields* (Moscow: Energoatomizdat, 1988, 288 p.).
- [2] Zel'dovich Ya.B., Starobinsky A.A. JETP (1971), **61**, No6, 2161–2175.
- [3] Pritomanov S.A. Izv. vuzov. (1984), **1**, No1, 35–40.
- [4] Grib A.A., Nesteruk .B. Nuclear Physics (1983), **38**, No5 (11), 1357–1366.
- [5] Moskaliuk S.S., Nesteruk A.V., Pritomanov S.A. – Kiev, 1990. – 20 p. (Preprint /AS UkrSSR, Inst. for Theoret. Phys., No90-40R).
- [6] Moskaliuk S.S. Quotient equations and integrals of motion for vector massless field // Hadronic Journal. – 2001. – **24**, No. 4. – P. 435–448.
- [7] Moskaliuk S.S. Quotient equations and integrals of motion for massive spinor field // Hadronic Journal. – 2001. – **24**, No. 6. – P. 699–674.