

A COCYCLE FOR THE COMULTIPLICATION ON THE QUANTUM $SU(2)$ GROUP

WOJCIECH SZYMAŃSKI

ABSTRACT. For the family Δ_q , $q \in [0, 1)$, of the $SU_q(2)$ -comultiplications on the C^* -algebra $A \cong C(SU_q(2))$, we show that there exist unitary operators Ω_q such that $\Delta_q(x) = \Omega_q \Delta_0(x) \Omega_q^*$ for $x \in A$.

One common way of producing Hopf $*$ -algebras is to twist the comultiplication of an already known Hopf $*$ -algebra by a unitary 2-cocycle or 2-pseudo-cocycle (for example, see [1]). In this note, we show that an analogous cocycle relation exists between the comultiplications on the quantum $SU_q(2)$ group corresponding to different values of the parameter $q \in [0, 1)$. However, our starting point is $q = 0$, at which $SU_q(2)$ is not a quantum group but only a quantum semigroup (with counit). We show that there exist unitary operators Ω_q such that $\Delta_q = \text{Ad}(\Omega_q)\Delta_0$. Thus, instead of twisting existing quantum group structure, Ω_q ‘regularizes’ the comultiplication of a quantum semigroup to a comultiplication of a quantum group.

We begin by recalling the definition of the quantum $SU(2)$ group [2]. For $q \in [0, 1]$, the C^* -algebra $C(SU_q(2))$ of continuous functions on the quantum $SU(2)$ group is defined as the universal C^* -algebra generated by two elements a and b , subject to the following relations.

$$\begin{aligned} a^*a + b^*b &= I, & ab &= qba, \\ aa^* + q^2b^*b &= I, & ab^* &= qb^*a, \\ & & b^*b &= bb^*. \end{aligned}$$

If $q = 1$ then this algebra is abelian and isomorphic with the C^* -algebra of continuous functions on the classical group $SU(2)$. If $q \in [0, 1)$ then the algebras $C(SU_q(2))$ are no longer abelian. However, they are all isomorphic as C^* -algebras [2, Theorem A2.2] to the universal C^* -algebra generated by two operators T and S , subject to the relations

$$\begin{aligned} T^*T &= I, \\ S^*S &= SS^*, \\ TT^* + S^*S &= I. \end{aligned}$$

We denote this C^* -algebra by A . By [2, Theorem A2.2] and [2, Formulae (A2.2) and (A2.3)], there exists an isomorphism $\phi_q : C(SU_q(2)) \rightarrow A$ given on the generators as follows. If $q \in (0, 1)$ then

$$\begin{aligned} \phi_q(a) &= \sum_{n=0}^{+\infty} \left(\sqrt{1 - q^{2(n+1)}} - \sqrt{1 - q^{2n}} \right) T^n (T^*)^{n+1}, \\ \phi_q(b) &= \sum_{n=0}^{+\infty} q^n T^n S (T^*)^n. \end{aligned}$$

If $q = 0$ then

$$\begin{aligned} \phi_0(a) &= T^*, \\ \phi_0(b) &= S. \end{aligned}$$

Date: 6 November, 2002.

This work was partially supported by Max-Planck-Institute für Mathematik, Bonn, by Universität Münster, and by Research Grants Committee of the University of Newcastle.

Here and in what follows we agree to write

$$T^0 = (T^*)^0 = I, \quad S^0 = (S^*)^0 = S^*S, \quad \text{and } S^{-1} = S^*.$$

We denote by J the closed 2-sided ideal of A generated by S . We have

$$(1) \quad J = \overline{\text{span}}\{T^m S^j T^{-n} : m, n \in \mathbb{N}, j \in \mathbb{Z}\}.$$

Thus J is isomorphic to $C(\mathbb{T}) \otimes \mathcal{K}$, with \mathcal{K} the compact operators on a separable Hilbert space. The quotient is generated by $T + J$ and isomorphic to $C(\mathbb{T})$, the C^* -algebra of continuous functions on the circle. Thus there is an exact sequence

$$0 \longrightarrow C(\mathbb{T}) \otimes \mathcal{K} \longrightarrow A \longrightarrow C(\mathbb{T}) \longrightarrow 0.$$

The ideal J has a natural faithful representation on the Hilbert space $\ell^2(\mathbb{N} \times \mathbb{Z})$, which extends to a representation ρ of A constructed by Woronowicz in [2]. Since the ideal J is essential, the representation ρ of A is faithful. If $\{\xi(k, m) \mid k \in \mathbb{N}, m \in \mathbb{Z}\}$ is an orthonormal basis of $\ell^2(\mathbb{N} \times \mathbb{Z})$ then we have

$$\begin{aligned} \rho(T) &: \xi(k, m) \mapsto \xi(k+1, m), \\ \rho(S) &: \xi(k, m) \mapsto \delta_{k,0} \xi(k, m+1), \\ \rho(T^k S (T^*)^k) &: \xi(k, m) \mapsto \xi(k, m+1), \\ \rho(\phi_q(a)) &: \xi(k, m) \mapsto \sqrt{1-q^{2k}} \xi(k-1, m), \\ \rho(\phi_q(b)) &: \xi(k, m) \mapsto q^k \xi(k, m+1), \end{aligned}$$

with δ being the Kronecker symbol, and the convention that $\xi(k, m) = 0$ whenever $k < 0$.

The irreducible representations of A consist of two families $\{\omega_t \mid t \in \mathbb{T}\}$ and $\{\rho_t \mid t \in \mathbb{T}\}$. Each ω_t is one dimensional and it is determined by $\omega_t(T) = t$, $\omega_t(S) = 0$. Each ρ_t acts on the infinite dimensional Hilbert space $\ell^2(\mathbb{N})$ with an orthonormal basis $\{\xi(n) \mid n \in \mathbb{N}\}$, and it is determined by $\rho_t(S)\xi(n) = \delta_{n,0} t \xi(0)$, $\rho_t(T)\xi(n) = \xi(n+1)$.

We recall the quantum group structure of $SU_q(2)$. The comultiplication

$$\Delta : C(SU_q(2)) \rightarrow C(SU_q(2)) \otimes C(SU_q(2))$$

is a coassociative, unital C^* -algebra homomorphism such that

$$\begin{aligned} \Delta(a) &= a \otimes a - qb^* \otimes b, \\ \Delta(b) &= b \otimes a + a^* \otimes b. \end{aligned}$$

We transport Δ to A via the isomorphism ϕ_q , and denote by

$$\Delta_q = (\phi_q \otimes \phi_q) \Delta \phi_q^{-1}$$

the resulting comultiplication on A . If $q = 0$ then we have

$$\begin{aligned} \Delta_0(T) &= T \otimes T, \\ \Delta_0(S) &= S \otimes T^* + T \otimes S. \end{aligned}$$

For $q > 0$ it is a non-trivial task to express $\Delta_q(T)$ and $\Delta_q(S)$ in terms of the generators T, S .

Lemma 1. *For each $x \in A$ the map $q \mapsto \Delta_q(x)$ is norm continuous.*

Proof. Since the homomorphism Δ_q is isometric and the C^* -algebra A is generated by S and T it suffices to show that the maps $q \mapsto \Delta_q(S)$ and $q \mapsto \Delta_q(T)$ are norm continuous.

Clearly, the maps $q \mapsto \phi_q(a)$ and $q \mapsto \phi_q(b)$ are continuous. It follows that the maps $q \mapsto \Delta_q(\phi_q(a))$ and $q \mapsto \Delta_q(\phi_q(b))$ are continuous. Since 1 is an isolated point of the spectrum of $\phi_q(bb^*)$ the spectral projection $E_{\phi_q(bb^*)}(\{1\})$ is in A . We have $S = \phi_q(b)E_{\phi_q(bb^*)}(\{1\})$ and hence $\Delta_q(S) = \Delta_q(\phi_q(b))E_{\Delta_q(\phi_q(bb^*))}(\{1\})$. It follows that the map $q \mapsto \Delta_q(S)$ is continuous. For each $q \in [0, 1)$ the element $|\phi_q(a^*)|$ is invertible and we have $T = \phi_q(a^*)|\phi_q(a^*)|^{-1}$. Hence $\Delta_q(T) = \Delta_q(\phi_q(a^*))|\Delta_q(\phi_q(a^*))|^{-1}$ and, consequently, the map $q \mapsto \Delta_q(T)$ is continuous. \square

The counit

$$\varepsilon : C(SU_q(2)) \rightarrow \mathbb{C}$$

is a character such that $\varepsilon(a) = 1$ and $\varepsilon(b) = 0$. Slightly abusing notation we write $\varepsilon = \varepsilon \circ \phi_q^{-1}$ for the counit on A . On the generators we have

$$\varepsilon(T) = 1 \quad \text{and} \quad \varepsilon(S) = 0.$$

In particular, ε does not depend on q . For all $q \in [0, 1)$ the following identity, characterizing the counit, holds

$$(\varepsilon \otimes \text{id})\Delta_q = \text{id} = (\text{id} \otimes \varepsilon)\Delta_q.$$

We denote by Pol_q the $*$ -subalgebra of A generated by $\phi_q(a)$ and $\phi_q(b)$. As shown in [2, 3], for each $q \in (0, 1)$ there exists the coinverse (or antipode)

$$\kappa_q : \text{Pol}_q \rightarrow \text{Pol}_q.$$

Pol_q is dense in A , while $\text{Pol}_q \cap J$ is dense in J (since the ideal J is generated by $\phi_q(b)$). We have $\kappa_q(\text{Pol}_q \cap J) = \text{Pol}_q \cap J$.

Let $\pi : A \rightarrow A/J$ be the canonical surjection and let $D = \ker(\pi \otimes \pi)$. It is not difficult to see that D coincides with the closure of $A \otimes J + J \otimes A$. We define B to be the C^* -subalgebra of $A \otimes A$ generated by D and $T \otimes T$. There exists an exact sequence

$$0 \longrightarrow D \longrightarrow B \longrightarrow C(\mathbb{T}) \longrightarrow 0.$$

For all $q \in [0, 1)$ we have

$$\Delta_q(A) \subseteq B \quad \text{and} \quad \Delta_q(J) \subseteq D.$$

Indeed, $\pi(\phi_q(a)) = \pi(T^*)$ and $\pi(\phi_q(b)) = 0$ yield $(\pi \otimes \pi)(\Delta_q(\phi_q(a))) = (\pi \otimes \pi)(T^* \otimes T^*)$ and $(\pi \otimes \pi)(\Delta_q(\phi_q(b))) = 0$. Consequently, both $\Delta_q(\phi_q(a))$ and $\Delta_q(\phi_q(b))$ belong to B and hence $\Delta_q(A) \subseteq B$. Since the ideal J of A is generated by $\phi_q(b)$ and $\Delta_q(\phi_q(b)) \in D$, we have $\Delta_q(J) \subseteq D$.

Also for all $q \in [0, 1)$ we see that

$$\Delta_q(x) - \Delta_0(x) \text{ belongs to } D \text{ for all } x \in A.$$

Indeed, this is easily checked on the generators $\phi_q(a)$ and $\phi_q(b)$ of a dense $*$ -subalgebra of A .

Lemma 2. $\overline{D\Delta_q(J)} = D$ for all $q \in [0, 1)$.

Proof. At first we observe that $\overline{D\Delta_q(J)}$ is a two-sided ideal of $A \otimes A$. We consider separately the cases $q \neq 0$ and $q = 0$. In the former, since κ_q is an antihomomorphism this follows from the fact that $(\kappa_q \otimes \kappa_q)(D \cap (\text{Pol}_q \otimes \text{Pol}_q)) = D \cap (\text{Pol}_q \otimes \text{Pol}_q)$ and $\kappa_q(J \cap \text{Pol}_q) = J \cap \text{Pol}_q$ are dense in D and J , respectively. In the latter, we notice that $D\Delta_0(J)$ is the left ideal of $A \otimes A$ generated by $\{\Delta_0(ST^{-n}) : n = 0, 1, \dots\}$. Thus it suffices to check that $\Delta_0(ST^{-n})x \in D\Delta_0(J)$ for $n = 0, 1, \dots$ and for all x in the set of generators for the algebra $A \otimes A$. This can be done directly. For example,

$$\Delta_0(ST^{-n})S \otimes T = \delta_{n,0}S^2 \otimes I = \delta_{n,0}(S \otimes T + S^2T^* \otimes S^*)\Delta_0(S).$$

Hence $\overline{D\Delta_q(J)}$ is a two-sided ideal of $A \otimes A$ for all $q \in [0, 1)$. Now if $\overline{D\Delta_q(J)} \neq D$ then there existed an irreducible representation of D vanishing on $\overline{D\Delta_q(J)}$. Such a representation of D could be extended to an irreducible representation of $A \otimes A$ unitarily equivalent to $\omega_t \otimes \rho_z$, $\rho_t \otimes \omega_z$ or $\rho_t \otimes \rho_z$, for some $t, z \in \mathbb{T}$. This, however, leads to a contradiction, as one can check directly that none of these representations vanishes on $D\Delta_q(\phi_q(b))$. \square

For $q \in [0, 1)$ we have a faithful representation $(\rho \otimes \rho)\Delta_q$ of A on $\ell^2(\mathbb{N} \times \mathbb{Z}) \otimes \ell^2(\mathbb{N} \times \mathbb{Z})$. Our first goal is to show that all of these representations are unitarily equivalent. To this end we need some preparation.

At first we observe that the restriction of $(\rho \otimes \rho)\Delta_q$ to the ideal J is still a nondegenerate representation. Indeed, otherwise there existed a subspace $\mathcal{H}_0 \neq \{0\}$ of $\ell^2(\mathbb{N} \times \mathbb{Z}) \otimes \ell^2(\mathbb{N} \times \mathbb{Z})$ on which $(\rho \otimes \rho)\Delta_q(J)$ were zero. If $q \neq 0$ then this is impossible since one can verify directly that the kernel of $(\rho \otimes \rho)\Delta_q(\phi_q(b))$ is trivial. In the case $q = 0$ this follows from Lemma 2.

Now for $n \in \mathbb{N}$ we define by induction on k numbers $\lambda_q(n, k) \in (0, +\infty)$ as

$$\begin{aligned}\lambda_q(n, 0) &= 1, \\ \lambda_q(n, k+1) &= \lambda_q(n, k)q^{n+2k+1}(1 - q^{2(n+k+1)})^{-1/2}(1 - q^{2(k+1)})^{-1/2}.\end{aligned}$$

Since

$$(1 - q^{2(n+k+1)})^{-1/2}(1 - q^{2(k+1)})^{-1/2} \leq (1 - q^2)^{-1}$$

it follows that

$$(2) \quad \lambda_q(n, k) \leq (q^{n+k}(1 - q^2)^{-1})^k.$$

In particular, the series $\sum_{k=0}^{+\infty} \lambda_q(n, k)^2$ converges. We set

$$\Lambda_q(n) = \left(\sum_{k=0}^{+\infty} \lambda_q(n, k)^2 \right)^{1/2}.$$

Now for $n, m \in \mathbb{N}$ such that either n or m (or both) equals 0 and for $i, j \in \mathbb{Z}$ we define a vector $f_{n,i,m,j}^q$ in $\ell^2(\mathbb{N} \times \mathbb{Z}) \otimes \ell^2(\mathbb{N} \times \mathbb{Z})$ as

$$f_{n,i,m,j}^q = \sum_{k=0}^{+\infty} \Lambda_q(n+m)^{-1} \lambda_q(n+m, k) \xi(n+k, i-k) \otimes \xi(m+k, j+k).$$

If $q = 0$ then simply $f_{n,i,m,j}^0 = \xi(n, i) \otimes \xi(m, j)$. On the other hand, if $q \in (0, 1)$ then for each vector $\xi(r, x) \otimes \xi(t, y)$, $r, t \in \mathbb{N}$ and $x, y \in \mathbb{Z}$, there exists exactly one vector $f_{n,i,m,j}^q$ so that $\langle \xi(r, x) \otimes \xi(t, y), f_{n,i,m,j}^q \rangle \neq 0$.

Lemma 3. *For each $q \in [0, 1)$ there exists a unique unitary operator U_q on $\ell^2(\mathbb{N} \times \mathbb{Z}) \otimes \ell^2(\mathbb{N} \times \mathbb{Z})$ such that*

$$(\rho \otimes \rho)\Delta_q(x) = U_q((\rho \otimes \rho)\Delta_0(x))U_q^*$$

for all $x \in A$, and

$$U_q : f_{n,i,m,j}^0 \mapsto f_{n,i,m,j}^q$$

for all $i, j \in \mathbb{Z}$ and $n, m \in \mathbb{N}$ with $nm = 0$.

Proof. For each $q \in [0, 1)$, 0 is an isolated point of the spectrum of $\phi_q(a^*a)$ and the corresponding spectral projection is S^*S . Thus, $(\rho \otimes \rho)\Delta_q(S^*S)$ is the projection onto the kernel of $(\rho \otimes \rho)\Delta_q(\phi_q(a))$. It is not difficult to verify that this kernel consists of all vectors $\sum c(n, i, m, j)\xi(n, i) \otimes \xi(m, j)$ in $\ell^2(\mathbb{N} \times \mathbb{Z}) \otimes \ell^2(\mathbb{N} \times \mathbb{Z})$ such that

$$c(n+1, i-1, m+1, j+1) = q^{n+m+1}(1 - q^{2(n+1)})^{-1/2}(1 - q^{2(m+1)})^{-1/2}c(n, i, m, j).$$

This identity implies that vectors $\{f_{n,i,m,j}^q \mid n, m \in \mathbb{N}, nm = 0, i, j \in \mathbb{Z}\}$ form an orthonormal basis of $(\rho \otimes \rho)\Delta_q(S^*S)$.

Since $\Delta_q(S) = \Delta_q(\phi_q(b))\Delta_q(S^*S)$ we find that

$$\begin{aligned}(\rho \otimes \rho)\Delta_q(S)f_{n,i,0,j}^q &= \Lambda_q(n)^{-1} \sum_{k=0}^{+\infty} \xi((n+1)+k, i-k) \otimes \xi(k, (j+1)+k) \\ &\quad \left(\lambda_q(n, k+1)q^{n+k+1}(1 - q^{2(k+1)})^{1/2} + \lambda_q(n, k)q^k(1 - q^{2(n+k+1)})^{1/2} \right).\end{aligned}$$

This implies that $\langle (\rho \otimes \rho)\Delta_q(S)f_{n,i,0,j}^q, f_{r,x,t,y}^q \rangle = 0$ if $(r, x, t, y) \neq (n+1, i, 0, j+1)$. Thus, since $(\rho \otimes \rho)\Delta_q(S)$ is a partial unitary with domain and range $(\rho \otimes \rho)\Delta_q(S^*S)$, we see that $(\rho \otimes \rho)\Delta_q(S)f_{n,i,0,j}^q$ is a scalar multiple of $f_{n+1,i,0,j+1}^q$. However, it is clear from the series expansions of $f_{n+1,i,0,j+1}^q$ and $(\rho \otimes \rho)\Delta_q(S)f_{n,i,0,j}^q$ that this scalar is a positive real number and,

hence, equal to 1. A similar reasoning yields the value of $(\rho \otimes \rho)\Delta_q(S)f_{0,i,m,j}^q$ for $m \neq 0$. We have

$$\begin{aligned}(\rho \otimes \rho)\Delta_q(S)f_{0,i,m,j}^q &= f_{0,i+1,m-1,j}^q, \\(\rho \otimes \rho)\Delta_q(S)f_{n,i,0,j}^q &= f_{n+1,i,0,j+1}^q,\end{aligned}$$

for $i, j \in \mathbb{Z}$ and $n, m \in \mathbb{N}$, $m \neq 0$. We now define an isometry \tilde{U}_q with domain $(\rho \otimes \rho)\Delta_0(S^*S)$ and range $(\rho \otimes \rho)\Delta_q(S^*S)$ by

$$\tilde{U}_q : f_{n,i,m,j}^0 \mapsto f_{n,i,m,j}^q.$$

By construction, we have

$$(\rho \otimes \rho)\Delta_q(S) = \tilde{U}_q((\rho \otimes \rho)\Delta_0(S))\tilde{U}_q^*.$$

Since the restriction of $(\rho \otimes \rho)\Delta_q$ to J is still a nondegenerate representation we can extend \tilde{U}_q to a unitary operator U_q on $\ell^2(\mathbb{N} \times \mathbb{Z}) \otimes \ell^2(\mathbb{N} \times \mathbb{Z})$ by setting

$$(3) \quad U_q = \sum_{k=0}^{+\infty} ((\rho \otimes \rho)\Delta_q(T)^k)\tilde{U}_q(\rho \otimes \rho)\Delta_0(T^*)^k.$$

It follows immediately from the construction and the property of \tilde{U}_q that

$$\begin{aligned}(\rho \otimes \rho)\Delta_q(T) &= U_q((\rho \otimes \rho)\Delta_0(T))U_q^*, \\(\rho \otimes \rho)\Delta_q(S) &= U_q((\rho \otimes \rho)\Delta_0(S))U_q^*.\end{aligned}$$

Furthermore, (3) is the unique extension of \tilde{U}_q which satisfies the above two identities. Since the C^* -algebra A is generated by T and S it follows that

$$(\rho \otimes \rho)\Delta_q(x) = U_q((\rho \otimes \rho)\Delta_0(x))U_q^*$$

for all $x \in A$, as desired. \square

Lemma 4. *For each $q \in [0, 1]$, the unitary U_q of (3) satisfies*

- (i) $U_q(\rho \otimes \rho)(D) \subseteq (\rho \otimes \rho)(D)$ and $(\rho \otimes \rho)(D)U_q \subseteq (\rho \otimes \rho)(D)$.
- (ii) $U_q(\rho \otimes \rho)(J \otimes J) \subseteq (\rho \otimes \rho)(J \otimes J)$ and $(\rho \otimes \rho)(J \otimes J)U_q \subseteq (\rho \otimes \rho)(J \otimes J)$.
- (iii) For each $w \in D$ the maps $q \mapsto U_q(\rho \otimes \rho)(w)$ and $q \mapsto (\rho \otimes \rho)(w)U_q$ are norm continuous.

Furthermore, the partial isometry $\tilde{U}_q : f_{n,i,m,j}^0 \mapsto f_{n,i,m,j}^q$ belongs to $(\rho \otimes \rho)(D)$.

Proof. An elementary calculation on the basis shows that \tilde{U}_q is equal to the sum of the series

$$(4) \quad (\rho \otimes \rho) \left(I \otimes I - TT^* \otimes TT^* + \sum_{\substack{n,m \in \mathbb{N} \\ nm=0}} (\Lambda_q(n+m)^{-1} - 1) T^n S^* ST^{-n} \otimes T^m S^* ST^{-m} \right. \\ \left. + \sum_{\substack{n,m \in \mathbb{N} \\ nm=0}} \Lambda_q(n+m)^{-1} \sum_{k=1}^{+\infty} \lambda_q(n+m, k) T^{n+k} S^{-k} T^{-n} \otimes T^{m+k} S^k T^{-m} \right),$$

and hence belongs to $(\rho \otimes \rho)(D)$. Inequality (2) implies that this series converges in norm and, furthermore, that the map $q \mapsto \tilde{U}_q$ is norm continuous.

By virtue of identity (1) and Lemma 2, to prove $U_q(\rho \otimes \rho)(D) \subseteq (\rho \otimes \rho)(D)$ it suffices to show that $U_q(\rho \otimes \rho)(\Delta_0(T^m S^i T^{-n})w)$ belongs to $(\rho \otimes \rho)(D)$ for all $w \in D$, $m, n \in \mathbb{N}$, $i \in \mathbb{Z}$. Indeed, formula (3) implies that

$$U_q(\rho \otimes \rho)(\Delta_0(T^m S^i T^{-n})w) = (\rho \otimes \rho)(\Delta_q(T^m))\tilde{U}_q(\rho \otimes \rho)(\Delta_0(T^{-m}))(\rho \otimes \rho)(\Delta_0(T^m S^i T^{-n})w).$$

This element belongs to $(\rho \otimes \rho)(D)$, since \tilde{U}_q is in $(\rho \otimes \rho)(D)$. Furthermore, the map $q \mapsto U_q(\rho \otimes \rho)(v)$ is norm continuous for $v = \Delta_0(T^m S^i T^{-n})w$, and consequently it is norm continuous for all $w \in D$. A similar argument yields the inclusion $(\rho \otimes \rho)(D)U_q \subseteq (\rho \otimes \rho)(D)$ and norm continuity of the map $q \mapsto (\rho \otimes \rho)(w)U_q$, $w \in D$. This proves (i) and (iii). Part (ii) follows from (i), since $J \otimes J$ is an ideal of D . \square

We denote by $M(J \otimes J)$ the multiplier C^* -algebra of $J \otimes J$ and by $\mathcal{U}(M(J \otimes J))$ the unitary group of $M(J \otimes J)$. Since $J \otimes J$ is an essential ideal of $A \otimes A$ there exists a canonical imbedding of $A \otimes A$ into $M(J \otimes J)$ and, hence, we may identify $A \otimes A$ with a subalgebra of $M(J \otimes J)$.

Since $\rho \otimes \rho$ is a faithful, non-degenerate representation of $J \otimes J$, it uniquely extends to a faithful representation of $M(J \otimes J)$, which we still denote $\rho \otimes \rho$. If U_q is the unitary operator defined in (3) then U_q belongs to $(\rho \otimes \rho)(M(J \otimes J))$ by Lemma 4. Thus we can define

$$(5) \quad \Omega_q = (\rho \otimes \rho)^{-1}(U_q),$$

$$(6) \quad \tilde{\Omega}_q = (\rho \otimes \rho)^{-1}(\tilde{U}_q).$$

It follows that Ω_q is a unitary in $M(J \otimes J)$ and $\tilde{\Omega}_q$ is a partial isometry in D .

Since $(\varepsilon \otimes \text{id})(D) = J = (\text{id} \otimes \varepsilon)(D)$, there exist unique homomorphism extensions (denoted by the same symbols as the original maps):

$$\begin{array}{ll} \varepsilon \otimes \text{id} : D \rightarrow J & \text{to} \quad \varepsilon \otimes \text{id} : M(D) \rightarrow M(J), \\ \text{id} \otimes \varepsilon : D \rightarrow J & \text{to} \quad \text{id} \otimes \varepsilon : M(D) \rightarrow M(J). \end{array}$$

By Lemma 2, $(\Delta_q \otimes \text{id})(J \otimes J) \cdot D \otimes J$ is dense in $D \otimes J$, and $(\text{id} \otimes \Delta_q)(J \otimes J) \cdot J \otimes D$ is dense in $J \otimes D$. Thus there exist unique homomorphism extensions:

$$\begin{array}{ll} \Delta_q \otimes \text{id} : J \otimes J \rightarrow D \otimes J & \text{to} \quad \Delta_q \otimes \text{id} : M(J \otimes J) \rightarrow M(D \otimes J), \\ \text{id} \otimes \Delta_q : J \otimes J \rightarrow J \otimes D & \text{to} \quad \text{id} \otimes \Delta_q : M(J \otimes J) \rightarrow M(J \otimes D). \end{array}$$

Theorem 5. *For $q \in [0, 1)$ let $\Omega_q \in \mathcal{U}(M(J \otimes J))$ be the unitary defined in (3). Then the following conditions hold true.*

- (i) $\text{Ad } \Omega_q$ is an automorphism of B .
- (ii) $\Delta_q(x) = \Omega_q \Delta_0(x) \Omega_q^*$ for all $x \in A$.
- (iii) The map $[0, 1) \ni q \mapsto \Omega_q \in \mathcal{U}(M(J \otimes J))$ is continuous in the strict topology.
- (iv) $(\varepsilon \otimes \text{id})(\Omega_q) = I = (\text{id} \otimes \varepsilon)(\Omega_q)$.

Proof. The unitary Ω_q satisfies (ii) and $\Omega_q D \Omega_q^* = D$ by Lemma 3. Since $\Omega_q(T \otimes T) \Omega_q^* = \Delta_q(T)$ equals $T \otimes T$ plus an element of D , it follows that $\text{Ad } \Omega_q(B) = B$. That is, condition (i) is satisfied. Condition (iii) follows from Lemma 4.

To prove $(\varepsilon \otimes \text{id})(\Omega_q) = I$ it suffices to show that this equality holds when multiplied on the right by an arbitrary element $x = (\varepsilon \otimes \text{id})(\Delta_0(x))$ of J . By (1), it is enough to take $x = T^m S^i T^{-n}$. We have

$$\begin{aligned} (\varepsilon \otimes \text{id})(\Omega_q)(\varepsilon \otimes \text{id})(\Delta_0(T^m S^i T^{-n})) &= (\varepsilon \otimes \text{id})(\Omega_q \Delta_0(T^m S^i T^{-n})) \\ &= (\varepsilon \otimes \text{id})\left(\Delta_q(T^m) \tilde{\Omega}_q \Delta_0(T^{-m}) \Delta_0(T^m S^i T^{-n})\right). \end{aligned}$$

Using (4) and (6) one can show that this element equals $T^m S^i T^{-n}$. Thus $(\varepsilon \otimes \text{id})(\Omega_q) = I$. Similarly, one shows that $I = (\text{id} \otimes \varepsilon)(\Omega_q)$, and (iv) is proved. \square

Remark 6. Since Δ_q is coassociative it easily follows from condition (ii) of Theorem 5 that the unitary element

$$(7) \quad (\text{id} \otimes \Delta_0)(\Omega_q^*)(I \otimes \Omega_q^*)(\Omega_q \otimes I)(\Delta_0 \otimes \text{id})(\Omega_q)$$

of $M(J \otimes J \otimes J)$ belongs to the relative commutant of $(\Delta_0 \otimes \text{id})\Delta_0(A)$. Thus, following [1], we may call Ω_q a 2-pseudo-cocycle for Δ_0 . We still do not know if Ω_q is a 2-cocycle for Δ_0 ; that is, if it satisfies

$$(8) \quad (\Omega_q \otimes I)(\Delta_0 \otimes \text{id})(\Omega_q) = (I \otimes \Omega_q)(\text{id} \otimes \Delta_0)(\Omega_q).$$

REFERENCES

- [1] M. Enock and L. Vainerman, *Deformation of a Kac algebra by an abelian subgroup*, Commun. Math. Phys. **178** (1996), 571–596.
- [2] S. L. Woronowicz, *Twisted $SU(2)$ group. An example of a non-commutative differential calculus*, Publ. Res. Inst. Math. Sci. **23** (1987), 117–181.
- [3] S. L. Woronowicz, *Compact matrix pseudogroups*, Commun. Math. Phys. **111** (1987), 613–665.

MATHEMATICS, THE UNIVERSITY OF NEWCASTLE, NSW 2308, AUSTRALIA

E-mail address: wojciech@frey.newcastle.edu.au