## Bernoulli Sieve

Alexander V. Gnedin

Department of Mathematics, Utrecht University Postbus 80010, 3508 TA Utrecht, The Netherlands

Abstract. Bernoulli sieve is a recursive construction of a random composition (ordered partition) of integer n. This composition can be induced by sampling from a random discrete distribution which has frequencies equal to the sizes of component intervals of a stick-breaking interval partition of [0, 1]. We exploit Markov property of the composition and its renewal representation to derive asymptotics of the moments and to prove a central limit theorem for the number of parts.

1. The Bernoulli sieve can be seen as a generalisation of the 'game' found in [3]. The first round of the game starts with n players and amounts to tossing a coin with probability  $X_1$ for tails. Each of the players tosses one time and the players flipping tails must drop out. If all n get heads the trial is disqualified and must be repeated completely with all n players, as many times as necessary until some players do quit. If at least one player remains after the first round, the second round continues with the remaining players, who must toss another coin with probability  $X_2$  for tails. The game lasts with probabilities  $X_3, X_4, \ldots$  for tails until all players are sorted out. It is assumed that the probabilities  $X_1, X_2, \ldots$  are independent random variables with a given distribution  $\omega$  on ]0, 1[, and that given  $X_j$  the individual outcomes at round j are conditionally independent. It follows readily that, as far as only the number of players is concerned, the outcome of a round depends on the past solely through the number of players which proceed that far.

A random composition  $C_n$  of integer *n* arises, with part *j* being the number of players dropping out at round *j*. In this paper we shall focus on some properties of  $C_n$ , in particular we are interested in the distribution of the number of parts of the composition, which may be thought of as the duration of the game.

There is a natural way to settle all  $C_n$ 's on the same probability space in a consistent fashion. Consider a random interval partition of [0, 1] by points  $1 - (1 - X_1)(1 - X_2) \dots (1 - X_j)$ ,  $j = 1, 2, \dots$  and assign each player a random uniform tag, independent of the  $X_j$ 's. The tags group within the intervals, and recording the cluster sizes, from the left to the right, yields a composition (intervals containing no tags are ignored). To establish equivalence with the coin-tossing construction we only need to note that the chance for a particular player to remain for at least j rounds in the game is precisely  $(1 - X_1)(1 - X_2) \dots (1 - X_j)$ .

The game in [3] corresponds to  $\omega$  supported by a single point, in which case the  $X_j$ 's are all equal. The composition is induced then by sampling from a geometric distribution and, of course, had appeared many times in the literature under different guises. Karlin distinguished this case in the context of a general occupation problem with infinitely many boxes and derived distributions for the number of parts, number of singletons, doubletons etc. The feature studied in [3], [14] was the probability that there is exactly one winner - a player remaining at the last round (that is to say, the last part of composition is 1).

When  $\omega$  is a Beta $(1,\theta)$  distribution the law of  $C_n$  is known as the ordered Ewens sampling formula (ESF). This structure is well understood, see [1] for a recent account and [18], [10] for generalisations.

Our interest in the construction arose in connection with the regenerative compositions [9]. Within this more general setting the Bernoulli sieve composition may be seen as discretisation of a subordinator with finite Lévy measure and zero drift. In what follows we shall treat general measures, with the only constraint that  $\omega$  is not supported by a geometric sequence like  $(1 - x^j)$  (in particular, sampling from the geometric distribution is ruled out) and such that  $\omega$  does not settle too much mass near the endpoints of [0, 1]. Our method relies on renewal theory and the analysis of 'divide-and-conquer' recurrences, the techniques intended to replace the independence-based tools available in the ESF case, see [1].

2. By exhangeability among the players the compositions  $C_n$  are sampling consistent for different values of n. That is to say, if a part of  $C_n$  is selected at random, in a size-biased fashion, and decremented by one unit then the resulting composition of n-1 (possibly with fewer parts) has the same distribution as  $C_{n-1}$ . The sequence  $(C_n)$  forms a composition structure in the sense of [7],[8] and determine a random exchangeable composition of a countable set.

There are two further constructions of  $C_n$  featuring renewal and Markov properties.

The renewal representation is obtained from the stick-breaking construction by applying transformation  $\phi(x) = -\log(1-x)$  which maps [0,1] onto  $[0,\infty]$ . Consider the range  $\mathcal{R}$  of a renewal process with initial state 0 and step distribution  $\Omega = \omega_{\phi}$ , and let  $E_1, \ldots, E_n$  be increasing order statistics from the standard exponential distribution (which correspond to exponentially distributed tags). The points of  $\mathcal{R}$  induce a partition of  $[0,\infty]$  into intervals making up the compliment  $\mathcal{R}^c = [0,\infty] \setminus \mathcal{R}$ , and the points  $E_j$  group within the intervals; in these terms composition  $C_n$  becomes a record of all nonzero cluster sizes, from the left to the right.

Markov chain representation of  $C_n$  stems from the following first-part deletion property of  $C_n$ . Given the first part of  $C_n$  is m, the composition of n - m obtained by removing this part has the same distribution as  $C_{n-m}$ . This property is obvious in the renewal context: it follows from the regenerative property of  $\mathcal{R}$  (applied at the leftmost point of  $\mathcal{R}$  to the right of  $E_1$ ) taken together with the memoryless property of the exponential distribution. The deletion property implies that the parts of  $C_n$  can be viewed as decrements of a decreasing Markov chain  $Q_n$  which has state-space  $\{0, 1, \ldots, n\}$ , starts at state n and eventually gets absorbed at 0. The one-step transition probability from n to n - m is

$$q(n,m) = \frac{w(n,m)}{1 - w(n,0)}, \qquad m = 1, \dots, n.$$
(1)

where

$$w(n,m) = \binom{n}{m} \int_0^1 x^m (1-x)^{n-m} \omega(\,\mathrm{d}x), \qquad m = 0, 1, \dots, n$$

and the binomial moments of  $\omega$ . Similar expression can be given in terms of  $\Omega$ , with  $1 - e^{-z}$  in place of x. The quantity  $1 - w(n, 0) = w(n, 1) + \ldots + w(n, n)$  will appear throughout as a normalising factor, we use therefore the shorter notation W(n) = 1 - w(n, 0). In other terms

$$W(n) = \int_0^\infty (1 - e^{-nz}) \,\Omega(\mathrm{d}z)$$

is the characteristic exponent of the measure  $\Omega$  thought of as a Lévy measure associated with  $\mathcal{R}$ .

For a given composition  $(n_1, \ldots, n_k)$  of n the probability that  $C_n$  assumes this value is of the product form

$$p(n_1, \dots, n_k) = q(n_1 + \dots + n_k, n_2 + \dots + n_k)q(n_2 + \dots + n_k, n_3 + \dots + n_k) \cdots q(n_k, n_k), \quad (2)$$

because this is the probability that the chain  $Q_n$  has decrements  $n_1, \ldots, n_k$  before absorption in 0.

**3.** We will be interested in the first instance in the number of parts  $K_n$  of composition  $C_n$ . It follows from  $\omega\{1\} = 0$  that q(n,m) > 0 for all  $n \ge m \ge 1$  and  $K_n$  goes to infinity with n.

Observe that the sizes of intervals comprising the partition of [0, 1] are  $Y_j = (1 - X_1) \dots (1 - X_{j-1})X_j$ . Rephrasing the stick-breaking interpretation,  $K_n$  is the number of boxes occupied by at least one of n balls, with probability  $Y_j$  of hitting the jth box. Karlin's paper [12] is a basic reference on the model with infinitely many boxes and nonrandom frequencies, and some information on  $K_n$  can be extracted from Karlin's results by conditioning on  $(Y_j)$ .

Consider two conditions on  $\omega$  which limit concentration of mass near 1 and 0

$$\mu := \int_0^1 |\log(1-x)| \,\omega(\mathrm{d}x) < \infty \,, \tag{3}$$

$$\int_0^1 |\log x| \,\omega(\mathrm{d}x) < \infty \,. \tag{4}$$

Reformulated, the condition (3) says that the first moment of  $\Omega$  is finite:

$$\mu = \int_0^\infty z \,\Omega(\mathrm{d}z) < \infty \,. \tag{5}$$

The unconditional law of large numbers from [12] implies

**Proposition 1** If conditions (3) and (4) are satisfied then, as  $n \to \infty$ ,

$$K_n \sim \frac{1}{\mu} \log n$$

with probability one.

*Proof.* By the strong law of large numbers we have for  $j \to \infty$ 

$$-\frac{1}{j}\log Y_j = -\frac{1}{j}\sum_{i=1}^{j-1}\log(1-X_j) - \frac{1}{j}\log X_j \to \frac{1}{\mu},$$

(condition (4) is necessary and sufficient to have the second term negligible). From this relation we have (for Karlin's alpha on p. 376 of [12])

$$\#\{j: Y_j > 1/x\} \sim \frac{1}{\mu} \log x, \quad \text{as} \quad x \to \infty$$

almost surely. By [12], Theorem 1',

$$\mathbb{E}(K_n|(Y_j)) \sim \frac{1}{\mu} \log n \,,$$

and by Theorem 8 from that paper the statement holds conditionally on  $(Y_j)$ , hence also unconditionally.  $\Box$ 

Note that 'deconditioning' itself does not allow to conclude about the asymptotics for  $\mathbb{E}K_n$  (see Proposition 2 to follow). Results of [12] could be used further to derive the asymptotics of the conditional variance of  $K_n$  and to obtain a conditional central limit theorem. We will not dwell on converting these results into their unconditional counterparts, rather will take an approach based on the renewal features of our model.

4. Let  $F_n$  be the first part of  $C_n$ , with distribution  $\mathbb{P}(F_n = m) = q(n, m)$ . Markov property of the composition implies that  $K_n$  satisfies a distributional equation

$$K_n \stackrel{d}{=} 1 + K'_{n-F_n} \tag{6}$$

where  $F_n, K'_1, K'_2, \ldots$  are independent and each  $K'_j$  has same distribution as  $K_j$ . Averaging in (6) we see that  $a_n = \mathbb{E} K_n$  satisfies a linear recursion

$$a_n = 1 + \sum_{m=1}^{n} q(n,m)a_{n-m}$$
(7)

with boundary value  $a_0 = 0$ .

Remark. Recursions akin to (7) are common in the average-case analysis of algorithms, see references in [19]. A recent dissertation by Bruhn [2] is devoted solely to them. Some results of Bruhn are reproduced in Rösler [20] along with distributional analysis of equations more general than (6). The class of recursions treated in the cited work relates to the assumption that the weights  $q(n, \cdot)$ , considered as measures with support  $\{1/n, 2/n, \ldots, n/n\}$ , satisfy an equiboundedness condition and converge weakly to some measure on [0, 1].

In our case the convergence of  $q(n, \cdot)$  to  $\omega$  is clear from the convergence of moments (which amounts to Bernstein's trick used to prove the Weierstrass uniform approximation theorem). Above that, the Bruhn-Rösler conditions certainly hold when  $\omega$  has a smooth density. However, we were unable to check their (very technical) conditions for the general measures  $\omega$  and will rely on the special structure (1). A specific feature of the class of recursions studied here is that we have a canonical renewal process as a part of the model, while Bruhn and Rösler needed to construct an auxiliary renewal process to 'mimic' the recursion. Remark. We formulate the next fact as  $L^1([0, 1], \omega)$ -approximability of the logarithm by the (ordinary) Bernstein polynomials, but in the sequel we will also make use of the formula (8). There is a variety of closely related results in the literature: the best known is the aforementioned argument due to Bernstein, then there is a number of  $L^1$ -results on generalised Bernstein polynomials [15], and pointwise asymptotic expansions found in [6] and [11]. Still, summation formulas (8), (18) seem to be new.

The Bernstein polynomial of degree n for  $\log(1-x)$  is

$$B_n(x) = \sum_{m=1}^{n-1} \binom{n}{m} x^m (1-x)^{n-m} \log\left(1-\frac{m}{n}\right)$$

**Lemma 1** If  $\omega$  satisfies (3) then

$$\lim_{n \to \infty} \int_0^1 |\mathbf{B}_n(x) - \log(1-x)| \,\omega(\mathrm{d}x) = 0 \,.$$

*Proof.* There is no simple formula for the expectation of the logarithm of binomial random variable but replacing the logarithms by the harmonic numbers, as  $\log(1 - m/n) = h_{n-m} - h_n + o((n-m)^{-1})$ , we have the explicit summation formula:

$$\sum_{m=0}^{n-1} \binom{n}{m} x^m (1-x)^{n-m} (h_{n-m} - h_n) - x^n h_n = -\frac{x}{1} - \frac{x^2}{2} - \dots - \frac{x^n}{n}.$$
 (8)

By monotone convergence the series in the RHS approaches  $\log(1-x)$  in the sense of  $L^1(\omega, [0, 1])$  whatever  $\omega$ . Getting back to  $B_n$  easily yields the claim.

Proposition 1 strongly suggests the logarithmic asymptotics for  $a_n$ . Our proof of this fact will rely on the following simple observation. Given  $n_0 > 1$  suppose  $(a_n)$  satisfies (7) for  $n \ge n_0$ , then  $(a_n + c)$  also satisfies the recursion for  $n \ge n_0$ , whatever constant c.

**Proposition 2** If  $\omega$  meets (3) then any sequence  $(a_n)$  satisfying (7) for  $n \ge n_0 \ge 1$  has asymptotics

$$a_n \sim \frac{\log n}{\mu}$$
.

In particular, this holds for the sequence  $a_n = \mathbb{E}K_n$  which is the unique solution which satisfies (7) for n > 0 and has the boundary value  $a_0 = 0$ .

*Proof.* Assume that there exists  $\epsilon > 0$  such that  $a_n > (1+\epsilon)\mu^{-1}\log n$  for infinitely many values of n. We will lead this to contradiction. Selecting  $\epsilon$  smaller, for any fixed c we could have inequality  $a_n > (1+\epsilon)\mu^{-1}\log n + c$  for infinitely many values of n. Let n(c) be the minimum such n, then  $n(c) \to \infty$  as  $c \to \infty$ . Thus for n < n(c) we have  $a_n < (1+\epsilon)\mu^{-1}\log n + c$  which implies

$$1 + \sum_{m=1}^{n(c)} q(n(c), m) a_{n(c)-m} < 1 + c + \frac{(1+\epsilon)}{\mu} \sum_{m=1}^{n(c)} q(n(c), m) \log(n(c) - m).$$

Now from (7) and the definition of n(c) we derive

$$(1+\epsilon)\frac{\log n(c)}{\mu} + c < 1 + c + \frac{(1+\epsilon)}{\mu} \sum_{m=1}^{n(c)} q(n(c),m)\log(n(c)-m)$$
(9)

where c itself cancels but n(c) can be taken arbitrarily large by the choice of c.

From Lemma 1 we see that

$$\sum_{m=1}^{n-1} q(n,m) \log(n-m) = \log n - \mu + o(1)$$

and substituting this formula into (9) and letting  $c \to \infty$  yields  $0 < -\epsilon$ , which is the promised contradiction. Thus the assumption was wrong and because  $\epsilon$  was arbitrary we have

$$\limsup \frac{a_n}{\mu^{-1}\log n} \le 1.$$

A symmetric argument proves the analogous lower bound, and the claim follows.  $\Box$ 

Turning to the variance of the number of parts  $v_n = \operatorname{Var} K_n$  we derive from (6) a recursion

$$v_n = \left(2a_n - 1 - a_n^2 + \sum_{m=1}^n q(n,m)a_{n-m}^2\right) + \sum_{m=1}^n q(n,m)v_{n-m}, \qquad v_0 = 0 \tag{10}$$

which involves  $a_n = \mathbb{E}K_n$ . Both (7) and (10) are instances of the general equation

$$b_n = r_n + \sum_{m=1}^n q(n,m)b_{n-m}, \qquad b_0 = 0,$$
 (11)

where  $(b_n)$  are unknowns, and  $(r_n)$  is given. The proof of Proposition 2 is easily extended to obtain

**Corollary 1** Assume (3). For any  $n_0$  and  $r \neq 0$ , if  $(b_n)$  satisfies (11) for  $n > n_0$  and if  $r_n \rightarrow r$  then  $b_n \sim r\mu^{-1} \log n$  as  $n \rightarrow \infty$ .

With a logarithmic asymptotics for  $v_n$  in mind, we aim to show the convergence of the bracketed inhomogeneous term in (10). It is easily seen that for this purpose we need more than just the principal-term asymptotics of the expectation, and it is exactly the point where the renewal theory provides indispensable tools.

5. It is well known that a renewal process starting at 0 admits a delayed version which has the expected number of renewals within [0, z] (the potential measure) growing linearly with z, see [5]. It turns out that the stationary renewal process induces a 'stationary' version of the Markov chain  $Q_n$ , which can be used for the asymptotic analysis of (6).

Let g(n, m) be the probability that  $Q_n$  ever visits state m (which means that at some round of the game there are exactly m players left). Since  $Q_n$  can visit each nonabsorbing state at most once g(n, m) is also the potential function, i.e. the expected number of visits to m. Interpreting  $r_m$  as a 'reward' collected at visit to state m, we can think of  $b_n$  satisfying (11) as the total expected reward of  $Q_n$ . The interpretation implies

$$b_n = \sum_{m=1}^{n-1} g(n,m) r_m$$
(12)

and reduces solving (7) to computation of the potential function. Explicit formula is complicated as it involves summation of products over a constrained set of compositions of n. Fortunately, there is a simple asymptotic formula.

Suppose  $\Omega$  is not supported by a lattice, and has finite first moment (5). For  $\omega$  this means (3) and that the support is not a geometric sequence like  $1 - x^j$  (in particular, the case of geometric frequencies, when  $\omega$  is supported by a single point, is excluded). Switching to the renewal representation, we introduce a probability distribution

$$\Omega_0[0,z] = \frac{1}{\mu} \int_0^z \Omega[\zeta,\infty] \,\mathrm{d}\zeta.$$

Let the overshoot B(z) be the distance from z to the leftmost point of  $\mathcal{R}$  to the right of z (B(z) is sometimes called the forward process, or forward recurrence time, or residual lifetime etc.). The renewal theory, as presented in vol. 2 of the Feller's textbook, says that  $\Omega_0$  is the limiting distribution of the overshoot as  $z \to \infty$ . Observe that  $Q_n$  visits m when there is a point of  $\mathcal{R}$  between  $E_{n-m-1}$  and  $E_{n-m}$  or, equivalently, when the overshoot at  $E_{n-m-1}$  does not exceed  $E_{n-m} - E_{n-m-1}$ . The spacing between the two order statistics is independent of  $E_{n-m-1}$  and its distribution is Exponential(m). By the renewal theorem the distribution of  $B(E_{n-m-1})$  converges to  $\Omega_0$  as  $n \to \infty$  because  $E_{n-m-1} \to \infty$  (in probability), thus

$$g(n,m) = \mathbb{P}(B(E_{n-m-1}) < E_{n-m} - E_{n-m-1}) \to \int_0^\infty e^{-mz} \,\Omega_0(\mathrm{d}z) = \frac{1}{\mu \, m} \int_0^\infty (1 - e^{-mz}) \,\Omega(\mathrm{d}z),$$

where the last step follows via integrating by parts. Changing measure back to  $\omega$  we obtain

**Proposition 3** If  $\omega$  is not supported by a geometric sequence and satisfies (3) then for any m

$$\lim_{n \to \infty} g(n, m) = \frac{W(m)}{\mu m}$$

The proposition suggests to modify chain  $Q_n$  so that the potential function becomes exactly

$$g_0(m) := \frac{W(m)}{\mu m}, \qquad m = 1, \dots, n-1.$$

We shall do this by assuming a special distribution for the first transition (which can be thought of as a qualifying round before the game).

Remark. Another possibility were to introduce a proper initial distribution on  $\{0, 1, ..., n\}$  so that the formula for potential function were valid also for m = n. But this would correspond to composition of a random integer, a model we wish to avoid.

Renewal theory offers construction of a stationary version of  $\mathcal{R}$ . Take  $Z_0$  independent of  $\mathcal{R}$ and with distribution  $\Omega_0$ . The shifted set  $\mathcal{R}_0 = Z_0 + \mathcal{R}$  is the range of the stationary (delayed) renewal process. For any  $z \ge 0$  the overshoot distribution for  $\mathcal{R}_0$  at z coincides with  $\Omega_0$ .

The points of  $\mathcal{R}_0$  induce an interval partition of  $[0, \infty]$  thus also a partition of the sequence of order statistics  $E_1, \ldots, E_n$ . Recording the sizes of blocks we obtain a *stationary* composition  $C_{0n}$  of n. The parts of  $C_{0n}$  are considered as decrements of a new Markov chain  $Q_{0n}$ . Repeating the argument which lead us to Proposition 3 we derive from invariance of the distribution of B(z) that  $g_0$  is the potential function of  $Q_{0n}$ .

For any reward function the solution of (11) satisfies

$$\sum_{m=1}^{n-1} g_0(m) r_m = \sum_{m=1}^{n-1} q_0(n,m) b_{n-m},$$
(13)

where  $q_0(n, \cdot)$  is the distribution of the first part of  $C_{0n}$ . This formula follows by computing the total expected reward of  $Q_{0n}$  upon departure from state n. Including state n leads to

$$r_n + \sum_{m=1}^{n-1} g_0(m) r_m = \sum_{m=0}^n w_0(n,m) a_{n-m},$$
(14)

where  $w_0(n, \cdot)$  is the distribution of the number of  $E_j$ 's to the left of  $Z_0$ . Explicitly,

$$w_0(n,m) = \binom{n}{m} \int_0^\infty (1 - e^{-z})^m e^{-(n-m)z} \Omega_0(\mathrm{d}z)$$

and

$$q_0(n,m) = \binom{n}{m} w_0(n,m) + w_0(n,0)q(n,m).$$

And when expressed via binomial moments of  $\omega$  this becomes

$$q_0(n,m) = \frac{1}{\mu} \binom{n}{m} \left( \sum_{k=0}^m (-1)^{m-k} \frac{W(n-k)}{n-k} + \frac{w(n,m)}{n} \right).$$

Remarks. The relation between compositions  $C_{0n}$  and  $C_n$  is that they are identically distributed given the size of the first part. The distribution of  $C_n$  is of the form (2) with the first factor replaced by  $q_0(n, n_1)$ .

The distributional identity  $(C_n) \stackrel{d}{=} (C_{0n})$  holds iff  $\Omega = \Omega_0$ , in which case  $\Omega$  is an exponential distribution,  $\mathcal{R}$  is a homogeneous Poisson point process and therefore  $C_n$  is governed by the ordered ESF. This explains, to an extent, the role of ESF as a 'central limit' because superposition of many rare renewal processes approaches the Poisson process.

For suitable choice of  $\Omega$  the sums in the RHS of (13) or (14) become Cesaro or Euler averages. The LHS is easy to analyse but concluding directly from these relations about the behaviour of  $(a_n)$  is only possible when  $(a_n)$  is known to satify certain regularity conditions (the Tauberian conditions). The direct approach seems hard to realise because the regularity conditions are very sensitive to the summability method. For  $r_n \equiv 1$  the LHS of (14) is the expected number of parts of the stationary composition, which is equal to

$$1 + \sum_{m=1}^{n-1} \frac{W(m)}{m\,\mu}$$

and, quite expectedly, is asymptotic to  $\mu^{-1} \log n$ .

Example. For ESF(1) we have  $W(n) = 1 - (n+1)^{-1}$  and  $\mu = 1$  whence the expected number of parts is the harmonic number  $h_n = 1 + (n-1)^{-1} + \ldots + 3^{-1} + 2^{-1}$  as is well-known [4].

We have seen that  $g(n,m) \to g_0(n,m)$  for  $n \to \infty$  and wish to obtain the asymptotics of (12) by substituting  $g_0$  instead of g. To this end we need a stronger assumption on  $\omega$ 

$$\nu := \int_0^1 (\log(1-x))^2 \,\omega(\mathrm{d}x) < \infty$$

which in terms of  $\Omega$  means finiteness of the second moment,  $\nu = \int_0^\infty x^2 \,\Omega(\mathrm{d}x)$ .

**Proposition 4** Suppose  $\omega$  is not supported by a geometric sequence and also  $\nu < \infty$ . Suppose  $(r_n)$  is such that  $|r_n| < r'_n$  where  $(r'_n)$  is a decreasing sequence satisfying  $\sum r'_n/n < \infty$ . Then for  $(b_n)$  solving (12) we have

$$\lim_{n \to \infty} b_n = \frac{1}{\mu} \sum_{n=1}^{\infty} \frac{W(n)r_n}{n}$$

*Proof.* Given integer J suppose  $(r_n)$  is such that  $r_n = 0$  for n < J, is decreasing for  $n \ge J$  and satisfies  $\sum r_n/n < \infty$ . We wish to show that for the sequence  $(b_n)$  solving (12) with such  $(r_n)$  there is a bound

$$\limsup b_n < \frac{1}{\mu} \sum_{n=1}^{\infty} \frac{r_n}{n} + \frac{r_J \nu}{\mu^2}.$$
 (15)

To this end we will make use of the renewal representation.

Recall that  $Q_n$  collects reward  $r_m$  if the chain visits state m. This occurs when  $\mathcal{R}$  has at least one point between  $E_{n-m}$  and  $E_{n-m+1}$ , in which case let us assign reward  $r_m$  to the rightmost such point (equivalently, given  $E_{n-m}$  is the leftmost point in a cluster, the point of  $\mathcal{R}$  in question is the left endpoint of the component interval  $\subset [0, \infty] \setminus \mathcal{R}$  containing  $E_{n-m}$ ). let U be the potential measure of  $\mathcal{R}$ , so that U[0, z] is the expected cardinality of  $\mathcal{R} \cap [0, z]$ . The total expected reward of  $Q_n$  may be written as

$$r_n W(n) + \int_0^\infty \Phi_n(z) U(\mathrm{d}z)$$

where the first term stands for the reward at  $0 \in \mathcal{R}$ , which is only due in the event that the first jump of the renewal process exceeds  $E_1$ , and the integrand is

$$\Phi_n(z) = \sum_{m=1}^n \binom{n}{m} e^{-zm} (1 - e^{-z})^{n-m} r_m W(m).$$
(16)

In the same manner, we associate the rewards collected by the stationary chain  $Q_{0n}$  with the separating points of  $\mathcal{R}_0$  and with 0 (exceptional point, not in  $\mathcal{R}_0$ ). The expected reward becomes

$$r_n \frac{W(n)}{n\mu} + \int_0^\infty \Phi_n(z) U_0(\mathrm{d}z)$$

where the first term stands for the event that  $E_1 < Z_0$  (i.e.,  $E_1$  falls to the left of  $\mathcal{R}_0$ ). This is, of course, yet another expression for the LHS of (14).

We modify now the reward processes for chains  $Q_n$  and  $Q_{0n}$  by deleting the first term (reward at 0) and by replacing  $\Phi_n$  with another function  $\tilde{\Phi}_n$  defined via (16) but with factors W(m) deleted. Deleting the first term has no asymptotic effect because  $r_n$  goes to 0 as  $n \to \infty$ . We also have  $\Phi_n(z) \leq \tilde{\Phi}_n(z)$ , thus  $\Phi_n(z)$  corresponds to a more generous reward structure, with reward at z being  $r_m$  if  $E_{n-m} < z < E_{n-m+1}$  (thus there is no other contraint on z except that  $z \in \mathcal{R}$ , respectively  $z \in \mathcal{R}_0$ ). The modified reward associated with  $\mathcal{R}_0$  is the sum in the LHS of (15).

The function  $\Phi_n(z)$  is unimodal, with the unique maximum attained at  $z^*$ , which is the unique positive solution of equation

$$-r_J \binom{n-1}{J} + \sum_{m=J}^{n-1} (r_m - r_{m+1}) \binom{n-1}{m} \left(\frac{e^{-z}}{1 - e^{-z}}\right)^{m-J} = 0$$

(the uniqueness follows from the monotonicity of  $(r_j), j > J$ ). For  $n \to \infty$  Poisson approximation provides asymptotics  $ne^{-z^*} \to \zeta$  where  $\zeta$  is the unique positive root of transcendental equation

$$-\frac{r_J}{J!} + \sum_{m=J}^{\infty} (r_m - r_{m+1}) \frac{\zeta^{m-J}}{m!} = 0.$$

In the following argument it is only important that  $z^* \to \infty$  as  $n \to \infty$ .

Because  $\mathcal{R}_0$  is  $\mathcal{R}$  shifted to the right,  $\mathcal{R}_0 = Z_0 + \mathcal{R}$ , there is a one-to-one correspondence between the sets  $\mathcal{R} \cap ](z^* - Z_0)_+, z^*]$  and  $\mathcal{R}_0 \cap ]0, z^*]$ . Furthermore, because  $\widehat{\Phi}_n$  is increasing on  $[0, z^*]$  the total (modified) reward of  $\mathcal{R}_0$  over  $]0, z^*]$  is larger than that of  $\mathcal{R}$  on  $]0, (z^* - Z_0)_+]$ . On the other hand, the expected reward of  $\mathcal{R}$  on  $](z^* - Z_0)_+, z^*]$  has an asymptotic bound  $r_J \nu/(2\mu^2)$ ; indeed  $r_J = \max r_j$  is an upper bound for the instantaneous reward and the potential  $U](z^* - Z_0)_+, z^*]$  is asymptotic to

$$\mathbb{E}\frac{Z_0}{\mu} = \frac{1}{\mu} \int_0^\infty z \,\Omega[z,\infty] \,\mathrm{d}z = \frac{1}{2\mu} \int_0^\infty z^2 \,\Omega(\mathrm{d}z)$$

as it follows from the two-term expansion in the renewal theorem, in the case  $\nu < \infty$  (see [5], Section 4, Chapter XI).

To the right of  $z^*$  the relation is reversed, since the function  $\widehat{\Phi}_n$  is decreasing. Shifting the origin to the leftmost point of  $\mathcal{R}_0 \cap [z^*, \infty]$  enables to view  $\mathcal{R}$  on the new scale as the range of a delayed renewal sequence. Thus the expected reward of  $\mathcal{R}_0$  on  $[z^*, \infty]$  is larger than that of  $\mathcal{R}$ , up to a term estimated by  $r_J \nu/(2\mu^2)$ , exactly as above. Putting the two parts together shows that the expected modified reward is bounded by the RHS of (15). The unmodified reward is smaller, hence (15).

Now suppose  $(r_n)$  is decreasing and satisfies  $\sum r_n/n < \infty$ . We split the sequence at J and decompose it in two:  $r_n = r_n \mathbb{1}_{\{n < J\}} + r_n \mathbb{1}_{\{n \ge J\}}$ . Since recursion (12) is linear, the decomposition forces the representation of solution as, say  $b_n = b'_n + b''_n$ . Applying the renewal theorem we get  $b'_n \to \mu^{-1} \sum_{n=1}^J r_n W(n)/n$ . As for the second part, lim sup  $b''_n$  is estimated with the help of (15) and approaches zero when  $J \to \infty$ , because both  $r_J$  and the tail-sum of the series vanish.

For arbitrary sequence satisfying the condition of proposition splitting at J yields one part converging to  $\mu^{-1} \sum_{n=1}^{J} r_n W(n)/n$  and another part estimated by a solution with reward sequence decreasing for n > J, thus going to 0 as J grows.

Now we are in a position to improve on the asymptotics of  $a_n = \mathbb{E}K_n$ . It will not be supererogatory to remind that asymptotic expansion of the harmonic number starts with  $h_n = \log n + \gamma + O(n^{-1})$ .

**Proposition 5** Suppose  $\omega$  is not supported by a geometric sequence, satisfies (4) and  $\nu < \infty$ . Then

$$a_n = \frac{\log n}{\mu} + \frac{\gamma}{\mu} + b + o(1)$$

where  $\gamma$  is the Euler constant and

$$b = \frac{1}{\mu} \int_0^1 \log x \ \omega(\mathrm{d}x) + \frac{\nu}{2\mu^2}.$$

*Proof.* Writing  $a_n = \mu^{-1}h_n + b_n$ , substituting this into (7) and using the summation formula (8) we find that  $(b_n)$  satisfies (11) with

$$r_n = 1 - \frac{1}{\mu W(n)} \int_0^1 \left(\frac{x}{1} + \frac{x^2}{2} + \dots + \frac{x^n}{n}\right) \,\omega(\mathrm{d}x),$$

which can also be written as

$$r_n W(n) = -\int_0^1 (1-x)^n \,\omega(\mathrm{d}x) + \frac{1}{\mu} \int_0^1 \left(\frac{x^{n+1}}{n+1} + \frac{x^{n+2}}{n+2} + \dots\right) \,\omega(\mathrm{d}x).$$

Using monotone convergence and manipulating the series we find

$$\sum_{n=1}^{\infty} \frac{r_n W(n)}{n\mu} = \frac{1}{\mu} \int_0^1 \log x \,\omega \,(\mathrm{d}x) + \frac{1}{2\mu^2} \int_0^1 (\log(1-x))^2 \,\omega(\mathrm{d}x) = b.$$

Since  $W(n) \to 1$  and  $r_n W(n)$  is the difference of two terms which decrease in n, application of Proposition 4 yields  $b_n \to b$ .  $\Box$ 

6. With no additional assumptions we will derive asymptotics of the variance  $v_n = \operatorname{Var} K_n$ . The key issue is the asymptotic evaluation of the inhomogeneous term of the recursion.

**Lemma 2** Under assumptions of Proposition 5 the expectation  $a_n = \mathbb{E}K_n$  satisfies

$$\lim_{n \to \infty} \left( 2a_n - 1 - a_n^2 + \sum_{m=1}^{n-1} q(n,m)a_{n-m}^2 \right) = \frac{\nu}{\mu^2} - 1.$$

*Proof.* For  $b_n, r_n$  bearing the same meaning as in Proposition 5, we have

$$b_n \to b$$
,  $W(n) \to 1$ ,  $b_n - \sum_{m=1}^n q(n,m)b_{n-m} = r_n$ ,

and integrating by parts yields

$$r_n W(n) = -n \int_0^1 \omega [0, x] (1 - x)^{n-1} dx + \frac{1}{\mu} \int_0^1 \frac{\omega [x, 1] x^n}{1 - x} dx.$$

Further useful estimates follow from the  $n \to \infty$  asymptotics

$$\int_0^1 x^n \,\omega(\mathrm{d}x) = o\left(\frac{1}{\log n}\right), \qquad \int_0^1 (1-x)^n \,\omega(\mathrm{d}x) = o\left(\frac{1}{\log n}\right), \qquad r_n = o\left(\frac{1}{\log n}\right). \tag{17}$$

To justify the first relation, observe that integrability and monotonicity of  $\log(1-x)$  imply that  $\omega[x,1] = o(|\log(1-x)|^{-1})$  for  $x \uparrow 1$  (in fact, the relation is equivalent to the integrability). Integrating by parts and using monotonicity we have

$$\int_0^1 x^n \,\omega(\mathrm{d}x) = \int_0^1 n x^{n-1} \omega[x, 1] \,\mathrm{d}x \,< \mathrm{const} \cdot \,\int_0^1 n x^{n-1} |\log x|^{-1} \,\mathrm{d}x$$

and by a Tauberian argument this is  $o(|\log n|^{-1})$ . The second relation follows in the same way from

$$\int_0^1 (1-x)^n \,\omega(\mathrm{d}x) = n \int_0^1 (1-x)^{n-1} \omega[0,x] \,\mathrm{d}x < \mathrm{const} \cdot \int_0^1 n(1-x)^{n-1} |\log(1-x)|^{-1} \,\mathrm{d}x$$

and  $\omega[0,x] = o(|\log x|^{-1})$  for  $x \downarrow 0$ . And the third relation follows from the first two.

Substituting  $a_n = \mu^{-1}h_n + b_n$  and grouping terms we have

$$2a_n - 1 - a_n^2 + \sum_{m=1}^n q(n,m)a_{n-m}^2 = T_1 + T_2 + T_3 - 1$$

with three to-be-evaluated terms

$$T_{1} = -b_{n}^{2} + \sum_{m=1}^{n} q(n,m)b_{n-m}^{2}$$

$$T_{2} = 2b_{n} - 2b_{n}\frac{h_{n}}{\mu} - \frac{2}{\mu}\sum_{m=1}^{n} q(n,m)b_{n-m}h_{n-m}$$

$$T_{3} = \frac{2h_{n}}{\mu} - \frac{h_{n}^{2}}{\mu^{2}} + \frac{1}{\mu^{2}}\sum_{m=1}^{n} q(n,m)h_{n-m}^{2}.$$

From  $b_n \to b$  it is obvious that  $T_1 \to 0$  as  $n \to \infty$ . To see that also  $T_2$  vanishes write

$$b_{n-m}h_{n-m} = b_{n-m}h_n + (b_{n-m} - b)(h_{n-m} - h_n) + b(h_{n-m} - h_n)$$

then from (8) and (17) we obtain

$$\sum_{m=1}^{n-1} q(n,m)(h_{n-m} - h_n) = \frac{1}{W(n)} \int_0^1 \left( h_n x^n - \sum_{j=1}^n \frac{x^j}{j} \right) \, \omega \mathrm{d}x \,,$$

hence by (17) and Lemma 1

$$b \sum_{m=1}^{n} q(n,m)(h_{n-m} - h_n) \to -b\mu$$
$$\sum_{m=1}^{n} q(n,m)(b_{n-m} - b)(h_{n-m} - h_n) \to 0$$
$$\sum_{m=1}^{n} q(n,m)b_{n-m}h_n = (b_n - r_n)h_n = b_nh_n + o(1)$$

which indeed implies  $T_2 \rightarrow 0$ .

To evaluate  $T_3$  we need a summation formula similar to (8), but this time we should take a combinatorial analogue of  $\log^2$  in place of log. To this end, introduce

$$s_n = \sum_{1 \le i \le j \le n} \frac{1}{i j}$$

then there is a summation formula

$$\sum_{m=0}^{n-1} \binom{n}{m} x^m (1-x)^{n-m} s_{n-m} = s_n - \sum_{j=1}^n \frac{x^j}{j} (h_n - h_{j-1})$$
(18)

where we recognise partial sum of the Taylor series

$$\frac{1}{2}(\log(1-x))^2 = \sum_{j=1}^{\infty} \frac{x^j}{j} h_{j-1}.$$

It follows that

$$\sum_{m=1}^{n-1} q(n,m) s_{n-m} = s_n - \frac{1}{W(n)} \int_0^1 \left( \sum_{j=1}^n \frac{x^j}{j} (h_n - h_{j-1}) \right) \, \omega(\mathrm{d}x)$$

and because  $h_n^2$  differs from  $2s_n$  by the partial sum of a converging series,

$$h_n^2 = 2s_n - \sum_{j=1}^n \frac{1}{j^2},$$

we conclude that

$$\sum_{m=1}^{n-1} q(n,m)h_{n-m}^2 = h_n^2 - \frac{2}{W(n)} \int_0^1 \left( \sum_{j=1}^n \frac{x^j}{j} (h_n - h_{j-1}) \right) + o(1) = h_n^2 - \frac{2\mu h_n}{W(n)} + \frac{\nu}{W(n)} + o(1) = h_n^2 - 2\mu h_n + \nu + o(1)$$

where we exploited monotone convergence and (17). Now it is easily seen that  $T_3 \rightarrow \nu/\mu^2$ .

Putting the terms together we arrive at  $T_1 + T_2 + T_3 - 1 \rightarrow 1 - \nu/\mu^2$ .

Remark. The summation formula (18) implies an analogue of Lemma 1: for arbitrary normalised weight  $\omega$  the square of logarithm is  $L^2([0, 1], \omega)$ -approximable by its Bernstein polynomial.

Appealing to Corollary 1 we obtain the desired asymptotics of variance. Define  $\sigma^2 = \nu - \mu^2$ , that is,  $\sigma^2 = \int_{[0,\infty]} (z-\mu)^2 \Omega(dz)$  is the variance of distribution  $\Omega$ .

**Proposition 6** Under assumptions of Proposition 5

$$\operatorname{Var} K_n \sim \frac{\sigma^2}{\mu^3} \log n$$
.

7. We turn next to the central limit theorem for  $K_n$ . Neininger and Rüschendorf [?] derived a general CLT for solutions of equations like (6). In our context, the assumptions of their Theorem 2.1 are easily checked, with the only exception that their CLT requires some expansion  $\operatorname{Var} K_n = \mu^{-1} \log n + O((\log n)^{1-\epsilon})$ , which is not guaranteed by the integrability of  $(\log(1-x))^2$  rather relies on integrability of a higher power of the logarithm. We shall see that in our situation no additional assumptions are necessary and the CLT follows by a simple comparison with the number of renewals.

Given n, define a *cell* to be a component interval of  $[0, \infty] \setminus \mathcal{R}$  containing at least one  $E_j$ ,  $j \leq n$ . Clearly, the total number of cells is  $K_n$ . Let  $L_n$  be the number of cells which have the left endpoint smaller  $\log n$ , and let  $R_n$  be the number of renewals on  $[0, \log n]$  (including 0), that is  $R_n = \#(\mathcal{R} \cap [0, \log n])$ . It is an easy matter to see that  $L_n \leq R_n$  and  $L_n \leq K_n$ . Moreover, since the expected number of order statistics that exceed  $\log n$  is 1, we have  $\mathbb{E}(K_n - L_n) < 1$ .

**Proposition 7** Under assumptions of Proposition 5,

$$\frac{K_n - \mu^{-1} \log n}{\sigma \mu^{-3/2} \log n} \tag{19}$$

converges weakly to the standard normal random variable.

Proof. By [5] (Section 5, Ch. XI),  $R_n$  is asymptotically normal with expectation  $\mu^{-1} \log n$  and variance  $\sigma^2 \mu^{-3} \log n$ . Furthermore,  $\mathbb{E}R_n = \mu^{-1} \log n + \nu(2\mu^2)^{-1} + o(1)$  ([5], Equation (4.5)). By asymptotics of moments (Propositions 5 and 6) and the above inequalities, the  $L^1$ -distance between any two of the three random variables  $(K_n - a_n)v_n^{-1/2}$ ,  $(L_n - a_n)v_n^{-1/2}$  and  $(R_n - a_n)v_n^{-1/2}$  goes to zero. It follows that  $L_n$  and  $K_n$  are also asymptotically normal.  $\Box$ 

In fact, the renewal theorem taken together with a Poisson limit for the number of  $E_j$ 's exceeding log n implies weak convergence of  $K_n - L_n$ . Asymptotics of the expectation involves the exponential integral function

$$I(z) = \int_{z}^{\infty} e^{-y} y^{-1} \,\mathrm{d}y$$

## Proposition 8 We have

$$\lim_{n \to \infty} \mathbb{E}(K_n - L_n) = \frac{\gamma}{\mu} + \frac{1}{\mu} \int_0^1 \log x \,\omega(\mathrm{d}x) + \int_0^1 I(x) \,\omega(\mathrm{d}x)$$

*Proof.* Recalling (16), using Poisson approximation and the renewal theorem, and changing the variable of integration for  $\zeta = ne^{-z}$  we compute

$$\mathbb{E}(K_n - L_n) = \frac{1}{\mu} \int_{\log n}^{\infty} \Phi_n(z) \, \mathrm{d}z + o(1) = \int_0^1 e^{-\zeta} \sum_{m=1}^{\infty} \frac{\zeta^m W(m)}{m!} \frac{\mathrm{d}\zeta}{\mu\zeta} + o(1) = \int_0^1 \int_0^1 \frac{1 - e^{-zx}}{\mu z} \, \mathrm{d}z \, \omega(\mathrm{d}x) + o(1) = \frac{\gamma}{\mu} + \frac{1}{\mu} \int_0^1 \log x \, \omega(\mathrm{d}x) + \frac{\gamma}{\mu} \int_0^1 I(x) \, \omega(\mathrm{d}x) + o(1)$$

where we also used

$$e^{-\zeta} \sum_{m=1}^{\infty} \frac{\zeta^m (1 - (1 - x)^m)}{m!} = 1 - e^{-\zeta x}$$

and the well-known formula

$$\int_0^x \frac{1 - e^{-y}}{y} \, \mathrm{d}y = I(x) + \log x + \gamma \, \Box$$

Now recalling

$$\mathbb{E}K_n = \frac{\log n}{\mu} + \frac{\gamma}{\mu} + \frac{1}{\mu} \int_0^1 \log x \,\omega \left( \mathrm{d}x \right) + \frac{\nu}{2\mu^2} + o(1)$$

and comparing the expectations

$$\mathbb{E}L_n = \frac{\log n}{\mu} + \frac{\nu}{2\mu^2} - \frac{1}{\mu} \int_0^1 I(x)\,\omega\,(\mathrm{d}x) + o(1)\,, \qquad \mathbb{E}R_n = \frac{\log n}{\mu} + \frac{\nu}{2\mu^2} + o(1)$$

we not only confirm 'by computation' the inequality  $\mathbb{E}L_n \leq \mathbb{E}R_n$  (=  $U[0, \log n]$ ) but also come to the conclusion that the number of component intervals of  $[0, \log n] \setminus \mathcal{R}$  which contain no  $E_j$ 's,  $j \leq n$ , remains bounded as  $n \to \infty$ . This conclusion is in good accord with the general point taken in [9], [10] that the composition  $C_n$  is a proper combinatorial analogue of the regenerative set  $\mathcal{R}.\Box$ 

Acknowledgement. I would like to thank Andrey Levin for help with (18) and further Euler-summation formulas.

## References

- [1] Arratia, R., Barbour, A.G., and Tavaré, S. (2002) Logarithmic Combinatorial Structures: a Probabilistic Approach, forthcoming book.
- [2] Bruhn, V. (1996) Eine Methode zur asymptotischen Behandlung einer Klasse von Rekursionsgleichungen mit einer Anwendung in der stochastischen Analyse des Quicksort-Algorithmus, Dissertation, Universität zu Kiel.

- [3] Bruss, T. and O'Cinneide, C. (1990) On the maximum and its uniqueness for geometric random samples, J. Appl. Prob. 27, 598-610.
- [4] Ewens, W.J. and Tavare, S. (1995) The Ewens sampling formula. In *Multivariate Discrete Distributions* (Johnson, N.S. et al eds), Wiley, NY.
- [5] Feller, W. (1966) An Introduction to Probability Theory and its Applications, vol. 2, Wiley, NY.
- [6] Flajolet, P. (1999) Singularity analysis and asymptotics of Bernoulli sums, *Theoretical Computer Science* 215, 371-381.
- [7] Gnedin, A.V. (1997) The representation of composition structures, Ann. Probab. 25, 1437-1450.
- [8] Gnedin, A.V. (1998) On the Poisson-Dirichlet limit, J. Multivariate Analysis 67, 90-98.
- [9] Gnedin, A.V. and Pitman, J. (2002) Regenerative compositions, (paper in progress).
- [10] Gnedin, A.V. (2002) Three sampling formulas, preprint.
- [11] Jacquet, P. and Szpankowski, W. (1999) Entropy computations via analytic depoissonization, *IEEE Trans. Inform. Theory* 45, 1072-1081.
- [12] Karlin, S. (1967) Central limit theorems for certain infinite urn schemes, J. Math. Mech. 17, 373-401.
- [13] Kingman, J.F.C. (1978) The representation of partition structures, J. London Math. Soc. 18, 374-380.
- [14] Kirschenhofer, P. and Prodinger, H. (1996) The number of winners in a discrete geometrically distributed sample, Ann. Appl. Prob. 6, 687-694.
- [15] Lorenz, G. (1953) Bernstein polynomials, University of Toronto Press, Toronto.
- [16] Neininger, R. and Rüschendorf, L. (2002) On the contraction method with degenerate limit equations, Preprint.
- [17] Pitman, J. (1997) Partition structures derived from Brownian motion and stable subordinators, *Bernoulli* 3, 79-96.
- [18] Pitman, J. (2002) Combinatorial stochastic processes, (Lecture Notes for St. Flour course, July 2002).
- [19] Rösler, U. and Rüschendorf, L. (2001) The contraction method for recursive algorithms, Algorithmica 29, 3-33.
- [20] Rösler, U. (2001) On the analysis of stochastic divide and conquer algorithms, Algorithmica 29, 238-261.

gnedin@math.uu.nl