TROPICAL R AND TAU FUNCTIONS

A. KUNIBA, M. OKADO, T. TAKAGI, AND Y. YAMADA

ABSTRACT. Tropical R is the birational map that intertwines products of geometric crystals and satisfies the Yang-Baxter equation. We show that the $D_n^{(1)}$ tropical R introduced by the authors and its reduction to $A_{2n-1}^{(2)}$ and $C_n^{(1)}$ are equivalent to a system of bilinear difference equations of Hirota type. Associated tropical vertex models admit solutions in terms of tau functions of the BKP and DKP hierarchies.

1. INTRODUCTION

Let $\mathcal{B} = \{x = (x_1, \dots, x_n)\}$ be the set of variables. The tropical R for $A_{n-1}^{(1)}$ [HHIKTT, Ki, Y] is the birational map $R : \mathcal{B} \times \mathcal{B} \to \mathcal{B} \times \mathcal{B}$ specified by

(1.1)

$$R(x,y) = (x',y'), \qquad x'_{i} = y_{i} \frac{P_{i}(x,y)}{P_{i-1}(x,y)}, \quad y'_{i} = x_{i} \frac{P_{i-1}(x,y)}{P_{i}(x,y)},$$

$$P_{i}(x,y) = \sum_{k=1}^{n} \prod_{j=k}^{n} x_{i+j} \prod_{j=1}^{k} y_{i+j},$$

where all the indices are considered to be in $\mathbb{Z}/n\mathbb{Z}$. It satisfies the inversion relation $R^2 = id$ on $\mathcal{B} \times \mathcal{B}$ and the Yang-Baxter equation $R_1R_2R_1 = R_2R_1R_2$ on $\mathcal{B} \times \mathcal{B} \times \mathcal{B}$, where $R_1(x, y, z) = (R(x, y), z)$ and $R_2(x, y, z) = (x, R(y, z))$. Given (x, y), it is characterized as the unique solution to the equations on (x', y'):

(1.2)
$$x_i y_i = x'_i y'_i, \qquad \frac{1}{x_i} + \frac{1}{y_{i+1}} = \frac{1}{x'_i} + \frac{1}{y'_{i+1}}$$

with an extra constraint $\prod_{i=1}^{n} (x_i/y'_i) = \prod_{i=1}^{n} (y_i/x'_i) = 1.$

A representation theoretical background for the tropical R is provided by the geometric crystals [BK] and their natural extrapolation into the affine setting as demonstrated in section 1 in [KOTY]. To recall it consider the matrix realization, where each element $x \in \mathcal{B}$ is associated with the matrix (see also [KNY])

$$M(x,\zeta) = \begin{pmatrix} x_1^{-1} & & -\zeta \\ -1 & x_2^{-1} & & \\ & -1 & \ddots & \\ & & -1 & x_n^{-1} \end{pmatrix}^{-1}$$

involving the spectral parameter ζ . Then the structure of $A_{n-1}^{(1)}$ geometric crystal on \mathcal{B} is realized as simple matrix operations. For instance the geometric Kashiwara operator $e_i^c : (x_1, \ldots, x_n) \mapsto (\ldots, x_{i-1}, cx_i, c^{-1}x_{i+1}, x_{i+2}, \ldots)$ is induced by a multiplication of M with certain unipotent matrices. The product of matrices $M(x^{(1)}, \zeta) \cdots M(x^{(j)}, \zeta)$ corresponds to the product of geometric crystals

 $(x^{(1)}, \ldots, x^{(j)}) \in \mathcal{B} \times \cdots \times \mathcal{B}$ in the sense of [BK]. Eq.(1.2) is equivalent to the matrix equation

(1.3)
$$M(x,\zeta)M(y,\zeta) = M(x',\zeta)M(y',\zeta).$$

Due to the presence of the spectral parameter ζ , its non-trivial solution is unique, which characterizes the tropical R as the intertwiner of the products of geometric crystals.

Geometric crystals are so designed that the relevant rational functions become totally positive [BK, BFZ, L]. There is no minus sign in $P_i(x, y)$ in (1.1) indeed. Such functions can consistently be transformed into piecewise linear ones by replacing +, × and / by max(min), + and -, respectively. Consequently, the structure of geometric crystals reduces to the one for crystal bases [K]. In the present case, the expression (1.1) leads to the piecewise linear formula [HHIKTT] for the combinatorial R [NY]. An analogous result is available in [KOTY] for the $D_n^{(1)}$ combinatorial R [HKOT]. The above transformation can be realized as a certain limiting procedure [TTMS] and is often called the ultradiscretization (UD). As related topics, we remark the tropical combinatorics [Ki, NoY] and the soliton cellular automata associated with crystal bases [TS, TNS, HKT1, FOY, HHIKTT, HKOTY, HKT2].

Now let us turn to the aspect of the tropical R as a classical integrable system, which is the main subject of this paper. Recall the discrete time Toda equation [HT, HTI] on the electric "current" I_i^t and "voltage" V_i^t :

$$I_i^{t+1}V_i^{t+1} = I_{i+1}^t V_i^t, \qquad I_i^{t+1} + V_{i-1}^{t+1} = I_i^t + V_i^t$$

with periodicity $I_i^t = I_{i+n}^t$, $V_i^t = V_{i+n}^t$. This difference equation is known to be integrable. Eq.(1.2) is identified with it via $x_i^{-1} = I_{i+1}^t$, $y_i^{-1} = V_i^t$, $x'_i^{-1} = V_i^{t+1}$ and $y'_i^{-1} = I_i^{t+1}$ [Y]. The matrix equation (1.3) is a Lax representation in this context. In fact the tropical R is equivalent to a system of bilinear difference equations of Hirota type, which was effectively the base of the analyses in [HHIKTT]. To see this, introduce the functions τ_i^J ($1 \leq J \leq 4, i \in \mathbb{Z}/n\mathbb{Z}$) and the parameters λ_i, κ_i , and make the change of variables

$$\begin{aligned} x_i^{-1} &= \lambda_i \delta \tau_i^3 / \delta \tau_i^2, \quad y_i^{-1} &= \kappa_i \delta \tau_i^2 / \delta \tau_i^1, \\ x_i'^{-1} &= \kappa_i \delta \tau_i^3 / \delta \tau_i^4, \quad y_i'^{-1} &= \lambda_i \delta \tau_i^4 / \delta \tau_i^1 \end{aligned}$$

with $\delta \tau_i^J = \tau_i^J / \tau_{i-1}^J$. Then the former relation in (1.2) is automatically satisfied and the latter is translated into

(1.4)
$$\lambda_i \tau_{i-1}^2 \tau_i^4 - \kappa_i \tau_i^2 \tau_{i-1}^4 = \alpha \tau_i^1 \tau_{i-1}^3$$

for any nonzero parameter α independent of *i*. The birational map $R : (x, y) \mapsto (x', y')$ is induced by an automorphism $\tau_i^2 \leftrightarrow \tau_i^4$, $\lambda_i \leftrightarrow \kappa_i$, $\alpha \to -\alpha$ of (1.4). Eq. (1.4) is a version of so-called Hirota-Miwa equation [H, M], a prototype discrete soliton equation to which the well-developed machinery of free fermions and infinite dimensional Lie algebras [DJM, JM] can be applied. The solutions are provided by tau functions of the KP hierarchy with a certain reduction and discretization of time variables.

In this paper we elucidate the classical integrability of such sort for the $D_n^{(1)}$ tropical R introduced in [KOTY] together with its reductions to $A_{2n-1}^{(2)}$ and $C_n^{(1)}$. We show that they are equivalent to a quartet of A type bilinear equations like (1.4), and construct solutions in terms of tau functions in soliton theory [JM]. The

cases $D_n^{(1)}$ and $A_{2n-1}^{(2)}$ are related to reductions of the DKP hierarchy associated to the algebras D_{∞} and D'_{∞} . A key is the relation between two kinds of tau functions that originate in the isomorphism $D_{\infty} \simeq D'_{\infty}$. See Lemmas B.1 and B.2. The $C_n^{(1)}$ case corresponds to a further reduction to the BKP hierarchy associated with the algebras $B_{\infty} \simeq B'_{\infty}$.

We shall formulate tropical vertex models, where the tropical R plays a role of local time evolution. Namely, it is the two dimensional system on a square lattice where each edge is assigned an element of a geometric crystal in such a way that those four surrounding a vertex are related by the tropical R. Our solutions to the bilinear equations are naturally extended to the tropical vertex models by duplicating the original lattice and attaching a Clifford group element to each face of it under a certain rule. The Yang-Baxter equation for the tropical R is naturally understood from such a point of view. See Remark 5.6.

We expect that the results in this paper are glimpses of deeper relations yet to be explored between geometric crystals, crystal bases, solvable lattice models, discrete and ultradiscrete solitons and so forth. A rough picture at present looks as follows.



In [D], Drinfeld proposed set-theoretical solutions of the Yang-Baxter equation as one of unsolved problems in quantum group theory. Since then a number of approaches have been made. See for example [ABS, ESS, JMY, MV, O, V1, V2, WX] and references therein. Our tropical R is a distinguishable example which is placed in the unique spot in the above picture and enjoys the total positivity. It should be noted that the combinatorial R is also a remarkable example of the set-theoretical solution which intertwines products of finite sets.

The paper is organized as follows. In Section 2 we recall the tropical R for $D_n^{(1)}$ following [KOTY]. In Section 3 the bilinear equations are introduced, which are divided into four groups corresponding to each vertex in Figure 1. Uniqueness and existence of the solution are shown in Proposition 3.5. In Section 4 we establish the bilinearization of the tropical R in Theorem 4.4. In Section 5 we construct solutions of the bilinear equations in terms of tau functions of the DKP hierarchy formulated with two component fermions. The final result is given in Theorem 5.5. In Section 6 we introduce the tropical R for $A_{2n-1}^{(2)}$ and $C_n^{(1)}$ as reductions of the $D_n^{(1)}$ case. Parallel results on the bilinearization and solutions are obtained. Appendix A provides a proof of Lemma 4.3. Appendix B contains elements of the free fermion approach [JM] as well as the key lemmas B.1 and B.2. We leave the calculation of the ultradiscrete limit of tau functions as in [HHIKTT] for $A_{n-1}^{(1)}$ as a future problem.

2. Tropical R for $D_n^{(1)}$

Let us recall the tropical R for $D_n^{(1)}$ from [KOTY]. Let $x = (x_1, \ldots, x_n, \overline{x}_{n-1}, \ldots, \overline{x}_1)$ be a set of variables. The geometric $D_n^{(1)}$ -crystal is the set $\mathcal{B} = \{x\}$ equipped with additional structures such as φ_i, ε_i and the geometric Kashiwara operators e_i^c . The product $\mathcal{B} \times \cdots \times \mathcal{B}$ again becomes a geometric crystal, which is an analogue of the tensor product of crystals. The tropical R is a birational map $\mathcal{B} \times \mathcal{B} \to \mathcal{B} \times \mathcal{B}$ commuting with the geometric Kashiwara operators. Leaving the precise definition to [KOTY], we concentrate here on its explicit form.

 Set

(2.1)
$$\ell(x) = x_1 x_2 \cdots x_n \overline{x}_{n-1} \cdots \overline{x}_2 \overline{x}_1,$$

and call it the *level* of x. On \mathcal{B} we introduce the involutions σ_1 and σ_n which preserve the level. Explicitly for $x = (x_1, \ldots, x_n, \overline{x}_{n-1}, \ldots, \overline{x}_1) \in \mathcal{B}$, they are defined by

(2.2)
$$\sigma_1: x_1 \longleftrightarrow \overline{x}_1,$$

(2.3)
$$\sigma_n: x_{n-1} \to x_{n-1} x_n, \quad \overline{x}_{n-1} \to \overline{x}_{n-1} x_n, \quad x_n \to 1/x_n,$$

where the variables not appearing in the above are unchanged. On $\mathcal{B} \times \mathcal{B}$ we introduce σ_1, σ_n analogously as

(2.4)
$$\sigma_a(x,y) = (\sigma_a(x), \sigma_a(y)) \quad a = 1, n.$$

Furthermore we introduce the involution σ_* by

(2.5)
$$\sigma_*: x_i \longleftrightarrow \overline{y}_i, \quad \overline{x}_i \longleftrightarrow y_i \quad (1 \le i \le n-1), \quad x_n \longleftrightarrow y_n$$

for the elements $x = (x_1, \ldots, \overline{x}_1)$ and $y = (y_1, \ldots, \overline{y}_1)$ of \mathcal{B} . Note that σ_1, σ_n and σ_* are all commutative, among which σ_* is the only one that mixes x and yand interchanges the levels. Given a function F = F(x, y), we will write $F^{\sigma_a} = F(\sigma_a(x, y))$ for a = 1, n, *.

We set

(2.6)
$$V_{0} = \ell(x)\frac{y_{1}}{\overline{y}_{1}} + \ell(x)\sum_{m=2}^{n-1} \left(\prod_{i=1}^{m-1} \frac{y_{i}}{x_{i}}\right) \left(1 + \frac{y_{m}}{\overline{y}_{m}}\right) + x_{n}y_{n}\prod_{i=1}^{n-1}\overline{x}_{i}y_{i}$$
$$+ \ell(y)\frac{\overline{x}_{1}}{x_{1}} + \ell(y)\sum_{m=2}^{n-1} \left(\prod_{i=1}^{m-1} \frac{\overline{x}_{i}}{\overline{y}_{i}}\right) \left(1 + \frac{\overline{x}_{m}}{x_{m}}\right) + \prod_{i=1}^{n-1}\overline{x}_{i}y_{i},$$

and define $V_i (1 \le i \le n-1)$ by the recursion relations:

(2.7)
$$V_{i} = \overline{y}_{i} \left(\frac{1}{\overline{x}_{i}} V_{i-1} + (\ell(x) - \ell(y)) \left(\frac{1}{x_{i}} + \frac{1}{\overline{y}_{i}} \right) \right), \quad (1 \le i \le n-2)$$
$$V_{n-1} = \frac{\overline{y}_{n-1}}{y_{n}} \left(\frac{y_{n}}{\overline{x}_{n-1}} V_{n-2} + (\ell(x) - \ell(y)) \left(\frac{y_{n}}{x_{n-1}} + \frac{1}{\overline{y}_{n-1}} \right) \right).$$

In terms of V_i we also define $U_i (1 \le i \le n-1)$ by

(2.8)
$$U_{1} = V_{0}V_{0}^{\sigma_{1}}, \quad U_{n-1} = V_{n-1}V_{n-1}^{\sigma_{*}},$$
$$U_{i} = \left(\frac{1}{x_{i}} + \frac{1}{\overline{y}_{i}}\right)^{-1} \left(\frac{1}{y_{i}}V_{i}V_{i-1}^{\sigma_{*}} + \frac{1}{\overline{x}_{i}}V_{i-1}V_{i}^{\sigma_{*}}\right) \quad (2 \le i \le n-2).$$

The tropical R is a birational map R(x, y) = (x', y') on $\mathcal{B} \times \mathcal{B}$ given by

$$\begin{split} x_1' &= y_1 \frac{V_0^{\sigma_1}}{V_1}, \quad \overline{x}_1' = \overline{y}_1 \frac{V_0}{V_1}, \\ x_i' &= y_i \frac{V_{i-1}U_i}{V_i U_{i-1}}, \quad \overline{x}_i' = \overline{y}_i \frac{V_{i-1}}{V_i}, \quad (2 \le i \le n-1) \\ x_n' &= y_n \frac{V_{n-1}}{V_{n-1}^{\sigma_n}}, \end{split}$$

(2.9)

$$\begin{split} y'_{1} &= x_{1} \frac{V_{0}}{V_{1}^{\sigma_{*}}}, \quad \overline{y}'_{1} = \overline{x}_{1} \frac{V_{0}^{\sigma_{1}}}{V_{1}^{\sigma_{*}}}, \\ y'_{i} &= x_{i} \frac{V_{i-1}^{\sigma_{*}}}{V_{i}^{\sigma_{*}}}, \quad \overline{y}'_{i} = \overline{x}_{i} \frac{V_{i-1}^{\sigma_{*}}U_{i}}{V_{i}^{\sigma_{*}}U_{i-1}}, \quad (2 \leq i \leq n-1) \\ y'_{n} &= x_{n} \frac{V_{n-1}^{\sigma_{n}}}{V_{n-1}}, \end{split}$$

In [KOTY], U_i here was denoted by W_i in Definition 4.9, and their transformation property under σ_a was summarized in Table 1. As shown therein, the tropical Ris subtraction-free (totally positive), interchanges the levels, commutes with the geometric Kashiwara operators, and satisfies the inversion relation R(R(x,y)) = (x, y) and the Yang-Baxter relation.

Remark 2.1. Let (x', y') = R(x, y) under the $D_n^{(1)}$ tropical R. It is easily seen that $x_n = y_n = 1$ is equivalent to $x'_n = y'_n = 1$. Similarly, $(x_1, y_1) = (\overline{x}_1, \overline{y}_1)$ is equivalent to $(x'_1, y'_1) = (\overline{x}'_1, \overline{y}'_1)$.

3. BILINEAR EQUATIONS

Here we introduce a system of bilinear equations and study its properties such as existence and uniqueness of the solution. It is a preparation for Section 4, where the equations are related to our tropical R.

In the rest of the paper we fix the elements $\lambda = (\lambda_1, \ldots, \overline{\lambda}_1)$ and $\kappa = (\kappa_1, \ldots, \overline{\kappa}_1)$ of \mathcal{B} , and set

(3.1)
$$l = \ell(\lambda), \qquad k = \ell(\kappa).$$

We assume $lk(l-k) \neq 0$ throughout. We introduce the variables which we call the *tau functions*:

$$S_i, W_i, N_i, E_i \quad (1 \le i \le n-2), \qquad \tau_i^0, \tau_i^1, \tau_i^2, \tau_i^3, \tau_i^4 \quad (0 \le i \le n).$$

They will be called *generic* if they are nonzero. Let α and β be arbitrary nonzero parameters. Our bilinear equations read as follows:

$$\overline{\kappa}_{1}N_{1}\tau_{1}^{4} - \overline{\lambda}_{1}E_{1}\tau_{1}^{2} = \alpha\tau_{0}^{0}\tau_{0}^{1},$$

$$\kappa_{1}N_{1}\tau_{0}^{4} - \lambda_{1}E_{1}\tau_{0}^{2} = \alpha\tau_{1}^{0}\tau_{1}^{1},$$

$$\kappa_{i}E_{i-1}N_{i} - \lambda_{i}N_{i-1}E_{i} = \alpha\tau_{i}^{0}\tau_{i}^{1} \quad (2 \le i \le n-2),$$

$$\kappa_{n-1}\kappa_{n}E_{n-2}\tau_{n}^{2} - \lambda_{n-1}\lambda_{n}N_{n-2}\tau_{n}^{4} = \alpha\tau_{n-1}^{0}\tau_{n}^{1},$$

$$\kappa_{n-1}E_{n-2}\tau_{n-1}^{2} - \lambda_{n-1}N_{n-2}\tau_{n-1}^{4} = \alpha\tau_{n}^{0}\tau_{n-1}^{1}.$$

The *i*-th equation here will be referred as $\langle 1, i \rangle$ $(0 \le i \le n)$.

$$\begin{split} \kappa_1 N_1 \tau_1^3 + \overline{\lambda}_1 W_1 \tau_1^1 &= \beta \tau_0^0 \tau_0^2, \\ \overline{\kappa}_1 N_1 \tau_0^3 + \lambda_1 W_1 \tau_0^1 &= \beta \tau_1^0 \tau_1^2, \\ \overline{\kappa}_i W_{i-1} N_i + \lambda_i N_{i-1} W_i &= \beta \tau_i^0 \tau_i^2 \quad (2 \le i \le n-2), \\ \kappa_n \overline{\kappa}_{n-1} W_{n-2} \tau_{n-1}^1 + \lambda_{n-1} N_{n-2} \tau_{n-1}^3 &= \beta \tau_{n-1}^0 \tau_{n-1}^2, \\ \overline{\kappa}_{n-1} W_{n-2} \tau_n^1 + \lambda_{n-1} \lambda_n N_{n-2} \tau_n^3 &= \beta \tau_n^0 \tau_n^2. \end{split}$$

The *i*-th equation here will be referred as $\langle 2, i \rangle$ $(0 \le i \le n)$.

$$\begin{aligned} \kappa_1 S_1 \tau_1^2 &- \lambda_1 W_1 \tau_1^4 = \alpha \tau_0^0 \tau_0^3, \\ \overline{\kappa}_1 S_1 \tau_0^2 &- \overline{\lambda}_1 W_1 \tau_0^4 = \alpha \tau_1^0 \tau_1^3, \\ \overline{\kappa}_i W_{i-1} S_i &- \overline{\lambda}_i S_{i-1} W_i = \alpha \tau_i^0 \tau_i^3 \quad (2 \le i \le n-2) \\ \overline{\kappa}_{n-1} W_{n-2} \tau_n^4 &- \overline{\lambda}_{n-1} S_{n-2} \tau_n^2 = \alpha \tau_{n-1}^0 \tau_n^3, \\ \kappa_n \overline{\kappa}_{n-1} W_{n-2} \tau_{n-1}^4 &- \lambda_n \overline{\lambda}_{n-1} S_{n-2} \tau_{n-1}^2 = \alpha \tau_n^0 \tau_{n-1}^3. \end{aligned}$$

The *i*-th equation here will be referred as (3, i) $(0 \le i \le n)$.

$$\begin{aligned} \overline{\kappa}_1 S_1 \tau_1^1 + \lambda_1 E_1 \tau_1^3 &= \beta \tau_0^0 \tau_0^4, \\ \kappa_1 S_1 \tau_0^1 + \overline{\lambda}_1 E_1 \tau_0^3 &= \beta \tau_1^0 \tau_1^4, \\ \kappa_i E_{i-1} S_i + \overline{\lambda}_i S_{i-1} E_i &= \beta \tau_i^0 \tau_i^4 \quad (2 \le i \le n-2), \\ \kappa_{n-1} E_{n-2} \tau_{n-1}^3 + \lambda_n \overline{\lambda}_{n-1} S_{n-2} \tau_{n-1}^1 &= \beta \tau_{n-1}^0 \tau_{n-1}^4, \\ \kappa_{n-1} \kappa_n E_{n-2} \tau_n^3 + \overline{\lambda}_{n-1} S_{n-2} \tau_n^1 &= \beta \tau_n^0 \tau_n^4. \end{aligned}$$

The *i*-th equation here will be referred as $\langle 4, i \rangle$ $(0 \le i \le n)$.

We call $\{\langle J, i \rangle \mid 1 \leq J \leq 4, 0 \leq i \leq n\}$ the $D_n^{(1)}$ bilinear equations.

Lemma 3.1. For generic input $N_1, \ldots, N_{n-2}, W_1, \ldots, W_{n-2}$ and $\tau_0^J, \ldots, \tau_n^J$ with J = 1, 2, 3, there exists a unique solution to the equations $\langle J, i \rangle (1 \le J \le 4, 1 \le i \le n-1)$ on the variables $S_1, \ldots, S_{n-2}, E_1, \ldots, E_{n-2}, \tau_1^0, \ldots, \tau_{n-1}^0$ and $\tau_0^4, \ldots, \tau_n^4$.

Proof. The equations $\langle 2, 1 \rangle - \langle 2, n-1 \rangle$ fix $\tau_1^0, \ldots, \tau_{n-1}^0$. Regard $\langle 1, i \rangle$ and $\langle 3, i \rangle$ with $1 \leq i \leq n-1$ as linear equations on the unknowns $S_1, \ldots, S_{n-2}, E_1, \ldots, E_{n-2}, \tau_0^4$ and τ_n^4 . Arrange them in a matrix form $A\mathbf{x} = \alpha \mathbf{b}$, where the column vectors \mathbf{x} and \mathbf{b} are specified by

$$\mathbf{x} = {}^{t}(\tau_{0}^{4}, E_{1}, \dots, E_{n-2}, \tau_{n}^{4}, S_{n-2}, \dots, S_{1}),$$

$$\mathbf{b} = {}^{t}(\tau_{1}^{0}\tau_{1}^{1}, \dots, \tau_{n-2}^{0}\tau_{n-2}^{1}, \tau_{n-1}^{0}\tau_{n}^{1}, \tau_{n-1}^{0}\tau_{n}^{3}, \tau_{n-2}^{0}\tau_{n-2}^{3}, \dots, \tau_{1}^{0}\tau_{1}^{3}).$$

Then one finds that the 2n-2 by 2n-2 coefficient matrix A is an upper triangular Jacobi matrix except the bottom left element $-\overline{\lambda}_1 W_1$. Hence it is easy to see det $A = (k-l)\tau_0^2\tau_n^2\prod_{i=1}^{n-2}(N_iW_i)$, which is nonzero for generic input. Therefore the unknowns in \mathbf{x} are uniquely determined. With $\tau_1^0, \ldots, \tau_{n-1}^0$ and \mathbf{x} at hand, one can determine the remaining ones $\tau_1^4, \ldots, \tau_{n-1}^4$ from $\langle 4, 1 \rangle - \langle 4, n-1 \rangle$.

According to Lemma 3.1, the bilinear equations $\langle J, 0 \rangle$ and $\langle J, n \rangle$ are not needed to determine the tau functions that will appear in the parameterization (4.2)-(4.3). However, they are essential in order to incorporate the involutions σ_a in the previous section into tau functions. The rest of this section concerns this point, leading to Propositions 3.5, 3.6, and ultimately to Proposition 4.2.

Lemma 3.2. Suppose that S_i, W_i, N_i, E_i $(1 \le i \le n-2)$ and τ_i^J $(0 \le J \le 4, 0 \le i \le n)$ satisfy $\langle J, i \rangle (1 \le J \le 4, 1 \le i \le n-1)$. Then we have

(3.2)
$$\alpha \overline{\lambda}_1 \overline{\kappa}_1 \tau_0^3 \tau_1^1 + \beta \lambda_1 \overline{\kappa}_1 \tau_0^2 \tau_1^4 - \alpha \lambda_1 \kappa_1 \tau_0^1 \tau_1^3 - \beta \overline{\lambda}_1 \kappa_1 \tau_0^4 \tau_1^2 = 0,$$

(3.3)
$$\alpha \tau_{n-1}^{3} \tau_{n}^{1} + \beta \lambda_{n} \tau_{n}^{4} \tau_{n-1}^{2} - \alpha \lambda_{n} \kappa_{n} \tau_{n-1}^{1} \tau_{n}^{3} - \beta \kappa_{n} \tau_{n}^{2} \tau_{n-1}^{4} = 0.$$

Proof. Write $\langle 1, 1 \rangle - \langle 4, 1 \rangle$ in the matrix form:

$$\begin{pmatrix} \kappa_1 \tau_0^4 & -\lambda_1 \tau_0^2 & 0 & 0 \\ 0 & \overline{\lambda}_1 \tau_0^3 & \kappa_1 \tau_0^1 & 0 \\ 0 & 0 & \overline{\kappa}_1 \tau_0^2 & -\overline{\lambda}_1 \tau_0^4 \\ \overline{\kappa}_1 \tau_0^3 & 0 & 0 & \lambda_1 \tau_0^1 \end{pmatrix} \begin{pmatrix} N_1 \\ E_1 \\ S_1 \\ W_1 \end{pmatrix} = \tau_1^0 \begin{pmatrix} \alpha \tau_1^1 \\ \beta \tau_1^4 \\ \alpha \tau_1^3 \\ \beta \tau_1^2 \end{pmatrix}.$$

The matrix on the left hand side is annihilated by multiplying the row vector $(\overline{\lambda}_1 \overline{\kappa}_1 \tau_0^3, \lambda_1 \overline{\kappa}_1 \tau_0^2, -\lambda_1 \kappa_1 \tau_0^1, -\overline{\lambda}_1 \kappa_1 \tau_0^4)$ from the left. So the same should happen on the right hand side, proving (3.2). The relation (3.3) is verified similarly by using $\langle 1, n-1 \rangle - \langle 4, n-1 \rangle$.

Lemma 3.3. Suppose that S_i, W_i, N_i, E_i $(1 \le i \le n-2), \tau_i^J$ $(1 \le J \le 4, 0 \le i \le n)$ and τ_i^0 $(1 \le i \le n-1)$ satisfy the equations $\langle J, i \rangle (1 \le J \le 4, 1 \le i \le n-1)$. Then there exists a unique solution to the equations $\langle J, 0 \rangle$ $(1 \le J \le 4)$ on τ_0^0 .

Proof. It suffices to check compatibility of the four equations $\langle J, 0 \rangle$ $(1 \leq J \leq 4)$, that is, they all lead to the same τ_0^0 . Multiply $\langle 1, 0 \rangle$ by τ_0^3 , and $\langle 3, 0 \rangle$ by τ_0^1 . The difference of the resulting left hand sides can be grouped as

$$\tau_1^4(\overline{\kappa}_1 N_1 \tau_0^3 + \lambda_1 W_1 \tau_0^1) - \tau_1^2(\overline{\lambda}_1 E_1 \tau_0^3 + \kappa_1 S_1 \tau_0^1).$$

Due to $\langle 2, 1 \rangle$ and $\langle 4, 1 \rangle$ this vanishes, proving the compatibility of $\langle 1, 0 \rangle$ and $\langle 3, 0 \rangle$. The compatibility of $\langle 2, 0 \rangle$ and $\langle 4, 0 \rangle$ is similarly shown by using $\langle 1, 1 \rangle$ and $\langle 3, 1 \rangle$. Let us check the compatibility of $\langle 2, 0 \rangle$ and $\langle 3, 0 \rangle$. We are to show

$$\alpha \tau_0^3 (\kappa_1 N_1 \tau_1^3 + \overline{\lambda}_1 W_1 \tau_1^1) - \beta \tau_0^2 (\kappa_1 S_1 \tau_1^2 - \lambda_1 W_1 \tau_1^4) = 0$$

Upon multiplying $\overline{\kappa}_1$, the first term contains the factor $\overline{\kappa}_1 N_1 \tau_0^3$ and the third does $\overline{\kappa}_1 S_1 \tau_0^2$. Eliminate them by using $\langle 2, 1 \rangle$ and $\langle 3, 1 \rangle$, respectively. After a cancellation the resulting expression becomes W_1 times the left hand side of (3.2).

Lemma 3.4. Suppose that S_i, W_i, N_i, E_i $(1 \le i \le n-2), \tau_i^J$ $(1 \le J \le 4, 0 \le i \le n)$ and τ_i^0 $(1 \le i \le n-1)$ satisfy the equations $\langle J, i \rangle (1 \le J \le 4, 1 \le i \le n-1)$. Then there exists a unique solution to the equations $\langle J, n \rangle$ $(1 \le J \le 4)$ on τ_n^0 .

Proof. This is verified similarly to Lemma 3.3 by using (3.3).

Since the compatibility of the two systems $\langle 1, 0 \rangle - \langle 4, 0 \rangle$ and $\langle 1, n \rangle - \langle 4, n \rangle$ is trivial, Lemmas 3.1, 3.3 and 3.4 lead to

Proposition 3.5 (Unique existence). For generic input $N_1, \ldots, N_{n-2}, W_1, \ldots, W_{n-2}$ and $\tau_0^J, \ldots, \tau_n^J$ with J = 1, 2, 3, there exists a unique solution to the bilinear equations $\langle J, i \rangle (1 \leq J \leq 4, 0 \leq i \leq n)$ on the remaining variables $S_1, \ldots, S_{n-2}, E_1, \ldots, E_{n-2}$ and $\tau_0^J, \ldots, \tau_n^J$ (J = 0, 4). Let us proceed to the automorphism of the bilinear equations. For brevity the array $\mathcal{T} = (\lambda, \kappa; S_i, W_i, N_i, E_i, \tau_i^J)$ consisting of all the tau functions together with $(\lambda, \kappa) \in \mathcal{B} \times \mathcal{B}$ will be called *data*. We extend the involutions σ_a (a = 1, n, *) on $\mathcal{B} \times \mathcal{B}$ defined in the previous section to the data as follows:

- (3.4) $\sigma_a \ (a = 1, n, *)$ acts on $(\lambda, \kappa) \in \mathcal{B} \times \mathcal{B}$ by (2.2)-(2.5) as before,
- (3.5) $\sigma_1: \tau_0^J \longleftrightarrow \tau_1^J \ (0 \le J \le 4),$
- (3.6) $\sigma_n: \ \tau_{n-1}^J \longleftrightarrow \tau_n^J \ (0 \le J \le 4),$

(3.7)
$$\sigma_*: \tau_{n-1}^0 \longleftrightarrow \tau_n^0, \quad \tau_{n-1}^2 \longleftrightarrow \tau_n^2, \quad \tau_{n-1}^4 \longleftrightarrow \tau_n^4, \\ \tau_i^1 \longleftrightarrow \tau_i^3 \quad (0 \le i \le n-2), \quad \tau_{n-1}^1 \longleftrightarrow \tau_n^3, \quad \tau_n^1 \longleftrightarrow \tau_{n-1}^3, \\ W_i \longleftrightarrow N_i, \qquad S_i \longleftrightarrow E_i \quad (1 \le i \le n-2).$$

In each involution, the tau functions not appearing in the above are unchanged. See also Proposition 4.2. It is immediate to check

Proposition 3.6 (Invariance). The involutions σ_a (a = 1, n, *) defined in (3.4)-(3.7) on the data \mathcal{T} are automorphism of the bilinear equations $\langle J, i \rangle (1 \leq J \leq 4, 0 \leq i \leq n)$.

In particular, the composition $\sigma_n \sigma_*$ interchanges $\langle 1, i \rangle$ and $\langle 3, i \rangle$ leaving $\langle 2, i \rangle$ and $\langle 4, i \rangle$ invariant.

4. Bilinearization of Tropical R

Given $\mu = (\mu_1, \ldots, \overline{\mu}_1) \in \mathcal{B}$ and the arrays

$$C = (C_1, \dots, C_{n-2}), \quad \tau = (\tau_0, \dots, \tau_n), \quad \tau' = (\tau'_0, \dots, \tau'_n),$$

we define

$$[\mu; \tau, C, \tau'] = (z_1, \dots, z_n, \overline{z}_{n-1}, \dots, \overline{z}_1) \in \mathcal{B},$$

$$z_1 = \mu_1^{-1} \frac{\tau_0 \tau_1'}{C_1}, \quad \overline{z}_1 = \overline{\mu}_1^{-1} \frac{\tau_1 \tau_0'}{C_1},$$

$$z_2 = \mu_2^{-1} \frac{C_1 \tau_2'}{C_2 \tau_0' \tau_1'}, \quad \overline{z}_2 = \overline{\mu}_2^{-1} \frac{C_1 \tau_2}{C_2 \tau_0 \tau_1},$$

$$z_i = \mu_i^{-1} \frac{C_{i-1} \tau_i'}{C_i \tau_{i-1}'}, \quad \overline{z}_i = \overline{\mu}_i^{-1} \frac{C_{i-1} \tau_i}{C_i \tau_{i-1}} \quad (3 \le i \le n-2),$$

$$z_{n-1} = \mu_{n-1}^{-1} \frac{C_{n-2} \tau_{n-1}'}{\tau_{n-1} \tau_{n-2}'}, \quad \overline{z}_{n-1} = \overline{\mu}_{n-1}^{-1} \frac{C_{n-2} \tau_n}{\tau_{n-2} \tau_n'},$$

$$z_n = \mu_n^{-1} \frac{\tau_{n-1} \tau_n'}{\tau_n \tau_{n-1}'}$$

in case $n \ge 4$. For n = 3 we slightly modify it into

$$z_{1} = \mu_{1}^{-1} \frac{\tau_{0} \tau_{1}'}{C_{1}}, \quad \overline{z}_{1} = \overline{\mu}_{1}^{-1} \frac{\tau_{1} \tau_{0}'}{C_{1}},$$

$$z_{2} = \mu_{2}^{-1} \frac{C_{1} \tau_{2}'}{\tau_{2} \tau_{0}' \tau_{1}'}, \quad \overline{z}_{2} = \overline{\mu}_{2}^{-1} \frac{C_{1} \tau_{3}}{\tau_{0} \tau_{1} \tau_{3}'},$$

$$z_{3} = \mu_{3}^{-1} \frac{\tau_{2} \tau_{3}'}{\tau_{3} \tau_{2}'}.$$

Note that $\ell([\mu; \tau, C, \tau']) = \ell(\mu)^{-1}$. From the data $\mathcal{T} = (\lambda, \kappa; S_i, W_i, N_i, E_i, \tau_i^J)$, we construct $x, y, x', y' \in \mathcal{B}$ as

(4.2)
$$x = [\lambda; \tau^3, W, \tau^2], \quad y = [\kappa; \tau^2, N, \tau^1],$$

(4.3) $x' = [\kappa; \tau^3, S, \tau^4], \quad y' = [\lambda; \tau^4, E, \tau^1].$

Note from (3.1) that

(4.4)
$$\ell(x) = \ell(y') = l^{-1}, \quad \ell(x') = \ell(y) = k^{-1}.$$

Given any pairs (x, y) and (λ, κ) such that $\ell(x) = \ell(\lambda)^{-1}$ and $\ell(y) = \ell(\kappa)^{-1}$, the parameterization (4.2) is always possible due to

Proposition 4.1. For any $z, \mu \in \mathcal{B}$ such that $\ell(z) = \ell(\mu)^{-1}$, the equation $[\mu; \tau, C, \tau'] = z$ with fixed τ , (resp. τ') on the variables (C, τ') (resp. (C, τ)) admits a one parameter family of solutions.

It is customary to attach a vertex diagram (cross) to the relation like R(x, y) = (x', y') for a quantum or combinatorial R, where the four edges correspond to x, y, x', y'. Figure 1 is an analogue of such a diagram, which may be of help to recognize the pattern (4.2)-(4.3). Elements of \mathcal{B} are represented there with double lines. Consequently the vertex diagram is separated into several domains where various tau functions live.





It is readily seen that the transformation $\mathcal{T} \to \sigma_a(\mathcal{T})$ of the data induces the change $(x, y) \to \sigma_a(x, y)$ and $(x', y') \to \sigma_a(x', y')$. More generally we have

Proposition 4.2. Let F = F(x, y) be any function on $\mathcal{B} \times \mathcal{B}$ and $\mathcal{T} = (\lambda, \kappa; S_i, W_i, N_i, E_i, \tau_i^J)$ be any data obeying all the bilinear equations. Denote by $F|_{\mathcal{T}}$ an expression of F resulting from the substitution of (4.2) followed by any possible application of the bilinear equations. Then one has $F^{\sigma_a}|_{\mathcal{T}} = \sigma_a(F|_{\mathcal{T}})$ (a = 1, n, *), where σ_a on the left hand side is specified by (2.2)-(2.5), whereas on the right hand side by (3.4)-(3.7). The same fact is valid also for (4.3) and F = F(x', y').

Proof. This is due to Propositions 3.5 and 3.6

Lemma 4.3. Suppose that the data $(\lambda, \kappa; S_i, W_i, N_i, E_i, \tau_i^J)$ satisfies all the bilinear equations. Then under the substitution (4.2), the functions U_i, V_i defined in (2.6)-(2.8) are expressed as follows:

$$\begin{split} V_{0} &= V_{0}^{\sigma_{*}} = \frac{\gamma \tau_{0}^{2} \tau_{0}^{4}}{\tau_{0}^{1} \tau_{0}^{3}}, \ V_{0}^{\sigma_{1}} = \frac{\gamma \tau_{1}^{2} \tau_{1}^{4}}{\tau_{1}^{1} \tau_{1}^{3}}, \ U_{1} = \frac{\gamma^{2} \tau_{0}^{2} \tau_{1}^{2} \tau_{0}^{4} \tau_{1}^{4}}{\tau_{0}^{1} \tau_{1}^{1} \tau_{0}^{3} \tau_{1}^{3}}, \\ V_{1} &= V_{1}^{\sigma_{1}} = \frac{\gamma S_{1} \tau_{0}^{2} \tau_{1}^{2}}{N_{1} \tau_{0}^{3} \tau_{1}^{3}}, \ V_{1}^{\sigma_{*}} = \frac{\gamma E_{1} \tau_{0}^{2} \tau_{1}^{2}}{W_{1} \tau_{0}^{1} \tau_{1}^{1}}, \\ V_{i} &= V_{i}^{\sigma_{1}} = \frac{\gamma S_{i} \tau_{i}^{2}}{N_{i} \tau_{i}^{3}}, \ V_{i}^{\sigma_{*}} = \frac{\gamma E_{i} \tau_{i}^{2}}{W_{i} \tau_{i}^{1}}, \ U_{i} = \frac{\gamma^{2} \tau_{i}^{2} \tau_{i}^{4}}{\tau_{i}^{1} \tau_{i}^{3}} \ (2 \leq i \leq n-2), \\ V_{n-1} &= V_{n-1}^{\sigma_{1}} = \frac{\gamma \tau_{n}^{2} \tau_{n}^{4}}{\tau_{n}^{1} \tau_{n}^{3}}, \ V_{n-1}^{\sigma_{*}} = V_{n-1}^{\sigma_{n}} = \frac{\gamma \tau_{n-1}^{2} \tau_{n-1}^{4}}{\tau_{n-1}^{1} \tau_{n-1}^{3}}, \ U_{n-1} = \frac{\gamma^{2} \tau_{n-1}^{2} \tau_{n}^{2} \tau_{n-1}^{4} \tau_{n}^{4}}{\tau_{n-1}^{1} \tau_{n}^{3} \tau_{n}^{3}}, \\ where \gamma = (k-l)\beta/(lk\alpha). \end{split}$$

The proof is available in Appendix A. The main result of this section is

Theorem 4.4 (Bilinearization of tropical R). Suppose that the data $(\lambda, \kappa; S_i, W_i, N_i, E_i, \tau_i^J)$ satisfies all the bilinear equations. Then x, y, x', y' specified in (4.2) and (4.3) obey the relation R(x, y) = (x', y').

Proof. Substitute (4.2) into the right hand sides of (2.9) and apply Lemma 4.3. Then the result agrees with (4.3).

Given (x, y), the tau functions $\tau^1, \tau^2, \tau^3, N, W$ satisfying (4.2) are not unique. Theorem 4.4 guarantees that (x', y') in (4.3) is independent of their choice under the bilinear equations. From Proposition 4.2 and the remark preceding it, we find that $R(\sigma_a(x, y)) = \sigma_a(R(x, y))$ in agreement with [KOTY]. We conclude that the tropical R acts on the data as the interchange

$$\lambda \longleftrightarrow \kappa, \ \tau_i^2 \longleftrightarrow \tau_i^4, \ W_i \longleftrightarrow S_i, \ N_i \longleftrightarrow E_i$$

for all *i*. Combined with $\alpha \to -\alpha$, this is another automorphism of our bilinear equations that interchanges $\langle 2, i \rangle$ and $\langle 4, i \rangle$ leaving $\langle 1, i \rangle$ and $\langle 3, i \rangle$ invariant. This is a complementary transformation with the $\sigma_n \sigma_*$ mentioned after Proposition 3.6.

5. Solutions of bilinear equations

There is a link between our bilinear equations and the soliton theory in terms of tau functions and infinite dimensional Lie algebras [JM]. We show in Section 5.1 that certain vacuum expectation values of the 2 component free fermions provide solutions to the bilinear equations. They are parametrized with the elements of the algebras $D'_{\infty} \simeq D_{\infty}$ and their reduction. Then in Section 5.2 the solutions for the local equations are naturally extended to the tropical vertex model on the two dimensional square lattice where the tropical R plays the role of local time evolution at each vertex. As for notations and basic facts on the free fermion approach, see Appendix B.

5.1. Tau functions as vacuum expectation values. In this subsection, the letters x, y, etc. stand for the array of infinitely many time variables as in Appendix B. They should not be confused with elements in \mathcal{B} , which have now been effectively replaced by the tau functions. Thus we refresh Figure 1 into the following:

FIGURE 2. Increment time arrays

Here we have introduced the parameters K, L, which will play a role of level in \mathcal{B} . Each line is assigned with the array ε or $\tilde{\varepsilon}$ determined from these parameters according to (B.2). We assign the time variables x^J on the 9 domains J = 0, 1, 2, 3, 4, N, S, W, E in Figure 2 so that the following recursion relations are satisfied:

(5.1)
$$\begin{aligned} x^{W} - x^{3} &= x^{0} - x^{S} = x^{E} - x^{4} = \tilde{\varepsilon}(L^{-1}), \\ x^{2} - x^{W} &= x^{N} - x^{0} = x^{1} - x^{E} = \varepsilon(L^{-1}), \\ x^{N} - x^{2} &= x^{0} - x^{W} = x^{S} - x^{3} = \tilde{\varepsilon}(K^{-1}), \\ x^{1} - x^{N} &= x^{E} - x^{0} = x^{4} - x^{S} = \varepsilon(K^{-1}). \end{aligned}$$

These relations determine all the x^{J} 's consistently upon specifying any one of them as an initial condition. We fix x^{1} to an arbitrary *odd* element in the sense of Appendix B.1. Then from the construction (5.1), all the time variables on the corner domains are odd, i.e.,

(5.2)
$$\widetilde{x^J} = x^J \quad J = 1, 2, 3, 4$$

On the other hand the variables in the domains N, W, S, E satisfy

(5.3)
$$\widetilde{x^{N}} = x^{N} + \varepsilon(K^{-1}) - \tilde{\varepsilon}(K^{-1}), \quad \widetilde{x^{S}} = x^{S} + \varepsilon(K^{-1}) - \tilde{\varepsilon}(K^{-1}),$$
$$\widetilde{x^{W}} = x^{W} + \varepsilon(L^{-1}) - \tilde{\varepsilon}(L^{-1}), \quad \widetilde{x^{E}} = x^{E} + \varepsilon(L^{-1}) - \tilde{\varepsilon}(L^{-1}).$$

For $i \geq 1$ define further

(5.4)
$$\begin{aligned} x_i^J &= x^J + z_i = (x_{i,1}^J, x_{i,2}^J, x_{i,3}^J, \ldots), \\ z_1 &= 0, \quad z_i = -\sum_{k=2}^i \varepsilon(a_k^{-1}) \quad (i \ge 2), \end{aligned}$$

where a_2, a_3, \ldots are nonzero parameters. Note that $x_1^J = x^J$ and $x_i^J + \varepsilon(a_i^{-1}) = x_{i-1}^J$.

Let $g' \in e^{D'_{\infty}}$ and $g \in e^{D_{\infty}}$ be the elements that are related as in (B.5). We shall use the vacuum expectation values (B.6) exclusively for odd $y = (y_1, 0, y_3, 0, \ldots)$ with fixed y_1, y_3, \ldots Thus we simply write them as $F_{l_1, l_2; l}(x; g)$ and $f_l(x; g')$. Our solution to the bilinear equations is constructed in two steps. In *Step 1*, we construct solutions in the infinite *n* limit. In *Step 2*, we will impose certain reduction condition on the elements g, g' to make the equations close for finite *n*.

Step 1. We identify the tau functions with the vacuum expectation values as

(5.5)
$$X_j = F_{1,1}(x_j^X; g) \ j \ge 1 \quad (X = S, W, N, E),$$

(5.6)
$$\tau_j^J = \begin{cases} f_j(x_1^J; g') & j = 0, 1, \\ F_{1,1}(x_j^J; g) & j \ge 2, \end{cases} \quad (J = 1, 2, 3, 4),$$

(5.7)
$$\tau_j^0 = \begin{cases} (F_{0,1} + (-1)^j i F_{0,1;1})(x_1^0; g) & j = 0, 1, \\ F_{1,1}(x_{j-1}^0; g) & j \ge 2. \end{cases}$$

In the bilinear equations $\langle J, i \rangle$ in Section 3, consider the *n* infinity limit, where one just forgets $\langle J, n-1 \rangle$ and $\langle J, n \rangle$ for all J = 1, 2, 3, 4. We call the resulting system the $D_{\infty}^{(1)}$ bilinear equations.

Lemma 5.1.

$$\begin{aligned} F_{1,1}(x_1^J) &= \tau_0^J \tau_1^J \quad J = 1, 2, 3, 4, \\ & 2F_{0,1}(x_1^N;g) \\ & 2iF_{0,1;1}(x_1^N;g) \\ & 2F_{0,1,1}(x_1^N;g) \\ & 2F_{0,1,1}(x_1^W;g) \\ & 2F_{0,1,1}(x_1^W;g) \\ & 2F_{0,1;1}(x_1^S;g) \\ & 2F_{0,1;1}(x_1^S;g) \\ & 2F_{0,1;1}(x_1^S;g) \\ & 2F_{0,1;1}(x_1^E;g) \\ \end{aligned} \right\} = \tau_1^4 \tau_1^1 \pm \tau_0^4 \tau_0^1. \end{aligned}$$

Proof. The first equality follows from Lemma B.1 and (5.2). The other relations are due to Lemma B.2 and (5.3). \Box

Proposition 5.2. The parameterization (5.5)-(5.7) solves the $D_{\infty}^{(1)}$ bilinear equations with $(a_1 = 0)$

$$\lambda_i = L - a_i, \ \overline{\lambda}_i = L + a_i \quad (i \ge 1),$$

$$\kappa_i = K - a_i, \ \overline{\kappa}_i = K + a_i \quad (i \ge 1),$$

$$\alpha = K - L, \ \beta = K + L.$$

Proof. First consider $\langle J, i \rangle$ with $i \ge 2$. Setting $l_1 = l_2 = 1, l = 0, x = x_i^0, b_3 = a_i$ in (B.13), we have

$$\begin{aligned} &(b_2 - a_i)F_{1,1}(x_{i-1}^0 + \varepsilon(b_2^{-1}))F_{1,1}(x_i^0 + \varepsilon(b_1^{-1})) \\ &- (b_1 - a_i)F_{1,1}(x_{i-1}^0 + \varepsilon(b_1^{-1}))F_{1,1}(x_i^0 + \varepsilon(b_2^{-1})) \\ &= (b_2 - b_1)F_{1,1}(x_{i-1}^0)F_{1,1}(x_i^0 + \varepsilon(b_1^{-1}) + \varepsilon(b_2^{-1})), \end{aligned}$$

where the dependence on $g \in e^{D_{\infty}}$ is suppressed. Due to (5.1) this yields $\langle J, i \rangle$ with $1 \leq J \leq 4$ upon taking $(b_1, b_2) = (L, K), (L, -K), (-L, -K), (-L, K)$. Next we treat $\langle J, 0 \rangle$ and $\langle J, 1 \rangle$. Setting $l_1 = l_2 = 1, l = 0, x = x_1^0, b_3^{-1} = 0$ in (B.12) $\mp i$ (B.11), we have

$$\begin{split} b_2 F_{1,1}(x_1^0 + \varepsilon(b_1^{-1}))(F_{0,1} \mp iF_{0,1;1})(x_1^0 + \varepsilon(b_2^{-1})) \\ &- b_1 F_{1,1}(x_1^0 + \varepsilon(b_2^{-1}))(F_{0,1} \mp iF_{0,1;1})(x_1^0 + \varepsilon(b_1^{-1})) \\ &= (b_2 - b_1)F_{1,1}(x_1^0 + \varepsilon(b_1^{-1}) + \varepsilon(b_2^{-1}))(F_{0,1} \mp iF_{0,1;1})(x_1^0) \end{split}$$

By setting $(b_1, b_2) = (L, K)$, (L, -K), (-L, -K), (-L, K), and applying Lemma 5.1, this becomes $\langle J, 1 \rangle \tau_0^J$ and $\langle J, 0 \rangle \tau_1^J$ for $1 \leq J \leq 4$.

Step 2. We choose the elements $g' \in e^{D'_{\infty}}$ and $g \in e^{D_{\infty}}$ related by (B.5) as

(5.8)
$$g' = \exp\left(\sum_{j=1}^{N} b_j \phi(p_j) \phi(q_j) + \sum_{j=1}^{M} c_j \phi(p'_j) \hat{\phi}(q'_j)\right),$$

(5.9)
$$g = \exp\left(\sum_{j=1}^{N} b_j \left(\psi^{(1)}(p_j)\psi^{(1)*}(-q_j) - \psi^{(1)}(q_j)\psi^{(1)*}(-p_j)\right) + \sum_{j=1}^{M} c_j \left(\psi^{(1)}(p_j')\psi^{(2)*}(-q_j') - \psi^{(2)}(q_j')\psi^{(1)*}(-p_j')\right)\right)$$

On these elements we impose the reduction condition:

(5.10)
$$p_j^2 A(p_j) A(-p_j) = q_j^2 A(q_j) A(-q_j), \qquad {p'}_j^2 A(p'_j) A(-p'_j) = {q'}_j^2$$

for all j, where the function A is defined by

$$A(p) = \prod_{k=2}^{n-1} (1 - \frac{p}{a_k})$$

in terms of a_2, a_3, \ldots introduced in (5.4).

Lemma 5.3. Under the condition (5.10), there exists $h' \in e^{D'_{\infty}}$ and $h \in e^{D_{\infty}}$ that are related by $h = \iota(h')\kappa\iota(h')$ as in (B.5) and satisfy

$$F_{l_1,l_2;l}(x+z_{n-1},y;\omega(g)) = F_{l_1,l_2;l}(x,y;h)$$

for any x and odd y.

Proof. Under the map ω specified in (B.3), $\psi^{(\alpha)}(p)$ and $\psi^{(\alpha)*}(q)$ are transformed into $p\psi^{(\alpha)}(p)$ and $q^{-1}\psi^{(\alpha)*}(q)$. Then under the time evolution $\operatorname{Ad} e^{H(z_{n-1},0)}$, they are further changed into $pA(p)^{\delta_{\alpha 1}}\psi^{(\alpha)}(p)$ and $q^{-1}A(q)^{-\delta_{\alpha 1}}\psi^{(\alpha)*}(q)$, respectively. Therefore if g in (5.9) is denoted by e^X , we get $\operatorname{Ad} e^{H(z_{n-1},0)}(\omega(g)) = e^{X'}$ with

$$\begin{aligned} X' &= \sum_{j} b_{j} \left(-\frac{p_{j} A(p_{j})}{q_{j} A(-q_{j})} \psi^{(1)}(p_{j}) \psi^{(1)*}(-q_{j}) + \frac{q_{j} A(q_{j})}{p_{j} A(-p_{j})} \psi^{(1)}(q_{j}) \psi^{(1)*}(-p_{j}) \right) \\ &+ \sum_{j} c_{j} \left(-\frac{p_{j}' A(p_{j}')}{q_{j}'} \psi^{(1)}(p_{j}') \psi^{(2)*}(-q_{j}') + \frac{q_{j}'}{p_{j}' A(-p_{j}')} \psi^{(2)}(q_{j}') \psi^{(1)*}(-p_{j}') \right). \end{aligned}$$

Under the condition (5.10), we find

$$\begin{split} X' &= X|_{b_j \to b'_j, c \to c'_j}, \\ b'_j &= -\frac{q_j A(q_j)}{p_j A(-p_j)} b_j, \quad c'_j = -\frac{p'_j A(p'_j)}{q'_j} c_j \end{split}$$

In view of the definition (B.6) and $H(x + z_{n-1}, y) = H(x, y) + H(z_{n-1}, 0)$, the elements h' and h in question are obtained by replacing b_j, c_j by b'_j, c'_j in g' and g given in (5.8) and (5.9), respectively.

Using h' and h in Lemma 5.3, we now modify (5.5)–(5.7) in *Step 1* into the finite n version as

(5.12) $X_{j} = F_{1,1}(x_{j}^{X};g) \ 1 \le j \le n-2 \quad (X = S, W, N, E),$ (5.13) $\tau_{j}^{J} = \begin{cases} f_{j}(x_{1}^{J};g') & j = 0, 1, \\ F_{1,1}(x_{j}^{J};g) & 2 \le j \le n-2, \quad (J = 1, 2, 3, 4), \\ f_{n-j}(x_{1}^{J};h') & j = n-1, n, \end{cases}$ (5.14) $0 \qquad \begin{cases} (F_{0,1} + (-1)^{j}iF_{0,1;1})(x_{1}^{0};g) & j = 0, 1, \\ F_{0,1} + (-1)^{j}iF_{0,1;1})(x_{1}^{0};g) & j = 0, 1, \end{cases}$

(5.14)
$$\tau_j^0 = \begin{cases} F_{1,1}(x_{j-1}^0;g) & 2 \le j \le n-2, \\ (F_{0,0} + \frac{(-1)^{n-j}ia_{n-1}}{KL}F_{-1,1;1})(x_{n-2}^0;\omega(g)) & j = n-1, n. \end{cases}$$

Lemma 5.4.

$$\begin{split} F_{0,0}(x_{n-1}^{J};\omega(g)) &= \tau_{n-1}^{J}\tau_{n}^{J} \quad J = 1, 2, 3, 4, \\ & 2F_{0,0}(x_{n-1}^{N};\omega(g)) \\ 2iK^{-1}F_{-1,1;1}(x_{n-1}^{N};\omega(g)) \\ & 2iL^{-1}F_{-1,1;1}(x_{n-1}^{N};\omega(g)) \\ & 2iL^{-1}F_{-1,1;1}(x_{n-1}^{W};\omega(g)) \\ & 2iL^{-1}F_{-1,1;1}(x_{n-1}^{S};\omega(g)) \\ & 2iK^{-1}F_{-1,1;1}(x_{n-1}^{S};\omega(g)) \\ & 2iK^{-1}F_{-1,1;1}(x_{n-1}^{S};\omega(g)) \\ & 2iK^{-1}F_{-1,1;1}(x_{n-1}^{S};\omega(g)) \\ & 2iL^{-1}F_{-1,1;1}(x_{n-1}^{S};\omega(g)) \\ & 2iL^{-1}F_{-1$$

Proof. The first relation follows from Lemma 5.3, Lemma B.1 and (5.2). From Lemma 5.3, the left hand side of the second equation $2F_{0,0}(x_{n-1}^N; \omega(g))$ is equal to $2F_{0,0}(x_1^N; h)$. From (5.1), (5.3) and Lemma B.2, we get

$$2F_{0,0}(x_1^N;h) = f_0(x_1^1;h')f_1(x_1^2;h') + f_0(x_1^2;h')f_1(x_1^1;h'),$$

which is the right hand side. The other relations can be checked similarly. $\hfill \Box$

Theorem 5.5. The parameterization (5.12)-(5.14) solves the $D_n^{(1)}$ bilinear equations with $(a_1 = 0)$

$$\begin{split} \lambda_i &= L - a_i, \ \overline{\lambda}_i = L + a_i \quad (1 \leq i \leq n - 1), \ \lambda_n = 1, \\ \kappa_i &= K - a_i, \ \overline{\kappa}_i = K + a_i \quad (1 \leq i \leq n - 1), \ \kappa_n = 1, \\ \alpha &= K - L, \ \beta = K + L. \end{split}$$

Proof. Thanks to Proposition 5.2 we are left to show $\langle J, n-1 \rangle$ and $\langle J, n \rangle$ only. Let us illustrate the proof for J = 1 since the equations for the other J follow from the same argument by taking the parameters b_1, b_2 in the proof similarly to Proposition

5.2. In (B.13) set $l_1 = l_2 = l = 0$, $(b_1, b_2) = (L, K)$, $b_3 = a_{n-1}$, $x = x_{n-1}^0$ and $g \to \omega(g)$. The result reads

$$\kappa_{n-1}F_{0,0}(x_{n-2}^E;\omega(g))F_{0,0}(x_{n-1}^N;\omega(g)) - \lambda_{n-1}F_{0,0}(x_{n-2}^N;\omega(g))F_{0,0}(x_{n-1}^E;\omega(g)) = (K-L)F_{0,0}(x_{n-1}^1;\omega(g))F_{0,0}(x_{n-2}^0;\omega(g)).$$

Owing to (B.8), we know $F_{0,0}(x_{n-2}^E;\omega(g)) = F_{1,1}(x_{n-2}^E;g) = E_{n-2}$ and similarly $F_{0,0}(x_{n-2}^N;\omega(g)) = N_{n-2}$. Rewriting $F_{0,0}(x_{n-1}^1;\omega(g)), F_{0,0}(x_{n-1}^N;\omega(g))$ and $F_{0,0}(x_{n-1}^E;\omega(g))$ by Lemma 5.4, we find that the above equation coincides with the combination $\langle 1, n-1 \rangle \tau_{n-1}^1 + \langle 1, n \rangle \tau_n^1$. The other combination $\langle 1, n-1 \rangle \tau_{n-1}^1 - \langle 1, n \rangle \tau_n^1$ can be shown by using (B.10) with $l_1 = l_2 = l = 0$ similarly.

5.2. Tropical vertex model. Consider the two dimensional square lattice \mathcal{L}' equipped with the space-time coordinates $(s,t) \in \mathbb{Z}^2$. The coordinate s (resp. t) increases rightward (resp. downward) and each vertical (resp. horizontal) line corresponds to s = constant (resp. t = constant). On each edge of \mathcal{L}' we assign an element in \mathcal{B} so that those four surrounding a vertex obey the relation R(x,y) = (x',y'), where x, y, x', y' are the ones on the east, north, south and west edges. We call the resulting two dimensional system the *tropical vertex model*. Unlike the usual vertex models in statistical mechanics [B], it is a deterministic system in the sense that all the edge variables are determined uniquely from their values on the northwest or southeast boundaries of \mathcal{L}' . The tropical vertex model reduces to the $D_n^{(1)}$ -soliton cellular automaton [HKT1, HKT2, HKOTY] in the ultradiscrete limit, where the tropical R is replaced by the combinatorial R [HKOT] and \mathcal{B} by the $D_n^{(1)}$ crystal [KKM]. From (4.4) it follows that all the elements in \mathcal{B} on the same line possess the same level. We let the levels be l_t^{-1} (resp. k_s^{-1}) on the *t*th horizontal (resp. *s*th vertical) line.

The bilinearization of the tropical R attained in Section 4 leads to another formulation of the tropical vertex model in terms of tau functions. In view of Figure 2 we duplicate each line in \mathcal{L}' into a pair of parallel lines to form a new lattice \mathcal{L} . Its unit structure looks as Figure 2 where we now replace (L, K) by (L(t), K(s)). To each face of \mathcal{L} we assign a coordinate (s, t) which now takes values in $(\mathbb{Z}/2)^2$ so that $\tau^0: (s,t) \in \mathbb{Z}^2$, $N: (s,t-\frac{1}{2})$, $W: (s-\frac{1}{2},t)$, $S: (s,t+\frac{1}{2})$, $E: (s+\frac{1}{2},t)$, $\tau^1: (s+\frac{1}{2},t-\frac{1}{2})$, $\tau^2: (s-\frac{1}{2},t-\frac{1}{2})$, $\tau^3: (s-\frac{1}{2},t+\frac{1}{2})$, $\tau^4: (s+\frac{1}{2},t+\frac{1}{2})$. There are three types of faces depending on whether the coordinate (s,t) belongs to \mathbb{Z}^2 , $(\mathbb{Z}+\frac{1}{2})^2$ or else. To the face of \mathcal{L} at (s,t), we associate a tau function $\tau_i(s,t)$ having the components $1 \leq i \leq n-2$ if $s+t \in \mathbb{Z}+\frac{1}{2}$ and $0 \leq i \leq n$ otherwise. Then up to a boundary condition, the tropical vertex model is equivalent to imposing the $D_n^{(1)}$ bilinear equations on the tau functions around each face at $(s,t) \in \mathbb{Z}^2$.

Let us construct such system of tau functions $\{\tau_i(s,t)\}$ by making use of the result in Section 5.1, which may be regarded as a solution of our tropical vertex model. To each face of \mathcal{L} at $(s,t) \in (\mathbb{Z}/2)^2$ we attach the time variables

$$x_{j}(s,t) = x(s,t) + z_{j},$$

$$x(s,t) = \eta + \sum_{t' \ge t} \varepsilon(L(t')^{-1}) + \sum_{t' > t} \tilde{\varepsilon}(L(t')^{-1}) + \sum_{s' < s} \varepsilon(K(s')^{-1}) + \sum_{s' \le s} \tilde{\varepsilon}(K(s')^{-1}),$$

where z_j is defined in (5.4) and $\eta = \tilde{\eta}$ is an arbitrary odd time. The sums extend over $s', t' \in \mathbb{Z}$. Let $g, h \in e^{D_{\infty}}, g', h' \in e^{D'_{\infty}}$ be as in (5.12)–(5.14). Set

$$\begin{aligned} \tau_j(s,t) &= F_{1,1}(x_j;g), \ 1 \le j \le n-2, \ s+t \in \mathbb{Z}+1/2, \\ &= \begin{cases} f_j(x_1;g') & j=0,1, \\ F_{1,1}(x_j;g) & 2 \le j \le n-2, \quad (s,t) \in (\mathbb{Z}+1/2)^2, \\ f_{n-j}(x_1;h') & j=n-1,n, \end{cases} \\ &= \begin{cases} (F_{0,1}+(-1)^j i F_{0,1;1})(x_1;g) & j=0,1, \\ F_{1,1}(x_{j-1};g) & 2 \le j \le n-2, \quad (s,t) \in \mathbb{Z}^2, \\ (F_{0,0}+\frac{(-1)^{n-j} i a_{n-1}}{K(s)L(t)}F_{-1,1;1})(x_{n-2};\omega(g)) & j=n-1,n, \end{cases} \end{aligned}$$

where $x_j = x_j(s,t)$, and the time variable y suppressed here is taken as explained in the paragraph preceding (5.5). Since x(s,t) obeys the recursion relation corresponding to (5.1), we conclude that $\{\tau_j(s,t)\}$ satisfies all the $D_n^{(1)}$ bilinear equations if the parameters $\lambda, \kappa, \alpha, \beta$ at $(s,t) \in \mathbb{Z}^2$ are identified with those in Theorem 5.5 with (L, K) replaced by (L(t), K(s)). In particular, the levels l_t^{-1} and k_s^{-1} of the edge variables are determined by

$$l_t = L(t)^2 \prod_{i=2}^{n-1} (L(t)^2 - a_i^2), \qquad k_s = K(t)^2 \prod_{i=2}^{n-1} (K(s)^2 - a_i^2).$$

Note that the formal sums over s', t' in x(s, t) make sense under a suitable choice of η .

Remark 5.6. Consider the composition of Figure 2 corresponding to $R_1R_2R_1(x, y, z) = (x', y', z')$ and $R_2R_1R_2(x, y, z) = (x'', y'', z'')$.



According to the bilinearization, assign the common time variables and tau functions to the faces $1, 2, \ldots, 7$ of the two diagrams to represent the incoming state $(x, y, z) \in \mathcal{B} \times \mathcal{B} \times \mathcal{B}$. Then the time variables on the faces a, b, \cdots, e are also the same in the two diagrams. Therefore one has $(x', y', z') = (x^*, y^*, z^*)$ for the associated outgoing state in $\mathcal{B} \times \mathcal{B} \times \mathcal{B}$, which implies the Yang-Baxter equation $R_1R_2R_1 = R_2R_1R_2$.

6. Reduction to $A_{2n-1}^{(2)}$ and $C_n^{(1)}$

Here we introduce the tropical R for $A_{2n-1}^{(2)}$ and $C_n^{(1)}$ based on Remark 2.1 and explain how the results on the $D_n^{(1)}$ case can be specialized to them. We note that these tropical R have more intrinsic characterization in terms of geometric crystals as argued in [KOTY] for $D_n^{(1)}$.

6.1. Tropical R for $A_{2n-1}^{(2)}$ and $C_n^{(1)}$. In this section we denote the tropical R for $D_n^{(1)}$ by $R_n : \mathcal{B}_n \times \mathcal{B}_n \to \mathcal{B}_n \times \mathcal{B}_n$. Let $\mathcal{B}' = \{x = (x_1, \dots, x_n, \overline{x}_n, \dots, \overline{x}_1)\}$ be the set of variables and $\rho : \mathcal{B}' \to \mathcal{B}_{n+1}$ be the embedding defined by

 $\rho((x_1,\ldots,x_n,\overline{x}_n,\ldots,\overline{x}_1)) = (x_1,\ldots,x_n,1,\overline{x}_n,\ldots,\overline{x}_1).$

The tropical R for $A_{2n-1}^{(2)}$ is the birational map $R'_n: \mathcal{B}' \times \mathcal{B}' \to \mathcal{B}' \times \mathcal{B}'$ defined by

$$R'_n(x,y) = (x',y') \Leftrightarrow R_{n+1}(\rho(x),\rho(y)) = (\rho(x'),\rho(y')).$$

Let $\mathcal{B}'' = \{x = (x_0, x_1, \dots, x_n, \overline{x}_n, \dots, \overline{x}_1)\}$ be the set of variables and $\rho' : \mathcal{B}'' \to \mathcal{B}_{n+2}$ be the embedding defined by

$$\rho'\big((x_0, x_1, \dots, x_n, \overline{x}_n, \dots, \overline{x}_1)\big) = (x_0, x_1, \dots, x_n, 1, \overline{x}_n, \dots, \overline{x}_1, x_0).$$

The tropical R for $C_n^{(1)}$ is the birational map $R_n'': \mathcal{B}'' \times \mathcal{B}'' \to \mathcal{B}'' \times \mathcal{B}''$ defined by

$$R''_{n}(x,y) = (x',y') \Leftrightarrow R_{n+2}(\rho'(x),\rho'(y)) = (\rho'(x'),\rho'(y')).$$

The maps R'_n and R''_n are well defined due to Remark 2.1. They satisfy the inversion relation and the Yang-Baxter equation. We shall write R'_n and R''_n simply as R' and R'' when the rank needs not be specified.

6.2. Bilinearization of $\mathbf{R'}$ and $\mathbf{R''}$. First consider the tropical R'_{n-1} for $A^{(2)}_{2n-3}$. The bilinear equation for it is given by setting $\lambda_n = \kappa_n = 1$ and specializing the tau functions as

(6.1)
$$\tau_{n-1}^J = \tau_n^J \qquad 0 \le J \le 4$$

in the $D_n^{(1)}$ bilinear equations. The constraint (6.1) implies $x_n = y_n = x'_n = y'_n = 1$ in (4.2)-(4.3), which is consistent with Remark 2.1 and Proposition 3.5. It makes the bilinear equations $\langle J, n-1 \rangle$ and $\langle J, n \rangle$ equivalent. We call the resulting subsystem $\{\langle J, i \rangle \mid 1 \leq J \leq 4, 0 \leq i \leq n-1\}$ of the $D_n^{(1)}$ case the $A_{2n-3}^{(2)}$ bilinear equations.

As for the tropical R''_{n-2} for $C^{(1)}_{n-2}$, the bilinear equation is obtained by further setting $\lambda_1 = \overline{\lambda}_1, \kappa_1 = \overline{\kappa}_1$ and imposing the constraint

(6.2)
$$\tau_0^J = \tau_1^J \qquad 0 \le J \le 4$$

on the $A_{2n-3}^{(2)}$ bilinear equations. The constraint implies $x_1 = \overline{x}_1$, $y_1 = \overline{y}_1$, $x'_1 = \overline{x}'_1$, $y'_1 = \overline{y}'_1$ in (4.2)-(4.3), which is again consistent with Remark 2.1 and Proposition 3.5. It makes the equations $\langle J, 0 \rangle$ and $\langle J, 1 \rangle$ equivalent. We call the resulting subsystem $\{\langle J, i \rangle \mid 1 \leq J \leq 4, 1 \leq i \leq n-1\}$ of the $A_{2n-3}^{(2)}$ case the $C_{n-2}^{(1)}$ bilinear equations. Under these specializations Theorem 4.4 is still valid for R'_{n-1} and R''_{n-2} .

6.3. Solutions of bilinear equations for R' and R''. Results in Section 5.1 can be specialized to make (6.1) and (6.2) hold. First we deal with (6.1) for R'. Let us restrict the elements (5.8) and (5.9) to the case M = 1 and take the limit

$$(6.3) q_1' \to 0, \quad p_1' \to a_m$$

for some $2 \le m \le n-1$ keeping $(1-p'_1/a_m)/{q'_1}^2$ to be an arbitrary nonzero constant. It makes the latter of (5.10) void, leaving a_1, \ldots, a_{n-1} as free parameters. All the tau functions in (5.12)-(5.14) are well defined in the limit as argued in the end of Appendix B.6. Moreover in (5.11), the coefficients $p'_1A(p'_1)/q'_1$ and $q'_1/(p'_1A(-p'_1))$ in the second sum tend to zero. Thus the element $h' \in e^{D'_{\infty}}$ in Lemma 5.3 actually belongs to $e^{B'_{\infty}}$. Then the property (B.15) ensures that $\tau^J_{n-1} = \tau^J_n$ for $1 \le J \le 4$ in (5.13) in agreement with (6.1). The remaining relation $\tau^0_{n-1} = \tau^0_n$ in (5.14) follows from (B.16).

Let us proceed to R'', where the further condition (6.2) should be satisfied. The reduction from $D_n^{(1)}$ case to $C_{n-2}^{(1)}$ is done by setting $\forall c_j = 0$ in (5.8)-(5.9). Namely we restrict $g' \in e^{D'_{\infty}}$ and $g \in e^{D_{\infty}}$ to $g' \in e^{B'_{\infty}}$ and $g \in e^{B_{\infty}}$, which reduces the relevant fermion theory essentially to the single component one. The former condition in (5.10) is still present although the latter becomes void. Then obviously $h' \in e^{B'_{\infty}}$ holds in Lemma 5.3, therefore (6.1) and (6.2) are valid due to (B.15), (B.9) and (B.8).

Appendix A. U_i, V_i in terms of tau functions

Let us prove Lemma 4.3. Our task is to substitute (4.2) into the functions U_i, V_i and simplify the result by means of the bilinear equations on the tau functions. Lemma A.1.

$$\begin{split} \frac{1}{x_i} + \frac{1}{\overline{y}_i} &= \frac{\beta \tau_1^0}{\tau_0^1 \tau_0^3} \; (i=1), \; \frac{\beta \tau_2^0 \tau_0^2 \tau_1^2}{N_1 W_1} \; (i=2), \; \frac{\beta \tau_i^0 \tau_{i-1}^2}{N_{i-1} W_{i-1}} \; (3 \le i \le n-2), \\ \frac{y_n}{x_{n-1}} + \frac{1}{\overline{y}_{n-1}} &= \frac{\beta \tau_{n-1}^0 \tau_n^1 \tau_{n-2}^2 \tau_{n-1}^2}{\kappa_n N_{n-2} W_{n-2} \tau_{n-1}^1 \tau_n^2}. \end{split}$$
Proof. Apply $\langle 2, 1 \rangle, \ldots, \langle 2, n-1 \rangle.$

It is straightforward to check

Lemma A.2. For
$$1 \le i \le n-1$$
, the ratio $(y_1 \cdots y_i)/(x_1 \cdots x_{i-1})$ is equal to
 $\frac{\tau_1^1 \tau_0^2}{\kappa_1 N_1}$ $(i = 1)$, $\frac{\lambda_1 W_1 \tau_0^2 \tau_2^1}{\kappa_1 \kappa_2 N_2 \tau_0^1 \tau_0^3 \tau_1^2}$ $(i = 2)$, $\frac{\lambda_1 \cdots \lambda_{i-1} W_{i-1} (\tau_0^2)^2 \tau_i^1}{\kappa_1 \cdots \kappa_i N_i \tau_0^1 \tau_0^3 \tau_{i-1}^2}$ $(3 \le i \le n-2)$,
 $\frac{\lambda_1 \cdots \lambda_{n-2} W_{n-2} (\tau_0^2)^2 \tau_{n-1}^1}{\kappa_1 \cdots \kappa_{n-1} \tau_0^1 \tau_0^3 \tau_{n-2}^2 \tau_{n-1}^2}$ $(i = n-1)$.

From Proposition 4.2, taking σ_* in Lemma A.2 implies

Lemma A.3. For $1 \le i \le n-2$, the ratio $(\overline{x}_1 \cdots \overline{x}_i)/(\overline{y}_1 \cdots \overline{y}_{i-1})$ is equal to

$$\frac{\tau_0^2 \tau_1^3}{\overline{\lambda}_1 W_1} \quad (i=1), \quad \frac{\overline{\kappa}_1 N_1 \tau_0^2 \tau_2^3}{\overline{\lambda}_1 \overline{\lambda}_2 W_2 \tau_0^1 \tau_0^3 \tau_1^2} \quad (i=2), \quad \frac{\overline{\kappa}_1 \cdots \overline{\kappa}_{i-1} N_{i-1} (\tau_0^2)^2 \tau_i^3}{\overline{\lambda}_1 \cdots \overline{\lambda}_i W_i \tau_0^1 \tau_0^3 \tau_{i-1}^2} \quad (3 \le i \le n-2).$$

In addition one has

$$\frac{\overline{x}_1\cdots\overline{x}_{n-1}}{\overline{y}_1\cdots\overline{y}_{n-2}y_n} = \frac{\overline{\kappa}_1\cdots\overline{\kappa}_{n-2}\kappa_n N_{n-2}(\tau_0^2)^2 \tau_{n-1}^1 \tau_n^3}{\overline{\lambda}_1\cdots\overline{\lambda}_{n-1}\tau_0^1 \tau_0^3 \tau_n^1 \tau_{n-2}^2 \tau_{n-1}^2}$$

Proof of Lemma 4.3. First we consider V_0 in (2.6). In view of (2.1) and (4.4), it is expressed as

(A.1)
$$V_{0} = l^{-1}v_{+} + k^{-1}v_{-},$$
$$v_{+} = \sum_{i=1}^{n-2} \frac{y_{1} \cdots y_{i}}{x_{1} \cdots x_{i-1}} \left(\frac{1}{x_{i}} + \frac{1}{\overline{y}_{i}}\right) + \frac{y_{1} \cdots y_{n-1}}{x_{1} \cdots x_{n-2}} \left(\frac{y_{n}}{x_{n-1}} + \frac{1}{\overline{y}_{n-1}}\right),$$
$$v_{-} = \sum_{i=1}^{n-2} \frac{\overline{x}_{1} \cdots \overline{x}_{i}}{\overline{y}_{1} \cdots \overline{y}_{i-1}} \left(\frac{1}{x_{i}} + \frac{1}{\overline{y}_{i}}\right) + \frac{\overline{x}_{1} \cdots \overline{x}_{n-1}}{\overline{y}_{1} \cdots \overline{y}_{n-2}y_{n}} \left(\frac{y_{n}}{x_{n-1}} + \frac{1}{\overline{y}_{n-1}}\right).$$

Using Lemmas A.1 and A.2 we get

$$\frac{\tau_0^1 \tau_0^3}{\beta \tau_0^2} v_+ = \frac{\tau_1^0 \tau_1^1}{\kappa_1 N_1} + \frac{\lambda_1 \tau_0^2 \tau_2^0 \tau_2^1}{\kappa_1 \kappa_2 N_1 N_2} + \tau_0^2 \sum_{i=3}^{n-2} \frac{\lambda_1 \cdots \lambda_{i-1} \tau_i^0 \tau_i^1}{\kappa_1 \cdots \kappa_i N_{i-1} N_i} + \frac{\lambda_1 \cdots \lambda_{n-2} \tau_0^2 \tau_{n-1}^0 \tau_n^1}{\kappa_1 \cdots \kappa_n N_{n-2} \tau_n^2}.$$

Rewrite $\tau_i^0 \tau_i^1 (1 \le i \le n-2)$ and $\tau_{n-1}^0 \tau_n^1$ appearing here by means of the bilinear equations $\langle 1, 1 \rangle - \langle 1, n-1 \rangle$. Then all but two terms cancel out, leading to

$$v_{+} = \frac{\beta \tau_0^2}{\alpha \tau_0^1 \tau_0^3} \left(\tau_0^4 - \frac{\lambda_1 \cdots \lambda_n \tau_0^2 \tau_n^4}{\kappa_1 \cdots \kappa_n \tau_n^2} \right).$$

Similarly one can derive

$$v_{-} = \frac{\beta \tau_0^2}{\alpha \tau_0^1 \tau_0^3} \left(-\tau_0^4 + \frac{\overline{\kappa}_1 \cdots \overline{\kappa}_{n-1} \tau_0^2 \tau_n^4}{\overline{\lambda}_1 \cdots \overline{\lambda}_{n-1} \tau_n^2} \right)$$

from Lemmas A.1, A.3 and the bilinear equations $\langle 3, 1 \rangle - \langle 3, n-1 \rangle$. Substituting these expressions of v_{\pm} into (A.1), we obtain $V_0 = \beta(k-l)\tau_0^2\tau_0^4/(lk\alpha\tau_0^1\tau_0^3)$ as desired. Starting from this result on V_0 , one can apply Lemma A.1 and the bilinear equations $\langle 3, 1 \rangle - \langle 3, n-1 \rangle$ to the recursion (2.7) successively to derive the formulas for V_1, \ldots, V_{n-1} . Then the formulas for $V_0^{\sigma_1}$ and $V_1^{\sigma_*}, \ldots, V_{n-1}^{\sigma_*}$ follow from (3.4)-(3.7) and Proposition 4.2. Finally substituting these results into (2.8) and using $\langle 4, 2 \rangle - \langle 4, n-2 \rangle$, one arrives at the desired expression for U_1, \ldots, U_{n-1} .

Appendix B. Tau functions and free fermions

Here we briefly recall the free fermion approach to the theory of tau functions based on [JM]. Lemma B.1 and Lemma B.2 play an important role in the main text.

B.1. **Single component fermions.** Single component charged fermions and the Fock states are given by

$$\begin{split} &[\psi_j, \psi_k^*]_+ = \delta_{jk}, \quad [\psi_j, \psi_k]_+ = [\psi_j^*, \psi_k^*]_+ = 0 \quad (j, k \in \mathbb{Z}), \\ &\psi(p) = \sum_j \psi_j p^j, \quad \psi(p)^* = \sum_j \psi_j^* p^{-j}, \\ &\langle l| = \langle \operatorname{vac} | \Psi_l^*, \quad | l \rangle = \Psi_l | \operatorname{vac} \rangle, \\ &\Psi_l = \begin{cases} \psi_{l-1} \cdots \psi_1 \psi_0, \quad l > 0 \\ 1 \quad l = 0, \quad \Psi_l^* = \begin{cases} \psi_0^* \psi_1^* \cdots \psi_{l-1}^*, \quad l > 0 \\ 1 \quad l = 0, \\ \psi_l^* \cdots \psi_{-2}^* \psi_{-1}^* \quad l < 0 \end{cases} \\ & \psi_j | \operatorname{vac} \rangle = 0, \ \langle \operatorname{vac} | \psi_j^* = 0 \ (j < 0), \quad \psi_j^* | \operatorname{vac} \rangle = 0, \ \langle \operatorname{vac} | \psi_j = 0 \ (j \ge 0) \end{cases} \end{split}$$

The neutral fermions are simple combinations thereof:

$$\phi_{j} = \frac{\psi_{j} + (-1)^{j} \psi_{-j}^{*}}{\sqrt{2}}, \quad \hat{\phi}_{j} = i \frac{\psi_{j} - (-1)^{j} \psi_{-j}^{*}}{\sqrt{2}},$$
$$[\phi_{j}, \phi_{k}]_{+} = [\hat{\phi}_{j}, \hat{\phi}_{k}]_{+} = (-1)^{k} \delta_{j,-k}, \quad [\phi_{j}, \hat{\phi}_{k}]_{+} = 0,$$
$$\phi(p) = \sum_{j} \phi_{j} p^{j}, \quad \hat{\phi}(p) = \sum_{j} \hat{\phi}_{j} p^{j},$$

(B.1) $\phi_j |l\rangle = \hat{\phi}_j |l\rangle = 0$ $(l = 0, 1, j < 0), \quad \langle l | \phi_j = \langle l | \hat{\phi}_j = 0 \quad (l = 0, 1, j > 0).$

We prepare time variables and their functions:

(B.2)
$$\begin{aligned} x &= (x_1, x_2, x_3, \ldots), \quad \tilde{x} = (x_1, -x_2, x_3, -x_4, \ldots), \\ \xi(x, p) &= \sum_{j \ge 1} x_j p^j, \\ \varepsilon(a) &= (a, \frac{a^2}{2}, \frac{a^3}{3}, \ldots) = -\tilde{\varepsilon}(-a). \end{aligned}$$

We say x is odd if $x = \tilde{x}$. With the Hamiltonians

$$H(x) = \sum_{m \ge 1} \sum_{j \in \mathbb{Z}} x_m \psi_j \psi_{j+m}^*,$$

$$H'(x,y) = \frac{1}{2} \sum_{l=1,3,5,\dots} \sum_{m \in \mathbb{Z}} (-1)^{m+1} (x_l \phi_m \phi_{-m-l} + y_l \hat{\phi}_m \hat{\phi}_{-m-l}),$$

the fermions exhibit the time evolutions

$$\begin{aligned} &\operatorname{Ad} e^{H(x)}\psi(p) = e^{\xi(x,p)}\psi(p), \ \operatorname{Ad} e^{H(x)}\psi^*(p) = e^{-\xi(x,p)}\psi^*(p), \\ &\operatorname{Ad} e^{H'(x,y)}\phi(p) = e^{\xi(x,p)}\phi(p), \ \operatorname{Ad} e^{H(x,y)}\hat{\phi}(p) = e^{\xi(y,p)}\hat{\phi}(p) \ \text{ for } (x,y) = (\tilde{x},\tilde{y}), \end{aligned}$$

where $\operatorname{Ad} g(\cdot) = g(\cdot)g^{-1}$.

B.2. 2 component fermions.

$$\begin{split} &[\psi_{j}^{(\alpha)},\psi_{k}^{(\beta)*}]_{+} = \delta_{\alpha\beta}\delta_{jk}, [\psi_{j}^{(\alpha)},\psi_{k}^{(\beta)}]_{+} = [\psi_{j}^{(\alpha)*},\psi_{k}^{(\beta)*}]_{+} = 0 \ (\alpha,\beta=1,2,\ j,k\in\mathbb{Z}), \\ &\psi^{(\alpha)}(p) = \sum_{j}\psi_{j}^{(\alpha)}p^{j},\ \psi^{(\alpha)*}(p) = \sum_{j}\psi_{j}^{(\alpha)*}p^{-j}, \\ &\phi_{j}^{(\alpha)} = \frac{\psi_{j}^{(\alpha)} + (-1)^{j}\psi_{-j}^{(\alpha)*}}{\sqrt{2}}, \quad \hat{\phi}_{j}^{(\alpha)} = i\frac{\psi_{j}^{(\alpha)} - (-1)^{j}\psi_{-j}^{(\alpha)*}}{\sqrt{2}}, \\ &\phi^{(\alpha)}(p) = \sum_{j}\phi_{j}^{(\alpha)}p^{j},\ \hat{\phi}^{(\alpha)}(p) = \sum_{j}\hat{\phi}_{j}^{(\alpha)}p^{j}, \\ &H(x,y) = \sum_{l\geq 1}\sum_{j}(x_{l}\psi_{j}^{(1)}\psi_{j+l}^{(1)*} + y_{l}\psi_{j}^{(2)}\psi_{j+l}^{(2)*}), \\ &\psi_{j}^{(\alpha)}|\mathrm{vac}\rangle = 0,\ \langle\mathrm{vac}|\psi_{j}^{(\alpha)*} = 0 \ (j<0), \quad \psi_{j}^{(\alpha)*}|\mathrm{vac}\rangle = 0,\ \langle\mathrm{vac}|\psi_{j}^{(\alpha)} = 0 \ (j\geq 0), \\ &\langle l_{1},l_{2}| = \langle\mathrm{vac}|\Psi_{l_{1}}^{(1)*}\Psi_{l_{2}}^{(2)*} \ |l_{2},l_{1}\rangle = \Psi_{l_{2}}^{(2)}\Psi_{l_{1}}^{(1)}|\mathrm{vac}\rangle, \end{split}$$

where $\Psi_l^{(\alpha)}$ and $\Psi_l^{(\alpha)*}$ stand for Ψ_l and Ψ_l^* with ψ_j, ψ_j^* replaced by $\psi_j^{(\alpha)}, \psi_j^{(\alpha)*}$. 20

B.3. Algebras $A_{\infty}, B'_{\infty}, B_{\infty}, D'_{\infty}$ and D_{∞} . The algebras $B'_{\infty} \subset D'_{\infty}$ are defined within the single component theory as $B'_{\infty} = \{\sum a_{jk} : \phi_j \phi_k : +d \mid \exists N, a_{jk} = 0 \text{ if } |j+k| > N\}$ and $D'_{\infty} = \{\sum a_{jk} : \phi_j \phi_k : +\sum b_{jk} : \hat{\phi}_j \hat{\phi}_k : +\sum c_{jk} : \phi_j \hat{\phi}_k : +d \mid \exists N, a_{jk} = b_{jk} = c_{jk} = 0 \text{ if } |j+k| > N\}$. The algebra A_{∞} in the 2 component theory is defined as $A_{\infty} = \{\sum a_{\alpha i,\beta j} : \psi_i^{(\alpha)} \psi_j^{(\beta)*} : +d \mid \exists N, a_{\alpha i,\beta j} = 0 \text{ if } |i-j| > N\}$. The automorphism σ, κ, ω of A_{∞} and the homomorphism $\iota : D'_{\infty} \to A_{\infty}$ are given by

$$(B.3) \qquad \begin{aligned} \sigma : \psi_j^{(\alpha)} \mapsto (-1)^j \psi_{-j}^{(\alpha)*}, \ \psi_j^{(\alpha)*} \mapsto (-1)^j \psi_{-j}^{(\alpha)} \\ \kappa : \psi_j^{(\alpha)} \mapsto i \psi_j^{(\alpha)}, \ \psi_j^{(\alpha)*} \mapsto -i \psi_j^{(\alpha)*}, \\ \omega : \psi_j^{(\alpha)} \mapsto \psi_{j-1}^{(\alpha)}, \ \psi_j^{(\alpha)*} \mapsto \psi_{j-1}^{(\alpha)*}, \\ \iota : \phi_j \mapsto \phi_j^{(1)}, \ \hat{\phi}_j \mapsto \phi_j^{(2)}. \end{aligned}$$

The algebras $B_{\infty} \subset D_{\infty}$ are defined as $D_{\infty} = \{X \in A_{\infty} \mid \sigma(X) = X\}$ and $B_{\infty} = \{X \in D_{\infty} \mid \pi(X) = X\}$, where π is the projection to the first component, i.e., $\pi(\psi_j^{(\alpha)}) = \delta_{\alpha 1} \psi_j^{(\alpha)}, \pi(\psi_j^{(\alpha)*}) = \delta_{\alpha 1} \psi_j^{(\alpha)*}$. Thus B_{∞} actually stays within the single component theory. The map

$$\iota + \kappa \iota : D'_{\infty} \longrightarrow D_{\infty}, \quad B'_{\infty} \longrightarrow B_{\infty}$$

is an isomorphism and one has

(B.4)
$$\iota(H'(x,y)) + \kappa\iota(H'(x,y)) = H(x,y) \quad \text{if } (x,y) = (\tilde{x},\tilde{y}).$$

The set of well-defined group elements corresponding to these algebras will be denoted, by abuse of notation, by

$$e^{D_{\infty}} := \{ e^{X_1} \cdots e^{X_k} \mid k \ge 0, X_1, \dots, X_k \in D_{\infty} \text{ are locally nilpotent} \},$$

and similarly for $A_{\infty}, B_{\infty}, B'_{\infty}$ and D'_{∞} . For $g' \in e^{D'_{\infty}}$, we write

(B.5)
$$g = e^{\iota(X') + \kappa\iota(X')} = \iota(g')\kappa\iota(g') \in e^{D_{\infty}}.$$

B.4. Tau functions. We concern two kinds of tau functions:

(B.6)
$$\begin{aligned} f_l(x,y;h) &= \langle l|e^{H'(x,y)}h|l\rangle \quad \text{for } (x,y) = (\tilde{x},\tilde{y}), \ h \in e^{D'_{\infty}}, \\ F_{l_1,l_2;l}(x,y;g) &= \langle l_1,l_2|e^{H(x,y)}g|l_2 - l,l_1 + l\rangle \quad g \in e^{A_{\infty}}. \end{aligned}$$

 $F_{l_1,l_2:0}(x,y;g)$ will simply be denoted by $F_{l_1,l_2}(x,y;g)$. They enjoy the symmetry

(B.7)
$$F_{l_1,l_2;l}(x,y;g) = (-1)^{(l_1+l_2)l} F_{1-l_1,1-l_2;-l}(\tilde{x},\tilde{y};\sigma(g)),$$

(B.8)
$$F_{l_1,l_2;l}(x,y;g) = F_{l_1-1,l_2-1;l}(x,y;\omega(g)).$$

Clearly we have

(B.9)
$$F_{l_1,l_2;l}(x,y;g) = 0$$
 if $l \neq 0$ and $g \in B_{\infty}$.

 $F_{l_1,l_2;l}(x,y;g)$ will be denoted by $F_{l_1,l_2;l}(x,y)$ or $F_{l_1,l_2;l}(x)$ for simplicity.

B.5. Bilinear identities. For any $g \in e^{A_{\infty}}$ one has the hierarchy of equations

$$\oint \frac{dk}{2\pi i k} (-)^{l_2+l_2'} k^{l_1-l_1'-1} e^{\xi(x-x',k)} F_{l_1-1,l_2;l+1}(x-\varepsilon(k^{-1}),y) F_{l_1'+1,l_2';l'-1}(x'+\varepsilon(k^{-1}),y') + \oint \frac{dk}{2\pi i k} k^{l_2-l_2'-1} e^{\xi(y-y',k)} F_{l_1,l_2-1;l}(x,y-\varepsilon(k^{-1})) F_{l_1',l_2'+1;l'}(x',y'+\varepsilon(k^{-1})) = 0,$$

where $l_1 - l'_1 \ge l' - l \ge l'_2 - l_2 + 2$ and the integration is taken along a small contour around $k = \infty$. This is eq.(4.4) in [JM]. Let b_1, b_2, b_3 be nonzero constants. We take y = y' and suppress the dependence on it. The above equations contain the following:

(B.10)
$$(b_2^{-1} - b_3^{-1})F_{l_1, l_2; l}(x + \varepsilon(b_2^{-1}) + \varepsilon(b_3^{-1}))F_{l_1 - 1, l_2 + 1; l + 1}(x + \varepsilon(b_1^{-1})) + \text{cyc} = 0,$$

(B.11)
$$(b_2^{-1} - b_3^{-1})F_{l_1, l_2; l}(x + \varepsilon(b_2^{-1}) + \varepsilon(b_3^{-1}))F_{l_1 - 1, l_2; l + 1}(x + \varepsilon(b_1^{-1})) + cyc = 0,$$

(B.12)
$$(b_2^{-1} - b_3^{-1})F_{l_1, l_2; l}(x + \varepsilon(b_2^{-1}) + \varepsilon(b_3^{-1}))F_{l_1 - 1, l_2; l}(x + \varepsilon(b_1^{-1})) + \text{cyc} = 0,$$

(B.13)
$$(b_2 - b_3)F_{l_1, l_2; l}(x + \varepsilon(b_2^{-1}) + \varepsilon(b_3^{-1}))F_{l_1, l_2; l}(x + \varepsilon(b_1^{-1})) + \text{cyc} = 0,$$

where +cyc means the additional two terms obtained by the cyclic permutations of b_1, b_2, b_3 . They also contain

(B.14)
$$F_{0,1}(x+\varepsilon(b_2^{-1}))F_{0,0}(x+\varepsilon(b_1^{-1})) - F_{0,1}(x+\varepsilon(b_1^{-1}))F_{0,0}(x+\varepsilon(b_2^{-1})) \\ = (b_1^{-1} - b_2^{-1})F_{1,0;-1}(x+\varepsilon(b_1^{-1}) + \varepsilon(b_2^{-1}))F_{-1,1;1}(x).$$

When both x and y are odd, $F_{l_1,l_2;l}(x,y;g)$ with $g \in e^{D_{\infty}}$ can be expressed in terms of $f_l(x,y;g')$ as noted in eq.(7.6) in [JM]. We recall this fact in

Lemma B.1. Let $g \in e^{D_{\infty}}$ and $g' \in e^{D'_{\infty}}$ be related as in (B.5). Suppose that x and y are both odd. Denoting $F_{l_1,l_2;l}(x,y;g)$ and $f_l(x,y;g')$ by $F_{l_1,l_2;l}(x)$ and $f_l(x)$, one has

$$F_{1,1}(x) = F_{0,0}(x) = f_0(x)f_1(x),$$

$$F_{0,1}(x) = F_{1,0}(x) = \frac{1}{2}(f_0(x)^2 + f_1(x)^2),$$

$$F_{0,1;1}(x) = -F_{1,0;-1}(x) = \frac{i}{2}(f_0(x)^2 - f_1(x)^2)$$

Proof. The first equality in each line is due to (B.7). Since $\phi_0^2 = \hat{\phi}_0^2 = 1/2$, we may write $\operatorname{Ad} e^{H'(x,y)} g' = X_0 + X_1 \phi_0 + X_2 \hat{\phi}_0 + X_3 \phi_0 \hat{\phi}_0$, where $X_j = X_j(\phi, \hat{\phi})$ contains $\phi_k, \hat{\phi}_k$ only for $k \neq 0$ and its order is even (resp. odd) for j = 0, 3 (resp. j = 1, 2). Due to (B.1) we find $f_l(x; g') = \langle X_0 \rangle + (-1)^l i \langle X_3 \rangle / 2$ for l = 0, 1, where $\langle X_j \rangle := \langle 0 | X_j | 0 \rangle = \langle 1 | X_j | 1 \rangle$. On the other hand from (B.4) and (B.5) it follows that $\operatorname{Ad} e^{H(x,y)} g = (X'_0 + X'_1 \phi_0^{(1)} + X'_2 \phi_0^{(2)} + X'_3 \phi_0^{(1)} \phi_0^{(2)}) (\hat{X}'_0 + \hat{X}'_1 \hat{\phi}_0^{(1)} + \hat{X}'_2 \hat{\phi}_0^{(2)} + \hat{X}'_3 \hat{\phi}_0^{(1)} \hat{\phi}_0^{(2)})$, where X'_j and \hat{X}'_j are obtained from X_j by substituting the 2 component fermions into X_j as $X'_j = X_j(\phi^{(1)}, \phi^{(2)}), \ \hat{X}'_j = X_j(\hat{\phi}^{(1)}, \hat{\phi}^{(2)}).$ Due to the property (B.1) for each component, we get $F_{0,0}(x; g) = \langle 0, 0 | (X'_0 \hat{X}'_0 + X'_3 \hat{X}'_3 \phi_0^{(1)} \phi_0^{(2)} \hat{\phi}_0^{(1)} \hat{\phi}_0^{(2)}) | 0, 0 \rangle = \langle X_0 \rangle^2 + \langle X_3 \rangle^2 / 4 = (\langle X_0 \rangle + i \langle X_3 \rangle / 2) (\langle X_0 \rangle - i \langle X_3 \rangle / 2)$, proving the first relation. The other relations can be shown similarly. □

For odd x and y, it is clear from the above proof that

(B.15)
$$f_0(x,y;g') = f_1(x,y;g')$$
 if $g' \in e^{B'_{\infty}}$.

Lemma B.1 admits a generalization to

Lemma B.2. Let $g \in e^{D_{\infty}}$ and $g' \in e^{D'_{\infty}}$ be as in (B.5). Suppose y is odd and $\tilde{x} = x + \varepsilon(c^{-1}) - \tilde{\varepsilon}(c^{-1})$. Denoting $F_{l_1, l_2; l}(x, y; g)$ and $f_l(x, y; g')$ by $F_{l_1, l_2; l}(x)$ and $f_l(x)$, one has

$$\begin{cases} 2F_{0,1}(x) \\ -2iF_{0,1;1}(x) \end{cases} = f_0(x + \varepsilon(c^{-1}))f_0(x - \tilde{\varepsilon}(c^{-1})) \pm f_1(x + \varepsilon(c^{-1}))f_1(x - \tilde{\varepsilon}(c^{-1})), \\ 2F_{0,0}(x) \\ 2ic^{-1}F_{-1,1;1}(x) \end{cases} = f_0(x + \varepsilon(c^{-1}))f_1(x - \tilde{\varepsilon}(c^{-1})) \pm f_1(x + \varepsilon(c^{-1}))f_0(x - \tilde{\varepsilon}(c^{-1})).$$

Proof. In (B.10)–(B.12), set $b_1 = -b_2 = c$ and $b_3^{-1} = 0$. Writing $x_+ = x + \varepsilon(c^{-1}), x_- = x - \tilde{\varepsilon}(c^{-1})$, one has

$$\begin{split} F_{1,1}(x_{-})F_{0,1}(x_{+}) + F_{1,1}(x_{+})F_{0,1}(x_{-}) &= 2F_{1,1}(\tilde{x})F_{0,1}(x), \\ F_{1,1}(x_{-})F_{0,1;1}(x_{+}) + F_{1,1}(x_{+})F_{0,1;1}(x_{-}) &= 2F_{1,1}(\tilde{x})F_{0,1;1}(x), \\ F_{1,0;-1}(x_{-})F_{0,1}(x_{+}) + F_{1,0;-1}(x_{+})F_{0,1}(x_{-}) &= 2F_{1,0;-1}(\tilde{x})F_{0,1}(x). \end{split}$$

By applying (B.7) together with $\sigma(g) = g$, the right hand sides here are equal to $2F_{0,0}(x)F_{0,1}(x), 2F_{0,0}(x)F_{0,1;1}(x)$ and $-2F_{0,1;1}(x)F_{0,1}(x)$, respectively. Since both x_+ and x_- are odd, the left hand sides are expressed in terms of $f_0(x_{\pm}), f_1(x_{\pm})$ by Lemma B.1. Solving the resulting three equations with respect to $F_{0,0}(x), F_{0,1}(x)$ and $F_{0,1;1}(x)$, we obtain the formulas in question up to an overall sign, which can be fixed by considering the smooth limit $g \to 1$. The formula for $F_{-1,1;1}(x)$ can be derived similarly by using (B.14).

Note that in the limit $c \to \infty,$ the first three relations in Lemma B.2 reduce to Lemma B.1.

B.6. **Explicit formula.** Let us write down the explicit formula for the vacuum expectation values (B.6). To simplify the result we introduce the parameters p_1, \ldots, p_{2N+M} and q_1, \ldots, q_M to modify the parameterization of $g' \in e^{D'_{\infty}}$ and $g \in e^{D_{\infty}}$ in (5.8) as

$$g' = \exp\left(\sum_{i=1}^{N} b_i \phi(p_i) \phi(p_{\bar{i}}) + \sum_{j=1}^{M} c_j \phi(p_{\bar{j}}) \hat{\phi}(q_j)\right),$$

$$g = \exp\left(\sum_{i=1}^{N} b_i \left(\psi^{(1)}(p_i) \psi^{(1)*}(-p_{\bar{i}}) - \psi^{(1)}(p_{\bar{i}}) \psi^{(1)*}(-p_i)\right)\right)$$

$$+ \sum_{j=1}^{M} c_j \left(\psi^{(1)}(p_{\bar{j}}) \psi^{(2)*}(-q_j) - \psi^{(2)}(q_j) \psi^{(1)*}(-p_{\bar{j}})\right)\right),$$

where $\overline{i} = 2N + 1 - i$ and $\tilde{j} = 2N + M + 1 - j$. Given $I \subseteq \{1, \ldots, N\}$ and $J \subseteq \{1, \ldots, M\}$, we shall write $\overline{I} = \{\overline{i} \mid i \in I\}$ and $\tilde{J} = \{\overline{j} \mid j \in J\}$. The cardinality of a set I is denoted by |I|. For any sets K and K' we set

$$\begin{split} \Delta_{K}^{\pm}(p) &= \prod_{\mu,\nu\in K, \; \mu<\nu} (p_{\mu} \pm p_{\nu}), \\ \Delta_{K,K'}^{m,l}(p) &= \frac{\Delta_{K}^{-}(p)\Delta_{K'}^{-}(-p)}{\prod_{\mu\in K,\nu\in K'} (p_{\mu} + p_{\nu})} \prod_{\mu\in K} p_{\mu}^{m+l} \prod_{\mu\in K'} (-p_{\mu})^{-m-l+1}, \end{split}$$

and similarly for q. Let x and y be odd time variables. Then we have

$$f_{l}(x, y; g') = \sum_{I,J} \varepsilon_{l,J} 2^{-|I| - |J|} b_{I}(x) c_{J}(x, y) \frac{\Delta_{K}^{-}(p) \Delta_{J}^{-}(q)}{\Delta_{K}^{+}(p) \Delta_{J}^{+}(q)} \qquad l = 0, 1,$$

$$K = I \sqcup \bar{I} \sqcup \tilde{J}, \qquad \varepsilon_{l,J} = \begin{cases} 1 & |J| \text{ even} \\ i(-1)^{l} & |J| \text{ odd,} \end{cases}$$

$$b_{I}(x) = \prod_{i \in I} b_{i} \exp(\xi(x, p_{i}) - \xi(x, -p_{\bar{i}})),$$

$$c_{J}(x, y) = \prod_{j \in J} c_{j} \exp(\xi(x, p_{\bar{j}}) + \xi(y, q_{j})),$$

where the sum $\sum_{I,J}$ extends over all the subsets $I \subseteq \{1, \ldots, N\}$ and $J \subseteq \{1, \ldots, M\}$. For any integers l_1, l_2 and l, $F_{l_1, l_2; l}(x, y; g)$ with odd y (x is not restricted to be odd) reads

$$\begin{split} F_{l_1,l_2;l}(x,y;g) &= \sum_{I,I',J,J'} (-1)^{|I'|+(|I|+|I'|)|J|} b_I(x) b'_{I'}(x) c_J(x,y) c'_{J'}(x,y) \Delta^{l_1,l}_{K,K'}(p) \Delta^{l_2,-l}_{J',J}(q) \\ K &= I \sqcup \bar{I'} \sqcup \tilde{J}, \qquad K' = I' \sqcup \bar{I} \sqcup \tilde{J'}, \\ b'_{I'}(x) &= \prod_{i \in I'} b_i \exp(\xi(x,p_{\bar{i}}) - \xi(x,-p_i)), \\ c'_{J'}(x,y) &= \prod_{j \in J'} c_j \exp(-\xi(x,-p_{\bar{j}}) + \xi(y,q_j)), \end{split}$$

where the sum $\sum_{I,I',J,J'}$ extends over all the subsets $I, I' \subseteq \{1, \ldots, N\}$ and $J, J' \subseteq \{1, \ldots, M\}$ such that |J'| - |J| = l.

Consider the dependence of $f_l(x, y; g')$ (l = 0, 1) and $F_{l_1, l_2; l}(x, y; g)$ on q_1 when M = 1. Apart from the factors $c_J(x, y)$ and $c'_{J'}(x, y)$, f_l is independent of q_1 and $F_{l_1, l_2; l}$ is proportional to the single power $q_1^{l_2-l}$. Therefore all the functions f_l and $F_{l_1, l_2; l}$ appearing in (5.12)-(5.14) are well defined in the limit $q_1 \to 0$ with M = 1. In particular from (B.8) one has

(B.16)
$$\lim_{q_1 \to 0} F_{-1,1;1}(x, y; \omega(g)) = \lim_{q_1 \to 0} F_{0,2;1}(x, y; g) = 0.$$

These facts are used in Section 6.3.

Acknowledgements

The authors thank R. Inoue for a careful reading of the manuscript. A.K. thanks M. Jimbo for explaining eq.(7.6) in [JM]. M.O. and Y.Y. were partially supported by Grant-in-Aid for Scientific Research JSPS No.14540026 and No.11440047, respectively.

References

- [ABS] V. E. Adler, A. I. Bobenko and Yu. B. Suris, Classification of integrable equations on quad-graphs. The consistency approach, math.QA/0202024.
- [B] R.J. Baxter, Exactly solved models in statistical mechanics, Academic Press, London (1982).
- [BFZ] A. Berenstein, S. Fomin and A. Zelevinsky, Parameterizations of canonical bases and totally positive matrices, Adv. in Math. 122 (1996) 49–149.

- [BK] A. Berenstein and D. Kazhdan, Geometric and unipotent crystals, GAFA 2000 (Tel Aviv, 1999), Geom. Funct. Anal. 2000, Special Volume, Part I, 188–236.
- [DJM] E. Date, M. Jimbo and T. Miwa, Method for generating discrete soliton equations, J. Phys. Soc. Japan. 51 (1982) 4116–4124, 4125–4131.
- [D] V. G. Drinfeld, On some unsolved problems in quantum group theory, in "Quantum groups" (Leningrad 1990), Lect. Note. in Math. 1510 pp.1–8, Springer, 1992.
- [ESS] P. Etingof, T. Schedler and A. Soloviev, Set-theoretical solutions to the quantum Yang-Baxter equation, Duke Math. J. 100 (1999) 169–209.
- [FOY] K. Fukuda, M. Okado, Y. Yamada, Energy functions in box ball systems, Int. J. Mod. Phys. A 15 (2000) 1379–1392.
- [HHIKTT] G. Hatayama, K. Hikami, R. Inoue, A. Kuniba, T. Takagi and T. Tokihiro, The A⁽¹⁾_M Automata related to crystals of symmetric tensors, J. Math. Phys. 42 (2001) 274-308.
- [HKOT] G. Hatayama, A. Kuniba, M. Okado and T. Takagi, Combinatorial R matrices for a family of crystals: B_n⁽¹⁾, D_n⁽¹⁾, A_{2n}⁽²⁾, and D_{n+1}⁽²⁾ cases, J. Alg. **247** (2002) 577–615.
 [HKOTY] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Scattering rules in
- [HKOTY] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Scattering rules in soliton cellular automata associated with crystal bases, Contemporary Math. 297 (2002) 151–182.
- [HKT1] G. Hatayama, A. Kuniba, and T. Takagi, Soliton cellular automata associated with crystal bases, Nucl. Phys. B577[PM] (2000) 619–645.
- [HKT2] G. Hatayama, A. Kuniba and T. Takagi Simple algorithm for factorized dynamics of g_n-automaton, J. Phys. A: Math. Gen. 34 (2001) 10697-10705.
- [H] R. Hirota, Discrete analogue of a generalized Toda equation, J. Phys. Soc. Japan 50 (1981) 3785–3791.
- [HT] R. Hirota and S. Tsujimoto, Conserved quantities of a class of nonlinear differencedifference equations, J. Phys. Soc. Japan 64 (1995) 3125–3127.
- [HTI] R. Hitota, S. Tsujimoto and T. Imai, Difference scheme of soliton equations, in Future directions of Nonlinear Dynamics in Physical and Biological Systems, ed. P.L. Christiansen et al, (Plenum, New York, 1993) p7.
- [JMY] L. Jiang-Hua, Y. Min, and Z. Yong-Chang, On the set-theoretical Yang-Baxter equation, Duke Math. J. 104 (2000) 1–18.
- [JM] M. Jimbo and T. Miwa, Solitons and infinite dimensional Lie algebras, Publ. RIMS. Kyoto Univ. 19 (1983) 943–1001.
- [KNY] K. Kajiwara, M. Noumi and Y. Yamada, Discrete dynamical systems with $W(A_{m-1}^{(1)} \times A_{n-1}^{(1)})$ symmetry, Lett. Math. Phys. **60** (2002) 211–219.
- [KKM] S-J. Kang, M. Kashiwara and K. C. Misra, Crystal bases of Verma modules for quantum affine Lie algebras, Compositio Math. 92 (1994) 299–325.
- [K] M. Kashiwara, Crystal bases of modified quantized enveloping algebra, Duke Math. J. 73 (1994) 383–413.
- [Ki] A. N. Kirillov, Introduction to tropical combinatorics, in "Physics and Combinatorics 2000" (Eds. A. N. Kirillov and N. Liskova) Proceedings of the Nagoya 2000 International Workshop, pp. 82–150, World Scientific, 2001.
- [KOTY] A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Geometric crystal and tropical R for $D_n^{(1)}$, preprint math.QA/0208239.
- [L] G. Lusztig, *Total positivity in reductive Groups*, in "Lie theory and geometry", Prog. in Math. 123, pp. 531–568, Birkhäuser, Boston, 1994.
- [M] T. Miwa, On Hirota's difference equations, Proc. Japan Acad. Ser. A Math. Sci. 58 (1982) 9–12.
- [MV] J. Moser and A. P. Veselov, Discrete versions of some classical integrable systems and factorization of matrix polynomials, Commun. Math. Phys. 139 (1991) 217–243.
- [NY] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, Selecta Mathematica, New Ser. 3 (1997) 547-599.
- [NoY] M. Noumi and Y. Yamada, Tropical Robinson-Schensted-Knuth correspondence and birational Weyl group actions, preprint (math-ph/0203030), to appear in Adv. Stud. in Pure Math.
- [O] A. Odesskii, Set-theoretical solutions to the Yang-Baxter relation from factorization of matrix polynomials and θ-functions, math.QA/0205051.

- [TNS] T. Tokihiro, A. Nagai and J. Satsuma, Proof of solitonical nature of box and ball systems by means of inverse ultra-discretization, Inverse Problems 15 (1999) 1639–1662.
- [TS] D. Takahashi and J. Satsuma, A soliton cellular automaton, J. Phys. Soc. Jpn. 59 (1990) 3514–3519.
- [TTMS] T. Tokihiro, D. Takahashi, J. Matsukidaira and J. Satsuma, From soliton equations to integrable cellular automata through a limiting procedure, Phys. Rev. Lett. 76 (1996) 3247– 3250.
- [V1] A. P. Veselov, Integrable mappings, Russian Math. Surveys 46 (1991) 1-51.
- [V2] A. P. Veselov, Yang-Baxter maps and integrable dynamics, math.QA/0205335.
- [WX] A. Weinstein and P. Xu, Classical solutions of the quantum Yang-Baxter equation, Commun. Math. Phys. 148 (1992) 309–343.
- [Y] Y. Yamada, A birational representation of Weyl group, combinatorial R-matrix and discrete Toda equation, in "Physics and Combinatorics 2000" (Eds. A. N. Kirillov and N. Liskova) Proceedings of the Nagoya 2000 International Workshop, pp. 305–319, World Scientific, 2001.

INSTITUTE OF PHYSICS, UNIVERSITY OF TOKYO, TOKYO 153-8902, JAPAN *E-mail address*: atsuo@gokutan.c.u-tokyo.ac.jp

DEPARTMENT OF INFORMATICS AND MATHEMATICAL SCIENCE, GRADUATE SCHOOL OF ENGI-NEERING SCIENCE, OSAKA UNIVERSITY, OSAKA 560-8531, JAPAN *E-mail address:* okado@sigmath.es.osaka-u.ac.jp

Department of Applied Physics, National Defense Academy, Kanagawa 239-8686, Japan

E-mail address: takagi@nda.ac.jp

Department of Mathematics, Faculty of Science, Kobe University, Hyogo 657-8501, Japan

E-mail address: yamaday@math.kobe-u.ac.jp