

# ASSOCIATIVE SUBMANIFOLDS OF A $G_2$ MANIFOLD

SELMAN AKBULUT AND SEMA SALUR

ABSTRACT. We study deformations of associative submanifolds of a  $G_2$  manifold. We show that deformation spaces can be perturbed to be smooth and finite dimensional, and they can be made compact by constraining them with an additional equation and reducing it to Seiberg-Witten theory. This allows us to associate invariants of certain associative submanifolds. More generally we apply this process to certain associated Grassmann bundles in order to assign invariants to  $G_2$  manifolds.

## 0. Introduction

McLean showed that, in a  $G_2$  manifold  $(M^7, \varphi)$  the space of associative submanifolds near a given one  $Y^3$ , can be identified with the harmonic spinors on  $Y$  twisted by a certain bundle  $E$  (the kernel of a twisted Dirac operator) [M]. But since we can not control the cokernel of the Dirac operator (it has index zero), the dimension of its kernel might vary. This is the obstruction to smoothness of the moduli space of associative submanifolds. It is generally expected that this problem should be remedied by deforming the ambient  $G_2$  structure (i.e. by deforming  $\varphi$ ), but unfortunately this process might move  $\varphi$  to a non-integrable  $G_2$  structure. If we are to view  $(M, \varphi)$  as an analogue of a symplectic manifold and  $\varphi$  a symplectic form, and view the associative submanifolds as analogues of holomorphic curves, deforming  $\varphi$  would be too destructive process. Ideally one would want to define auxiliary objects analogous to almost complex structures ‘compatible with  $\varphi$ ’, and then deform the associative submanifolds by the help of these objects in a larger class of manifolds, just like deforming the holomorphic curves in the larger class of manifolds (pseudo-holomorphic curves). We will take approach to obtain the smoothness and the finite dimensionality of the moduli space. To get compactness of the moduli space we will relate the deformation equation to the Seiberg-Witten equations. Metric deformations also comes handy at the end in defining an invariant for  $G_2$  manifolds since they help to increase the dimension of the relevant moduli space.

In [AS] in order to carry this program we introduced *complex associative submanifolds* of  $G_2$  manifolds, they are associative submanifolds whose normal bundles carry

---

*Date:* February 17, 2019.

1991 *Mathematics Subject Classification.* 53C38, 53C29, 57R57.

*Key words and phrases.* deformation of calibrated manifolds.

First named author is partially supported by NSF grant DMS 9971440.

a  $U(2)$  structure. This is no restriction, since every associative submanifold has this structure, but if we require that their deformations to be compatible with the background connection we must have an integrability condition, i.e. the condition that the connection on their normal bundles (induced by the  $G_2$  background metric) to reduce to a  $U(2)$  connection. We call such manifolds *integrable complex associative submanifolds*. Prototype of these manifolds are holomorphic curves crossed by circles inside of Calabi Yau  $\times S^1$ . We don't know how abundant these manifolds are. In this paper we circumvent this restriction by considering a more general class of submanifolds, namely the associative submanifolds whose normal bundles carry a  $Spin^c(4)$  structure. This structure allows us to break the normal bundle into pair of complex bundles, define necessary Dirac operator between them, and perform the deformations in these bundles. In this case we don't have a connection compatibility problem, because we perform the deformation in a larger space by using the whole background connection with the help of an auxiliary connection. By applying this process globally to the 15 dimensional associative Grassmannian bundles over a  $G_2$  manifold  $(M, \varphi)$  we define an invariant of  $(M, \varphi)$ . In a future paper we hope to study this in a more general context by relating it to a similar theory for Cayley submanifolds of a  $Spin(7)$ .

We would like to thank G. Tian and R. Bryant for many valuable comments, and also thank R. Kirby for constant encouragement.

## 1. Preliminaries

Here we give some basic definitions and known facts about the manifolds with special holonomies, the reader can find all these interesting facts about them in [B2], [B3], [HL]. The exceptional Lie group  $G_2$  can be defined as the subgroup of  $GL(7, \mathbb{R})$  fixing a particular 3-form  $\varphi_0 \in \Omega^3(\mathbb{R}^7)$ . Denote  $e^{ijk} = dx^i \wedge dx^j \wedge dx^k \in \Omega^3(\mathbb{R}^7)$ , then

$$G_2 = \{A \in GL(7, \mathbb{R}) \mid A^* \varphi_0 = \varphi_0\}$$

$$\varphi_0 = e^{123} + e^{145} + e^{167} + e^{246} - e^{257} - e^{347} - e^{356}$$

**Definition 1.** A smooth 7-manifold  $M^7$  has a  $G_2$  structure if its tangent frame bundle reduces to  $G_2$  bundle. Equivalently  $M^7$  has a  $G_2$  structure if it has a 3-form  $\varphi \in \Omega^3(M)$ , such that at each  $x \in M$  the pair  $(T_x(M), \varphi(x))$  is isomorphic to  $(\mathbb{R}^7, \varphi_0)$ .

In fact  $G_2$  is a 14-dimensional subgroup of  $SO(7)$ . Since  $GL(7, \mathbb{R})$  acts on  $\Omega^3(\mathbb{R}^7)$  with the stabilizer  $G_2$ , its orbit is open by dimension reasons, so the choice of  $\varphi_0$  in the above definition is generic. In fact the action of  $GL^+(7, \mathbb{R})$  has two orbits containing  $\pm\varphi_0$ . The set of smooth 7-manifolds with  $G_2$ -structures coincides with the set of 7-manifolds with spin structure, though this correspondance is not 1 – 1. This

is because  $Spin(7)$  acts on  $S^7$  with stabilizer  $G_2$  inducing the fibrations

$$G_2 \rightarrow Spin(7) \rightarrow S^7 \rightarrow BG_2 \rightarrow BSpin(7)$$

and hence there is no obstruction to lifting maps  $M^7 \rightarrow BSpin(7)$  to  $BG_2$ . The cotangent frame bundle  $\mathcal{P}^*(M) \rightarrow M$  of a manifold with  $G_2$  structure  $(M, \varphi)$  can be expressed as  $\mathcal{P}^*(M) = \cup_{x \in M} \mathcal{P}_x^*(M)$ , where each fiber is given by:

$$(1) \quad \mathcal{P}_x^*(M) = \{u \in Hom(T_x(M), \mathbb{R}^7) \mid u^*(\varphi_0) = \varphi(x)\}$$

It turns out that any  $G_2$  structure  $\varphi$  on  $M^7$  gives an orientation  $\mu \in \Omega^3(M)$  on  $M$ , and in turn  $\mu$  determines a metric  $g = \langle \cdot, \cdot \rangle$  on  $M$ , and a cross product structure  $\times$  on its tangent bundle of  $M$  as follows: Let  $i_v$  denote the interior product with a vector  $v$ , then

$$(2) \quad g(u, v) = [i_u(\varphi) \wedge i_v(\varphi) \wedge \varphi] / \mu$$

$$(3) \quad \varphi(u, v, w) = g(u \times v, w)$$

To emphasize dependency on  $\varphi$  sometimes  $g$  is denoted by  $g_\varphi$ . There is a notion of a  $G_2$  structure  $\varphi$  on  $M^7$  to be integrable, which corresponds to  $\varphi$  being harmonic.

**Definition 2.** A manifold with  $G_2$  structure  $(M, \varphi)$  is called a  $G_2$  manifold if the holonomy group of the Levi-Civita connection (of the metric  $g_\varphi$ ) lies inside of  $G_2$ . Equivalently  $(M, \varphi)$  is a  $G_2$  manifold if  $d\varphi = d^*\varphi = 0$  (i.e.  $\varphi$  harmonic).

In short one can define  $G_2$  manifold to be any Riemannian manifold  $(M^7, g)$  whose holonomy group is contained in  $G_2$ , then  $\varphi$  and the cross product  $\times$  comes as a consequence. It turns out that the condition  $\varphi$  being harmonic is equivalent to the condition that at each point  $x \in M$  there is a chart such that  $\varphi(x) = \varphi_0 + O(|x|^2)$ .

For example if  $(X, \omega, \Omega)$  is a complex 3-dimensional Calabi-Yau manifold with the Kähler form  $\omega$  and a nowhere vanishing holomorphic 3-form  $\Omega$ , then  $X^6 \times S^1$  has holonomy group  $SU(3) \subset G_2$ , hence is a  $G_2$  manifold. In this case  $\varphi = \text{Re } \Omega + \omega \wedge dt$ .

**Definition 3.** A 3-dimensional submanifold  $Y$  of a  $G_2$  manifold  $(M, \varphi)$  is called *associative submanifold* if  $\varphi|_Y \equiv \text{vol}(Y)$ . This condition is equivalent to  $\chi|_Y \equiv 0$ , where  $\chi \in \Omega^3(M, TM)$  is a tangent bundle valued 3-form defined by the identity:

$$(4) \quad \langle \chi(u, v, w), z \rangle = * \varphi(u, v, w, z)$$

The equivalence of these conditions follows from the ‘associator equality’ of [HL]

$$\varphi(u, v, w)^2 + |\chi(u, v, w)|^2 / 4 = |u \wedge v \wedge w|^2$$

We will denote the tangent bundle valued 3-form in  $\mathbb{R}^7$  corresponding to  $\varphi_0$  by  $\chi_0$ . Throughout this paper we will denote the sections of a bundle  $\xi \rightarrow Y$  by  $\Omega^0(Y, \xi)$  or

simply by  $\Omega^0(\xi)$ , and the bundle valued  $p$ -forms by  $\Omega^p(\xi) = \Omega^0(\Lambda^p T^*Y \otimes \xi)$ . We will denote the coframe bundle by  $\mathcal{P}^*(M) \rightarrow M$  and its adapted frame bundle by  $\mathcal{P}(M)$ . They can be  $G_2$  or  $SO(7)$  frame bundles, when needed we will specify them by the notations  $\mathcal{P}_{SO(7)}(M)$  or  $\mathcal{P}_{G_2}(M)$ .

## 2. Grassmann bundles

Let  $G(3, 7)$  be the Grassmann manifold of oriented 3-planes in  $\mathbb{R}^7$ . Let  $M^7$  be any smooth 7-manifold, and let  $\tilde{M}$  be the 3-planes in  $T(M)$ , i.e.  $\tilde{M} \rightarrow M$  is the bundle of Grassmannians over  $M$  defined by

$$\tilde{M} = \mathcal{P}_{SO(7)}(M) \times_{SO(7)} G(3, 7) \rightarrow M$$

Let  $\xi \rightarrow G(3, 7)$  be the universal  $\mathbb{R}^3$  bundle, and  $\nu = \xi^\perp \rightarrow G(3, 7)$  be the dual  $\mathbb{R}^4$  bundle. Hence  $Hom(\xi, \nu) = \xi^* \otimes \nu \rightarrow G(3, 7)$  is just the tangent bundle  $TG(3, 7)$ .  $\xi, \nu$  extend fiberwise to give bundles  $\Xi \rightarrow \tilde{M}, \mathbb{V} \rightarrow \tilde{M}$  respectively, and let  $\Xi^*$  denote the dual of  $\Xi$ . Notice that  $Hom(\Xi, \mathbb{V}) = \Xi^* \otimes \mathbb{V} \rightarrow \tilde{M}$  is just the bundle of vertical vectors  $T_{vert}(\tilde{M})$  of  $T(\tilde{M}) \rightarrow M$ , i.e. it is the bundle of tangent vectors to the fibers of  $\pi : \tilde{M} \rightarrow M$ . Hence

$$(5) \quad T\tilde{M} = T_{vert}(\tilde{M}) \oplus \Xi \oplus \mathbb{V} = (\Xi^* \otimes \mathbb{V}) \oplus \pi^*TM$$

Let  $\mathcal{P}(\mathbb{V}) \rightarrow \tilde{M}$  be the  $SO(4)$  frame bundle of the vector bundle  $\mathbb{V}$ , and identify  $\mathbb{R}^4$  with the quaternions  $\mathbb{H}$ , and identify  $SU(2)$  with the unit quaternions  $Sp(1)$ . Recall that  $SO(4)$  is the equivalence classes of pairs  $[q, \lambda]$  of unit quaternions

$$SO(4) = (SU(2) \times SU(2))/\mathbb{Z}_2$$

Hence  $\mathbb{V} \rightarrow \tilde{M}$  is the associated vector bundle to  $\mathcal{P}(\mathbb{V})$  via the  $SO(4)$  representation

$$(6) \quad x \mapsto qx\lambda^{-1}$$

There is a pair of  $\mathbb{R}^3 = im(\mathbb{H})$  bundles over  $\tilde{M}$  corresponding to the left and right  $SO(3)$  reductions of  $SO(4)$ , so they are given by the  $SO(3)$  representations

$$(7) \quad \begin{aligned} \lambda_+(\mathbb{V}) &: x \mapsto qxq^{-1} \\ \lambda_-(\mathbb{V}) &: y \mapsto \lambda y \lambda^{-1} \end{aligned}$$

The map  $x \otimes y \mapsto xy$  gives actions  $\lambda_+(\mathbb{V}) \otimes \mathbb{V} \rightarrow \mathbb{V}$  and  $\mathbb{V} \otimes \lambda_-(\mathbb{V}) \rightarrow \mathbb{V}$ , by combining we can think of it as one conjugation action

$$(8) \quad (\lambda_+(\mathbb{V}) \otimes \lambda_-(\mathbb{V})) \otimes \mathbb{V} \rightarrow \mathbb{V}$$

If the  $SO(4)$  bundle  $\mathbb{V} \rightarrow \tilde{M}$  lifts to a  $Spin(4) = SU(2) \times SU(2)$  bundle (locally it does), we get two additional bundles over  $\tilde{M}$

$$(9) \quad \begin{aligned} \mathcal{S} &: y \mapsto qy \\ \mathbb{E} &: y \mapsto y\lambda^{-1} \end{aligned}$$

which gives  $\mathbb{V}$  as a tensor product of two quaternionic line bundles  $\mathbb{V} = \mathcal{S} \otimes_{\mathbb{H}} \mathbb{E}$ . In particular  $\lambda_+(\mathbb{V}) = ad(\mathcal{S})$  and  $\lambda_-(\mathbb{V}) = ad(\mathbb{E})$ , i.e. they are the  $SO(3)$  reduction of the  $SU(2)$  bundles  $\mathcal{S}$  and  $\mathbb{E}$ . In particular there is multiplication map  $\mathcal{S} \otimes \mathbb{E} \rightarrow \mathbb{V}$ .

### 3. Associative Grassmann bundles

Now consider the *Grassmannian of associative 3-planes*  $G^\varphi(3, 7)$  in  $\mathbb{R}^7$ , consisting of elements  $L \in G(3, 7)$  with the property  $\varphi_0|_L = vol(L)$  (or equivalently  $\chi_0|_L = 0$ ).  $G_2$  acts on  $G^\varphi(3, 7)$  transitively with the stabilizer  $SO(4)$ , giving the the identification  $G^\varphi(3, 7) = G_2/SO(4)$ . Identify the imaginary octanions  $\mathbb{R}^7 = \text{Im}(\mathbb{O}) \cong im(\mathbb{H}) \oplus \mathbb{H}$ , then the action of the subgroup  $SO(4) \subset G_2$  on  $\mathbb{R}^7$  is given by:

$$(10) \quad \begin{pmatrix} A & 0 \\ 0 & \rho(A) \end{pmatrix}$$

where  $\rho : SO(4) = (SU(2) \times SU(2))/\mathbb{Z}_2 \rightarrow SO(3)$  is the projection of the first factor ([HL]). Now let  $M^7$  be a  $G_2$  manifold. As above, this time we can construct the bundle of associative Grassmannians over  $M$

$$\tilde{M}_\varphi = \mathcal{P}_{G_2}(M) \times_{G_2} G^\varphi(3, 7) \rightarrow M$$

which is just the quotient bundle  $\tilde{M}_\varphi = \mathcal{P}(M)/SO(4) \rightarrow \mathcal{P}(M)/G_2 = M$ . As in previous section, the universal bundles  $\xi, \nu = \xi^\perp \rightarrow G^\varphi(3, 7)$  induce 3 and 4 plane bundles  $\Xi \rightarrow \tilde{M}_\varphi$  and  $\mathbb{V} \rightarrow \tilde{M}_\varphi$  (i.e. restricting the universal bundles from  $\tilde{M}$ ). Also

$$(11) \quad T\tilde{M}_\varphi = T_{vert}(\tilde{M}_\varphi) \oplus \Xi \oplus \mathbb{V}$$

So in this associative case, we have an important identification  $\Xi = \lambda_+(\mathbb{V})$  (as bundles over  $\tilde{M}_\varphi$ ). For simplicity we will denote  $\mathbb{Q} = \lambda_-(\mathbb{V})$ . As before we have an action  $\mathbb{V} \otimes \mathbb{Q} \rightarrow \mathbb{V}$ , and the similar action of  $\lambda_+(\mathbb{V})$  on  $\mathbb{V}$ . We will brake the dual of this last action into a pair of Clifford multiplications on  $\mathbb{V}^\pm = \mathbb{V}$

$$(12) \quad \Xi^* \otimes \mathbb{V}^\pm \rightarrow \mathbb{V}^\mp$$

given by  $x \otimes y \mapsto -\bar{x}y$  and  $x \otimes y \mapsto xy$ , on  $\mathbb{V}^+$  and  $\mathbb{V}^-$  respectively. By writing  $x_1 \wedge x_2 = \frac{1}{2}(x_1 \otimes x_2 - x_2 \otimes x_1)$  we can extend this to an action of the whole exterior algebra  $\Lambda^*(\Xi^*)$ . For example  $(x_1 \wedge x_2) \otimes y \mapsto \frac{1}{2}(-x_1\bar{x}_2 + x_2\bar{x}_1)y = Im(x_2\bar{x}_1)y$  gives

$$(13) \quad \Lambda^2(\Xi^*) \otimes \mathbb{V} \rightarrow \mathbb{V}$$

Recall  $T_{vert}(\tilde{M}) = \Xi^* \otimes \mathbb{V}$  is the the subbundle of vertical vectors of  $T(\tilde{M}) \rightarrow M$ . The total space  $E(\nu_\varphi)$  of the normal bundle of the imbedding  $\tilde{M}_\varphi \subset \tilde{M}$  should be thought of an open tubular neighborhood of  $\tilde{M}_\varphi$  in  $\tilde{M}$ , and it has a nice description:

**Lemma 1.** ([M]) *The normal bundle  $\nu_\varphi$  of  $\tilde{M}_\varphi \subset \tilde{M}$  is isomorphic to  $\mathbb{V}$ , and the bundle of vertical vectors  $T_{vert}(\tilde{M}_\varphi)$  of  $T(\tilde{M}_\varphi) \rightarrow M$  is the kernel of the Clifford multiplication  $c : \Xi^* \otimes \mathbb{V} \rightarrow \mathbb{V}$ .*

Hence we have  $\Xi^* \otimes \mathbb{V}|_{\tilde{M}_\varphi} = T_{vert}(\tilde{M}_\varphi) \oplus \nu_\varphi$  from the exact sequence over  $\tilde{M}_\varphi$

$$(14) \quad T_{vert}(\tilde{M}_\varphi) \rightarrow \Xi^* \otimes \mathbb{V}|_{\tilde{M}_\varphi} \xrightarrow{c} \mathbb{V}|_{\tilde{M}_\varphi}$$

the quotient bundle,  $T_{vert}(\tilde{M})/T_{vert}(\tilde{M}_\varphi)$  is isomorphic to  $\nu_\varphi$ . Also, if  $\{e^i\}$  is a local ON basis for  $\Xi^*$  then projection  $\pi_\varphi : T(\tilde{M})|_{\tilde{M}_\varphi} \rightarrow T(\tilde{M}_\varphi)$  is given by

$$(15) \quad \pi_\varphi(a \otimes v) = a \otimes v + (1/3) \sum_{j=1}^3 e^j \otimes e^j(a.v)$$

**Remark 1.** *The lemma holds since the Lie algebra inclusion  $\mathfrak{g}_2 \subset \mathfrak{so}(7)$  is given by*

$$\begin{pmatrix} a & \beta \\ -\beta^t & \rho(a) \end{pmatrix}$$

where  $a \in \mathfrak{so}(4)$  is  $y \mapsto qy - yq$ , and  $\rho(a) \in \mathfrak{so}(3)$  is by  $x \mapsto qx - xq$ . So the tangent space inclusion of  $G_2/SO(4) \subset SO(7)/SO(4) \times SO(3)$  is given by the matrix  $\beta$ . And if we write  $\beta$  as column vectors  $\beta = (\beta_1, \beta_2, \beta_3)$ , then  $\beta_1 i + \beta_2 j + \beta_3 k = 0$  ([M], [Mc])

#### 4. Associative submanifolds

Any imbedding of a 3-manifold  $f : Y^3 \hookrightarrow M^7$  induces a bundle  $\tilde{Y} \rightarrow Y$  with fibers  $G(3, 7)$ , and the Gauss map of  $f$  canonically lifts to an imbedding  $\tilde{f} : Y \hookrightarrow \tilde{M}$ :

$$(16) \quad \begin{array}{ccc} & \tilde{M} \supset \tilde{M}_\varphi & \\ \tilde{f} \nearrow & \downarrow & \\ Y & \xrightarrow{f} & M \end{array}$$

Also the pull backs  $\tilde{f}^*\Xi = T(Y)$  and  $\tilde{f}^*\mathbb{V} = \nu(Y)$  give tangent and normal bundles of  $Y$ . Furthermore, if  $f$  is an imbedding as an associative submanifold in a  $G_2$  manifold  $(M, \varphi)$ , then the image of  $\tilde{f}$  lands in  $\tilde{M}_\varphi$ . We will denote this canonical lifting of any 3-manifold  $Y \subset M$  by  $\tilde{Y} \subset \tilde{M}$ .

$\tilde{M}_\varphi$  can be thought of a universal space parametrizing associative submanifolds of  $M$ . In particular if  $\tilde{f} : Y \hookrightarrow \tilde{M}_\varphi$  is the lifting of an associative submanifold, by pulling

back we see that the principal  $SO(4)$  bundle  $\mathcal{P}(\mathbb{V}) \rightarrow \tilde{M}_\varphi$  induces an  $SO(4)$ -bundle  $\mathcal{P}(Y) \rightarrow Y$ , and gives the following vector bundles over  $Y$  via the representations:

$$(17) \quad \begin{aligned} \nu(Y) &: y \mapsto qy\lambda^{-1} \\ T(Y) &: x \mapsto qxq^{-1} \\ Q &: y \mapsto \lambda y\lambda^{-1} \end{aligned}$$

where  $[q, \lambda] \in SO(4)$ . Note that  $T(Y) = \lambda_+(\nu)$  and  $Q = \lambda_-(\nu)$ , where  $\nu = \nu(Y)$ . Also we can identify  $T^*Y$  with  $TY$  by the induced metric. We have the actions  $T^*Y \otimes \nu \rightarrow \nu$ ,  $\Lambda^*(T^*Y) \otimes \nu \rightarrow \nu$ ,  $\nu \otimes Q \rightarrow \nu$  from above. By combining them we can define an action of the bundle valued differential forms

$$(18) \quad \Omega^*(Y; Q) \otimes \Omega^0(Y, \nu) \rightarrow \Omega^0(Y, \nu)$$

For example, if  $F = (x_1 \wedge x_2) \otimes y$ , and  $y \in \Omega^0(Y, Q)$ , and  $z \in \Omega^0(Y, \nu)$  then

$$(19) \quad F \otimes z \mapsto \text{Im}(x_2 \bar{x}_1)zy$$

Let  $\mathbb{L} = \Lambda^3(\Xi) \rightarrow \tilde{M}$  be the determinant (real) line bundle. Note that the definition (4) implies that if  $\chi$  maps every oriented 3-plane in  $T_x(M)$  to its complementary subspace, so  $\chi$  gives a bundle map  $\mathbb{L} \rightarrow \mathbb{V}$  over  $\tilde{M}$ , which is a section of  $\mathbb{L}^* \otimes \mathbb{V} \rightarrow \tilde{M}$ . Also,  $\Xi$  being an oriented bundle  $\mathbb{L}$  is trivial so  $\chi$  actually gives a section

$$(20) \quad \chi \in \Omega^0(\tilde{M}, \mathbb{V})$$

Clearly  $\tilde{M}_\varphi \subset \tilde{M}$  is the codimension 4 submanifold given as the zeros of this section. So associative submanifolds  $Y \subset M$  are characterized by the condition  $\chi|_{\tilde{Y}} = 0$  where  $\tilde{Y} \subset \tilde{M}$  is the canonical lifting of  $Y$ . Similarly  $\varphi$  defines a map  $\varphi : \tilde{M} \rightarrow \mathbb{R}$ .

## 5. Dirac operator

In general, the normal bundle  $\nu = \nu(Y)$  of any orientable 3-manifold  $Y$  in a  $G_2$  manifold  $(M, \varphi)$  has a  $\text{Spin}(4)$  structure (e.g. [B2]). Hence we have  $SU(2)$  bundles  $S$  and  $E$  over  $Y$  (the pull-backs of  $\mathcal{S}$  and  $\mathbb{E}$  in (9)) such that  $\nu = S \otimes_{\mathbb{H}} E$ , with  $SO(3)$  reductions  $adS = \lambda_+(\nu)$  and  $adE = \lambda_-(\nu)$ . Note that  $adE$  is also the bundle of endomorphisms  $\text{End}(E)$ . If  $Y$  is associative, then the  $SU(2)$  bundle  $S$  reduces to  $TY$ , i.e.  $S$  becomes the spinor bundle of  $Y$ , and so  $\nu$  becomes a twisted spinor bundle on  $Y$ .

The Levi-Civita connection of the  $G_2$  metric of  $(M, \varphi)$  induces connections on the associated bundles  $\mathbb{V}$  and  $\Xi$  on  $\tilde{M}$ . In particular it induces connections on the tangent and normal bundles of any submanifold  $Y^3 \subset M$ . We will call these connections the *background connections*. Let  $\mathbb{A}_0$  be the induced connection on the normal bundle  $\nu = S \otimes E$ . From the Lie algebra decomposition  $so(4) = so(3) \oplus so(3)$ , we can write  $\mathbb{A}_0 = A_0 \oplus B_0$ , where  $A_0$  and  $B_0$  are connections on  $S$  and  $E$  respectively.

Let  $\mathcal{A}(S)$  and  $\mathcal{A}(E)$  be the set of connections on the bundles  $S$  and  $E$ . Hence connections  $A \in \mathcal{A}(S)$ ,  $B \in \mathcal{A}(E)$  are in the form  $A = A_0 + a$ ,  $B = B_0 + b$ , where  $a \in \Omega^1(Y, adS)$  and  $b \in \Omega^1(Y, adE)$ . So  $\Omega^1(Y, \lambda_{\pm}(\nu))$  parametrizes connections on  $S$  and  $E$ , and the connections on  $\nu$  are in the form  $\mathbb{A} = A \oplus B$ . To emphasize the dependency on  $a$  and  $b$  we will sometimes denote  $\mathbb{A} = \mathbb{A}(a, b)$ , and  $\mathbb{A}_0 = \mathbb{A}(0, 0)$ .

Now, let  $Y^3 \subset M$  be any smooth manifold. We can express the covariant derivative  $\nabla_{\mathbb{A}} : \Omega^0(Y, \nu) \rightarrow \Omega^1(Y, \nu)$  on  $\nu$  by  $\nabla_{\mathbb{A}} = \sum e^i \otimes \nabla_{e_i}$ , where  $\{e_i\}$  and  $\{e^i\}$  are orthonormal tangent and cotangent frame fields of  $Y$ , respectively. Furthermore if  $Y$  is an associative submanifold, we can use the Clifford multiplication (12) to form the twisted Dirac operator  $\not{D}_{\mathbb{A}} : \Omega^0(Y, \nu) \rightarrow \Omega^0(Y, \nu)$

$$(21) \quad \not{D}_{\mathbb{A}} = \sum e^i \cdot \nabla_{e_i}$$

The sections lying in the kernel of this operator are usually called harmonic spinors twisted by  $(E, \mathbb{A})$ . Elements of the kernel of  $\not{D}_{\mathbb{A}_0}$  are called the harmonic spinors twisted by  $E$ , or just the twisted harmonic spinors.

## 6. Complex associative submanifolds

There is an interesting class of associative submanifolds which we call *complex associative*, they are defined as follows: The subgroups  $U(2) \subset SO(4) \subset G_2$ , from

$$(SU(2) \times S^1)/\mathbb{Z}_2 \subset (SU(2) \times SU(2))/\mathbb{Z}_2 \subset G_2$$

give a  $U(2)$ -principal bundle  $\mathcal{P}_{G_2}(M) \rightarrow \bar{M}_{\varphi} = \mathcal{P}_{G_2}(M)/U(2)$ . Note that  $\bar{M}_{\varphi}$  is the total space of an  $S^2$  bundle  $\bar{M}_{\varphi} \rightarrow \tilde{M}_{\varphi} = \mathcal{P}(M)/SO(4)$ . This is just the sphere bundle of the  $\mathbb{R}^3$ -bundle  $\lambda_-(\mathbb{V}) \rightarrow \tilde{M}_{\varphi}$ . So we can identify the unit sections  $j \in \lambda_-(\mathbb{V})$  with complex structures on  $\mathbb{V}$ , i.e. the right reductions of  $SO(4)$  to  $U(2)$ , where

$$U(2) = \text{Spin}^c(3) = (SU(2) \times S^1)/\mathbb{Z}_2$$

Just as  $SO(4)$  is the stabilizer subgroup of the action of  $G_2$  on  $G^{\varphi}(3, 7)$ , the subgroup  $U(2) \subset SO(4)$  is the stabilizer of the action of  $G_2$  on the "framed" version of  $G^{\varphi}(3, 7)$ . That is, if  $\nu \rightarrow G^{\varphi}(3, 7)$  is the dual universal bundle, then the action of  $G_2$  on  $G^{\varphi}(3, 7)$  extends to an action to the sphere bundle of  $\lambda_-(\nu)$  with stabilizer  $U(2)$ .

By the representations of  $U(2)$  induced from  $SO(4)$  we can form the same vector bundles  $\mathbb{V}, \Xi$  on  $\bar{M}_{\varphi}$  as in Sections (2) and (3). Except in this case we have  $\lambda \in S^1$  and hence  $\lambda_-(\mathbb{V})$  becomes the trivial bundle, and  $\mathbb{V}$  becomes a  $\mathbb{C}^2$  bundle. The reduction  $j$  gives an extra line bundle  $\mathbb{L}$  via the representation  $\rho_2(q, \lambda)(y) = y\lambda^{-2}$  (the determinant line bundle of  $\mathbb{V}$ ). Similarly we have the same actions of (12), (13).



Since  $\lambda_-(\mathbb{V})$  is trivial its action becomes the multiplication with complex numbers. In this case we have a useful quadratic bundle map  $\sigma : \mathbb{V} \otimes \mathbb{V} \rightarrow \Lambda^2(\Xi^*)_{\mathbb{C}}$  given by

$$(22) \quad \sigma(x, y) = -\frac{1}{2}(xi\bar{y})i$$

In particular if  $x = z + jw \in \mathbb{O}$  (octanions), we have  $\sigma(x, x) = \frac{|z|^2 - |w|^2}{2} + j\bar{z}w$

So if  $f : Y^3 \hookrightarrow M$  is an associative submanifold, and  $\tilde{f} : Y^3 \hookrightarrow \tilde{M}_\varphi$  is the lifting of its Gauss map, then the normal bundle of  $f$  has a  $U(2)$  structure if and only if  $\tilde{f}$  lifts

$$(23) \quad \begin{array}{ccc} & & \bar{M}_\varphi \\ & \nearrow \bar{f} & \downarrow \\ Y & \xrightarrow{\tilde{f}} & \tilde{M}_\varphi \end{array}$$

As before we get  $\bar{f}^*\Xi = T(Y)$  and  $\bar{f}^*\mathbb{V} = \nu(Y)$ , plus a line bundle  $L = \bar{f}^*(\mathbb{L})$ , and  $\nu$  becomes a complex  $U(2)$  bundle, which we will call  $W$ . So  $W$  is the spinor bundle  $W(L) \rightarrow Y$  of a  $Spin^c(3)$  structure on  $Y$ , and  $L$  is its determinant bundle. If this  $Spin^c(3)$  structure comes from a  $Spin(3)$  structure, then  $W = S \otimes L^{1/2}$  where  $S$  is the spinor bundle on  $Y$ . So in concrete terms:

**Definition 4.** A *complex associative submanifold*  $(Y, j)$  of a  $G_2$  manifold  $M$  is an associative manifold in  $M$  with a unit section  $j \in \Omega^0(\lambda_-(\nu))$ , where  $\nu = \nu(Y)$ .

Note that the normal bundle  $\nu(Y)$  of any associative submanifold  $Y$  has a  $U(2)$  structure (in fact it is trivial [BSh]), so liftings to  $\tilde{M}_\varphi$  always exists. We will call any  $Y$  with such a lifting a *complex associative submanifold*. But for such a lifting the background  $SO(4)$  connection  $\mathbb{A}_0$  on  $\nu = \nu(Y)$  may not reduce to a  $U(2)$  connection. Let  $j \in \lambda_-(\nu)$  be the unit section describing the  $U(2)$  structure on  $\nu$ . Then  $\mathbb{A}_0$  reduces to a  $U(2)$  connection if  $\nabla_{A_0}(j) = 0$ , where  $A_0$  is the connection on  $\lambda_-(\nu)$  induced from the background connection  $\mathbb{A}_0$ , and  $\nabla_{A_0}$  is the covariant differentiation.

$$(24) \quad \Omega^0(Y, \lambda_-(\nu)) \xrightarrow{\nabla_{A_0}} \Omega^1(Y, \lambda_-(\nu))$$

**Definition 5.** An *integrable complex associative submanifold*  $(Y, j)$  of a  $G_2$  manifold  $M$  is an associative manifold in  $M$  with a unit section  $j \in \Omega^0(\lambda_-(\nu))$  with  $\nabla_{A_0}(j) = 0$ .

Let  $Y$  be an integrable complex associative submanifold of a  $G_2$  manifold  $M$ , and  $\mathbb{A}_0$  be the background  $U(2)$  connection on its normal bundle  $W$  induced by the  $G_2$  metric of  $M$ . Then  $\mathbb{A}_0 = S_0 \oplus A_0$  where  $S_0$  is a connection on  $\lambda_+(W) = T(Y)$  and  $A_0$  is a connection on the line bundle  $L$ . In particular, we can deform the background connection  $\mathbb{A}_0 = \mathbb{A}(0) \mapsto \mathbb{A}(a)$ , where  $\mathbb{A}(a) = S_0 \oplus (A_0 + a)$ , where  $a \in \Omega^1(Y)$ . We will use this deformation in a coming section.

**Remark 2.** Recall that if  $(X, \omega, \Omega)$  is a complex 3-dimensional Calabi-Yau manifold then  $(X^6 \times S^1, \varphi)$  is a  $G_2$  manifold, where  $\varphi = \text{Re } \Omega + \omega \wedge dt$ . The submanifolds of the form  $C \times S^1$  where  $C \subset X$  a holomorphic curve, are examples of integrable complex associative submanifolds of  $X \times S^1$ .

## 7. Deformations

In [M] McLean showed that, in a neighborhood of an associative submanifold  $Y$  in a  $G_2$  manifold  $(M, \varphi)$  the space of associative submanifolds can be identified with the harmonic spinors on  $Y$  twisted by  $E$ . Since the cokernel of the Dirac operator can vary, we can not determine the dimension of its kernel (it always has zero index on  $Y^3$ ). We will remedy this by deforming  $Y$  in a larger class of submanifolds. To motivate our approach let us sketch a proof of the McLean's theorem by adapting the explanation in [B3]. Let  $Y \subset M$  be an associative submanifold,  $Y$  will determine a lifting  $\tilde{Y} \subset \tilde{M}_\varphi$ . Let us recall that the  $G_2$  structure  $\varphi$  gives a metric connection on  $M$ , hence it gives a covariant differentiation in the normal bundle  $\nu(Y) = \nu$

$$\nabla_{\mathbb{A}_0} : \Omega^0(Y, \nu) \rightarrow \Omega^1(Y, \nu) = \Omega^0(Y, T^*Y \otimes \nu)$$

Recall that we identified  $T_y^*(Y) \otimes \nu_y(Y)$  with a copy of  $TG(3, 7)$ , more precisely we identified it by the tangent space of the Grassmannian of 3-planes in  $T_y(M)$ . So the covariant derivative lifts normal vector fields  $v$  of  $Y \subset M$  to vertical vector fields  $\tilde{v}$  in  $T(\tilde{M})|_{\tilde{Y}}$ . We want the normal vector fields  $v$  of  $Y$  always move  $Y$  in the class of associative submanifolds of  $M$ , i.e. we want the liftings  $\tilde{Y}_v$  of the nearby copies  $Y_v$  of  $Y$  (pushed off by the vector field  $v$ ) to lie in  $T(\tilde{M}_\varphi) \subset T(\tilde{M})$  upstairs, i.e. we want the component of  $\tilde{v}$  in the direction of the normal bundle  $\tilde{M}_\varphi \subset \tilde{M}$  vanish. By Lemma 1, this means  $\nabla_{\mathbb{A}_0}(v)$  should be in the kernel of the Clifford multiplication  $c : \Omega^0(T^*(Y) \otimes \nu) \rightarrow \Omega^0(\nu)$ , i.e.

$$(25) \quad \mathcal{D}_{\mathbb{A}_0}(v) = c(\nabla_{\mathbb{A}_0}(v)) = 0$$

Here  $\mathcal{D}_{\mathbb{A}_0}$  is the Dirac operator induced by the background connection  $\mathbb{A}_0$ , which is the composition.

$$(26) \quad \Omega^0(Y, \nu) \xrightarrow{\nabla_{\mathbb{A}_0}} \Omega^0(Y, T^*Y \otimes \nu) \xrightarrow{c} \Omega^0(Y, \nu)$$

To sum up what we have done: Let  $f : Y^3 \hookrightarrow M$  be an associative manifold in a  $G_2$  manifold, and  $\tilde{f} : Y^3 \hookrightarrow \tilde{M}_\varphi$  be its canonical lifting. By using the metric connection, the map on the space of immersions  $\Phi : \text{Im}(Y, M) \rightarrow \text{Im}(Y, \tilde{M})$  given by  $f \mapsto \tilde{f}$ , induces a map on the tangent spaces

$$(27) \quad d\Phi : T_f \text{Im}(Y, M) \longrightarrow T_{\tilde{f}} \text{Im}(Y, \tilde{M}) \supset T_{\tilde{f}} \text{Im}(Y, \tilde{M}_\varphi)$$

and the condition that  $d\Phi(v)$  to lie in  $T_{\tilde{f}} \text{Im}(Y, \tilde{M}_\varphi)$  is given by  $\mathcal{D}_{\mathbb{A}_0}(v) = 0$ .




FIGURE 1.

Now  $\Phi$  may not be transversal to  $Im(Y, \tilde{M}_\varphi)$ , i.e. cokernel of  $\mathcal{D}_{\mathbb{A}_0}$  maybe non zero, but we can make it zero by deforming the connection in the normal bundle (i.e. in  $E$  or  $S$  as indicated in Section 5)  $\mathbb{A}_0 = \mathbb{A}(0, 0) \rightarrow A = \mathbb{A}(a, b)$ , where  $a \in \Omega^1(Y, \lambda_+(\nu))$  and  $b \in \Omega^1(Y, \lambda_-(\nu))$ , and  $\nu = \nu(Y)$ . Then (26) becomes parametrized

$$(28) \quad \Omega^0(\nu) \times \Omega^1(\lambda_\pm(\nu)) \xrightarrow{\mathcal{D}_{\mathbb{A}}} \Omega^0(\nu)$$

that is (25) becomes a twisted Dirac equation  $\mathcal{D}_{\mathbb{A}}(v) = c(\nabla_{\mathbb{A}}(v)) = \mathcal{D}_{\mathbb{A}_0}(v) + \alpha v = 0$ , where  $\alpha = (a, b)$ . This gives a smooth moduli space parametrized by  $\Omega^0(\lambda_\pm(\nu))$ . Then by a generic choice of  $\alpha \in \Omega^0(\lambda_\pm(\nu))$  we get a zero dimensional perturbed moduli space, whose elements perhaps should be called  $A$ -associative submanifolds (to make an analogy with  $J$ -holomorphic curves). Perturbation of  $A_0$  may be achieved by deforming background  $G_2$  structure (i.e. by changing  $\varphi$ ), but we will not pursue this line, instead we will deform the associative submanifolds in a larger class of objects. To get compactness we will couple the Dirac equation with another equation to get Seiberg-Witten equations. We will first treat the complex associative case.

## 8. Deforming Complex Associative Submanifolds

Let  $\mathcal{M}_{\mathbb{C}}(M, \varphi)$  be the set of integrable complex associative submanifolds of a  $G_2$  manifold  $(M, \varphi)$ . We wish to study the local structure of  $\mathcal{M}_{\mathbb{C}}(M, \varphi)$  near a particular  $Y$ . The normal bundle of  $Y$  is a  $U(2)$  bundle  $\nu(Y) = W$ , and  $L \rightarrow Y$  is its determinant line bundle. Let  $\mathbb{A}_0 = S_0 \oplus A_0$  be the background connection on  $W$ , where  $S_0$  is a

connection on  $\lambda_+(W) = T(Y)$ , and  $A_0$  is a connection on  $L$ . As before we will denote  $\mathbb{A} = \mathbb{A}(a) = S_0 \oplus (A_0 + a)$ , where  $a \in \Omega^1(Y) = T_{\mathbb{A}_0} \mathcal{A}(L)$  (tangent space to space of connections). So (28) becomes

$$(29) \quad \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R}) \xrightarrow{\mathcal{D}_{\mathbb{A}}} \Omega^0(Y, W)$$

which can be thought of the derivative of a similarly defined map

$$(30) \quad \Omega^0(Y, W) \times \mathcal{A}(L) \rightarrow \Omega^0(Y, W)$$




FIGURE 2.

In each slice  $A$ , we are deforming along normal vector fields by the connection  $\mathbb{A}$  (which is a perturbation of the background connection  $\mathbb{A}_0$ ). To get compactness we can cut the moduli space by an additional equation as in Seiberg-Witten theory, i.e.

$$(31) \quad \Phi : \Omega^0(Y, W) \times \mathcal{A}(L) \rightarrow \Omega^0(Y, W) \times \Omega^2(Y, i\mathbb{R})$$

$$(32) \quad \begin{aligned} \mathcal{D}_{\mathbb{A}}(v) &= 0 \\ F_A &= \sigma(v, v) \end{aligned}$$

where  $F_A$  is the curvature of the connection  $A = A_0 + a$  in  $L$ . For convenience we will replace the second equation above by  $*F_A = \mu(v, v)$ , where  $*$  is the star operator on  $Y$  and  $\mu(v, v) = *\sigma(v, v)$ . Now we proceed exactly as in Seiberg-Witten theory of 3-manifolds (e.g. [C], [Ma], [W]). To obtain smoothness on the zeros of  $\Phi$ , we perturb the equations by 1-forms  $\delta \in \Omega^1(Y)$ . This gives a new parametrized equation  $\Phi = 0$

$$(33) \quad \Phi : \Omega^0(Y, W) \times \mathcal{A}(L) \times \Omega^1(Y) \rightarrow \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R})$$

$$(34) \quad \begin{aligned} \mathcal{D}_A(v) &= 0 \\ *F_A + i\delta &= \mu(v, v) \end{aligned}$$

It follows that in particular  $\delta$  must be coclosed.  $\Phi$  has a linearization:

$$(35) \quad D\Phi_{(v_0, A_0, 0)} : \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R}) \times \Omega^1(Y) \rightarrow \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R})$$

$$D\Phi_{(v_0, A_0, 0)}(v, a, \delta) = (\mathcal{D}_{A_0}(v) + a \cdot v_0, *da + i\delta - 2\mu(v_0, v))$$

We see that  $\Phi^{-1}(0)$  is smooth and the projection  $\Phi^{-1}(0) \rightarrow \Omega^1(Y)$  is onto, so by Sard's theorem for generic choice of  $\delta$  we can make  $\Phi_\delta^{-1}(0)$  smooth, where  $\Phi_\delta(v, A) = \Phi(v, A, \delta)$ . The normal bundle of  $Y$  has a complex structure, so the gauge group  $\mathcal{G}(Y) = \text{Map}(Y, S^1)$  acts on the solution set  $\Phi_\delta^{-1}(0)$ , and makes quotient  $\Phi_\delta^{-1}(0)/\mathcal{G}(Y)$  a smooth zero dimensional manifold. This is because the infinitesimal action of  $\mathcal{G}(Y)$  on the complex  $\Phi_\delta : \Omega^0(Y, W) \times \mathcal{A}(L) \rightarrow \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R})$  is given by the map

$$\Omega^0(Y, i\mathbb{R}) \xrightarrow{G} \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R})$$

where  $G(f) = (fv_0, df)$ . So after dividing by  $\mathcal{G}$ , tangentially the complex  $\Phi_\delta$  becomes

$$(36) \quad \Omega^0(Y; i\mathbb{R}) \xrightarrow{G} \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R}) \rightarrow \Omega^0(Y, W) \times \Omega^1(Y, i\mathbb{R})/\mathcal{G}$$

Hence the index of this complex is the sum of the indices of the Dirac operator  $\mathcal{D}_{A_0} : \Omega^0(Y, W) \rightarrow \Omega^0(Y, W)$  (which is zero), and the index of the following complex

$$\Omega^0(Y, i\mathbb{R}) \times \Omega^1(Y, i\mathbb{R}) \rightarrow \Omega^0(Y, i\mathbb{R}) \times \Omega^1(Y; i\mathbb{R})$$

given by  $(f, a) \mapsto (d^*(a), df + *da)$ , which is also zero, since  $Y^3$  has zero Euler characteristic. Furthermore  $\Phi_\delta^{-1}(0)/\mathcal{G}(Y)$  is compact and oriented (by Seiberg-Witten theory). Hence we get a number  $\chi(Y)$ . Here we don't worry about metric dependence of  $\chi(Y)$  since we have a fixed background metric induced from the  $G_2$  structure. Hence we associated a number to a complex associative submanifold  $Y$  in a  $G_2$ . In particular such a  $Y$  moves unobstructed way after perturbation, not in the sections of its normal bundle  $\Omega^0(Y, W)$ , but in  $\Omega^0(Y, W) \times \mathcal{A}(L)$  that lies over it.

**Remark 3.** *We can use the complex structures  $j$  of a complex associative manifold  $(Y, j)$  as parameters to obtain families of Seiberg-Witten invariants parametrized by  $S^2$ . It is convenient to work with the whole pull back  $S^2$  bundle  $\bar{Y}_f = \{(Y, j)\} \cong Y \times S^2$*

$$(37) \quad \begin{array}{ccc} \bar{Y}_f & \longrightarrow & \bar{M}_\varphi \\ \downarrow & \nearrow & \downarrow \\ Y & \xrightarrow{\tilde{f}} & \tilde{M}_\varphi \end{array}$$

*In particular for each  $j$  we get a complex structure  $W_j$  on  $\nu(Y)$ , and a family of Dirac actions  $c_j : T^*Y \otimes W_j \rightarrow W_j$ , and get family of Seiberg-Witten equations parametrized by  $S^2$ . More generally, we can write family of Seiberg-Witten equations defined directly on the bundles  $\mathbb{V} \rightarrow \bar{M}_\varphi$ . In this general case we will run into difficulty controlling*

the dimension Seiberg-Witten solutions since the  $*$  operator does not behave nicely under restriction. We will discuss more general version of this in Section 10.

## 9. Deformation as a parallel translation

We can obtain a version of the deformation equations (32) by an invariant geometric way. That is, we can count the normal vector fields of  $Y$ , such that moving along them preserves the connection coordinate as the covariantly constant sections:

$$(38) \quad \nabla_A(v) = 0$$

In particular this implies  $\mathcal{D}_A(v) = 0$ . Recall the Weitzenböck formula ([LM], pp 398)

$$(39) \quad \mathcal{D}_A^2(v) = \nabla_A^* \nabla_A(v) + \frac{s}{4} v + \frac{1}{2} F_A \cdot v$$

where  $s$  is the scalar curvature of  $Y$ , and  $F_A \in i\Omega^2(Y)$  is the curvature of  $A$ . We get

$$(40) \quad F_A \cdot v = -(s/2)v$$

If we write  $F_A = \Omega \cdot i$ , then from (19) the action is  $\Omega v i = -(s/2)v$ , hence by (22)

$$\Omega |v|^2 = (s/2) v i \bar{v} = s \sigma(v, v) i$$

$$(41) \quad F_A = -s \sigma(v, v) / |v|^2$$

Hence (38) is equivalent to the following equations

$$(42) \quad \begin{aligned} \mathcal{D}_A(v) &= 0 \\ F_A |v|^2 &= -s \sigma(v, v) \end{aligned}$$

Note that if  $Y$  is a hyperbolic manifold (i.e.  $s$  is a negative constant) then, in the unit normal bundle these equations coincide with the Seiberg-Witten equations. Also note that in the zero slice  $A(0) = A_0$  the equations (42) gives the McLean's deformation space  $\ker(\mathcal{D}_{A_0})$  cut down with an extra equation  $F_{A_0} = -s \sigma(v, v) / |v|^2$ . Notice that

$$v \mapsto \sigma(v, v) / |v|^2$$

is just the cone on the Hopf map  $S^3 \rightarrow S^2$ . So in the zero slice we are only including sections  $v \in \ker(\mathcal{D}_{A_0})$  in  $W$  that live in a certain complex line bundle.

### 10. $Spin^c(4)$ Structures (general case)

Now suppose that the  $SO(4)$  bundle  $\mathcal{P}(\mathbb{V}) \rightarrow \tilde{M}$  lifts to a  $Spin^c(4)$  bundle  $\mathbf{V}$ , since

$$Spin^c(4) = (SU(2) \times SU(2) \times S^1)/\mathbb{Z}_2$$

a lifting exists if the Stiefel-Whitney class  $w_2(\mathbb{V}) \in H^2(\tilde{M}, \mathbb{Z}_2)$  is an integral class, and the number of different liftings, i.e.  $Spin^c$  structures, on  $\mathbb{V}$  is given by  $H^2(\tilde{M}, \mathbb{Z})$ . These facts follow from the fibrations:

$$S^1 \rightarrow Spin^c(4) \rightarrow SO(4) \rightarrow K(\mathbb{Z}, 2) \rightarrow BSpin^c(4) \rightarrow BSO(4) \rightarrow K(\mathbb{Z}, 3)$$

$\mathbf{V}$  gives the following vector bundles by the representations, for  $[q, \lambda, t] \in Spin^c(4)$ :

$$(43) \quad \begin{aligned} \mathbb{V} &: & y &\mapsto qy\lambda^{-1} \\ \mathbb{V}^+ &: & y &\mapsto qyt^{-1} \\ \mathbb{V}^- &: & y &\mapsto \lambda yt^{-1} \\ ad(\mathbb{V}^+) &: & y &\mapsto qyq^{-1} \\ ad(\mathbb{V}^-) &: & y &\mapsto \lambda y\lambda^{-1} \\ \mathbf{L} &: & y &\mapsto yt^2 \end{aligned}$$

$\mathbb{V}^\pm$  are pair of  $\mathbb{C}^2$  bundles that reduce to  $\mathbb{R}^3$  bundles  $ad(\mathbb{V}^\pm)$ , and  $\mathbf{L}$  is a complex line bundle, and  $\mathbb{V}_{\mathbb{C}} = \mathbb{V}^+ \otimes_{\mathbb{C}} \bar{\mathbb{V}}^-$ . Clearly  $\lambda_+(\mathbb{V}) = ad(\mathbb{V}^+)$ , and  $\lambda_-(\mathbb{V}) = ad(\mathbb{V}^-)$ . Recall that if we restrict to  $\tilde{M}_\varphi$  we get an additional identifications  $\Xi = ad(\mathbb{V}^+)$ . Also the  $Spin^c$  structure on  $\mathbb{V}$  gives a  $Spin^c$  structure on the  $\mathbb{R}^3$  bundle  $ad(\mathbb{V}^+)$ , which can be identified as an equivalence class of a non-vanishing vector field  $\xi_0$  on  $ad(\mathbb{V}^+)$ , call this the basic section of  $ad(\mathbb{V}^+)$ .

Recall that  $T(\tilde{M}_\varphi) = T_{vert}(\tilde{M}_\varphi) \oplus \Xi \oplus \mathbb{V}$ , and  $T_{vert}(\tilde{M}_\varphi) \subset \Xi^* \otimes \mathbb{V}$ . Let  $\xi_0 \in \Xi$  be the basic section. We can define an action  $\rho : T(\tilde{M}_\varphi) \rightarrow Hom(\mathbb{V}^+ \oplus \mathbb{V}^-)$ , i.e.

$$T(\tilde{M}_\varphi) \otimes \mathbb{V}^\pm \rightarrow \mathbb{V}^\mp$$

For  $w = a \otimes v + x + y \in T(\tilde{M}_\varphi)$ ,  $a \otimes v \in \Xi^* \otimes \mathbb{V}$ ,  $x \in \Xi$ ,  $y \in \mathbb{V}$ , let  $v_0 = a(\xi_0)v$  and

$$w.(z_1, z_2) = (\bar{a}vz_2 + \bar{x}v_0z_2 + yz_2, -\bar{v}az_1 - \bar{v}_0xz_1 - \bar{y}z_1)$$

where  $(z_1, z_2) \in \mathbb{V}^+ \oplus \mathbb{V}^-$ . From (43) it is easy to check that this is a Dirac action, i.e.  $\rho(w) \circ \rho(w) = -|w|^2 I$ . As usual we can dualize and extend this to an action

$$\Lambda^2(T^*\tilde{M}_\varphi) \otimes \mathbb{V}^+ \rightarrow \mathbb{V}^+$$

Also as in (22) we have the quadratic bundle map  $\sigma : \mathbb{V}^+ \otimes \mathbb{V}^+ \rightarrow \Lambda^2(\Xi^*)_{\mathbb{C}}$  defined by

$$(44) \quad \sigma(x, y) = -\frac{1}{2}(xi\bar{y})i$$

Now let  $\mathcal{A}(\mathbb{L})$  be the space of connections on  $\mathbb{L}$ . Let  $\mathbb{A}_0$  be a given background connection on  $\mathbb{V}$  (e.g. the one induced from the  $G_2$  metric on  $M$ ). Then any  $A \in \mathcal{A}(\mathbb{L})$  along with  $\mathbb{A}_0$  determines a connection on  $\mathbb{V}$  and hence connections all the above associated bundles. Also if we fix  $A_0 \in \mathcal{A}(\mathbb{L})$ , any other  $A \in \mathcal{A}(\mathbb{L})$  can be written as  $A = A_0 + a$  where  $a \in \Omega^1(\tilde{M}_\varphi)$ . Hence for each  $A \in \mathcal{A}(\mathbb{L})$  we get the corresponding Dirac operator  $\mathcal{D}_A(v) = \mathcal{D}_{A_0}(v) + a.v$  on bundles  $\mathbb{V}^\pm \rightarrow \tilde{M}_\varphi$  which is the composition:

$$\Omega^0(\tilde{M}_\varphi, \mathbb{V}^+) \xrightarrow{\nabla_A} \Omega^0(\tilde{M}_\varphi, T^*\tilde{M}_\varphi \otimes \mathbb{V}^+) \xrightarrow{c} \Omega^0(\tilde{M}_\varphi, \mathbb{V}^-)$$

We can now write generalized Seiberg-Witten equations on  $\tilde{M}_\varphi$  in the usual way

$$\phi : \Omega^0(\tilde{M}_\varphi, \mathbb{V}^+) \times \mathcal{A}(\mathbb{L}) \rightarrow \Omega^0(\tilde{M}_\varphi, \mathbb{V}^-) \times \Omega^2(\tilde{M}_\varphi)$$

$$(45) \quad \begin{aligned} \mathcal{D}_A(v) &= 0 \\ F_A &= \sigma(v, v) \end{aligned}$$

and proceed as in Section 8. Equations (45) imply  $F_A - \sigma(v, v)$  is closed. By perturbing the equations  $\phi = 0$  by closed 2-forms  $\mathcal{Z}^2(\tilde{M}_\varphi)$  of  $\tilde{M}_\varphi$  (c.f. [A]), i.e. by changing  $\phi$  to

$$\Phi : \Omega^0(\tilde{M}_\varphi, \mathbb{V}^+) \times \mathcal{A}(\mathbb{L}) \times \mathcal{Z}^2(\tilde{M}_\varphi) \rightarrow \Omega^0(\tilde{M}_\varphi, \mathbb{V}^-) \times \mathcal{Z}^2(\tilde{M}_\varphi)$$

where  $\Phi(A, v, \delta) = (F_A - \sigma(v, v) + i\delta, \mathcal{D}_{A_0}(v) + a.v)$  we can make  $\Phi^{-1}(0)$  smooth. That is, by applying Sard's theorem to the projection  $\Phi^{-1}(0) \rightarrow \mathcal{Z}^2(\tilde{M}_\varphi)$  for generic choice of  $\delta$  we can make  $\Phi_\delta^{-1}(0)$  smooth, where  $\Phi_\delta(v, A) = \Phi(v, A, \delta)$ . The normal bundle of  $Y$  has a complex structure, so the gauge group  $\mathcal{G}(\tilde{M}_\varphi) = \text{Map}(\tilde{M}_\varphi, S^1)$  acts on the solution set  $\Phi^{-1}(0)$ , and makes quotient  $\Phi^{-1}(0)/\mathcal{G}(\tilde{M}_\varphi)$  smooth of Fredholm index  $d = b_1 - b_2 - 1$ , where  $b_i = b_i(\tilde{M}_\varphi)$  denote the betti numbers. This is because the index of the linearization of  $\Phi$  (modulo the gauge group)

$$\Omega^0(\tilde{M}_\varphi) \rightarrow \Omega^1(\tilde{M}_\varphi) \times \Omega^0(\tilde{M}_\varphi, \mathbb{V}^+) \rightarrow \mathcal{Z}^2(\tilde{M}_\varphi) \times \Omega^0(\tilde{M}_\varphi, \mathbb{V}^-)$$

is the sum of the index of the Dirac operator  $\mathcal{D}_A$  (which is zero being a self adjoint operator on an odd dimensional manifold) and index of

$$\Omega^0(\tilde{M}_\varphi) \xrightarrow{d} \Omega^1(\tilde{M}_\varphi) \xrightarrow{d} \mathcal{Z}^2(\tilde{M}_\varphi)$$

(the first map is the derivative of the gauge group action), Furthermore  $\Phi_\delta^{-1}(0)/\mathcal{G}(Y)$  is compact and orientable (the same proof as in Seiberg-Witten theory [A]).



**Remark 4.** We can localize these universal constructions to submanifolds of  $M$ . The normal bundle of any 3-manifold  $f : Y^3 \hookrightarrow M$  in a  $G_2$  manifold  $(M, \varphi)$  has a  $Spin^c(4)$  structure. Let  $\tilde{f} : Y \rightarrow \tilde{M}$  be the lifting of its Gauss map. By pulling back  $\tilde{f}$  we can transform all the bundles and their Clifford multiplication information to  $Y$ . That is, we get a pair of  $\mathbb{C}^2$  bundles  $V^\pm = \tilde{f}^*(\mathbb{V}^\pm)$  and a line bundle  $L \rightarrow Y$  describing a  $Spin^c(4)$  structure on  $\nu(Y) = \tilde{f}^*(\mathbb{V})$  with  $\nu(Y)_\mathbb{C} = V^+ \otimes V^-$ . Furthermore if  $Y$  is associative we also have  $T^*(Y) = \tilde{f}^*ad(\mathbb{V}^+)$ . Also any  $A \in \mathcal{A}(L)$  with the help of the background connection  $A_0$  gives connections on  $\mathbb{V}^\pm$ . We can then proceed as in Section 8 to do Seiberg-Witten theory on  $Y$ , and associate an invariant to  $Y$ .

In order to get a counting invariant for  $M$  we need to have positive dimensional Seiberg-Witten moduli space  $\mathcal{M}_d(M)$ . Unfortunately  $d = b_1 - b_2 - 1$  is often negative (for example any  $G_2$  manifold, whose holonomy group is the whole  $G_2$ , necessarily has  $b_1=0$  [J]). But we can use the Seiberg-Witten invariants for families to increase the dimension  $d$ . For example, we can vary the metric of  $(M, \varphi)$ . Recall that  $G_2$  metrics  $\mathcal{M}et(M)$  on  $M$  are parametrized by an open subset of 3-forms  $\Omega_+^3(M, \mathbb{R}) \subset \Omega^3(M, \mathbb{R})$ , and the harmonic version  $H_+^3(M, \mathbb{R}) \subset H^3(M, \mathbb{R})$  for the integrable ones. So at least we can increase the dimension of  $\mathcal{M}_d(M)$  to  $d = b_3 + b_1 - b_2 - 1$ . We can then proceed as in [Li-Liu], i.e. by taking a closed submanifold  $B \subset \mathcal{M}$  and doing the families of Seiberg-Witten over  $B$ , then pairing  $\langle [\mathcal{M}_d], Z \rangle$  where  $Z$  is a  $d$ -dimensional cohomology class of the moduli space  $\mathcal{B} = [\Omega^0(\tilde{M}_\varphi, \mathbb{V}^+) \times \mathcal{A}(\tilde{L})] / \mathcal{G}(\tilde{M}_\varphi)$ . Note that when  $M$  is simply connected then  $\mathcal{B} = \mathbb{C}\mathbb{P}^\infty$  (for irreducibles).

## REFERENCES

- [A] S. Akbulut, *Lectures on Seiberg-Witten Invariants*, Turkish Jour. of Math 20 (1996) 1329-1355.
- [AS] S. Akbulut and S. Salur, *Calibrated manifolds and gauge theory*, GT/0402368.
- [B1] R.L. Bryant, *Metrics with exceptional holonomy*, Ann of Math 126 (1987) 525-576.
- [B2] R.L. Bryant, *Some remarks on  $G_2$ -structures*, DG/0305124 v3
- [B3] R.L. Bryant *Manifolds with  $G_2$ -holonomy* (Duke Lectures).
- [BSh] R. L. Bryant and E. Sharpe, *D-Branes and  $Spin^c$  structures* hep-th/9812084
- [BSa] R.L. Bryant, R.L. and M.S. Salamon, *On the construction of some complete metrics with exceptional holonomy*, Duke Mathematical Journal, vol. 58, no 3 (1989), 829-850.
- [C] Weimin Chen, *The Seiberg-Witten theory of homology spheres*, arXiv: dg-ga/9703009
- [HL] F.R. Harvey, and H.B. Lawson, *Calibrated geometries*, Acta. Math. 148 (1982), 47-157.
- [J] D.D. Joyce, *Compact Manifolds with Special Holonomy*, OUP, Oxford, 2000.
- [Li-Liu] Tian-Jun Li and Ai-Ko Liu, *Family of Seiberg-Witten invariants and wall crossing formulas*, GT/0107211.
- [LM] H.B. Lawson and M.L. Michelson, *Spin geometry*, Princeton University Press (1989)
- [Ma] M. Marcolli, *Seiberg-Witten Floer homology and Heegaard splittings*, arXiv: dg-ga/9601011
- [M] R.C. McLean, *Deformations of calibrated submanifolds*, Comm. Anal. Geom. 6 (1998), 705-747.
- [Mc] B. McKay,  *$G_2$  Manifolds of cohomogeneity two*, DG/0311441

- [W] Bai-Liang Wang, *Seiberg-Witten-Floer homology for homology 3-spheres*, arXiv: dg-ga/9602003

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, MI, 48824  
*E-mail address:* `akbulut@math.msu.edu`

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, IL, 60208  
*E-mail address:* `salur@math.northwestern.edu`