

Exotic smooth structures on $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P^2}$

JONGIL PARK
ANDRÁS I. STIPSICZ
ZOLTÁN SZABÓ

*Department of Mathematical Sciences
Seoul National University
San 56-1, Shinlim-dong, Gwanak-gu
Seoul 151-747, Korea*

*Rényi Institute of Mathematics
Hungarian Academy of Sciences
H-1053 Budapest*

*Reáltanoda utca 13–15, Hungary and
Institute for Advanced Study, Princeton, NJ
and*

*Department of Mathematics
Princeton University,
Princeton, NJ, 08540*

Email: jipark@math.snu.ac.kr stipsicz@renyi.hu and szabo@math.princeton.edu

Abstract Motivated by a construction of Fintushel and Stern, we show that the topological 4–manifold $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P^2}$ supports infinitely many distinct smooth structures.

AMS Classification 53D05, 14J26; 57R55, 57R57

Keywords exotic smooth 4–manifolds, Seiberg–Witten invariants, rational blow–down, rational surfaces

1 Introduction

It is a basic problem in 4–dimensional topology to find exotic smooth structures on rational surfaces. The first such structures were found by Donaldson [4]; these examples were homeomorphic to $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P^2}$. While in this homeomorphism type many exotic examples were constructed [9, 6, 18], the cases of $\mathbb{C}P^2 \# k\overline{\mathbb{C}P^2}$ with $k < 9$ were more elusive. The Barlow surface [1] provided the first exotic structure on $\mathbb{C}P^2 \# 8\overline{\mathbb{C}P^2}$, see [13]. More recently, an exotic smooth structure on $\mathbb{C}P^2 \# 7\overline{\mathbb{C}P^2}$ has been constructed [15]. After this example many new exotic 4–manifolds with small Euler characteristic have been found. In [16] symplectic 4–manifolds homeomorphic but not diffeomorphic to $\mathbb{C}P^2 \# 6\overline{\mathbb{C}P^2}$ were constructed, implying the existence of an exotic smooth struc-

ture on $\mathbb{C}\mathbb{P}^2\#6\overline{\mathbb{C}\mathbb{P}^2}$. In a beautiful recent paper [7] Fintushel and Stern showed the existence of infinitely many distinct smooth structures on $\mathbb{C}\mathbb{P}^2\#k\overline{\mathbb{C}\mathbb{P}^2}$ with $k = 6, 7, 8$. Combining their technique of knot surgery in a double node neighborhood with a particular form of generalized rational blow-down, in this note we prove

Theorem 1.1 *There exist infinitely many pairwise nondiffeomorphic 4-manifolds all homeomorphic to $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$.*

In Section 2 various constructions of 4-manifolds homeomorphic to $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$ are described. In Section 3 we use Seiberg–Witten theory to show that many of these examples are mutually nondiffeomorphic, leading us to the proof of Theorem 1.1.

Acknowledgement: We would like to thank Ron Fintushel and Ron Stern for sending us an early version of [7] which provided an essential ingredient to the construction given in this paper. We also thank Mustafa Korkmaz for useful e-mail exchange. JP was supported by the Korea Research Foundation Grant (KRF-2004-013-C00002), AS by OTKA T34885 and ZSz by NSF grant number DMS 0107792.

2 The constructions

We construct our examples using knot surgery (in a double node neighborhood, as in [7]) when applied to particular elliptic fibrations. The special properties of the chosen elliptic fibration allow us to find a configuration in the result of the knot surgery such that after blowing it down we arrive to a 4-manifold homeomorphic, but not diffeomorphic to $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$. By using a suitable infinite set of knots (the ones already encountered in [7], cf. also [6, 18]), we get an infinite family of 4-manifolds all homeomorphic to $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$.

2.1 Elliptic fibrations

Singular fibers of elliptic fibrations have been classified [12] (cf. also [11]). In this note we will consider fibrations containing only singular fibers of type I_n ($n \geq 1$). Recall that the singular fiber I_1 (also known as the fishtail fiber) is an immersed 2-sphere with one positive double point, and it is created from a regular torus fiber by collapsing a homologically essential simple closed curve (the *vanishing cycle* of the singular fiber). The I_n -fiber ($n \geq 2$) is a collection

of n 2-spheres of self-intersection (-2) , intersecting each other in a circular pattern, see [11, page 35]. An elliptic fibration with singular fibers only of type I_n are *Lefschetz fibrations* in the sense of [10, Chapter 8]. The only subtlety we have to keep in mind is that here we allow a singular fiber to contain more than one singular points as well.

Lefschetz fibrations can be conveniently described by the monodromy factorization induced by the singular fibers of the fibration, that is, by a word involving right-handed Dehn twists which is equal to 1 in the mapping class group of the regular fiber. The mapping class group Γ_1 of the 2-torus T^2 can be presented as

$$\Gamma_1 = \{a, b \mid aba = bab, (ab)^6 = 1\},$$

where $a, b \in \Gamma_1$ denote the right-handed Dehn twists along the two standard simple closed curves A, B in T^2 intersecting each other transversally in a unique point. This group can be identified with $\mathrm{SL}(2; \mathbb{Z})$ by mapping a to $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and b to $\begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$. For example, the standard elliptic fibration we get by blowing up nine base points of a generic elliptic pencil in \mathbb{CP}^2 results the monodromy factorization $(ab)^6$. Using the braid relation $aba = bab$ it can be shown that $(a^3b)^3$ also defines an elliptic fibration on $\mathbb{CP}^2 \# 9\mathbb{CP}^2$. Furthermore, it is easy to see that for any expression $x \in \Gamma_1$ the mapping class $a^x = xax^{-1}$ can be identified with the right-handed Dehn twist along the image of a under a map giving x . Note, for example, that the braid relation implies that $b = a^{ab}$.

The monodromy of a fishtail fiber can be shown to be equal to the right-handed Dehn twist along the vanishing cycle corresponding to the given singular fiber. An I_n -fiber can be created by collapsing n parallel (homologically essential) simple closed curves, therefore the monodromy of such a fiber is equal to the n th power of the right-handed Dehn twist along one of the parallel curves.

In our constructions we will need the existence of a section, which can be also read off from the monodromy factorization. In general, a Lefschetz fibration admits a section if the monodromy factorization induced by it can be lifted from the mapping class group of its generic fiber to the mapping class group of the fiber with one marked point. In the case of a genus-1 Lefschetz fibration, however, the forgetful map $f: \Gamma_1^1 \rightarrow \Gamma_1$ mapping from the mapping class group Γ_1^1 of T^2 with one marked point to Γ_1 is an isomorphism, implying in particular

Lemma 2.1 *Any genus-1 Lefschetz fibration over S^2 admits a section. \square*

2.2 The definition of the 4-manifold X_n

Our first construction of exotic 4-manifolds relies on the following existence result. (For a schematic picture of the fibration see Figure 1.)




Figure 1: The schematic diagram of the fibration with an I_7 fiber

Proposition 2.2 *There exists an elliptic fibration $\mathbb{C}\mathbb{P}^2 \# 9\overline{\mathbb{C}\mathbb{P}^2} \rightarrow \mathbb{C}\mathbb{P}^1$ with five fishtail fibers, an I_7 -fiber and a section. Furthermore, we can assume that two of the five fishtail fibers have isotopic vanishing cycles.*

Proof We will give the fibration by finding an appropriate factorization of 1 in the mapping class group Γ_1 of the torus. Start with the fibration on $\mathbb{C}\mathbb{P}^2 \# 9\overline{\mathbb{C}\mathbb{P}^2}$ defined by the factorization

$$(a^3b)^3$$

of $1 \in \Gamma_1$. Notice that

$$(a^3b)^3 = a^7(a^{-1}(a^{-3}ba^3)a)(a^{-1}ba)a^2b = a^7b^{a^{-4}}b^{a^{-1}}a^2b.$$

Since a^7 is the monodromy of an I_7 -fiber, its existence in the above fibration is verified. The term a^2 gives rise to two fishtail fibers with isotopic vanishing cycles in the complement of the I_7 -fiber. Finally, Lemma 2.1 shows the existence of a section in the fibration. \square

Suppose now that $p > q > 0$ are relatively prime integers. Let us define the 4-manifold $C_{p,q}$ as the result of the linear plumbing with weights specified by the continued fraction coefficients of $-\frac{p^2}{pq-1}$. It is known [2] (cf. also [14, 16, 17]) that the boundary $\partial C_{p,q} = L(p^2, pq-1)$ is a lens space which bounds a rational ball $B_{p,q}$. The replacement of an embedded copy of $C_{p,q} \subset X$ with $B_{p,q}$ is called the (*generalized*) *rational blow-down* of X along $C_{p,q}$. This operation was introduced and successfully applied by Fintushel and Stern [5] in the case of $q = 1$ and studied in [14, 17] in the above generality.

Now we are ready to turn to the construction of the 4-manifolds homeomorphic but not diffeomorphic to $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P^2}$. Let K_n denote the n -twist knot as it is depicted in [7]. Let F_3, F_4 of Figure 1 denote the fishtail fibers with isotopic vanishing cycles. Following the convention of [7] we denote the result of the knot surgery in a double node neighborhood containing F_3, F_4 and with knot K_n by Y_n . Fintushel and Stern [7] proves the existence of a “pseudo-section” $S \subset Y_n$ which is an immersed sphere with one positive double point, self-intersection -1 , and which transversally intersects F_1, F_2 and one of the spheres in the I_7 -fiber. Let us blow up Y_n in the double point of the pseudo-section, and in the double points of the fishtail fibers F_1 and F_2 . After smoothing the intersections P_1, P_2 , we get a sphere of self-intersection -9 intersecting the I_7 -fiber transversally at one point. Now we apply eight infinitely close blow-ups at the point Q as it is shown by Figure 2. (The small circles indicate the




Figure 2: Infinitely close blow-ups at Q

locations of the blow-ups.) This construction results a chain of 2-spheres, with a neighborhood diffeomorphic to the 4-manifold C we get by plumbing along

a linear chain with weights

$$(-9, -10, -2, -2, -2, -2, -2, -3, -2, -2, -2, -2, -2, -2)$$

in the eleven-fold blow-up of Y_n . Simple computation identifies C with $C_{71,8}$. Define X_n as the (generalized) rational blow-down of Y_n along C , that is,

$$X_n = (Y_n \# 11\overline{\mathbb{C}\mathbb{P}^2} - \text{int } C) \cup B_{71,8}.$$

Theorem 2.3 X_n is homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$.

Proof The 4-manifold Y_n has trivial fundamental group, since the fibration admits a section and two different vanishing cycles in the complement of the double node neighborhood. Simple connectivity of X_n follows from the fact that the complement of C in $Y_n \# 11\overline{\mathbb{C}\mathbb{P}^2}$ is simply connected, since the generator of $\pi_1(\partial C)$ can be contracted along the fishtail fiber F_0 present in the fibration but not used in constructing the configuration C . Now simple Euler characteristic and signature computation together with Freedman's Theorem on the classification of topological 4-manifolds [8] imply the result. \square

2.3 Further constructions

Many similar constructions can be carried out using different elliptic fibrations or different sets of knots. Below we outline constructions relying on various types of elliptic fibrations.

2.3.1 A configuration using the I_7 -fiber

A similar argument provides an embedding of $C_{212,55}$ into $Y_n \# 12\overline{\mathbb{C}\mathbb{P}^2}$ by smoothing only at P_2 and keeping the transverse intersection P_1 . In this case one further blow-up of a (-2) -sphere is necessary, leading to the chain

$$(-4, -7, -10, -2, -2, -2, -2, -2, -3, -2, -2, -2, -2, -3, -2, -2)$$

in $Y_n \# 12\overline{\mathbb{C}\mathbb{P}^2}$. Blowing this configuration down we get a sequence of 4-manifolds with the same properties as X_n . (The hemisphere originated from the exceptional sphere of the last blow-up can be used to show that the resulting configuration of spheres in the twelve-fold blow-up of Y_n has simply connected complement.)

2.3.2 Configurations using the I_8 -fiber

Many other examples can be given using the I_8 -fiber. To see the existence of the required fibration, we need a result similar to Proposition 2.2.

Proposition 2.4 *There exists an elliptic fibration $\mathbb{C}\mathbb{P}^2 \# 9\overline{\mathbb{C}\mathbb{P}^2} \rightarrow \mathbb{C}\mathbb{P}^1$ with four fishtail fibers, an I_8 -fiber and a section. Furthermore, we can assume that two of the four fishtail fibers have isotopic vanishing cycles.*

Proof Using the braid relation it is fairly easy to see that the expression

$$a^3ba^2b^2a^2ba$$

is equal to 1 in Γ_1^1 , hence defines an elliptic fibration with a section. Since it can be written as

$$a^8(b^{a^{-2}})b^2(b^{a^2}),$$

the resulting fibration can be chosen to have an I_8 -fiber and two fishtails in its complement with isotopic vanishing cycles. \square

Our further constructions rely on

- Proposition 2.5** (1) *The 4-manifold $C_{44,9}$ embeds into $Y_n \# 8\overline{\mathbb{C}\mathbb{P}^2}$;*
(2) *$C_{79,44}$ admits an embedding into $Y_n \# 9\overline{\mathbb{C}\mathbb{P}^2}$;*
(3) *$C_{89,9}$ embeds into $Y_n \# 13\overline{\mathbb{C}\mathbb{P}^2}$;*
(4) *$C_{169,89}$ can be embedded into $Y_n \# 14\overline{\mathbb{C}\mathbb{P}^2}$;*
(5) *$C_{301,62}$ admits an embedding into $Y_n \# 14\overline{\mathbb{C}\mathbb{P}^2}$; and finally*
(6) *$C_{540,301}$ is a submanifold of $Y_n \# 15\overline{\mathbb{C}\mathbb{P}^2}$.*

The complements of these configurations are simply connected.

Remark 2.6 Recall that the above 4-manifolds can be given by the linear plumbings as follows:

$$C_{44,9} = (-5, -11, -2, -2, -2, -2, -2, -2, -3, -2, -2, -2),$$

$$C_{79,44} = (-2, -5, -11, -2, -2, -2, -2, -2, -2, -3, -2, -2, -3),$$

$$C_{89,9} = (-10, -11, -2, -2, -2, -2, -2, -2, -3, -2, -2, -2, -2, -2, -2, -2),$$

$$C_{169,89} = (-2, -10, -11, -2, -2, -2, -2, -2, -2, -3, -2, -2, -2, -2, -2, -2, -3),$$

$$C_{301,62} = (-5, -7, -11, -2, -2, -2, -2, -2, -2, -3, -2, -2, -2, -2, -3, -2, -2, -2)$$

and finally

$$C_{540,301} = (-2, -5, -7, -11, -2, -2, -2, -2, -2, -2, -3, -2, -2, -2, -2, -3, -2, -2, -3).$$

Proof We use the configuration of Figure 3 to indicate the embeddings given




Figure 3: A fibration with an I_8 -fiber

above. First of all, perform the knot surgery in the neighborhood of the fishtail fibers F_3, F_4 with isotopic vanishing cycles and blow up the two double points of the remaining two fishtail fibers F_1, F_2 together with the double point of the pseudo-section. To get the first embedding, smooth the transverse intersections P_2, P_3 and apply four infinitely close blow-ups at Q , resulting the configuration

$$(-4, -11, -2, -2, -2, -2, -2, -2, -3, -2, -2, -2).$$

One further blow-up of the (-4) -sphere provides the first embedding. If we blow up this sphere as instructed by Figure 4, a final blow-up of the last (-2) -sphere in the chain gives the second embedding.

If we smooth the intersections P_1 and P_2 then eight infinitely close blow-ups at Q , together with a final blow-up on any of the former fishtail fibers F_1 or F_2 results the third embedding. Once again, the last blow-up can be performed as in Figure 4, in which case we need to blow up the other end of the chain, resulting the fourth embedding. Finally, resolving only P_2 , eight infinitely close blow-ups at Q , one further blow-up on the appropriate (-2) -sphere in the I_8 -fiber and one more on the fishtail passing through P_1 gives the fifth configuration. If this last blow-up is performed as in Figure 4, by blowing up the last (-2) -sphere of the configuration we get the last promised embedding. Since in any of the above constructions the last blow-up provides an exceptional




Figure 4: Infinitely close blow-up at the double point of the fishtail fiber

divisor transversally intersecting the first or last sphere of the configuration, the complements of the configurations are obviously simply connected. \square

Simple Euler characteristic computation and Freedman's Theorem implies that after rationally blowing down any of the configurations presented in Proposition 2.5 we get further interesting examples of 4-manifolds homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$.

3 Seiberg–Witten computations

We will prove Theorem 1.1 by computing Seiberg–Witten invariants of the 4-manifolds constructed above. We will only give details of the computation for the first construction, resulting the manifolds X_n . The argument sketched below is closely modeled on the argument encountered in [7].

It is shown in [6, 18] that Y_n has two Seiberg–Witten basic classes $\pm K$, moreover $|SW_{Y_n}(\pm K)| = n$. Furthermore, we can choose the sign of K so that it evaluates on the pseudo-section S as -1 . Consequently

$$(K - e_1 - \dots - e_{11})(u_i) = u_i \cdot u_i + 2$$

for each sphere u_i appearing in the plumbing C . Let L be the extension of $K|_{Y_n - C}$ to X_n . Using the blow-up and the rational blow-down formula together with the wall-crossing formula we get

Proposition 3.1 *The Seiberg–Witten invariant $SW_{X_n}(L)$ is an element of the set $\{\pm n, \pm n \pm 1\}$. Therefore the 4–manifold X_n with $n \geq 2$ admits a Seiberg–Witten basic class. \square*

This computation leads us to

Corollary 3.2 *There exists an exotic smooth structure on $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$.*

Proof Since the Seiberg–Witten function is a diffeomorphism invariant for manifolds with $b_2^+ = 1$ and $b_2^- \leq 9$, and by the existence of a positive scalar curvature metric we have $SW_{\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}} \equiv 0$, the corollary follows from Proposition 3.1. \square

The same computation as above actually shows

Lemma 3.3 *The Seiberg–Witten function of X_n takes its values in a subset of $\{0, \pm 1, \pm n, \pm n \pm 1\}$. \square*

Proof of Theorem 1.1 Combining Proposition 3.1 with Lemma 3.3 it follows that X_n and X_{n+3k} are not diffeomorphic once $n \geq 2$ and $k > 0$. This observation proves the existence of infinitely many distinct smooth structures on $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$. \square

References

- [1] **R. Barlow**, *A simply connected surface of general type with $p_g = 0$* , Invent. Math. **79** (1985), 293–301.
- [2] **A. Casson and J. Harer**, *Some homology lens spaces which bound rational homology balls*, Pacific J. Math. **96** (1981), 23–36.
- [3] **S. Donaldson**, *An application of gauge theory to four dimensional topology*, J. Diff. Geom. **18** (1983), 279–315.
- [4] **S. Donaldson**, *Irrationality and the h-cobordism conjecture*, J. Diff. Geom. **26** (1987), 141–168.
- [5] **R. Fintushel and R. Stern**, *Rational blowdowns of smooth 4–manifolds*, J. Diff. Geom. **46** (1997), 181–235.
- [6] **R. Fintushel and R. Stern**, *Knots, links and 4–manifolds*, Invent. Math. **134** (1998), 363–400.
- [7] **R. Fintushel and R. Stern**, *Double node neighborhoods and families of simply connected 4–manifolds with $b^+ = 1$* , arXiv:math.GT/0412126

- [8] **M. Freedman**, *The topology of four-dimensional manifolds*, J. Diff. Geom. **17** (1982), 357–453.
- [9] **R. Friedman and J. Morgan**, *On the diffeomorphism types of certain algebraic surfaces I., II.* J. Diff. Geom. **27** (1988), 297–369. and 371–398.
- [10] **R. Gompf and A. Stipsicz**, *4-manifolds and Kirby calculus*, AMS Grad. Studies in Math. **20**, 1999.
- [11] **J. Harer, A. Kas and R. Kirby**, *Handlebody decompositions of complex surfaces*, Memoirs of the AMS, vol. **62**, 1986.
- [12] **K. Kodaira**, *On compact analytic surfaces: II*, Ann. Math. **77** (1963), 563–626.
- [13] **D. Kotschick**, *On manifolds homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 8\overline{\mathbb{C}\mathbb{P}^2}$* , Invent. Math. **95** (1989), 591–600.
- [14] **J. Park**, *Seiberg–Witten invariants of generalized rational blow-downs*, Bull. Austral. Math. Soc. **56** (1997), 363–384.
- [15] **J. Park**, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , to appear in Invent. Math., arXiv:math.GT/0311395
- [16] **A. Stipsicz and Z. Szabó**, *An exotic smooth structure on $\mathbb{C}\mathbb{P}^2 \# 6\overline{\mathbb{C}\mathbb{P}^2}$* , arXiv.math:GT/0411258
- [17] **M. Symington**, *Generalized symplectic rational blowdowns*, Algebr. Geom. Topol. **1** (2001), 503–518.
- [18] **Z. Szabó**, *Exotic 4-manifolds with $b_2^+ = 1$* , Math. Res. Letters **3** (1996), 731–741.