REIDEMEISTER TORSION OF A SYMPLECTIC COMPLEX

YAŞAR SÖZEN

ABSTRACT. We consider a claim mentioned in [5] pp 187 about the relation between a symplectic chain complex with ω -compatible bases and Reidemeister Torsion of it. This is an explanation of it.

INTRODUCTION

Even though, we approach Reidemeister torsion as a linear algebraic object, it actually is a combinatorial invariant for the space of representations of a compact surface into a fixed gauge group [5],[2].

More precisely, let S be a compact surface with genus at least 2 and without boundary, G be a gauge group with its (semi-simple) Lie algebra \mathfrak{g} . Then, for a representation $\rho : \pi_1(S) \to G$, we can associate the corresponding adjoint bundle

 $\begin{pmatrix} \widetilde{S} \times_{\rho} \mathfrak{g} \\ \downarrow \\ S \end{pmatrix} \text{ over } S, \text{ i.e. } \widetilde{S} \times_{\rho} \mathfrak{g} = \widetilde{S} \times \mathfrak{g} / \sim, \text{ where } (x,t) \text{ is identified with all }$

the elements in its orbit i.e. $(\gamma \bullet x, \gamma \bullet t)$ for all $\gamma \in \pi_1(S)$, and where in the first component the element $\gamma \in \pi_1(S)$ of the fundamental group of S acts as a deck transformation, and in the second component by conjugation by $\rho(\gamma)$.

Suppose K is a cell-decomposition of S so that the adjoint bundle $\widetilde{S} \times_{\rho} \mathfrak{g}$ on S is trivial over each cell. Let \widetilde{K} be the lift of K to the universal covering \widetilde{S} of S. With the action of $\pi_1(S)$ on \widetilde{S} as deck transformation, $C_*(\widetilde{K}; \mathbb{Z})$ can be considered a left $-\mathbb{Z}[\pi_1(S)]$ module and with the action of $\pi_1(S)$ on \mathfrak{g} by adjoint representation, \mathfrak{g} can be considered as a left $-\mathbb{Z}[\pi_1(S)]$ module, where $\mathbb{Z}[\pi_1(S)]$ is the integral group

$$\operatorname{ring}\left\{\sum_{i=1}^{p} m_{i}\gamma_{i} ; m_{i} \in \mathbb{Z}, \ \gamma_{i} \in \pi_{1}(S), \ p \in \mathbb{N}\right\}.$$
More explicitly, if $\sum_{i=1}^{p} m_{i}\gamma_{i}$ is in $\mathbb{Z}[\pi_{1}(S)], \ t$ is in \mathfrak{g} , and $\sum_{j=1}^{q} n_{j} \ \sigma_{j} \in C_{*}(\widetilde{S};\mathbb{Z}),$

then
$$\left(\sum_{i=1}^{p} m_i \gamma_i\right) \bullet \left(\sum_{j=1}^{q} n_j \sigma_j\right) \stackrel{\text{defn}}{=} \sum_{i,j} n_j m_i \ (\gamma_i \bullet \sigma_j), \text{ where } \gamma_i \text{ acts on } \sigma_j \subset \widetilde{S}$$

by deck transformation, and $\left(\sum_{j=1}^{q} m_j \gamma_j\right) \bullet t \stackrel{\text{defn}}{=} \sum_{j=1}^{q} m_j \ (\gamma_j \bullet t)$, where $\gamma_j \bullet t =$

Date: December 4, 2004.

¹⁹⁹¹ Mathematics Subject Classification. 15A90; 57M99.

Key words and phrases. Reidemeister torsion, Symplectic Complex.

 $Ad_{\rho(\gamma_j)}(t) = \rho(\gamma_j)t\rho(\gamma_j)^{-1}.$

To talk about the tensor product $C_*(\widetilde{K};\mathbb{Z}) \otimes \mathfrak{g}$, we should consider the left $\mathbb{Z}[\pi_1(S)]$ -module $C_*(\widetilde{K};\mathbb{Z})$ as a right $\mathbb{Z}[\pi_1(S)]$ -module as $\sigma \bullet \gamma \stackrel{defn}{=} \gamma^{-1} \bullet \sigma$, where the action of γ^{-1} is as a deck-transformation. Note that the relation $\sigma \bullet \gamma \otimes t = \sigma \otimes \gamma \bullet t$ becomes $\gamma^{-1} \bullet \sigma \otimes t = \sigma \otimes \gamma \bullet t$, equivalently $\sigma' \otimes t = \gamma \bullet \sigma' \otimes \gamma \bullet t$, where σ' is $\gamma^{-1} \bullet \sigma$. We may conclude that tensoring with $\mathbb{Z}[\pi_1(S)]$ has the same effect as factoring with $\pi_1(S)$. Thus, $C_*(K; Ad_\rho) \stackrel{\text{defn}}{=} C_*(\widetilde{K}; \mathbb{Z}) \otimes_\rho \mathfrak{g}$ is defined as the quotient $C_*(\widetilde{K}; \mathbb{Z}) \otimes \mathfrak{g} / \sim$, where the elements of the orbit $\{\gamma \bullet \sigma \otimes \gamma \bullet t; \text{ for all } \gamma \in \pi_1(S)\}$ of $\sigma \otimes t$ are identified.

In this way, we obtain the following complex:

$$0 \to C_2(K; Ad_\rho) \stackrel{\partial_2 \otimes \mathrm{id}}{\longrightarrow} C_1(K; Ad_\rho) \stackrel{\partial_1 \otimes \mathrm{id}}{\longrightarrow} C_0(K; Ad_\rho) \to 0,$$

where ∂_i is the usual boundary operator. For this complex, we can associate the homologies $H_*(K; Ad_{\rho})$. Similarly, the twisted cochains $C^*(K; Ad_{\rho})$ will result the cohomologies $H^*(K; Ad_{\rho})$, where $C^*(K; Ad_{\rho}) \stackrel{\text{defn}}{=} \text{Hom}_{\mathbb{Z}[\pi_1(S)]}(C_*(\widetilde{K}; \mathbb{Z}), \mathfrak{g})$ is the set of $\mathbb{Z}[\pi_1(S)]$ -module homomorphisms from $C_*(\widetilde{K}; \mathbb{Z})$ into \mathfrak{g} . For more information, we refer [2],[3],[5].

If $\rho, \rho' : \pi_1(S) \to G$ are conjugate, i.e. $\rho'(\cdot) = A\rho(\cdot)A^{-1}$ for some $A \in G$, then $C_*(K; Ad_{\rho})$ and $C_*(K; Ad_{\rho'})$ are isomorphic. Similarly, the twisted cochains $C^*(K; Ad_{\rho})$ and $C^*(K; Ad_{\rho'})$ are isomorphic. Moreover, the homologies $H_*(K; Ad_{\rho})$ are independent of the cell-decomposition. For details, see [3],[5],[2].

If $\{e_1^i, \dots, e_{m_i}^i\}$ is a basis for the $C_i(K; \mathbb{Z})$, then $c_i := \{\widetilde{e_1^i}, \dots, \widetilde{e_{m_i}^i}\}$ will be a $\mathbb{Z}[\pi_1(S)]$ -basis for $C_i(\widetilde{K}; \mathbb{Z})$, where $\widetilde{e_j^i}$ is a lift of e_j^i . If we choose a basis \mathcal{A} of \mathfrak{g} , then $c_i \otimes_{\rho} \mathcal{A}$ will be a \mathbb{C} -basis for $C_i(K; Ad_{\rho})$, called a *geometric* basis for $C_i(K; Ad_{\rho})$. Recall that $C_i(K; Ad_{\rho}) = C_i(\widetilde{K}; \mathbb{Z}) \otimes_{\rho} \mathfrak{g}$, is defined as the quotient $C_i(\widetilde{K}; \mathbb{Z}) \otimes_{\mathfrak{g}} \mathfrak{g} / \sim$, where we identify the orbit $\{\gamma \bullet \sigma \otimes \gamma \bullet t; \gamma \in \pi_1(S)\}$ of $\sigma \otimes t$, and where the action of the fundamental group in the first slot by deck-transformations, and in the second slot by the conjugation with $\rho(\cdot)$.

In this set-up, one can also define $\operatorname{Tor}(C_*(K; Ad_{\rho}), \{c_i \otimes_{\rho} \mathcal{A}\}_{i=0}^2, \{\mathfrak{h}_i\}_{i=0}^2)$ the *Reidemiester torsion* of the triple K, Ad_{ρ} , and $\{\mathfrak{h}_i\}_{i=0}^2$, where \mathfrak{h}_i is a \mathbb{C} -basis for $H_i(K; Ad_{\rho})$. Moreover, one can easily prove that this definition does not depend on the lifts \tilde{e}_j^i , conjugacy class of ρ , and cell-decomposition K of the surface S. Details can be found in [3],[2],[5].

Let K, K' be dual cell-decompositions of S so that $\sigma \in K \sigma' \in K'$ meet at most once and moreover the diameter of each cell has diameter less than, say, half of the injectivity radius of S. If we denote $C_* = C_*(K; Ad_\rho), C'_* = C_*(K'; Ad_\rho)$, then by the invariance of torsion under subdivision, $\operatorname{Tor}(C_*) = \operatorname{Tor}(C'_*)$. Let D_* denote the complex $C_* \oplus C'_*$. Then, easily we have the short-exact sequence

$$0 \to C_* \to D_* = C_* \oplus C'_* \to C'_* \to 0.$$

The complex $D_* = C_* \oplus C'_*$ can also be considered as a symplectic complex. Moreover, in the case of irreducible representation $\rho : \pi_1(S) \to G$, torsion $\operatorname{Tor}(C_*)$ gives a two-form on $H^1(S; Ad_{\rho})$. See [5],[3].

In this article, we will consider Reidemeister torsion as a linear algebraic object and try to rephrase a statement mentioned in [5].

The main result of the article is as stated in [5] pp 187 "the torsion of a symplectic complex (C_*, ω) computed using a compatible set of measures is "trivial" in the sense that"

Theorem 0.0.1. For a general symplectic complex C_* , if \mathfrak{c}_p , \mathfrak{h}_p are bases for C_p , H_p , respectively, then

$$Tor(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n) = \left(\prod_{p=0}^{\frac{n}{2}-1} (\det[\omega_{p,n-p}])^{(-1)^p}\right) \cdot \left(\sqrt{\det[\omega_{\frac{n}{2},\frac{n}{2}}]}\right)^{(-1)^{\frac{n}{2}}},$$

where det $[\omega_{p,n-p}]$ is the determinant of the matrix of the non-degenerate pairing $[\omega_{p,n-p}]: H_p(C) \times H_{n-p}(C) \to \mathbb{R}$ in bases $\mathfrak{h}_p, \mathfrak{h}_{n-p}$.

For topological application of this, we refer [3], [5].

The plan of paper is as follows. In section §1, we will give the definition of Reidemeister torsion for a general complex C_* and recall some properties. See [1],[2] for more information. In section §2, we will explain torsion using Witten's notation [5]. Finally, symplectic complex will be explained in section §3 and also the proof of main result Theorem 0.0.1.

1. Reidemeister Torsion of a general Chain Complex

Let $C_* = (C_n \xrightarrow{\partial_n} C_{n-1} \longrightarrow \cdots \longrightarrow C_1 \xrightarrow{\partial_1} C_0 \longrightarrow 0)$ be a chain complex of a finite dimensional vector spaces over \mathbb{R} or \mathbb{C} . Let $H_p = Z_p/B_p$ denote the homologies of the complex, where $B_p = \text{Im}\{\partial_{p+1} : C_{p+1} \to C_p\}, Z_p = \text{ker}\{\partial_p : C_p \to C_{p-1}\}$, respectively.

If we start with bases $\mathfrak{b}_p = \{b_p^1, \cdots, b_p^{m_p}\}$ for B_p , and $\mathfrak{h}_p = \{h_p^1, \cdots, h_p^{n_p}\}$ for H_p , a new basis for C_p can be obtained by considering the following short-exact sequences:

$$(1.0.1) 0 \to Z_p \hookrightarrow C_p \to B_{p-1} \to 0$$

$$(1.0.2) 0 \to B_p \hookrightarrow Z_p \to H_p \to 0$$

where the first row is a result of 1^{st} -Isomorphism Theorem and the second follows simply from the definition of H_p .

Starting with 1.0.2 and a section $\overline{s_p} : H_p \to Z_p$, then Z_p will have a basis $\mathfrak{b}_p \oplus \overline{s_p}(\mathfrak{h}_p)$. Using 1.0.1 and a section $s_p : B_{p-1} \to C_p$, C_p will have a basis $\mathfrak{b}_p \oplus \overline{s_p}(\mathfrak{h}_p) \oplus s_p(\mathfrak{b}_{p-1})$.

If V is a vector space with bases \mathfrak{e} and \mathfrak{f} , then we will denote $[\mathfrak{f}, \mathfrak{e}]$ for the determinant of the change-base-matrix $T_{\mathfrak{e}}^{\mathfrak{f}}$ from \mathfrak{e} to \mathfrak{f} .

Definition 1.0.2. For $p = 1, \dots, n$, let $\mathfrak{c}_p, \mathfrak{b}_p$, and \mathfrak{h}_p be bases for C_p, B_p and H_p , respectively. Tor $(C_*, {\mathfrak{c}_p}_{p=0}^n, {\mathfrak{h}_p}_{p=0}^n) = \prod_{p=0}^n [\mathfrak{b}_p \oplus \overline{s_p}(\mathfrak{h}_p) \oplus s_p(\mathfrak{b}_{p-1}), \mathfrak{c}_p]^{(-1)^{(p+1)}}$ is called the *torsion of the complex* C_* with respect to bases ${\mathfrak{c}_p}_{p=0}^n, {\mathfrak{h}_p}_{p=0}^n,$

Milnor [1] proved that torsion does not depend on neither the bases \mathfrak{b}_p , nor the sections $s_p, \overline{s_p}$. In other words, it is well-defined.

Remark 1.0.3. If we choose another bases $\mathfrak{c}'_p, \mathfrak{h}'_p$ respectively for C_p and H_p , then an easy computation shows that

$$\operatorname{Tor}(C_*, \{\mathfrak{c}'_p\}_{p=0}^n, \{\mathfrak{h}'_p\}_{p=0}^n) = \prod_{p=0}^n \left(\frac{[\mathfrak{c}'_p, \mathfrak{c}_p]}{[\mathfrak{h}'_p, \mathfrak{h}_p]}\right)^{(-1)^p} \cdot \operatorname{Tor}(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n).$$

This follows easily from the fact that torsion is independent of \mathfrak{b}_p and sections $s_p, \overline{s_p}$. For example, if $[\mathfrak{c}'_p, \mathfrak{c}_p] = 1$, and $[\mathfrak{h}'_p, \mathfrak{h}_p] = 1$, then they produce the same torsion.

If we have a short-exact sequence of chain complexes $0 \to A_* \xrightarrow{i} B_* \xrightarrow{\pi} D_* \to 0$, then we also have a long-exact sequence of vector space C_*

$$\cdots \to H_p(A) \stackrel{\iota_*}{\to} H_p(B) \stackrel{\pi_*}{\to} H_p(D) \stackrel{\Delta}{\to} H_{p-1}(A) \to \cdots$$

i.e. an acyclic (or exact) complex C_* of length 3n + 2 with $C_{3p} = H_p(D_*)$, $C_{3p+1} = H_p(A_*)$ and $C_{3p+2} = H_p(B_*)$. In particular, the bases $\mathfrak{h}_p(D_*)$, $\mathfrak{h}_p(A_*)$, and $\mathfrak{h}_p(B_*)$ will serve as bases for C_{3p}, C_{3p+1} , and C_{3p+2} , respectively.

Theorem 1.0.4. (Milnor [1]) Using the above setup, let $\mathbf{c}_p^A, \mathbf{c}_p^B, \mathbf{c}_p^D$ be bases for A_p, B_p, D_p , respectively, and let $\mathfrak{h}_p^A, \mathfrak{h}_p^B, \mathfrak{h}_p^D$ be bases for the corresponding homologies $H_p(A), H_p(B)$, and $H_p(D)$. If, moreover, the bases $\mathbf{c}_p^A, \mathbf{c}_p^B, \mathbf{c}_p^D$ are compatible in the sense that $[\mathbf{c}_p^B, \mathbf{c}_p^A \oplus \widetilde{\mathbf{c}_p}^D] = \pm 1$ where $\pi\left(\widetilde{\mathbf{c}_p^D}\right) = \mathbf{c}_p^D$, then $Tor(B_*, \{\mathbf{c}_p^B\}_{p=0}^n, \{\mathbf{b}_p^B\}_{p=0}^n) = Tor(A_*, \{\mathbf{c}_p^A\}_{p=0}^n, \{\mathbf{b}_p^A\}_{p=0}^n) \cdot Tor(D_*, \{\mathbf{c}_p^D\}_{p=0}^n, \{\mathbf{b}_p^D\}_{p=0}^n) \cdot Tor(C_*, \{\mathbf{c}_{3p}\}_{p=0}^{3n+2}, \{0\}_{p=0}^{3n+2})$

2. Reidemeister Torsion Using Witten's notations

Let V be a vector space of dimension k over \mathbb{R} . Let $\det(V)$ denote the top exterior power $\bigwedge^k V$ of V. A *measure* on V is a non-zero functional $\alpha : \det(V) \to \mathbb{R}$ on $\det(V)$, i.e. $\alpha \in \det(V)^{-1}$, -1 denotes the dual space.

Recall that the isomorphism between $\det(V)^{-1}$ and $\det(V^*)$ is given by the pairing $\langle \cdot, \cdot \rangle : \det(V^*) \times \det(V) \to \mathbb{R}$, defined by

$$\langle f_1^* \wedge \cdots \wedge f_k^*, e_1 \wedge \cdots \wedge e_k \rangle = \det \left[f_i^*(e_j) \right],$$

i.e. $[\mathfrak{f},\mathfrak{e}]$ the determinant of the change-base-matrix from basis $\mathfrak{e} = \{e_1, \dots, e_k\}$ to $\mathfrak{f} = \{f_1, \dots, f_k\}$, where f_i^* is the dual element corresponding to f_i , namely,

$$f_i^*(f_j) = \delta_{ij}$$
. Below $(v_1 \wedge \cdots \wedge v_k)^{-1}$ will denote $(v_1)^* \wedge \cdots \wedge (v_k)^*$

Note also that $\langle f_1^* \wedge \cdots \wedge f_k^*$, $e_1 \wedge \cdots e_k \rangle = \langle e_1^* \wedge \cdots \wedge e_k^*$, $f_1 \wedge \cdots f_k \rangle^{-1}$, i.e. $[\mathfrak{f}, \mathfrak{e}] = [\mathfrak{e}, \mathfrak{f}]^{-1}$. So, using the pairing, $[\mathfrak{f}, \bullet]$ can be considered a linear functional on det(V) and $[\bullet, \mathfrak{e}]$ can be considered a linear functional on det (V^*) .

Let $C_*: 0 \to C_n \xrightarrow{\partial_n} C_{n-1} \to \cdots \to C_1 \xrightarrow{\partial_1} C_0 \to 0$ be a chain complex of finite dimensional vector spaces with volumes $\alpha_p \in \det(V)^{-1}$, i.e. $\alpha_p = (c_1^p)^* \wedge \cdots \wedge (c_{m_p}^p)^*$ for some basis $\{c_1^p, \cdots, c_{m_p}^p\}$ for C_p . If, moreover, we assume that C_* is exact (or acyclic), then $H_p(C_*) = 0$ for all p. In particular, we have the short exact sequence

$$0 \to \underbrace{\operatorname{Im}\{\partial_{p+1}: C_{p+1} \to C_p\}}_{B_p} \xrightarrow{i_p} C_p \xrightarrow{\partial_p} \underbrace{\operatorname{Im}\{\partial_p: C_p \to C_{p-1}\}}_{B_{p-1}} \to 0.$$

Let $\{b_1^p, \dots, b_{k_p}^p\}$, $\{b_1^{p-1}, \dots, b_{k_{p-1}}^{p-1}\}$ be bases for B_p, B_{p-1} , respectively. Then, $\{b_1^p, \dots, b_{k_p}^p, \widetilde{b_1^{p-1}}, \dots, \widetilde{b_{k_{p-1}}^{p-1}}\}$ is a basis for C_p , where $\partial_p(\widetilde{b_{p-1}^i}) = b_{p-1}^i$ and thus $b_1^p \wedge \dots \wedge b_{k_p}^p \wedge \widetilde{b_1^{p-1}} \wedge \dots \wedge \widetilde{b_{k_{p-1}}^{p-1}}$ is a basis for $\det(C_p)$.

If u denotes $\bigotimes_{p=0}^{n} (b_1^p \wedge \dots \wedge b_{k_p}^p \wedge \widetilde{b_1^{p-1}} \wedge \dots \wedge \widetilde{b_{k_{p-1}}^{p-1}})^{(-1)^p}$, then u is an element of

 $\bigotimes_{p=0}^{p=0} (\det(C_p))^{(-1)^p}$, where the exponent (-1) denotes the dual of the vector space. Then, E. Witten describes the torsion as:

$$\operatorname{Tor}(F_{*}) = \langle u, \bigotimes_{p=0}^{n} \alpha_{p}^{(-1)^{p}} \rangle$$

= $\prod_{p=0}^{n} \langle b_{1}^{p} \wedge \dots \wedge b_{k_{p}}^{p} \wedge \widetilde{b_{1}^{p-1}} \wedge \dots \wedge \widetilde{b_{k_{p-1}}^{p-1}}, (c_{1}^{p})^{*} \wedge \dots \wedge (c_{m_{p}}^{p})^{*} \rangle^{(-1)^{p}},$
which is nothing but $\prod_{p=0}^{n} [\{c_{1}^{p}, \dots, c_{m_{p}}^{p}\}, \{b_{1}^{p}, \dots, b_{k_{p}}^{p}, \widetilde{b_{1}^{p-1}}, \dots, \widetilde{b_{k_{p-1}}^{p-1}}\}]^{(-1)^{p}}$ or

 $\prod_{p=0}^{n} ([\{b_1^p, \cdots, b_{k_p}^p, \widetilde{b_1^{p-1}}, \cdots, \widetilde{b_{k_{p-1}}^{p-1}} \}, \{c_1^p, \cdots, c_{m_p}^p\}]^{(-1)})^{(-1)^p}.$ The last term coincides with the Tor $(C_*, \{c_p\}_{p=0}^n, \{0\}_{p=0}^n)$ defined in previous section.

We will now explain how a general chain complex can be (unnaturally) as a direct sum of an two chain complexes, one of which is exact and the other is ∂ -zero.

Theorem 2.0.5. If $C_* : 0 \to C_n \xrightarrow{\partial_n} C_{n-1} \to \cdots \to C_1 \xrightarrow{\partial_1} C_0 \to 0$ is any chain complex, then it can be splitted as $C_* = C'_* \oplus C''_*$, where C'_* is exact, and C''_* is ∂ -zero.

Proof. Consider the short-exact sequences

If $\ell_p : \operatorname{Im}\partial_p \to C_p$, and $s_p : H_p \to \ker \partial_p$ are sections, i.e. $\partial_p \circ \ell_p = id_{\operatorname{Im}\partial_p}$, and $\pi_p \circ s_p = id_{H_p(C)}$, then C_p is equal to $\ker \partial_p \oplus \ell_p(\operatorname{Im}\partial_p)$ or $\operatorname{Im}\partial_{p+1} \oplus s_p(H_p) \oplus \ell_p(\operatorname{Im}\partial_p)$. Define $C'_p := \operatorname{Im}\partial_{p+1} \oplus \ell_p(\operatorname{Im}\partial_p)$ and $C''_p := s_p(H_p)$. Restricting $\partial_p : C_p \to C_{p-1}$ to these, we obtain two chain complexes $(C'_*, \partial'_*) (C''_*, \partial''_*)$.

As C''_p is a subspace of ker ∂_p , $\partial''_p : C''_p \to C''_{p-1}$ is the zero map, i.e. C''_* is ∂ -zero chain complex. Note also ker $\{\partial''_p : C''_p \to C''_{p-1}\}$ equals to C''_p and $\operatorname{Im}\{\partial''_{p+1} : C''_{p+1} \to C''_p\}$ is $\{0\}$. Then, $H_p(C''_*) = C''_p/\{0\}$ is isomorphic to $H_p(C)$, because $C''_p = s_p(H_p(C))$ is isomorphic to $H_p(C)$.

The exactness of (C'_*, ∂'_*) can be seen as follows: Since $\operatorname{Im}\partial_{p+1}$ is a subspace of ker ∂_p , the image of $\operatorname{Im}\partial_{p+1}$ under ∂'_p is zero. Hence, ker $\{\partial'_p : C'_p \to C'_{p-1}\}$ equals to $\operatorname{Im}\{\partial_{p+1} : C_{p+1} \to C_p\}$. Since $\partial_p \circ \ell_p = id_{\operatorname{Im}\partial_p}$, and $\partial'_p : C'_p \to C'_{p-1}$ is the restriction of $\partial_p : C_p \to C_{p-1}$, then $\operatorname{Im}\{\partial'_p : C'_p \to C'_{p-1}\}$ equals to $\operatorname{Im}\{\partial_p : C_p \to C_{p-1}\}$. Similarly, $\operatorname{Im}\{\partial'_{p-1} : C'_{p-1} \to C'_{p-2}\} = \operatorname{Im}\{\partial_{p-1} : C_{p-1} \to C_{p-2}\}$ and ker $\{\partial'_{p-1} : C'_{p-1} \to C'_{p-2}\} = \operatorname{Im}\{\partial_p : C_p \to C_{p-1}\}$, because $\operatorname{Im}\partial_p$ is a subspace of ker ∂_{p-1} and ℓ_{p-1} is a section of $\partial_{p-1} : C_{p-1} \to \operatorname{Im}\partial_{p-1}$. Consequently, $\operatorname{Im}\{\partial'_p : C'_p \to C'_{p-1}\} = \operatorname{ker}\{\partial'_{p-1} : C_{p-1} \to C_{p-2}\} = \operatorname{Im}\partial_p$ and we have the exactness of C'_* .

This concludes Theorem 2.0.5.

In the next result, we will explain Witten's remark on ([5] pp 185) how torsion $\operatorname{Tor}(C_*)$ of a general complex can be interpreted as an element of the dual of the one dimensional vector space $\bigotimes_{p=0}^n (\det(H_p(C)))^{(-1)^p}$.

Theorem 2.0.6. $Tor(C_*)$ of a general complex is as an element of the dual of the one dimensional vector space $\bigotimes_{p=0}^n (\det(H_p(C)))^{(-1)^p}$.

Proof. Let C_* be a general chain complex of finite dimensional vector spaces with volumes $\alpha_p \in (\det C_p)^{-1}$, i.e. $\alpha_p = (c_p^1)^* \wedge \cdots \wedge (c_p^{i_p})^*$, for some basis $\mathfrak{c}_p = \{c_p^1, \cdots, c_p^{i_p}\}$ of C_p . Let $C_* = C'_* \oplus C''_*$ be the above unnatural splitting of C_* i.e. $C'_p = \operatorname{Im} \partial_{p+1} \oplus \ell_p(\operatorname{Im} \partial_p)$ and $C''_p = s_p(H_p(C))$, where $\ell_p : \operatorname{Im} \partial_p \to C_p$ is the lift of $\partial_p : C_p \to \operatorname{Im} \partial_p$ and $s_p : H_p \to \ker \partial_p$ is the lift of $\pi_p : \ker \partial_p \to H_p(C)$ used in Theorem 2.0.5.

Since $C_p = \operatorname{Im}\partial_{p+1} \oplus s_p(H_p) \oplus \ell_p(\operatorname{Im}\partial_p)$, we can break the basis \mathfrak{c}_p of C_p into three blocks as $\mathfrak{c}_1^p \sqcup \mathfrak{c}_2^p \sqcup \mathfrak{c}_3^p$, where \mathfrak{c}_1^p generates $\operatorname{Im}\partial_{p+1}$, \mathfrak{c}_2^p is basis for $s_p(H_p(C))$ i.e. $[\mathfrak{c}_2^p] = \pi_p(\mathfrak{c}_2^p)$ generates $H_p(C)$, and \mathfrak{c}_3^p is a basis for $\operatorname{Im}\partial_p$. As the determinant of change-base-matrix from \mathfrak{c}_p to \mathfrak{c}_p is 1, the bases \mathfrak{c}_2^p , $\mathfrak{c}_p = \mathfrak{c}_1^p \sqcup \mathfrak{c}_2^p \sqcup \mathfrak{c}_3^p$, and $\mathfrak{c}_1^p \sqcup \mathfrak{c}_3^p$ for C''_p, C_p, C'_p , will be compatible with the short-exact sequence of complexes

$$0 \to C_*'' \hookrightarrow C_* = C_*'' \oplus C_*' \to C_*' \to 0,$$

where we consider the inclusion as section $C'_p \to C_p$. Note also that $H_p(C'') = C''_p/0$ i.e. $s_p(H_p(C))$ which is isomorphic to $H_p(C)$. By Milnor's result Theorem 1.0.4, we have $\operatorname{Tor}(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n)$ is the product of $\operatorname{Tor}(C_*'', \{\mathfrak{c}_p^2\}_{p=0}^n, \{s_p(\mathfrak{h}_p)\}_{p=0}^n)$, $\operatorname{Tor}(C_*', \{\mathfrak{c}_p^1 \sqcup \mathfrak{c}_p^3\}_{p=0}^n, \{0\}_{p=0}^n)$, and $\operatorname{Tor}(\mathcal{H}_*)$, where \mathcal{H}_* is the long-exact sequence obtained from the above short-exact of chain complexes.

More precisely, $\mathcal{H}_* : 0 \to H_n(C'') \to H_n(C) \to H_n(C') \to H_{n-1}(C'') \to H_{n-1}(C') \to H_{n-1}(C') \to \dots \to H_0(C'') \to H_0(C) \to H_0(C') \to 0$. Since C'_* is exact, then \mathcal{H}_* is the long exact-sequence $0 \to H_n(C'') \to H_n(C) \to 0 \to H_{n-1}(C'') \to H_{n-1}(C) \to 0 \to \dots \to 0 \to H_0(C'') \to H_0(C) \to 0 \to 0$. Using the isomorphism $H_p(C) \to H_p(C'')$, namely s_p as section, we conclude that $\operatorname{Tor}(\mathcal{H}_*, \{s_p(\mathfrak{h}_p), \mathfrak{h}_p, 0\}_{p=0}^n, \{0\}_{p=0}^{3n+2}) = 1$.

Moreover, we can also verify that $\operatorname{Tor}(C'_*, \{\mathfrak{c}^1_p \sqcup \mathfrak{c}^3_p\}_{p=0}^n, \{0\}_{p=0}^n) = 1$ as follows:

$$0 \to \ker\{\partial'_p; C'_p \to C'_{p-1}\} \hookrightarrow C'_p \stackrel{\partial'_p \stackrel{i.e.}{=} \partial_p}{\to} \operatorname{Im}\{\partial'_p: C'_p \to C'_{p-1}\} \to 0,$$

where ker{ $\partial'_p : C'_p \to C'_{p-1}$ } is Im{ $\partial_{p+1} : C_{p+1} \to C_p$ } and Im{ $\partial'_p : C'_p \to C'_{p-1}$ } is Im{ $\partial_p : C_p \to C_{p-1}$ }. If we consider the section ℓ_p , then we also have $\operatorname{Tor}(C'_*, \{\mathfrak{c}^1_p \sqcup \mathfrak{c}^3_p\}_{p=0}^n, \{0\}_{p=0}^n) = 1.$

Therefore, $\operatorname{Tor}(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n)$ is equal to $\operatorname{Tor}(C_*'', \{\mathfrak{c}_p^2\}_{p=0}^n, \{s_p(\mathfrak{h}_p)\}_{p=0}^n)$ i.e. $\prod_{n=0}^n [s_p(\mathfrak{h}_p), \mathfrak{c}_p^2]^{(-1)^p}$, where $[s_p(\mathfrak{h}_p), \mathfrak{c}_p^2]$ is the determinant of the change-base-

matrix from \mathfrak{c}_p^2 to $s_p(\mathfrak{h}_p)$ of $C_p'' = s_p(H_p(C))$. Recall that $s_p : H_p(C) \to \ker \partial_p$ is the lift of π_p : $\ker \partial_p \to H_p(C)$. So, $[\mathfrak{c}_p^2]$, i.e. $\pi_p(\mathfrak{c}_p)$, and $\mathfrak{h}_p = [s_p(\mathfrak{h}_p)]$ are bases for $H_p(C)$. Since s_p is isomorphism onto its image, change-base-matrix from \mathfrak{c}_p^2 to $s_p(\mathfrak{h}_p)$ coincides with the one from $[\mathfrak{c}_p^2]$ to \mathfrak{h}_p .

As a result, we obtained that

$$\operatorname{Tor}(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n) = \prod_{p=0}^n [\mathfrak{h}_p, \, [\mathfrak{c}_p^2] \,]^{(-1)^p} \\ = [\mathfrak{h}_0, \, [\mathfrak{c}_0^2] \,] \cdot [\mathfrak{h}_1, \, [\mathfrak{c}_1^2] \,]^{-1} \cdots [\mathfrak{h}_n, \, [\mathfrak{c}_n^2] \,]^{(-1)^n}.$$

For p even, $[\mathfrak{h}_p, \, [\mathfrak{c}_p^2] \,]^{(-1)^p}$ is $[\mathfrak{h}_p, \, [\mathfrak{c}_p^2] \,]$, for p odd, it is $[\mathfrak{h}_p, \, [\mathfrak{c}_p^2] \,]^{-1}$ or $[\, [\mathfrak{c}_p^2] \,, \, \mathfrak{h}_p \,].$

By the remark at the beginning of §2, for even p's, $[\bullet, [\mathfrak{c}_p^2]]$ is linear functional on det $(H_p(C))$, and for odd p's, $[[\mathfrak{c}_p^2], \bullet]$ is linear functional on det $(H_p(C)^*) \equiv$

 $\det(H_p(C))^{-1}$, where the exponent -1 denotes the dual of the space.

This finishes the proof of Theorem 2.0.6.

3. Symplectic Chain Complex

Definition 3.0.7. $C_*: 0 \to C_n \xrightarrow{\partial_n} C_{n-1} \to \cdots \to C_{\frac{n}{2}} \to \cdots \to C_1 \xrightarrow{\partial_1} C_0 \to 0$ is a symplectic chain complex, if

• $n \equiv 2 \pmod{4}$ and

• there exist non-degenerate anti-symmetric ∂ -compatible bilinear maps i.e. $\omega_{p,n-p}: C_p \times C_{n-p} \to \mathbb{R}$ s.t. $\omega_{p,n-p}(a,b) = (-1)^{p(n-p)}\omega_{n-p,p}(b,a)$ and $\omega_{p,n-p}(\partial_{p+1}a,b) = (-1)^{p+1}\omega_{p+1,n-(p+1)}(a,\partial_{n-p}b).$

In the definition, since $n \equiv 2 \pmod{4}$ i.e. *n* is even and $\frac{n}{2}$ is odd, $\omega_{p,n-p}(a,b) = (-1)^p \omega_{n-p,p}(b,a)$.

Using the ∂ -compatibility of the non-degenerate anti-symmetric bilinear maps $\omega_{p,n-p}: C_p \times C_{n-p} \to \mathbb{R}$, one can easily extend these to homologies. Namely,

Lemma 3.0.8. The bilinear map $[\omega_{p,n-p}] : H_p(C) \times H_{n-p}(C) \to \mathbb{R}$ defined by $[\omega_{p,n-p}]([x], [y]) = \omega_{p,n-p}(x, y)$ is anti-symmetric and non-degenerate.

Proof. For the well-definiteness, let x, x' be in ker ∂_p with $x - x' = \partial_{p+1}x''$ for some $x'' \in C_{p+1}$ and let y, y' be in ker ∂_{n-p} with $y - y' = \partial_{n-p+1}y''$ for some $y'' \in C_{n-p+1}$. Then from the bilinearity and ∂ -compatibility, $[\omega_{p,n-p}]([x], [y])$ is equal to $\omega_{p,n-p}(x', y') + (-1)^p \omega_{p-1,n-p+1}(\partial_p x', y'') + (-1)^{p+1} \omega_{p+1,n-p-1}(x'', \partial_{n-p}y') + (-1)^{p+1} \omega_{p+1,n-p-1}(x'', \partial_{n-p} \circ \partial_{n-p+1}y'') = \omega_{p,n-p}(x', y').$

Assume for some $[y_0] \in H_{n-p}(C), \ [\omega_{p,n-p}]([x], [y_0]) = 0$ for all $[x] \in H_p(C)$.

Lemma 3.0.9. y_0 is in $Im\partial_{n-p+1}$.

Proof. Let $\varphi : \frac{C_p}{Z_p} \to \mathbb{R}$ be defined by $\varphi(x + Z_p) = \omega_{p,n-p}(x, y_0)$. This is a welldefined linear map because if $x - x' \in Z_p$, then $\omega_{p,n-p}(x, y_0) - \omega_{p,n-p}(x', y_0) = [\omega_{p,n-p}]([x - x'], [y_0])$ equals to 0. By the 1st isomorphism theorem, $\frac{C_p}{Z_p} \stackrel{\overline{\partial_p}}{\cong} \operatorname{Im} \partial_p = B_{p-1}$, where $\overline{\partial_p}(x + Z_p)$ is $\partial_p(x)$.

Consider the linear functional $\widetilde{\varphi} := \varphi \circ (\overline{\partial_p})^{-1}$ on B_{p-1} , where $(\overline{\partial_p})^{-1}(\partial_p y) = y + Z_p$. Extend $\widetilde{\varphi}$ to $\widehat{\varphi} : C_{p-1} = B_{p-1} \oplus \frac{C_{p-1}}{B_{p-1}} \to \mathbb{R}$ as zero on complement. Since $\omega_{p-1,n-p+1} : C_{p-1} \times C_{n-p+1} \to \mathbb{R}$ is non-degenerate, it induces an isomorphism between the dual space C_{p-1}^* of C_{p-1} and C_{n-p+1} . Therefore, there exists a unique $u_0 \in C_{n-p+1}$ such that $\widehat{\varphi}(\cdot) = \omega_{p-1,n-p+1}(\cdot, u_0)$.

For $x \in C_p$, $v = \partial_p(x)$ is in B_{p-1} Then, on one hand, $\widehat{\varphi}(v)$ is $\omega_{p-1,n-p+1}(\partial_p x, u_0)$ or $(-1)^p \omega_{p,n-p}(x, \partial_{n-p+1}u_0)$ by the ∂ -compatibility. On the other hand, by the construction of $\widehat{\varphi}, \widehat{\varphi}(v) = \omega_{p,n-p}(x, y_0)$. So, $\omega_{p,n-p}(x, y_0)$ is $\omega_{p,n-p}(x, (-1)^p \partial_{n-p+1}u_0)$ for all $x \in C_p$.

The nondegeneracy of $\omega_{p,n-p}$ finishes the proof of Lemma 3.0.9.

This concludes the proof of Lemma 3.0.8

We will define ω -compatibility for bases in a symplectic chain complex.

8

Definition 3.0.10. Let $C_* : 0 \to C_n \xrightarrow{\partial_n} C_{n-1} \to \cdots \to C_{\frac{n}{2}} \to \cdots \to C_1 \xrightarrow{\partial_1} C_0 \to 0$ be a symplectic chain complex. Bases $\mathfrak{o}_p, \mathfrak{o}_{n-p}$ of C_p, C_{n-p} are ω -compatible if the matrix of $\omega_{p,n-p}$ in bases $\mathfrak{o}_p, \mathfrak{o}_{n-p}$ is

$$\begin{cases} Id_{k\times k} & ;p\neq \frac{n}{2}\\ \begin{bmatrix} O_{m\times m} & Id_{m\times m}\\ -Id_{m\times m} & 0_{m\times m} \end{bmatrix} & ;p=\frac{n}{2} \end{cases}$$

where k is dim $C_p = \dim C_{n-p}$ and $2m = \dim C_{\frac{n}{2}}$. In the same way, considering $[\omega_{p,n-p}] : H_p(C) \times H_{n-p}(C) \to \mathbb{R}$, we can also define $[\omega_{p,n-p}]$ -compatibility of bases $\mathfrak{h}_p, \mathfrak{h}_{n-p}$ of $H_p(C), H_{n-p}(C)$.

In the next result, we will explain how a general symplectic chain complex C_* can be splitted ω -orthogonally as a direct sum of an exact and ∂ -zero symplectic complexes.

Theorem 3.0.11. Let $C_* : 0 \to C_n \xrightarrow{\partial_n} C_{n-1} \to \cdots \to C_1 \xrightarrow{\partial_1} C_0 \to 0$ be a symplectic chain complex. Assume $\mathfrak{o}_p, \mathfrak{o}_{n-p} \omega$ -compatible. Then C_* can be splitted as a direct sum of symplectic complexes C'_*, C''_* , where C'_* is exact, C''_* is ∂ -zero and C'_* is perpendicular to C''_* .

Proof. Start with the following short-exact sequence

Consider the section $\ell_p : \operatorname{Im} \partial_p \to C_p$ defined by $\ell_p(\partial_p x) = x$ for $\partial_p x \neq 0$, and $s_p : H_p \to \ker \partial_p$ by $s_p([x]) = x$.

As C_p disjoint union of $\operatorname{Im}\partial_{p+1}$, $s_p(H_p)$, and $\ell_p(\operatorname{Im}\partial_p)$, the basis \mathfrak{o}_p of C_p has three blocks $\mathfrak{o}_p^1, \mathfrak{o}_p^2, \mathfrak{o}_p^3$, where \mathfrak{o}_p^1 is a basis for $\operatorname{Im}\partial_{p+1}, \mathfrak{o}_p^2$ generates $s_p(H_p)$ the rest of ker ∂_p , i.e. $[\mathfrak{o}_p^2]$ generates $H_p(C)$, and $\partial_p \mathfrak{o}_p^3$ is a basis for $\operatorname{Im}\partial_p$. Similarly, $\mathfrak{o}_{n-p} = \mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^2 \sqcup \mathfrak{o}_{n-p}^3$. Because $[\omega]_{p,n-p} : H_p(C) \times H_{n-p}(C) \to \mathbb{R}$ defined by $[\omega]_{p,n-p}([a], [b]) = \omega_{p,n-p}(a, b)$ is non-degenerate and bases $\mathfrak{o}_p \mathfrak{o}_{n-p}$ of C_p, C_{n-p} are ω -compatible, $\omega_{p,n-p}(\cdot, s_{n-p}(H_{n-p})) : C_p \to \mathbb{R}$ vanishes on $\operatorname{Im}\partial_{p+1} \oplus \ell_p(\operatorname{Im}\partial_p)$. Likewise, $\omega_{p,n-p}(s_p(H_p(C)), \cdot) : C_{n-p} \to \mathbb{R}$ vanishes on $\operatorname{Im}\partial_{n-p+1} \oplus \ell_{n-p}(\operatorname{Im}\partial_{n-p})$.

Set $C'_p = \operatorname{Im} \partial_{p+1} \oplus \ell_p(\operatorname{Im} \partial_p)$ and $C''_p = s_p(H_p)$. Note that C'_p with basis $\mathfrak{o}_p^1 \sqcup \mathfrak{o}_p^3$ and C''_{n-p} with basis \mathfrak{o}_{n-p}^2 are ω -orhogonal to each other. Hence, (C'_*, ∂) , (C''_*, ∂) will be the desired splitting, where we consider the corresponding restrictions of $\omega_{p,n-p} : C_p \times C_{n-p} \to \mathbb{R}$.

Clearly, (C''_*, ∂) is ∂ -zero for C''_p being subspace of ker ∂_p . Since $[\omega_{p,n-p}]$: $H_p(C) \times H_{n-p}(C) \to \mathbb{R}$ is non-degenerate, the restriction $\omega_{p,n-p} : C''_p \times C''_{n-p} \to \mathbb{R}$ is also non-degenerate. Being the restriction of $\omega_{p,n-p}$, it is also ∂ -compatible. Hence C''_* becomes symplectic chain complex with ∂ -zero.

In the sequence $C'_{p+1} \xrightarrow{\partial_{p+1}} C'_p \xrightarrow{\partial_p} C'_{p-1}$, first map ∂_{p+1} sends $\operatorname{Im}\partial_{p+2}$, $\ell_{p+1}(\operatorname{Im}\partial_{p+1})$ to zero and $\operatorname{Im}\partial_{p+1}$, respectively. Hence, $\operatorname{ker}\{\partial_{p+1} : C'_{p+1} \to C'_p\}$ equals to $\operatorname{Im}\{\partial_{p+2} : C_{p+2} \to C_{p+1}\}$ and $\operatorname{Im}\{\partial_{p+1} : C'_{p+1} \to C'_p\}$ is $\operatorname{Im}\{\partial_{p+1} : C_{p+1} \to C_p\}$. Similarly, $\operatorname{ker}\{\partial_p : C'_p \to C'_{p-1}\} = \operatorname{Im}\{\partial_{p+1} : C_{p+1} \to C_p\}$ and $\operatorname{Im}\{\partial_p : C'_p \to C'_{p-1}\}$

is $\operatorname{Im}\{\partial_p: C_p \to C_{p-1}\}$. Thus, C'_* is exact.

Moreover, since $\omega_{p,n-p}: C_p \times C_{n-p} \to \mathbb{R}$ is non-degenerate, and C'_p, C'_{n-p} are ω -perpendicular to C''_{n-p}, C''_p , respectively, $\omega_{p,n-p}: C'_p \times C'_{n-p} \to \mathbb{R}$ is non-degenerate. Also, it is ∂ -compatible for being restriction of the ∂ -compatible map $\omega_{p,n-p}: C_p \times C_{n-p} \to \mathbb{R}$.

This concludes the proof of Theorem 3.0.11

Above theorem is a special case of Theorem 2.0.5. The only only difference is using ω -compatible bases \mathfrak{o}_p the splitting is ω -orthogonal, too.

We will now explain how the torsion of a symplectic complex with ∂ -zero is connected with Pfaffian of the anti-symmetric $[\omega_{\frac{n}{2},\frac{n}{2}}] : H_{\frac{n}{2}}(C) \times H_{\frac{n}{2}}(C) \to \mathbb{R}$. Then, Pfaffian will be defined. After that, we will give the relation for a general symplectic complex.

Theorem 3.0.12. Let C_* be symplectic chain complex with ∂ -zero. Let \mathfrak{h}_p be a basis for H_p . Assume the bases $\mathfrak{o}_p, \mathfrak{o}_{n-p}$ of C_p, C_{n-p} are ω -compatible with the property that the bases $\mathfrak{o}_{\frac{n}{2}}$ and $h_{\frac{n}{2}}$ of $H_{\frac{n}{2}}(C)$ are in the same orientation class. Then,

$$Tor(C_*, \{\mathfrak{o}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n) = \left(\prod_{p=0}^{\frac{n}{2}-1} (\det[\omega_{p,n-p}])^{(-1)^p}\right) \cdot \left(\sqrt{\det[\omega_{\frac{n}{2},\frac{n}{2}}]}\right)^{(-1)^{\frac{n}{2}}},$$

where det $[\omega_{p,n-p}]$ is the determinant of the matrix of the non-degenerate pairing $[\omega_{p,n-p}]: H_p(C) \times H_{n-p}(C) \to \mathbb{R}$ in bases $\mathfrak{h}_p, \mathfrak{h}_{n-p}$.

Proof. C_* is ∂ -zero complex, so all B_p 's are zero and $Z_p = C_p$. In particular, H_p is equal to $C_p/\{0\}$ and hence the basis \mathfrak{h}_p of H_p can also be considered as a basis in C_p . Recall $\operatorname{Tor}(C_*, \{\mathfrak{o}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n)$ is defined as the alternating product

$$\prod_{p=0}^{n} [\mathfrak{o}_{p},\mathfrak{h}_{p}]^{(-1)^{p}} = [\mathfrak{o}_{0},\mathfrak{h}_{0}]^{(-1)^{0}} \cdots [\mathfrak{o}_{\frac{n}{2}},\mathfrak{h}_{\frac{n}{2}}]^{(-1)^{\frac{n}{2}}} \cdots [\mathfrak{o}_{n},\mathfrak{h}_{n}]^{(-1)^{n}},$$

of the determinants $[\mathfrak{o}_p, \mathfrak{h}_p]$ of the change-base-matrices from \mathfrak{h}_p to \mathfrak{o}_p . If we combine the terms symmetric with the middle term $[\mathfrak{o}_{\frac{n}{2}}, \mathfrak{h}_{\frac{n}{2}}]^{(-1)^{\frac{n}{2}}}$, torsion becomes

$$\left(\prod_{p=0}^{\frac{n}{2}-1} [\mathfrak{o}_p,\mathfrak{h}_p]^{(-1)^p} [\mathfrak{o}_{n-p},\mathfrak{h}_{n-p}]^{(-1)^{n-p}}\right) [\mathfrak{o}_{\frac{n}{2}},\mathfrak{h}_{\frac{n}{2}}]^{(-1)^{\frac{n}{2}}}$$

Moreover, note that $[\mathfrak{o}_p, \mathfrak{h}_p]^{(-1)^p} [\mathfrak{o}_{n-p}, \mathfrak{h}_{n-p}]^{(-1)^{n-p}} = \{ [\mathfrak{o}_p, \mathfrak{h}_p] [\mathfrak{o}_{n-p}, \mathfrak{h}_{n-p}] \}^{(-1)^p}$ for *n* being even. Let $T^{\mathfrak{o}_p}_{\mathfrak{h}_p}, T^{\mathfrak{o}_{n-p}}_{\mathfrak{h}_{n-p}}$ denote the change-base-matrices from \mathfrak{h}_p to \mathfrak{o}_p of C_p , and from \mathfrak{h}_{n-p} to \mathfrak{o}_{n-p} of C_{n-p} respectively, i.e. $h^i_p = \sum_{\alpha} (T^{\mathfrak{o}_p}_{\mathfrak{h}_p})_{\alpha i} o^{\alpha}_p$ and $h^j_{n-p} = \sum_{\beta} (T^{\mathfrak{o}_{n-p}}_{\mathfrak{h}_{n-p}})_{\beta j} o^{\beta}_{n-p}$, where h^i_p is the *i*th-element of the basis \mathfrak{h}_p .

If A and B are the matrices of $\omega_{p,n-p}$ in the bases $\mathfrak{h}_p, \mathfrak{h}_{n-p}$, and in the bases $\mathfrak{o}_p, \mathfrak{o}_{n-p}$, respectively, then $A = (T^{\mathfrak{o}_p}_{\mathfrak{h}_p})^{\text{transpose}} B T^{\mathfrak{o}_{n-p}}_{\mathfrak{h}_{n-p}}$. By the ω -compatibility

11

of the bases $\mathfrak{o}_p, \mathfrak{o}_{n-p}$, the matrix B is equal to $Id_{k\times k}$, $\begin{bmatrix} 0_{m\times m} & Id_{m\times m} \\ -Id_{m\times m} & 0_{m\times m} \end{bmatrix}$ for $p \neq \frac{n}{2}$, $p = \frac{n}{2}$, respectively, where k is dim $C_p = \dim C_{n-p}$ and $2m = \dim C_{\frac{n}{2}}$. Clearly, determinant of B is $1^k = (-1)^m (-1)^m$ or 1.

Hence, det A equals det $T_{\mathfrak{h}_p}^{\mathfrak{o}_p} \det T_{\mathfrak{h}_{n-p}}^{\mathfrak{o}_{n-p}}$ or $[\mathfrak{o}_p, \mathfrak{h}_p][\mathfrak{o}_{n-p}, \mathfrak{h}_{n-p}]$ for all p. In particular, for $p = \frac{n}{2}$, it is $[\mathfrak{o}_{\frac{n}{2}}, \mathfrak{h}_{\frac{n}{2}}]^2$. Since 2m is even, and $\omega_{\frac{n}{2}, \frac{n}{2}}$ is non-degenerate skew-symmetric, the determinant det $A_{\frac{n}{2}}$ is positive actually equals to $\mathrm{Pfaf}(\omega_{\frac{n}{2}, \frac{n}{2}})^2$, and thus $[\mathfrak{o}_{\frac{n}{2}}, \mathfrak{h}_{\frac{n}{2}}] = \pm \sqrt{\det A_{\frac{n}{2}}}$. Because $\mathfrak{o}_{\frac{n}{2}}, \mathfrak{h}_{\frac{n}{2}}$ are in the same orientation class, then $[\mathfrak{o}_{\frac{n}{2}}, \mathfrak{h}_{\frac{n}{2}}] = \sqrt{\det A_{\frac{n}{2}}}$.

The proof is finished by the fact $\omega_{p,n-p}(h_p^i,h_{n-p}^j) = [\omega_{p,n-p}](h_p^i,h_{n-p}^j).$

Theorem 3.0.13. Let C_* be an exact symplectic chain complex. If $\mathfrak{c}_p, \mathfrak{c}_{n-p}$ are bases for C_p, C_{n-p} , respectively, then $Tor(C_*, {\mathfrak{c}_p}_{p=0}^n, {\{0\}}_{p=0}^n) = 1$.

Proof. From the exactness of C_* , we have $H_p = 0$ or ker $\partial_p = \text{Im}\partial_{p+1}$. Using the short-exact sequence

$$0 \to \ker \partial_p \hookrightarrow C_p \to \operatorname{Im} \partial_p \to 0,$$

we also have $C_p = \ker \partial_p \oplus \ell_p(\operatorname{Im}\partial_p)$, where we consider the section $\ell_p(\partial_p x) = x$ for $\partial_p x \neq 0$.

Let $\mathfrak{o}_p, \mathfrak{o}_{n-p}$ be ω -compatible bases for C_p, C_{n-p} , respectively. We can break $\mathfrak{o}_p = \mathfrak{o}_p^1 \sqcup \mathfrak{o}_p^3$, where \mathfrak{o}_p^1 generates ker $\partial_p = \operatorname{Im}\partial_{p+1}$, and $\partial_p \mathfrak{o}_p^3$ generates $\operatorname{Im}\partial_p$. Similarly, $\mathfrak{o}_{n-p} = \mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^3$, where \mathfrak{o}_{n-p}^1 generates ker $\partial_{n-p} = \operatorname{Im}\partial_{n-p+1}$, and $\partial_{n-p}\mathfrak{o}_{n-p}^3$ generates $\operatorname{Im}\partial_{n-p}$. Since $\omega_{p,n-p} : C_p \times C_{n-p} \to \mathbb{R}$ is non-degenerate, ∂ -compatible, then $\omega_{p,n-p}(\mathfrak{o}_p^1,\mathfrak{o}_{n-p}^1) = 0$, and $\omega_{p,n-p}(\mathfrak{o}_p^1,\mathfrak{o}_{n-p}^3)$ does not vanish. Also by the ω -compatibility of $\mathfrak{o}_p, \mathfrak{o}_{n-p}$, for every *i* there is unique j_i such that $\omega_{p,n-p}((\mathfrak{o}_p^1)_i, (\mathfrak{o}_{n-p}^3)_{\alpha}) = \delta_{j_i,\alpha}$. Likewise, for every *k* there is unique q_k such that $\omega_{p,n-p}((\mathfrak{o}_p^3)_k, (\mathfrak{o}_{n-p}^1)_{\beta}) = \delta_{q_k,\beta}$.

Recall torsion is independent of bases \mathfrak{b}_p for $\operatorname{Im}\partial_p$ and section $\operatorname{Im}\partial_p \to C_p$. Let A_p be the determinant of the matrix of $\omega_{p,n-p}$ in bases $\mathfrak{c}_p \mathfrak{c}_{n-p}$, and let O_p be the determinant of the matrix of $\omega_{p,n-p}$ in bases $\mathfrak{o}_p^1 \sqcup \mathfrak{o}_p^3$, $\mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^3$. Since the set $\partial_p \mathfrak{o}_p^3 = \{\partial_p((\mathfrak{o}_p^3)_1), \cdots, \partial_p((\mathfrak{o}_p^3)_\alpha)\}$ generates $\operatorname{Im}\partial_p$, so does the set $\{\partial_p(A_pO_p(\mathfrak{o}_p^3)_1), \partial_p((\mathfrak{o}_p^3)_2) \cdots, \partial_p((\mathfrak{o}_p^3)_\alpha)\}$. Hence, image of the latter set under ℓ_p , namely, $\widetilde{\mathfrak{o}_p^3} = \{A_p \cdot O_p \cdot (\mathfrak{o}_p^3)_1, (\mathfrak{o}_p^3)_2 \cdots, (\mathfrak{o}_p^3)_\alpha\}$ will also be basis for $\ell_p(\operatorname{Im}\partial_p)$. Keeping $\widetilde{\mathfrak{o}_{n-p}^3}$ as \mathfrak{o}_{n-p}^3 , we have

$$\begin{bmatrix} \omega_{p,n-p} & \text{in} \\ \\ \mathfrak{o}_p^1 \sqcup \widetilde{\mathfrak{o}_p^3} & \mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^3 \end{bmatrix} = (T^{\mathfrak{c}_p}_{\mathfrak{o}_p^1 \sqcup \widetilde{\mathfrak{o}_p^3}})^{\text{transpose}} \begin{bmatrix} \omega_{p,n-p} & \text{in} \\ \\ \\ \mathfrak{c}_p & \mathfrak{c}_{n-p} \end{bmatrix} T^{\mathfrak{c}_{n-p}}_{\mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^3}.$$

Determinant of left-hand-side is $A_p \cdot O_p \cdot O_p$, or A_p because of the determinant of $\omega_{p,n-p}$ in the ω -compatible bases $\mathfrak{o}_p, \mathfrak{o}_{n-p}$. Thus, for $p \neq \frac{n}{2}$, we obtained that $[\mathfrak{c}_p, \mathfrak{o}_p^1 \sqcup \widetilde{\mathfrak{o}_p^3}][\mathfrak{c}_{n-p}, \mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^3] = 1$.

For $p = \frac{n}{2}$, we can prove the same property as follows. Since $\frac{n}{2}$ is odd, $\omega_{\frac{n}{2},\frac{n}{2}}$: $C_{\frac{n}{2}} \times C_{\frac{n}{2}} \to \mathbb{R}$ is non-degenerate and alternating, then the matrix of $\omega_{\frac{n}{2},\frac{n}{2}}$ in any basis of $C_{\frac{n}{2}}$ will be an invertible $2m \times 2m$ skew-symmetric matrix X with real entries, where $2m = \dim C_{\frac{n}{2}}$. Actually, we can find an orthogonal $2m \times 2m$ matrix Q with real entries so that

$$QXQ^{-1} = \operatorname{diag}\left(\left(\begin{array}{cc} 0 & a_1 \\ -a_1 & 0 \end{array}\right), \cdots, \left(\begin{array}{cc} 0 & a_m \\ -a_m & 0 \end{array}\right)\right).$$

So, the determinant of $\omega_{\frac{n}{2},\frac{n}{2}}$ in any basis will be positive, in particular, the determinants $A_{\frac{n}{2}}, O_{\frac{n}{2}}$ of $\omega_{\frac{n}{2},\frac{n}{2}}$ in basis $\mathfrak{c}_{\frac{n}{2}}, \mathfrak{o}_{\frac{n}{2}}^1 \sqcup \mathfrak{o}_{\frac{n}{2}}^3$ respectively will be positive. Having noticed that, let $\widetilde{\mathfrak{o}_{\frac{n}{2}}^3} = \{\sqrt{A_{\frac{n}{2}}} \cdot \sqrt{O_{\frac{n}{2}}} \cdot (\mathfrak{o}_{\frac{n}{2}}^3)_1, (\mathfrak{o}_{\frac{n}{2}}^3)_2 \cdots, (\mathfrak{o}_{\frac{n}{2}}^3)_{\alpha}\}.$

As explained above, on one side, we have that det $\begin{bmatrix} \omega_{\frac{n}{2},\frac{n}{2}} & \text{in} \\ \mathfrak{o}_{\frac{1}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3} \end{bmatrix}$ is equal to $\sqrt{A_{\frac{n}{2}}} \cdot \sqrt{A_{\frac{n}{2}}} \sqrt{O_{\frac{n}{2}}} \cdot \sqrt{O_{\frac{n}{2}}} \det \begin{bmatrix} \omega_{\frac{n}{2},\frac{n}{2}} & \text{in} \\ \mathfrak{o}_{\frac{1}{2}}^1 \sqcup \mathfrak{o}_{\frac{n}{2}}^3 \end{bmatrix}$ or $A_{\frac{n}{2}}$. On the other side, it is the product $[\mathfrak{c}_{\frac{n}{2}}, \mathfrak{o}_{\frac{1}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3}] \cdot A_{\frac{n}{2}} \cdot [\mathfrak{c}_{\frac{n}{2}}, \mathfrak{o}_{\frac{1}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3}]$. Consequently, $[\mathfrak{c}_{\frac{n}{2}}, \mathfrak{o}_{\frac{n}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3}]^2$ is equal to 1.

If $\mathfrak{o}_{\frac{n}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3}$ and $\mathfrak{c}_{\frac{n}{2}}$ are already in the same orientation class, then $[\mathfrak{c}_{\frac{n}{2}}, \mathfrak{o}_{\frac{n}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3}] = 1$. If not, considering $\widetilde{\mathfrak{o}_{\frac{n}{2}}^3}$ as $\{-\sqrt{A_{\frac{n}{2}}} \cdot \sqrt{O_{\frac{n}{2}}} \cdot (\mathfrak{o}_{\frac{n}{2}}^3)_1, (\mathfrak{o}_{\frac{n}{2}}^3)_2 \cdots, (\mathfrak{o}_{\frac{n}{2}}^3)_\alpha\}$, we still have $[\mathfrak{c}_{\frac{n}{2}}, \mathfrak{o}_{\frac{n}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3}] = 1$.

As a result, we proved that

$$\operatorname{Tor}(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{0\}_{p=0}^n) = \prod_{p=0}^n [\mathfrak{c}_p, \mathfrak{o}_p^1 \sqcup \widetilde{\mathfrak{o}_p^3}]^{(-1)^p}$$
$$= \prod_{p=0}^{\frac{n}{2}-1} \left([\mathfrak{c}_p, \mathfrak{o}_p^1 \sqcup \widetilde{\mathfrak{o}_p^3}] [\mathfrak{c}_{n-p}, \mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^3] \right)^{(-1)^p} \cdot [\mathfrak{c}_{\frac{n}{2}}, \mathfrak{o}_{\frac{n}{2}}^1 \sqcup \widetilde{\mathfrak{o}_{\frac{n}{2}}^3}]^{(-1)^{\frac{n}{2}}} = 1$$

Before explaining the corresponding result for a general symplectic complex, we would like to recall the Pfaffian of a skew-symmetric matrix.

Let V be an even dimensional vector space over reals. Let $\omega : V \times V \to \mathbb{R}$ be a bilinear and anti-symmetric. If we fix a basis for V, then ω can be represented by a $2m \times 2m$ skew-symmetric matrix.

If A any $2m \times 2m$ skew-symmetric matrix with real entries then, by the spectral theorem of normal matrices, one can easily find an orthogonal $2m \times 2m$ -real matrix Q so that $QAQ^{-1} = \text{diag}\left(\begin{pmatrix} 0 & a_1 \\ -a_1 & 0 \end{pmatrix}, \cdots, \begin{pmatrix} 0 & a_m \\ -a_m & 0 \end{pmatrix}\right)$, where

 a_1, \dots, a_m are positive real. Thus, in particular, determinant of A is non-negative.

Definition 3.0.14. For $2m \times 2m$ real skew-symmetric matrix A, Pfafian of A will be $\sqrt{\det A}$.

Actually, if $A = [a_{ij}]$ is any $2m \times 2m$ skew-symmetric matrix and if we let $\omega_A = \sum_{i < j} a_{ij} \overrightarrow{e_i} \wedge \overrightarrow{e_j}$, then we can also define $\operatorname{Pfaf}(A)$ as the coefficient of $\overrightarrow{e_1} \wedge \cdots \wedge \overrightarrow{e_{2m}}$

in the product $\frac{\widetilde{\omega_A \wedge \cdots \wedge \omega_A}}{m!}$.

For example, if A is the matrix diag $\begin{pmatrix} 0 & a_1 \\ -a_1 & 0 \end{pmatrix}, \dots, \begin{pmatrix} 0 & a_m \\ -a_m & 0 \end{pmatrix} \end{pmatrix}$, then ω_A is $\sum_{i=1}^m a_i \cdot \overrightarrow{e_{2i-1}} \wedge \overrightarrow{e_{2i}}$. An easy computation shows that $\underbrace{\omega_A \wedge \omega_A \wedge \dots \wedge \omega_A}_{m-times}$ equals to $m! \underbrace{(a_1 \cdots a_m)}_{\text{Pfaffian of A}} \overrightarrow{e_1} \wedge \dots \wedge \overrightarrow{e_{2m}}$.

For a general $2m \times 2m$ skew-symmetric A, we can find an orthogonal matrix Q such that $QAQ^{-1} = \operatorname{diag}\left(\begin{pmatrix} 0 & a_1 \\ -a_1 & 0 \end{pmatrix}, \cdots, \begin{pmatrix} 0 & a_m \\ -a_m & 0 \end{pmatrix}\right)$. As a result, $\underbrace{\omega_{QAQ^{-1}} \wedge \omega_{QAQ^{-1}} \wedge \cdots \wedge \omega_{QAQ^{-1}}}_{m-times}$ equals to m! $\underbrace{(a_1 \cdots a_m)}_{\operatorname{Pfaffian of } QAQ^{-1}} \stackrel{i.e.}{e_1} \wedge \cdots \wedge \stackrel{i.e.}{e_{2m}}$ i.e.

On the other hand, one can easily prove that for any $2m \times 2m$ skew-symmetric matrix X and any $2m \times 2m$ matrix Y, $Pfaf(YXY^t)$ is equal to $Pfaf(A) \det(B)$. Consequently, since Q is orthogonal matrix, we can conclude that $Pfaf(A)^2 = \det(A)$ for any skew-symmetric $2m \times 2m$ real matrix A. In other words, both definitions coincide.

Using Pfafian, we can rephrase Theorem 3.0.12 as follows.

If C_* is a symplectic chain complex with ∂ -zero, \mathfrak{h}_p is a basis for H_p , $\mathfrak{o}_p, \mathfrak{o}_{n-p}$ ω -compatible bases for C_p, C_{n-p} so that $\mathfrak{h}_{\frac{n}{2}}$ and $[\mathfrak{o}_{\frac{n}{2}}]$ are in the same orientation class, then

$$\operatorname{Tor}(C_*, \{\mathfrak{o}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n) = \left(\prod_{p=0}^{\frac{n}{2}-1} (\operatorname{det}[\omega_{p,n-p}])^{(-1)^p}\right) \cdot \left(\operatorname{Pfaf}\left[\omega_{\frac{n}{2},\frac{n}{2}}\right]\right)^{(-1)^{\frac{n}{2}}},$$

where Pfaf $[\omega_{\frac{n}{2},\frac{n}{2}}]$ is the Pfafian of the matrix of the non-degenerate pairing $[\omega_{\frac{n}{2},\frac{n}{2}}]$: $H_{\frac{n}{2}}(C) \times H_{\frac{n}{2}}(C) \to \mathbb{R}$ in bases $\mathfrak{h}_{\frac{n}{2}}, \mathfrak{h}_{\frac{n}{2}}$. **Theorem 3.0.15.** For a general symplectic complex C_* , if \mathfrak{c}_p , \mathfrak{h}_p are bases for C_p , H_p , respectively, then

$$Tor(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n) = \left(\prod_{p=0}^{\frac{n}{2}-1} (\det[\omega_{p,n-p}])^{(-1)^p}\right) \cdot \left(\sqrt{\det[\omega_{\frac{n}{2},\frac{n}{2}}]}\right)^{(-1)^{\frac{n}{2}}},$$

where det $[\omega_{p,n-p}]$ is the determinant of the matrix of the non-degenerate pairing $[\omega_{p,n-p}]: H_p(C) \times H_{n-p}(C) \to \mathbb{R}$ in bases $\mathfrak{h}_p, \mathfrak{h}_{n-p}$.

Proof. Since C_p is disjoint union $\operatorname{Im}\partial_{p+1} \sqcup s_p(H_p) \sqcup \ell_p(\operatorname{Im}\partial_p)$, any basis \mathfrak{a}_p of C_p has three parts $\mathfrak{a}_p^1, \mathfrak{a}_p^2, \mathfrak{a}_p^3$, where \mathfrak{a}_p^1 is basis for $\operatorname{Im}\partial_{p+1}, \mathfrak{a}_p^2$ generates $s_p(H_p)$ the rest of ker ∂_p i.e. $[\mathfrak{a}_p^2]$ generates $H_p(C)$, and $\partial_p \mathfrak{a}_p^3$ is basis for $\operatorname{Im}\partial_p$, where $\ell_p : \operatorname{Im}\partial_p \to C_p$ is the section defined by $\ell_p(\partial_p x) = x$ for $\partial_p x \neq 0$, and $s_p : H_p \to \ker \partial_p$ by $s_p([x]) = x$.

If \mathfrak{o}_p , \mathfrak{o}_{n-p} are ω -compatible bases for C_p and C_{n-p} , then we can also write $\mathfrak{o}_p = \mathfrak{o}_p^1 \sqcup \mathfrak{o}_p^2 \sqcup \mathfrak{o}_p^3$ and $\mathfrak{o}_{n-p} = \mathfrak{o}_{n-p}^1 \sqcup \mathfrak{o}_{n-p}^2 \sqcup \mathfrak{o}_{n-p}^3$. We may assume $[\mathfrak{o}_{\frac{n}{2}}]$ and $\mathfrak{h}_{\frac{n}{2}}$ are in the same orientation class. Otherwise, switch, say, the first element $(\mathfrak{o}_{\frac{n}{2}})^1$ and the corresponding ω -compatible element $(\mathfrak{o}_{\frac{n}{2}})^{m+1}$ i.e. $\omega_{\frac{n}{2},\frac{n}{2}}((\mathfrak{o}_{\frac{n}{2}})^1, (\mathfrak{o}_{\frac{n}{2}})^{m+1}) = 1$, where $2m = \dim H_{\frac{n}{2}}(C)$. In this way, we still have ω -compatibility and moreover we can guarantee that $[\mathfrak{o}_{\frac{n}{2}}]$, $\mathfrak{h}_{\frac{n}{2}}$ are in the same orientation class.

Using these ω -compatible bases \mathfrak{o}_p , as in Theorem 3.0.11, we have the ω -orthogonal splitting $C_* = C'_* \oplus C''_*$, where C'_p and C''_p are $\operatorname{Im}(\partial_{p+1}) \oplus \ell_p(\operatorname{Im}\partial_p)$, $s_p(H_p)$, and $\ell_p : \operatorname{Im}\partial_p \to C_p$ is the section defined by $\ell_p(\partial_p x) = x$ for $\partial_p x \neq 0$, and $s_p : H_p \to \ker \partial_p$ by $s_p([x]) = x$.

 C_p is the disjoint union $\operatorname{Im}\partial_{p+1} \sqcup s_p(H_p) \sqcup \ell_p(\operatorname{Im}\partial_p)$, so the basis \mathfrak{c}_p of C_p has also three blocks $\mathfrak{c}_p^1, \mathfrak{c}_p^2, \mathfrak{c}_p^3$, where \mathfrak{c}_p^1 is a basis for $\operatorname{Im}\partial_{p+1}, \mathfrak{c}_p^2$ generates $s_p(H_p)$ the rest of ker ∂_p , i.e. $[\mathfrak{c}_p^2]$ generates $H_p(C)$, and $\partial_p \mathfrak{c}_p^3$ is a basis for $\operatorname{Im}\partial_p$.

Consider the ∂ -zero symplectic C''_* with the ω -compatible bases $\mathfrak{o}_p^2, \mathfrak{o}_{n-p}^2$. Note that by the ∂ -zero property of $C''_*, H_p(C'')$ is $C''_p/0$ or $s_p(H_p(C))$. Hence $s_p(\mathfrak{h}_p)$ will be a basis $H_p(C'')$. Recall also that $[\mathfrak{o}_{\frac{n}{2}}^2]$ and $\mathfrak{h}_{\frac{n}{2}}^2$ are in the same orientation class. Therefore, by Theorem 3.0.12, we can conclude that

$$\operatorname{Tor}(C_*'', \{\mathfrak{o}_p^2\}_{p=0}^n, \{s_p(\mathfrak{h}_p)\}_{p=0}^n) = \left(\prod_{p=0}^{\frac{n}{2}-1} (\det[\omega_{p,n-p}])^{(-1)^p}\right) \cdot \left(\sqrt{\det[\omega_{\frac{n}{2},\frac{n}{2}}]}\right)^{(-1)^{\frac{n}{2}}}$$

where det $[\omega_{p,n-p}]$ is the determinant of the matrix of the non-degenerate pairing $[\omega_{p,n-p}]: H_p(C) \times H_{n-p}(C) \to \mathbb{R}$ in bases $\mathfrak{h}_p, \mathfrak{h}_{n-p}$.

On the other hand, if \mathfrak{c}'_p is any basis for C'_p , then by Theorem 3.0.13 the torsion $\operatorname{Tor}(C'_*, \{c'_p\}_{p=0}^n, \{0\}_{p=0}^n)$ of the exact symplectic complex C'_* is equal to 1.

Let A_p be the determinant of the change-base-matrix from \mathfrak{o}_p^2 to \mathfrak{c}_p^2 . If we consider the basis $\mathfrak{c}_p^1 \sqcup (\frac{1}{A_p}\mathfrak{c}_p^3)$ for the C'_p , then for the short-exact sequence

$$0 \to C''_* \to C_* = C'_* \oplus C''_* \to C'_* \to 0$$

the bases $\mathfrak{o}_p^2, \mathfrak{c}_p, \mathfrak{c}_p^1 \sqcup (\frac{1}{A_p}\mathfrak{c}_p^3)$ of C_p'', C_p, C_p' respectively will be compatible i.e. the determinant of the change-base-matrix from basis $\mathfrak{c}_p^1 \sqcup \mathfrak{o}_p^2 \sqcup (\frac{1}{A_p}\mathfrak{c}_p^3)$ to $\mathfrak{c}_p = \mathfrak{c}_p^1 \sqcup \mathfrak{c}_p^2 \sqcup \mathfrak{c}_p^3$ is 1.

Thus, by Milnor's result Theorem 1.0.4, $\operatorname{Tor}(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n)$ is equal to the product of $\operatorname{Tor}(C_*'', \{\mathfrak{o}_p^2\}_{p=0}^n, \{s_p(\mathfrak{h}_p)\}_{p=0}^n)$, $\operatorname{Tor}(C_*', \{\mathfrak{c}_p^1 \sqcup (\frac{1}{A_p}\mathfrak{c}_p^3)\}_{p=0}^n, \{0\}_{p=0}^n)$, and $\operatorname{Tor}(\mathcal{H}_*, \{s_p(\mathfrak{h}_p), \mathfrak{h}_p, 0\}_{p=0}^n, \{0\}_{p=0}^{3n+2})$, where \mathcal{H}_* is the long-exact sequence $0 \to H_n(C'') \to H_n(C) \to H_n(C') \to H_{n-1}(C'') \to \cdots \to H_0(C'') \to H_0(C) \to H_0(C') \to 0$ obtained from the short-exact sequence of complexes. Since C_*' is exact, $H_p(C')$ are all zero. So, using the isomorphisms $H_p(C) \to H_p(C'') = C_p''/0$, namely s_p as section, we can conclude that $\operatorname{Tor}(\mathcal{H}_*, \{s_p(\mathfrak{h}_p), \mathfrak{h}_p, 0\}_{p=0}^n, \{0\}_{p=0}^{3n+2}) = 1$. From Theorem 3.0.13, we also obtain $\operatorname{Tor}(C_*', \{\mathfrak{c}_p^1 \sqcup (\frac{1}{A_p}\mathfrak{c}_p^3)\}_{p=0}^n, \{0\}_{p=0}^n) = 1$.

Therefore, we verified that

$$\operatorname{Tor}(C_*, \{\mathfrak{c}_p\}_{p=0}^n, \{\mathfrak{h}_p\}_{p=0}^n) = \operatorname{Tor}(C_*'', \{\mathfrak{o}_p^2\}_{p=0}^n, \{s_p(\mathfrak{h}_p)\}_{p=0}^n).$$

This finishes the proof of Theorem 3.0.15.

References

- [1] J. Milnor, Whitehead torsion, Bull. Amer. Soc. 72 (1966), pp 358-426.
- [2] J. Porti, *Torsion de Reidemeister pour les Varieties Hyperboliques*, Memoirs of the Amer. Math. Soc. 128 (1997) Number 612.
- [3] Y. Sözen, Reidemeister torsion and Pfaffian of Thurston symplectic form, in preparation.
- [4] Y. Sözen, F. Bonahon, Weil-Petersson and Thurston symplectic forms, Duke Math. Jour. Vol 108 No.3 (2001), pp 581-597.
- [5] E. Witten, On quantum gauge theories in two dimension, Commun. in Math.Phys. 141 (1991), pp 153-209.

Rowan University, Department of Mathematics, Glassboro NJ 08028 $E\text{-}mail\ address:\ sozen@rowan.edu$