A Bayes method for a Monotone Hazard Rate via S-paths^{1,2}

Man-Wai Ho

National University of Singapore (March 21, 2005)

Abstract

A class of random hazard rates, that is defined as a mixture of an indicator kernel convoluted with a completely random measure, is of interest. We provide an explicit characterization of the posterior distribution of this mixture hazard rate model via a finite mixture of **S**-paths. A closed and tractable Bayes estimator for the hazard rate is derived to be a finite sum over **S**-paths. The path characterization or the estimator is proved to be a Rao-Blackwellization of an existing partition characterization or partition-sum estimator. This accentuates the importance of **S**-path in Bayesian modeling of monotone hazard rates. An efficient Markov chain Monte Carlo (MCMC) method is proposed to approximate this class of estimates. It is shown that **S**-path characterization also exists in modeling with covariates by a proportional hazard model, and the proposed algorithm again applies. Numerical results of the method are given to demonstrate its practicality and effectiveness.

1 Introduction

Estimation of monotone hazard rate (or hazard function) is important in reliability theory and survival analysis. The hazard rate $\lambda(t)$ is interpreted as the propensity of failure of a system (or an item) in the instant future given that it has survived until time t. In general, the function has a wide variety of shapes. The simplest case of a constant hazard rate corresponds to an exponential lifetime distribution for the system. Cases of increasing or decreasing hazard rate, broadly speaking, correspond to lifetime distributions that are of a lighter or heavier tail, respectively, compared to an exponential distribution. There is a substantial amount of literature about estimation of monotone hazard rates from a frequentist viewpoint. They include, for example, the pioneer work of Grenander (1956) and Prakasa Rao (1970), extensions of their works to different censoring schemes by Padgett and Wei (1980) and Mykytyn and Santner (1981), a constrained spline smoothing technique by Villalobos and Wahba (1987), works of Lo and Phadia (1992) and Huang and Wellner (1995)

¹This research was in part supported by National University of Singapore research grant R-155-000-047-112 and Hong Kong RGC Competitive Earmarked Research Grant HKUST6159/02P.

²AMS 2000 subject classifications. Primary 62G05; secondary 62F15.

Keywords and phrases. Hazard rates; Completely random measure; Gamma process; Weighted gamma process; Rao-Blackwellization; Markov chain Monte Carlo; Proportional hazard model; Gibbs sampler;

based on the least concave/greatest convex minorants, and a kernel-based method by Hall, Huang, Gifford and Gijbels (2001). The last article also provides a good recent survey on hazard rate estimations.

By modeling an increasing hazard rate on the half line $\mathcal{R} = (0, \infty)$ as indicator functions convolved with a gamma process, Dykstra and Laud (1981) first obtained, as a nonparametric Bayes estimate of an increasing hazard rate, a posterior mean of the hazard rate that is expressible as a finite sum over, what they call, e-vectors. Lo and Weng (1989) consider a more general mixture hazard rate model of which the hazard rate is a mixture of an arbitrary kernel k and a weighted gamma process μ [Lo (1982)],

$$\lambda(t|\mu) = \int_{\mathcal{R}} k(t|u)\mu(du), \qquad t \in \mathcal{R}.$$
 (1)

Assuming a multiplicative intensity model [see Aalen (1975, 1978)] under right censoring, they show that the posterior law of the mixing measure μ is a mixture of weighted gamma processes, depending on partitions of *n* integers where *n* is the number of complete observations. It follows that the posterior mean of the mixture hazard rate is expressible as a finite sum over partitions. In case the kernel is an indicator, they obtain that the posterior mean reduces to a coarser sum over **S**-paths, which are equivalent to **e**-vectors. This includes the Bayes estimate by Dykstra and Laud (1981) as a special case. For Bayes inference of general hazard rates with presence of covariates, see Kalbfleisch (1978), Ibrahim, Chen and MacEachern (1999), James (2003), Ishwaran and James (2004) and among others.

This paper discusses a more general class of random monotone hazard rates for Bayes inference, which extends both works of Dykstra and Laud (1981) and Lo and Weng (1989). This work is motivated by a recent work of James (2002, 2004), which extends Lo and Weng's work by considering a mixture hazard rate model represented by (1) of which μ is modeled by a, more general, completely random measure prior [Kingman (1967, 1993)]. One of his interesting results is that the posterior distribution of the mixing measure is again a mixture of partitions. This preservation of partition structure in the posterior distribution inspires a study of the fine structure of the class of random monotone hazard rates defined by (1) of which k is an indicator function and μ is a completely random measure.

A major contribution of this paper is to provide an explicit expression of the posterior distribution of the model as a significantly coarser mixture over **S**-paths, compared to the general result of James (2002, 2004) as a mixture over partitions. In particular, the posterior distribution is nicely described as a three-step hierarchical experiment, which appears in neither of the previous works. The basis, which is the first step, of the hierarchy depends on a posterior (discrete) probability distribution of **S**-paths. The hierarchical description of the posterior distribution allows us to have a better understanding of the model; it entails the importance of this **S**-path characterization as it is shown to be a Rao-Blackwellization of the partition characterization. Analogously, a posterior mean of the monotone hazard rate

expressible as a finite sum over **S**-paths should be preferred to the specialized "partitionsum" estimator of James (2002, 2004), regarding estimation errors. More importantly, these accentuate that in general **S**-path should deserve to have more attention due to its nice statistical properties, as well as its existences in Bayes inference with a much larger class of random processes, but not be restricted to only Dirichlet processes [Brunner and Lo (1989, 1994) and Ho (2005a, 2005b)] and gamma processes.

As "path-sum" estimators appear not only in modeling monotone hazard rates, but also in many other statistical problems under monotonicity constraints, such as monotone or symmetric unimodal densities [Brunner and Lo (1989)], and rotationally symmetric and unimodal densities on the sphere [Brunner and Lo (1994)], there have been several attempts to sample **S**-paths in order to approximate finite sums over **S**-paths [see Brunner and Lo (1989, 1994) and Brunner (1995)]. However, they are all far from successful due to their incapability of sampling from desired posterior distributions of **S**-paths in the respective models. To facilitate practical usage of the path-sum estimates of monotone hazard rates (and those of many other models), this paper proposes an efficient MCMC computational procedure to evaluate finite sums over **S**-paths. The algorithm is designed in view of accelerating a straightforward Gibbs sampler [Geman and Geman (1984)] with the target stationary distribution on the space of all **S**-paths, in the sense that it allows more efficient movements among different **S**-paths based on an idea of Hastings (1970).

To sum up, this paper provides a Bayes method that can serve as a viable alternative to frequentist methods for estimating monotone hazard rates. Hall *et al.* (2001) proposed a kernel technique based on biased-bootstrap for estimating a monotone hazard rate as an alternative of the explicit, but rough, nonparametric likelihood estimates in Huang and Wellner (1995). The Bayes method is comparable to their kernel technique; it results in smooth hazard estimates under various censoring schemes via a computationally straightforward algorithm that does not rely on any choice of bandwidth or bootstrap weights. In addition, Section 6 presents that the method can be easily extended to cases where covariates exist. Under a proportional hazard model, the random vector, **S**-path, analogously plays an important role in characterizing the posterior distribution. Evaluation of posterior quantities of the semi-parametric model can be done by applying the proposed MCMC algorithm with slight modifications. Last but not least, consistency of the posterior distribution of this class of monotone hazard rates follows from Drăgichi and Ramamoorthi (2003), who establish the Bayesian consistency of a large class of monotone hazard rates under wide choice of μ .

The paper is organized as follows: Section 2 introduces the class of random hazard rates that are represented as a mixture of an indicator kernel convolved with a completely random measure. The posterior structure of the model is described in terms of **S**-paths as a three-step hierarchical experiment. An explicit Bayes estimator, which is a finite sum over **S**-paths, of the monotone hazard rate is presented. Section 2.1 provides a detailed description of the RaoBlackwellized result of the path characterization over James' partition characterization in the model. Section 3 discusses a straightforward Gibbs sampler that can be used to approximate the sum over **S**-paths, and presents an efficient MCMC computational procedure. Section 4 demonstrates that our Bayes estimator generalizes corresponding results in other mixture hazard rate models. Section 5 gives numerical examples to illustrate the MCMC algorithms.

2 A posterior distribution of a monotone hazard rate model

This section is concerned with the Bayes estimation of a decreasing hazard rate on the half line $\mathcal{R} = (0, \infty)$. Analogous results exist for estimations of the increasing counterpart. Our interest is in the class of random decreasing hazard rates defined by

$$\lambda(t|\mu) = \int_{\mathcal{R}} \mathbb{I}(t < u)\mu(du), \tag{2}$$

where $\mathbb{I}(A)$ is the indicator function of a set A and μ is taken to be a *completely random* measure [Kingman (1967, 1993)] on the half line, *characterized* by an intensity measure $\rho(dz|u)\eta(du)$. That is, μ can be represented in a distributional sense as

$$\mu(du) = \int_{\mathcal{R}} z \mathcal{N}(dz, du),$$

where $\mathcal{N}(dz, du)$ is a Poisson random measure, taking on points (z, u) in $\mathcal{R} \times \mathcal{R}$, with mean intensity

$$\mathbb{E}[\mathcal{N}(dz, du)] = \rho(dz|u)\eta(du),\tag{3}$$

such that for any bounded set B on the half line, $\int_B \int_{\mathcal{R}} \min(z, 1)\rho(dz|u)\eta(du) < \infty$. In other words, the law of μ is uniquely characterized by the Laplace functional

$$\mathcal{L}_{\mu}(g|\rho,\eta) = \exp\left[-\int_{\mathcal{R}}\int_{\mathcal{R}}\left(1 - e^{-g(u)z}\right)\rho(dz|u)\eta(du)\right],\tag{4}$$

where g is a non-negative function on \mathcal{R} . Suppose we collect failure time observations from N items with hazard rates given by (2) until time τ . Denote the completely observed failure times by $T_1 < \cdots < T_n < \tau$, and right-censored times by $T_{n+1} = \cdots = T_N \equiv \tau$. Define

$$g_N(u) = \int_0^\tau \left[\sum_{i=1}^N \mathbb{I}(T_i \ge t)\right] \mathbb{I}(t < u) dt, \tag{5}$$

and write $\mu(g_N) = \int_{\mathcal{R}} g_N(u)\mu(du) = \int_0^{\tau} \left[\sum_{i=1}^N \mathbb{I}(T_i \ge t)\right] \left[\int_{\mathcal{R}} \mathbb{I}(t < u)\mu(du)\right] dt$. The sum $\sum_{i=1}^N \mathbb{I}(T_i \ge t)$ is called the total time transform [Barlow, Bartholomew, Bremner and Brunk (1972)]. Assume a multiplicative intensity model, the likelihood of the data $\mathbf{T} = (T_1, \ldots, T_N)$ is

$$\frac{N!}{(N-n)!} \left[\prod_{i=1}^{n} \int_{\mathcal{R}} \mathbb{I}(t_i < u_i) \mu(du_i) \right] \exp\left[-\mu(g_N)\right].$$
(6)

Monotone hazard

The posterior law of $\mu | \mathbf{T}$ can be described in terms of **S**-paths in Theorem 2.1. Define an integer-valued vector $\mathbf{S} = (S_0, S_1, \ldots, S_{n-1}, S_n)$, referred to as an **S**-path (of n + 1coordinates), which satisfies (i) $S_0 = 0$ and $S_n = n$; (ii) $S_j \leq j, j = 1, \ldots, n-1$; and (iii) $S_j \leq S_{j+1}, j = 1, \ldots, n-1$. That is, an **S**-path is a nondecreasing integer-valued vector of n + 1 coordinates, starting from 0 to n, bounded above by the diagonal line [Dykstra and Laud (1981), Brunner and Lo (1989)]. Denote the increment (or jump size) at location j by $m_j = S_j - S_{j-1}, j = 1, \ldots, n$, and the number of positive increments/jumps for a path **S** by $n(\mathbf{S}) = \sum_{j=1}^n \mathbb{I}(m_j > 0)$. Define $f_N(z, u) = g_N(u)z$. Given the data **T**, for each integer $i \leq n$ and fixed u > 0, assume that

$$\kappa_i(e^{-f_N}\rho|u) = \int_{\mathcal{R}} z^i e^{-g_N(u)z} \rho(dz|u) < \infty.$$
(7)

This assumption and the meaning of the function will be discussed in Remark 2.1 and Remark 2.2, respectively. Write $\sum_{\mathbf{S}}$ as summing over all paths \mathbf{S} , and $\prod_{\{j:m_j>0\}}$ and $\sum_{\{j:m_j>0\}}$ as $\prod_{j=1:m_j>0}^{n}$ and $\sum_{j=1:m_j>0}^{n}$ conditioned on \mathbf{S} , respectively.

Theorem 2.1. Suppose the likelihood of the data is (6), and μ is a completely random measure characterized by the Laplace functional (4). Then given **T**, the posterior distribution of μ can be described by a three-step experiment:

(i) An **S**-path $\mathbf{S} = (0, S_1, \dots, S_{n-1}, n)$ has a distribution $Z(\mathbf{S}) = \phi(\mathbf{S}) / \sum_{\mathbf{S}} \phi(\mathbf{S})$, where

$$\phi(\mathbf{S}) = \prod_{\{j:m_j>0\}} \left(\begin{array}{c} j-1-S_{j-1} \\ j-S_j \end{array} \right) \int_{T_j}^{\infty} \kappa_{m_j} (e^{-f_N} \rho | y) \eta(dy).$$
(8)

(ii) Given **S**, there exist $n(\mathbf{S})$ independent pairs of (y_j, Q_j) , denoted by $(\mathbf{y}, \mathbf{Q}) = \{(y_j, Q_j) : m_j > 0, j = 1, ..., n\}$, where $y_j | \mathbf{S}, \mathbf{T}$ is distributed as

$$\eta_j(dy_j|\mathbf{S},\mathbf{T}) \propto \mathbb{I}(T_j < y_j)\kappa_{m_j}(e^{-f_N}\rho|y_j)\eta(dy_j),\tag{9}$$

and

$$\Pr\{Q_j \in dz | \mathbf{S}, y_j, \mathbf{T}\} \propto z^{m_j} e^{-g_N(y_j)z} \rho(dz | y_j).$$
(10)

(iii) Given $(\mathbf{S}, \mathbf{y}, \mathbf{Q})$, μ has a distribution as

$$\mu_N^* + \sum_{\{j:m_j > 0\}} Q_j \delta_{y_j}, \tag{11}$$

where μ_N^* is a completely random measure characterized by an intensity measure

$$e^{-g_N(u)z}\rho(dz|u)\eta(du).$$

A proof of this theorem is given in the appendix.

Remark 2.1. The finiteness condition in (7) guarantees the existences of the posterior distributions of $Q_j|\mathbf{S}, y_j$, for j = 1, ..., n such that $m_j > 0$.

Corollary 2.1. Theorem 2.1 implies that the posterior mean of the decreasing hazard rate (2) given **T** is given by, for $t \in [0, \tau]$,

$$\mathbb{E}[\lambda(t|\mu)|\mathbf{T}] = \int_{t}^{\infty} \kappa_{1}(e^{-f_{N}}\rho|y)\eta(dy) + \sum_{\mathbf{S}} Z(\mathbf{S})\sum_{j=1}^{n} \lambda_{j}(t|\mathbf{S}),$$
(12)

where $\kappa_i(e^{-f_N}\rho|y), i = 1, ..., n$, is defined in (7), $Z(\mathbf{S})$ is given in Theorem 2.1, and

$$\lambda_j(t|\mathbf{S}) = \frac{\int_{\max(t,T_j)}^{\infty} \kappa_{m_j+1}(e^{-f_N}\rho|y)\eta(dy)}{\int_{T_j}^{\infty} \kappa_{m_j}(e^{-f_N}\rho|y)\eta(dy)}$$
(13)

if $m_j > 0$; otherwise 0.

Proof. It follows from (11) that the posterior mean of μ |**y**, **S**, **T** is given by

$$\mathbb{E}[\mu_N^*(dy)|\mathbf{y}, \mathbf{S}, \mathbf{T}] = \kappa_1(e^{-f_N}\rho|y)\eta(dy) + \sum_{\{j:m_j>0\}} \mathbb{E}[Q_j|y_j]\delta_{y_j}(dy),$$

where

$$\mathbb{E}[Q_j|y_j] = \frac{\kappa_{m_j+1}(\mathrm{e}^{-f_N}\rho|y_j)}{\kappa_{m_j}(\mathrm{e}^{-f_N}\rho|y_j)}.$$

This implies that the posterior mean of the decreasing hazard rate $\lambda(t)|\mathbf{y}, \mathbf{S}, \mathbf{T}$ is given by

$$\mathbb{E}[\lambda(t)|\mathbf{y}, \mathbf{S}, \mathbf{T}] = \int_{\mathcal{R}} \mathbb{I}(t < y) \kappa_1(e^{-f_N}\rho|y)\eta(dy) + \sum_{\{j:m_j>0\}} \mathbb{I}(t < y_j)\mathbb{E}[Q_j|y_j].$$

Marginalizing over (\mathbf{S}, \mathbf{y}) gives the result. \Box

Remark 2.2. It is convenient to view $\eta_j(dy_j|\mathbf{S}, \mathbf{T})$, given by (9), as a "posterior" distribution of y_j given the cluster of maximum element j and with number of elements m_j , where $\eta(dy_j)$ is the "prior" and $\kappa_{m_j}(e^{-f_N}\rho|y_j)\mathbb{I}(T_j < y_j)$ is the "likelihood" of the data given y_j .

Remark 2.3. As discussed in Section 4 of James (2004), the multiplicative intensity model captures a large variety of models that appear in event history analysis. Bayesian analysis for models under different censoring schemes, such as left truncation together with right censorship, and random censoring at different time points, follow similarly as the likelihoods differ slightly from (6) [see Andersen, Borgan, Gill and Keiding (1993, Section III.2) and James (2004)].

Remark 2.4. Often, a machine part is likely to fail in its early period, and then the failure rate decreases gradually to a constant. Later on, the machine becomes more likely to fail again. Estimation for a "bathtub" or unimodal hazard rate is an important statistical problem. A posterior characterization via **S**-paths also appears in Bayes estimation of a "bathtub" or U-shaped hazard rate with a minimum at a, that can be modeled by [Ishwaran and James (2004)]

$$\lambda(t|a,\mu) = \int \mathbb{I}(|t-a| \ge u) \mu(du).$$

The posterior distribution of (a, μ) can then be jointly described by a posterior distribution of a and a posterior distribution of $\mu|a$, where the latter follows naturally as a path characterization for a fixed a. It turns out that the posterior distribution of the (scalar) location parameter a can also be described in terms of finite sums over **S**-paths; see Ho (2005b) for the Bayes estimation of a similarly defined semi-parametric family of asymmetric unimodal densities with mode at zero, which is parametrized by a scalar parameter and a mixing distribution.

2.1 Rao-Blackwellization – relation between S-path and partition

By definition, an **S**-path carries less information than a partition \mathbf{p} in clustering a set of n integers; an **S**-path records in the form of a jump, in each cluster, only the number of elements and the maximum elements, but not the remaining elements that a partition also takes into account. As such, an **S**-path corresponds to possibly many partitions; therefore, the space of **S**-paths of n + 1 coordinates is much coarser/smaller than that of partitions of n integers for a moderate n (see Table 1 for illustration). Given an **S**-path of n + 1 coordinates, denote $\mathbb{C}_{\mathbf{S}}$ as a collection of partitions $\mathbf{p} = \{C_1, \ldots, C_{n(\mathbf{p})}\}$ of integers $\{1, \ldots, n\}$ that correspond to the path **S**, in the sense that (i) the number of clusters is identical to the number of positive increments, that is, $n(\mathbf{S}) = n(\mathbf{p})$, and (ii) **p** has a cluster C_i of e_i number of elements, and of a maximum j if and only if **S** has a jump at location j of an increment $m_j = e_i$. The total number of these partitions [due to Dr. Peisen Zhang; see the proof of Lemma 2.1 in Brunner and Lo (1989)] is given by

$$|\mathbb{C}_{\mathbf{S}}| = \prod_{\{j:m_j>0\}} \left(\begin{array}{c} j-1-S_{j-1}\\ j-S_j \end{array}\right).$$

This number, appearing in the posterior distribution $Z(\mathbf{S})$, plays an important role in deriving the **S**-path characterization [see the proof of Theorem 2.1 in the appendix]. Furthermore, it yields the following statistically important (posterior) structure in model (2).

Lemma 2.1. Suppose $S|T \sim Z(S)$. Then, there exists a conditional distribution

$$\pi(\mathbf{p}|\mathbf{S},\mathbf{T}) = \frac{1}{|\mathbb{C}_{\mathbf{S}}|}, \qquad \mathbf{p} \in \mathbb{C}_{\mathbf{S}}.$$
(14)

That is, $\mathbf{p}|\mathbf{S}, \mathbf{T}$ is uniformly distributed over the $|\mathbb{C}_{\mathbf{S}}|$ different partitions that correspond to the given \mathbf{S} .

A proof is available in the appendix. Concerning model (2), the path characterization of the posterior distribution of μ , stated in Theorem 2.1, is always preferred to the one in terms of partitions in two-folds:

- (a) the space of **S**-paths is coarser than that of partitions for a moderate size of number of complete observations; and
- (b) posterior estimates in terms of **S**-paths are always less variable than estimates in terms of partitions.

Provided that exact sampling from posterior distribution $Z(\mathbf{S})$ is available [Propp and Wilson (1996)], the Bayes hazard estimate (12) should be used instead of that in terms of partitions due to reason (b). Though exact sampling methods of neither partitions nor **S**-paths from the appropriate posterior distributions are available, reason (a) above on difference between complexities of spaces suggests that one might be able to design more efficient Markov chain Monte Carlo (MCMC) methods for sampling **S**-paths than existing MCMC methods for sampling partitions.

\overline{n}	# of paths	# of partitions	Ratio in $\%$
1	1	1	100.000
3	5	5	100.000
5	42	52	80.769
7	429	877	48.917
10	16,796	$115,\!975$	14.482
15	$9,\!694,\!845$	$1,\!382,\!958,\!545$	0.701
20	$6,\!564,\!120,\!420$	51,724,158,235,372	0.013

Table 1: Total numbers of paths and partitions versus sample sizes (n).

Remark 2.5. Lemma 2.1 extends an analogous Rao-Blackwellization result, established in an unpublished Ph.D. thesis of Ho (2002), regarding the specialized model considered by Lo and Weng (1989), as the same conditional law of $\mathbf{p}|\mathbf{S}, \mathbf{T}$, which does not depend on the data, applies.

Remark 2.6. Similar posterior structures related to **S**-paths and partitions do appear in an analogous problem of Bayes estimation of a monotone density, in which the monotone density on the half line is represented as a mixture of uniform densities and the mixing distribution is assumed to be a Dirichlet processes [Ferguson (1973)]. Results in Lo (1984) specialize

and yield a characterization for the posterior distribution in terms of partitions. The same reduction by **S**-paths and analogous Rao-Blackwellization results, as in the hazard rate estimation, exist in the monotone case. On one hand, a coarser characterization of the posterior distribution in terms of **S**-paths exists. On the other hand, the same discrete uniform distribution applies to the posterior (conditional) law of **p** given **S** and the data [Ho (2002)]. Recently, Ho (2005a) also obtained an analogous result by considering monotone densities defined by uniform densities convolved with the more general two-parameter Poisson-Dirichlet process [Pitman and Yor (1997)].

3 The Markov chain Monte Carlo method

This section introduces a MCMC path-sampler to approximate the monotone hazard estimate (12), and in general sums over **S**-paths. The Markov chain has a state space as the collection of **S**-paths of n+1 coordinates with a stationary distribution $Z(\mathbf{S})$. The algorithm is named as *accelerated path (AP) sampler*, which is designed to accelerate a straightforward Gibbs sampler in the sense that the algorithm allows more efficient movements among different **S**-paths.

A straightforward Gibbs sampler, which has a stationary distribution proportional to $\phi(\mathbf{S})$ [see equation (8)], can be defined [Ho (2002)]: Each Gibbs cycle consists of sampling $S_r|\mathbf{S}_{-r}$, where $\mathbf{S}_{-r} = (S_1, \ldots, S_{r-1}, S_{r+1}, \ldots, S_{n-1})$ is the "deleted-r" vector, and cycling through $r = 1, \ldots, n-1$. The conditional probabilities are

$$\Pr\{S_r = j | \mathbf{S}_{-r}\} \propto \phi(0, S_1, \dots, S_{r-1}, j, S_{r+1}, \dots, S_{n-1}, n),$$

for $j = S_{r-1}, S_{r-1} + 1, \ldots, S_{r+1} - 1, S_{r+1}$. The conditional move of S_r given \mathbf{S}_{-r} only depends on the two most neighboring coordinates S_{r-1} and S_{r+1} . In case if $S_{r-1} = S_{r+1}$ at step r, S_r would remain unchanged (subject to the nondecreasing property of an \mathbf{S} -path). This retards the convergence of the chain to its equilibrium state, and thus results in poor approximations of averages of \mathbf{S} -paths. The above phenomenon motivates us to accelerate this naïve chain in accordance with an increasing number of possible movements among the state space within any step. Notice that each step of the Gibbs sampler is equivalent to re-determinations of the two increments m_r and m_{r+1} at locations r and r+1, respectively. The idea is to replace m_{r+1} by some other m_q such that, at any step, it is relatively less likely that the resulting chain is bounded to remain unchanged by the definition of \mathbf{S} -paths. [See Remark 3.1.]

The accelerated path sampler, like the Gibbs sampler, consists of n-1 steps within one cycle. Suppose q > r denotes the next location (to the right of r) that the path jumps.

Algorithm 3.1 (Accelerated path sampler). The accelerated path (AP) sampler is a MCMC scheme that samples a Markov chain of S-paths of n + 1 coordinates which has a distribution $Z(\mathbf{S}) = \phi(\mathbf{S}) / \sum_{\mathbf{S}} \phi(\mathbf{S})$. A cycle of it is defined by two steps:

(i) At the r-th step of a cycle, suppose $\mathbf{S}^* = (0, S_1, \dots, S_{r-1}, c, \dots, c, S_q, \dots, S_{n-1}, n)$ where $S_{r-1} \leq c \leq \min(r, S_q - 1)$. \mathbf{S}^* moves to $(0, S_1, \dots, S_{r-1}, j, \dots, j, S_q, \dots, S_{n-1}, n)$ with conditional probability proportional to

$$\frac{r - S_{r-1}}{S_q - 1 - S_{r-1}} \int_{T_q}^{\infty} \kappa_{S_q - S_{r-1}} (e^{-f_N} \rho | y) \eta(dy),$$
(15)

if $j = S_{r-1}$; otherwise, if $j \in \{S_{r-1} + 1, \dots, \min(r, S_q - 1)\}$, with probability proportional to

$$\begin{pmatrix}
S_{q} - S_{r-1} - 2 \\
S_{q} - j - 1
\end{pmatrix} \prod_{i=r+1}^{q-1} \left(\frac{i-j}{i-S_{r-1}}\right)$$

$$\times \int_{T_{r}}^{\infty} \kappa_{j-S_{r-1}} \left(e^{-f_{N}}\rho|y\rangle\eta(dy) \times \int_{T_{q}}^{\infty} \kappa_{S_{q}-j} \left(e^{-f_{N}}\rho|z\rangle\eta(dz),\right)$$
(16)

(ii) repeat steps r = 1, 2, ..., n - 1 to complete a cycle.

Starting with an arbitrary path $\mathbf{S}^{(0)}$, and repeating M cycles according to the above scheme give a Markov chain $\mathbf{S}^{(0)}, \mathbf{S}^{(1)}, \ldots, \mathbf{S}^{(M)}$ with a unique stationary distribution $Z(\mathbf{S})$, the ergodic average

$$\widehat{\lambda}_M(t) = \int_t^\infty \kappa_1(e^{-f_N}\rho|y)\eta(dy) + \frac{1}{M}\sum_{i=1}^M\sum_{j=1}^n \lambda_j(t|\mathbf{S}^{(i)})$$
(17)

approximates the hazard estimate (12).

10

Remark 3.1. At step r of a cycle, the re-determinations of m_r and m_q can be viewed as moving the chain by assigning a single value to every element in the whole block/subvector $(S_r, S_{r+1}, \ldots, S_{q-1})$, subject to the definition of **S**-paths. This "block" change is more favorable to a "single" change of only S_r in the Gibbs sampler, in terms of efficiency of movements among different **S**-paths, as it is relatively less likely that $S_q = S_{r-1} + 1$ happens at step rso that our chain is bound to remain unchanged too often.

The validity of the AP sampler is justified by an idea in Hastings (1970) [see also Tierney (1994) and Gelfand and Smith (1990)]. One could always define a sequence of reducible transition kernels $\mathbb{P}^{(r)}$, r = 1, 2, ..., d, that all have the target stationary distribution. Multiplying them in series gives a transition kernel $\mathbb{P} = \mathbb{P}^{(1)} \times \mathbb{P}^{(2)} \times \cdots \times \mathbb{P}^{(d)}$ with the target stationary distribution (from construction). If the chain defined by \mathbb{P} is irreducible as all states communicate, the target stationary distribution will be unique. In the case of the AP sampler, we have d = n - 1. At each step r, kernel $\mathbb{P}^{(r)}$ is defined by the probability that the chain moves from the path after step r-1, $\mathbf{S}_0 = (0, S_1, \dots, S_{r-1}, c, \dots, c, S_q, \dots, S_{n-1}, n)$, to $\mathbf{S}^* = (0, S_1, \dots, S_{r-1}, j, \dots, j, S_q, \dots, S_{n-1}, n)$ for $j \in \{S_{r-1}, S_{r-1}+1, \dots, \min(r, S_q-1)\}$. The probability is proportional to $\phi(\mathbf{S}^*)$. That is, \mathbf{S}_0 communicates only with paths in the collection defined by $\{\mathbf{S} : \mathbf{S} = (0, S_1, \dots, S_{r-1}, j, \dots, j, S_q, \dots, S_{n-1}, n), j \in \{S_{r-1}, S_{r-1} + 1, \dots, \min(r, S_q - 1)\}\}$. With this construction, kernel $\mathbb{P}^{(r)}$ decomposes the state space of all \mathbf{S} -paths of n + 1 coordinates into a finite collection of mutually exclusive communication classes [see Theorem 3 in Feller (1968), P.392]. One can easily check that each kernel $\mathbb{P}^{(r)}$, though not irreducible, has a stationary distribution $Z(\mathbf{S})$. In addition, the chain defined by $\mathbb{P} = \mathbb{P}^{(1)} \times \mathbb{P}^{(2)} \times \cdots \times \mathbb{P}^{(n-1)}$ is irreducible, as all states can communicate with the path $\mathbf{S} = (0, 0, \dots, 0, n)$ within one cycle. Hence, the AP sampler gives a Markov chain of \mathbf{S} -paths with a unique stationary distribution $Z(\mathbf{S})$ in the space of all \mathbf{S} -paths of n + 1 coordinates.

4 Examples

One can model μ in (2) by a variety of completely random measures. Corresponding posterior analysis follows from the results in the previous sections. This section looks at two explicit examples of posterior distribution of μ where μ is characterized by the mean measure,

$$\rho_{\alpha,\beta}(dz|u)\eta(du) = \frac{1}{\Gamma(1-\alpha)} z^{-\alpha-1} \exp[-z/\beta(u)] dz \,\eta(du).$$
(18)

This class of random measures generalizes the generalized gamma random measure proposed by Brix (1999), for $0 < \alpha < 1$ and $0 \le \beta < \infty$ or $-\infty < \alpha \le 0$ and $0 < \beta < \infty$. Random measures defined by (18) include the gamma process (when $\alpha = 0$), a stable law with index $0 < \alpha < 1$ (when $\beta = \infty$), and the inverse-Gaussian process (when $\alpha = 1/2$ and $\beta > 0$).

4.1 The weighted gamma random measure

If $\alpha = 0$ in (18), μ is the weighted gamma random measure with shape measure η and scale measure β . Corollary 2.1 gives the Bayes estimate of the decreasing hazard rate (2), of which

$$\kappa_i(e^{-f_N}\rho|u) = (i-1)! \times \left[\beta(u)^{-1} + g_N(u)\right]^{-i}, \qquad i = 1, 2, \dots, n.$$
(19)

An implementation of the AP sampler relies on conditional probabilities (15) and (16) being proportional to

$$(r - S_{r-1}) \times \xi_{S_q - S_{r-1}}(T_q | \mathbf{T}),$$

and

$$\left[\prod_{i=r+1}^{q-1} \left(\frac{i-j}{i-S_{r-1}}\right)\right] \times \xi_{j-S_{r-1}}(T_r | \mathbf{T}) \times \xi_{S_q-j}(T_q | \mathbf{T}),$$

respectively, where $\xi_i(t|\mathbf{T}) = \int_t^\infty \left[\beta(v)^{-1} + g_N(v)\right]^{-i} \eta(dv), i = 1, \dots, n.$

4.2 The stable law

The stable law with index $0 < \alpha < 1$ appears when $\beta = \infty$ in (18). The posterior mean of the decreasing hazard rate (2) defined by this class of random measures follows from Corollary 2.1 and it can be evaluated by implementing the AP sampler based on

$$\kappa_i(e^{-f_N}\rho|y) = \frac{\Gamma(i-\alpha)}{\Gamma(1-\alpha)[g_N(y)]^{i-\alpha}}, \qquad i = 1, 2, \dots, n$$

5 Numerical results

This section concerns the effectiveness of the proposed AP sampler for evaluation of Bayes decreasing hazard estimate (12). In particular, a special case of the Bayes decreasing hazard estimate (12) according to a gamma process, which follows from Section 4.1 with $\beta(\cdot) = 1$, is selected. The posterior mean reduces to

$$\mathbb{E}[\lambda(t|\mu)|\mathbf{T}] = \xi_1(t|\mathbf{T}) + \sum_{\mathbf{S}} Z^*(\mathbf{S}) \sum_{j=1}^n \left[m_j \frac{\xi_{m_j+1}(\max(t, T_j)|\mathbf{T})}{\xi_{m_j}(T_j|\mathbf{T})} \right],\tag{20}$$

where $Z^*(\mathbf{S}) \propto \prod_{\{j:m_j>0\}} (j-1-S_{j-1})!/(j-S_j)! \times \xi_{m_j}(T_j|\mathbf{T})$. The complexity in evaluating $\xi_j(t|\mathbf{T})$ can be reduced by assuming a uniform shape probability from 0 to $6(\geq \tau)$ for $\eta(\cdot)$, even though the closed-form expression is tedious [see Ho and Lo (2001) for its exact expression]. The methodology is tested by data from a piecewise constant hazard rate model, of which the hazard rate of an item is

$$\lambda(t) = \begin{cases} 1 & 0 \le t < 1\\ 0.5 & t \ge 1. \end{cases}$$
(21)

Data are generated subject to a termination time $\tau = 3$, such that the censoring rate is about 15%. All simulation results that follow are based on the following settings: The Monte Carlo size is M = 1000, and the initial path is set at $\mathbf{S}^{(0)} = (0, 1, \dots, n-1, n)$.

Remark 5.1. In practice, the implementation of the AP sampler depends heavily on evaluations of double integrals,

$$\int_{y}^{\infty} \int_{\mathcal{R}} z^{i} e^{-g_{N}(u)z} \rho(dz|u) \eta(du), \qquad y > 0, \quad i = 1, 2, \dots, n$$

where, according to (5),

$$g_N(u) = \begin{cases} \sum_{i=1}^{j-1} T_i + (N-j+1)u, & T_{j-1} < u \le T_j, j = 1, \dots, n+1\\ \sum_{i=1}^n T_i + (N-n)\tau, & u > \tau, \end{cases}$$

12

is a piecewise linear function of u. Apart from the specific gamma process chosen in this simulation study, here we suggest at least another two random measures that result in practically convenient computations. They are a weighted gamma process with a uniform η and a scale measure $\beta(u) = 1/u$, and a stable law with index $0 < \alpha < 1$ together with a uniform η , under the context of Section 4.1 and Section 4.2, respectively. In both cases, the double integrals reduce to closed-form and piecewise expressions similar to that in the gamma case.

5.1 Resolution of the AP sampler

This section focuses on the convergence property of the hazard estimate (20) approximated by the AP sampler as the sample size N increases. Based on nested samples of sizes N = 100, 500, and 1000, MCMC estimates (17) according to (19) are displayed in Figure 1. The graphs echo the fact that the approximated posterior mean of the decreasing hazard function, $\hat{\lambda}_M(t)$, tends to the "true" hazard rate (21) as sample size increases. We remark that the drop of the hazard estimates after t = 3 results from the fact that the estimates are mainly constructed based on the prior information owing to no complete data observed after that time point.



Figure 1: MCMC hazard estimates $\hat{\lambda}_{1000}(t)$ produced by the accelerated path sampler (with no "burn-in" or "warm-up" period, and Monte Carlo size M = 1000) of the posterior mean (20) based on data generated from model (21).

5.2 Comparison with other methods

Path-sum Bayes estimates of monotone hazard rates, though they appeared two decades ago, have not received much attention and are not commonly used, probably due to un14

availability of efficient numerical methods. On the contrary, corresponding partition-sum monotone hazard estimates have been used as a substitute [Hayakawa, Zukerman, Paul and Vignaux (2001) and Ho and Lo (2001)] as there are many well-developed numerical methods for sampling partitions, such as weighted Chinese restaurant algorithms [Lo, Brunner and Chan (1996) and Ishwaran and James (2003)], and blocked Gibbs sampling methods [Ishwaran and James (2001, 2004)]. This section aims at comparing MCMC hazard estimates produced by the AP sampler, the Gibbs path sampler defined in Ho (2002), and a Gibbs sampler for partitions introduced by MacEachern (1994), which is a special case of the gWCR sampler in Lo *et al.* (1996). See Ho and Lo (2001) for an exact description of the algorithm being applied to this gamma model. Standard errors of MCMC hazard estimates by the three different methods are estimated by repetitions of experiment, and are used as the standard of comparison.

Here, the sample size is fixed at N = 100, and there are n = 82 complete observations in our simulated data set. Markov samples from all the three different Markov chain experiments are collected after a "burn-in" or "warm-up" period of 10000 cycles. By producing 1000 repetitions of each experiment, we have 1000 independent hazard estimates by each of the methods. These are used to estimate the average and the standard error of the hazard estimates by the three corresponding MCMC methods in the usual manner. The hazard rates $\lambda(t)$ at time t = 0.5, 0.99, 1.01, and 2.0, are studied. The points, 0.99 and 1.01, are near to the change point at 1, and they seem to reflect well the effectiveness and the efficiency of the MCMC methods in the worst case. Table 2 displays the averages and the standard errors of 1000 realizations of MCMC hazard estimates produced by the three methods. At different time points, the three averages are close to each other, yet the standard errors vary substantially. The standard error of hazard estimate produced by the AP sampler is the smallest among the three. On one hand, the AP sampler definitely outweighs the naive Gibbs path sampler. This shows that our "acceleration" scheme works extremely well. On the other hand, the AP sampler beats the closest competitor, the gWCR sampler, by a comfortable margin.

6 Proportional Hazards

The Cox regression model [Cox (1972)] is an important example of a multiplicative intensity model that can allow incorporation of covariates, together with right independent censoring, in survival analysis. In particular, if we assume that the underlying hazard defined on the half line \mathcal{R} is modeled by

$$\lambda(t | \mathbf{Z}, \boldsymbol{\theta}, \mu) = \int_{\mathcal{R}} \exp(\boldsymbol{\theta}^T \mathbf{Z}) \mathbb{I}(t < u) \mu(du),$$

t	MCMC method	average of	standard error of
		hazard estimates	hazard estimates
		at time t	at time t
	AP	0.9667340	0.0038426
0.5	gP	0.9677822	0.0517161
	gWCR	0.9668593	0.0080103
	AP	0.8815065	0.0065156
0.99	gP	0.8812967	0.0594452
	gWCR	0.8820966	0.0097827
	AP	0.8530503	0.0067767
1.01	$_{\rm gP}$	0.8524295	0.0541091
	gWCR	0.8537771	0.0099268
	AP	0.3708132	0.0055500
2.0	$_{\rm gP}$	0.3692810	0.0440281
	gWCR	0.3707660	0.0106327

Table 2: A large-sample study of MCMC hazard estimates based on N = 100 samples (of which n = 82 are completely observed) from model (21) according to 1000 independent replications of the accelerated path (AP) sampler, of the Gibbs path (gP) sampler, and of the weighted Chinese restaurant process (gWCR) sampler. All estimates are evaluated based on Markov chains run with settings: 10000 cycles of "warm-up" period, and Monte Carlo size M = 1000.

where **Z** is a covariate vector with parameter vector $\boldsymbol{\theta}$, and $\lambda_0(t|\mu) = \int_{\mathcal{R}} \mathbb{I}(t < u)\mu(du)$, same as (2), is a decreasing baseline hazard rate. Suppose we observe data until time τ and the data $\mathbf{D} = ((T_1, \mathbf{Z}_1), \dots, (T_N, \mathbf{Z}_N))$ can be summarized as failure times $T_1 < \dots < T_n$ and right-censored times $T_i = \tau$, $i = n + 1, \dots, N$, associated with covariate vector \mathbf{Z}_i , $i = 1, \dots, N$ with unknown parameter vector $\boldsymbol{\theta}$. Define $f_{N,\boldsymbol{\theta}}(z, u) = g_{N,\boldsymbol{\theta}}(u)z$, where

$$g_{N,\boldsymbol{\theta}}(u) = \int_{0}^{\tau} \left[\sum_{i=1}^{N} \mathbb{I}(T_{i} \ge t) \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{i}) \right] \mathbb{I}(t < u) dt$$
$$= \begin{cases} \sum_{i=1}^{j-1} \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{i}) T_{i} + \left[\sum_{i=j}^{N} \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{i}) \right] u, & T_{j-1} < u \le T_{j}, j \le n+1 \\ \sum_{i=1}^{n} \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{i}) T_{i} + \left[\sum_{i=n+1}^{N} \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{i}) \right] \tau, & u > \tau \end{cases}$$
(22)

Then, the Cox proportional hazards likelihood may be written as

$$\left[\prod_{i=1}^{n} \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{i}) \lambda_{0}(T_{i}|\boldsymbol{\mu})\right] \exp\left[-\boldsymbol{\mu}(g_{N,\boldsymbol{\theta}})\right], \qquad (23)$$

where $\mu(g_{N,\boldsymbol{\theta}}) = \int_{\mathcal{R}} g_{N,\boldsymbol{\theta}}(u)\mu(du) = \int_{0}^{\tau} \left[\sum_{i=1}^{N} \mathbb{I}(T_{i} \ge t) \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{i})\right] \lambda_{0}(t|\mu) dt.$

Proposition 6.1. Suppose the likelihood of the data is given by (23). Let $\pi(d\theta)$ denote our prior for θ and independently assume μ is a completely random measure characterized by the Laplace functional (4). Then, the posterior distribution of $\mu|\theta$, **D** can be described as an analogous three-step hierarchical experiment of that in Theorem 2.1, of which $f_N(\cdot, \cdot)$ and $g_N(\cdot)$ are replaced by $f_{N,\theta}(\cdot, \cdot)$ and $g_{N,\theta}(\cdot)$, respectively.

To evaluate any posterior quantities of model (23), such as the posterior mean of the underlying monotone baseline hazard rate and the posterior mean of the covariate parameters θ , run the following Gibbs sampler:

- 1. Draw $\mathbf{S}|\mathbf{Q}, \mathbf{y}, \boldsymbol{\theta}, \mathbf{D}$ by implementing Algorithm 3.1 with $f_N(\cdot, \cdot)$ and $g_N(\cdot)$ replaced by $f_{N,\boldsymbol{\theta}}(\cdot, \cdot)$ and $g_{N,\boldsymbol{\theta}}(\cdot)$, respectively.
- 2. Draw $\mathbf{Q}, \mathbf{y} | \mathbf{S}, \boldsymbol{\theta}, \mathbf{D}$ according to the analogues of the conditional distributions (9) and (10) in Theorem 2.1 with $f_N(\cdot, \cdot)$ and $g_N(\cdot)$ replaced by $f_{N,\boldsymbol{\theta}}(\cdot, \cdot)$ and $g_{N,\boldsymbol{\theta}}(\cdot)$, respectively.
- 3. Draw $\theta | \mathbf{Q}, \mathbf{y}, \mathbf{S}, \mathbf{D}$ from the density proportional to

$$\pi(d\boldsymbol{\theta})B(\boldsymbol{\theta})\prod_{\{j:m_j>0\}} e^{-g_{N,\boldsymbol{\theta}}(y_j)Q_j},$$

where $B(\boldsymbol{\theta})$ is given by (31).

Appendix

Proof of Theorem 2.1 and Lemma 2.1

Proof. The proof relies on the following three key points.

(i) Suppose p is a partition (of integers {1,...,n}) that corresponds to a path S (of n+1 coordinates). We have the following identity,

$$\prod_{i=1}^{n(\mathbf{p})} \mathbb{I}(\max_{j \in C_i} T_j < v_i) = \prod_{\{j: m_j > 0\}} \mathbb{I}(T_j < y_j), \qquad \mathbf{p} \in \mathbb{C}_{\mathbf{S}},$$
(24)

where $\mathbf{v} = (v_1, \dots, v_{n(\mathbf{p})})$ represents the distinct values of (u_1, \dots, u_n) in (6) and $\mathbf{y} = \{y_j : m_j > 0, j = 1, \dots, n\}$ is a permutation of \mathbf{v} according to $\mathbf{p} \in \mathbb{C}_{\mathbf{S}}$. (ii) The total number of partitions that correspond to a given **S** equals

$$\sum_{\mathbf{p}\in\mathbb{C}_{\mathbf{S}}} 1 = |\mathbb{C}_{\mathbf{S}}|,\tag{25}$$

where $\sum_{\mathbf{p}\in\mathbb{C}_{\mathbf{S}}}$ represents summing over all partitions corresponding to the path \mathbf{S} .

- (iii) The posterior law of μ |**T** in terms of partitions obtained by James (2002, 2004), specialized in our model, can be described as a three-step experiment:
 - (a) A partition **p** has a discrete posterior distribution $W(\mathbf{p}) = \zeta(\mathbf{p}) / \sum_{\mathbf{p}} \zeta(\mathbf{p})$, where $\zeta(\mathbf{p}) = \prod_{i=1}^{n(\mathbf{p})} \int_{\max_{j \in C_i} T_j}^{\infty} \kappa_{e_i} (e^{-f_N} \rho | y) \eta(dy)$ and $\sum_{\mathbf{p}}$ represent summing over all partitions **p**.
 - (b) Given \mathbf{p} , there exist $n(\mathbf{p})$ independent pairs of (v_i, J_i) , denoted by $(\mathbf{v}, \mathbf{J}) = \{(v_1, J_1), \dots, (v_{n(\mathbf{p})}, J_{n(\mathbf{p})})\}$, where for $i = 1, \dots, n(\mathbf{p})$, $v_i | \mathbf{p}, \mathbf{T}$ has a distribution proportional to $\mathbb{I}(\max_{j \in C_i} T_j < v_i) \kappa_{e_i}(e^{-f_N}\rho | v_i) \eta(dv_i)$, and $\Pr\{J_i \in dz | \mathbf{p}, v_i, \mathbf{T}\} = z^{e_i} e^{-g_N(v_i)z} \rho(dz | v_i) / \kappa_{e_i}(e^{-f_N}\rho | v_i)$.
 - (c) Given $(\mathbf{p}, \mathbf{v}, \mathbf{J})$ and the data \mathbf{T} , μ has a distribution as $\mu_N^* + \sum_{i=1}^{n(\mathbf{p})} J_i \delta_{v_i}$, where μ_N^* is defined in Theorem 2.1.

The three-step experiment in (iii) states that the posterior law of $\mu | \mathbf{T}$ is equivalent to the distribution of a random measure $\mu_N^* + \sum_{i=1}^{n(\mathbf{p})} J_i \delta_{v_i}$, which is determined by the joint conditional law of μ_N^* , $\mathbf{J}, \mathbf{v}, \mathbf{p} | \mathbf{T}$. The conditional distribution of μ_N^* , $\mathbf{J}, \mathbf{v}, \mathbf{p} | \mathbf{T}$ is proportional to

$$\mathbb{P}(d\mu_N^*) \prod_{i=1}^{n(\mathbf{p})} J_i^{e_i} e^{-g_N(v_i)J_i} \rho(dJ_i|v_i) \prod_{i=1}^{n(\mathbf{p})} \mathbb{I}(\max_{j \in C_i} T_j < v_i) \eta(dv_i),$$
(26)

where $\mathbb{P}(d\mu_N^*)$ is a completely random measure characterized by an intensity measure $e^{-g_N(u)z}\rho(dz|u)\eta(du)$.

Notice that (26), due to its irrelevance to the remaining members other than the maximum elements of the clusters in a partition, may be rewritten in terms of the intrinsic characteristics of a path **S** provided that $\mathbf{p} \in \mathbb{C}_{\mathbf{S}}$ based on (24) as

$$\mathbb{P}(d\mu_N^*) \prod_{\{j:m_j>0\}} Q_j^{m_j} e^{-g_N(y_j)Q_j} \rho(dQ_j|y_j) \prod_{\{j:m_j>0\}} \mathbb{I}(T_j < y_j)\eta(dy_j),$$
(27)

where $\mathbf{Q} = \{Q_j : m_j > 0, j = 1, ..., n\}$ is a relabelling of \mathbf{J} according to the correspondence $\mathbf{p} \in \mathbb{C}_{\mathbf{S}}$. That is, the conditional law of $\mu_N^*, \mathbf{J}, \mathbf{v}, \mathbf{p} | \mathbf{T}$ only depends on \mathbf{p} through \mathbf{S} -paths.

Additionally, we have the following equivalence in distribution relation between the two random measures,

$$\mathcal{L}\left\{ \left. \mu_{N}^{*} + \sum_{i=1}^{n(\mathbf{p})} J_{i} \delta_{v_{i}} \right| \mathbf{T} \right\} \stackrel{d}{=} \mathcal{L}\left\{ \left. \mu_{N}^{*} + \sum_{\{j:m_{j}>0\}} Q_{j} \delta_{y_{j}} \right| \mathbf{T} \right\}.$$
(28)

Equations (27) and (28) imply that the posterior law of $\mu | \mathbf{T}$ can be expressed in terms of $\mu_N^* | \mathbf{Q}, \mathbf{y}, \mathbf{S}, \mathbf{T}$ and $\mathbf{Q}, \mathbf{y}, \mathbf{S} | \mathbf{T}$. Note that the distribution of $\mathbf{Q}, \mathbf{y}, \mathbf{S} | \mathbf{T}$ is determined by integrating out μ_N^* and then summing over all $\mathbf{p} \in \mathbb{C}_{\mathbf{S}}$ in (27), and thus it is proportional to

$$\prod_{\{j:m_j>0\}} Q_j^{m_j} e^{-g_N(y_j)Q_j} \rho(dQ_j|y_j) \prod_{\{j:m_j>0\}} \mathbb{I}(T_j < y_j)\eta(dy_j) \left[\sum_{\mathbf{p}\in\mathbb{C}_{\mathbf{S}}} 1\right],$$
(29)

where $\sum_{\mathbf{p}\in\mathbb{C}_{\mathbf{S}}} 1 = |\mathbb{C}_{\mathbf{S}}|$ by (25). Now, the laws of $\mathbf{Q}|\mathbf{S}, \mathbf{y}, \mathbf{T}, \mathbf{y}|\mathbf{S}, \mathbf{T}$, and $\mathbf{S}|\mathbf{T}$ follow from Bayes' theorem and multiplication rule. Hence, the result in Theorem 2.1 follows.

The conditional distribution of $\mathbf{p}|\mathbf{S}, \mathbf{T}$ in Lemma 2.1 is obtained by dividing (26) without the leading term $\mathbb{P}(d\mu_N^*)$ by (29). \Box

Proof of Proposition 6.1

Proof. Following from the same arguments as in James (2004), the posterior law of $\mu | \mathbf{D}$ is equivalent to the distribution of a random measure $\mu_N^* + \sum_{i=1}^{n(\mathbf{p})} J_i \delta_{v_i}$. It is determined by the joint distribution of $\mu_N^*, \mathbf{J}, \mathbf{v}, \mathbf{p}, \boldsymbol{\theta} | \mathbf{D}$, which is proportional to

$$\mathbb{P}(d\mu_N^*)\pi(d\boldsymbol{\theta})B(\boldsymbol{\theta})\prod_{i=1}^{n(\mathbf{p})}J_i^{e_i}e^{-g_{N,\boldsymbol{\theta}}(v_i)J_i}\rho(dJ_i|v_i)\prod_{i=1}^{n(\mathbf{p})}\mathbb{I}(\max_{j\in C_i}T_j < v_i)\eta(dv_i),$$
(30)

where $\mathbb{P}(d\mu_N^*)$ is a completely random measure characterized by an intensity measure $e^{-g_{N,\theta}(u)z}\rho(dz|u)\eta(du)$, and

$$B(\boldsymbol{\theta}) = \exp\left[-\int_{\mathcal{R}} \int_{\mathcal{R}} \left(1 - e^{-g_{N,\boldsymbol{\theta}}(u)z}\right) \rho(dz|u)\eta(du)\right] \prod_{i=1}^{n} \exp(\boldsymbol{\theta}^{T} \mathbf{Z}_{j}).$$
(31)

As in the proof of Theorem 2.1, $\mu_N^* | \mathbf{Q}, \mathbf{y}, \mathbf{S}, \boldsymbol{\theta}, \mathbf{D}$ is equivalent to $\mathbb{P}(d\mu_N^*)$. Summing over all $\mathbf{p} \in \mathbb{C}_{\mathbf{S}}$ in (30) yields that the conditional distribution of $\mathbf{Q}, \mathbf{y}, \mathbf{S}, \boldsymbol{\theta} | \mathbf{D}$ is proportional to

$$\pi(d\boldsymbol{\theta})B(\boldsymbol{\theta})\prod_{\{j:m_j>0\}}Q_j^{m_j}e^{-g_{N,\boldsymbol{\theta}}(y_j)Q_j}\rho(dQ_j|y_j)\prod_{\{j:m_j>0\}}\mathbb{I}(T_j < y_j)\eta(dy_j) \times |\mathbb{C}_{\mathbf{S}}|.$$
(32)

Ho

Hence, by Bayes' theorem and multiplication rule, the result follows from the conditional dis-

tribution of $\mathbf{Q}, \mathbf{y}, \mathbf{S}|\boldsymbol{\theta}, \mathbf{D}$, which is proportional to (32) without the leading term $\pi(d\boldsymbol{\theta})B(\boldsymbol{\theta})$.

References

- AALEN, O. O. (1975) Statistical inference for a family of counting processes. Unpublished Ph.D. thesis. University of California, Berkeley.
- [2] AALEN, O. O. (1978) Nonparametric inference for a family of counting processes. Ann. Statist., 6, 701–726.
- [3] ANDERSEN, P. K., BORGAN, O., GILL, R. D. AND KEIDING, N. (1993) Statistical Models Based On Counting Processes, Springer-Verlag, New York.
- [4] BARLOW, R. E., BARTHOLOMEW, D. J., BREMNER, J. M. AND BRUNK, H. D. (1972). Statistical Inference Under Order Restrictions, John Wiley & Sons.
- [5] BRIX, A. (1999). Generalized gamma measures and shot-noise Cox processes. Adv. in Appl. Probab., 31, 929–953.
- [6] BRUNNER, L. J. (1995). Bayesian linear regression with error terms that have symmetric unimodal densities. Nonpara. Statist., 4, 335–348.
- [7] BRUNNER, L. J. AND LO, A. Y. (1989). Bayes methods for a symmetric unimodal density and its mode. Ann. Statist., 17, 1550–1566.
- [8] BRUNNER, L. J. AND LO, A. Y. (1994). Nonparametric Bayes methods for directional data. *Canadian J. Statist.*, 22, 401–412.
- [9] Cox, D. R. (1972). Regression models and life-tables (with discussion). J. Royal Statist. Soc., Series B, 34, 187–220.
- [10] DRĂGICHI, L. AND RAMAMOORTHI, R. V. (2003). Consistency of Dykstra-Laud priors. Sankhyā, 65, 464–481.
- [11] DYKSTRA, R. L. AND LAUD, P. (1981). A Bayesian nonparametric approach to reliability. Ann. Statist., 9, 356–367.
- [12] FELLER, W. (1968). An Introduction to Probability Theory and Its Applications 1, 3rd ed., John Wiley & Sons, New York.
- [13] FERGUSON, T. S. (1973). A Bayesian analysis of some nonparametric problems. Ann. Statist., 1, 209–230.
- [14] GELFAND, A. E. AND SMITH, A. F. M. (1990). Sampling based approaches to calculating marginal densities. J. Amer. Statist. Assoc., 85, 398–409.
- [15] GEMAN, A. AND GEMAN, D. (1984). Stochastic relaxation, Gibbs distributions, and Bayesian restoration of images. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 6, 721–741.

- [16] GRENANDER, U. (1956). On the theory of mortality measurement, Part II. Skand. Aktuar., 39, 125–153.
- [17] HALL, P., HUANG, L.-S., GIFFORD, J. A. AND GIJBELS, I. (2001). Nonparametric estimation of hazard rate under the constraint of monotonicity. J. Comp. Graph. Statist., 10, 592–614.
- [18] HASTINGS, W. K. (1970). Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57, 97–109.
- [19] HAYAKAWA, Y., ZUKERMAN, J., PAUL, S. AND VIGNAUX, T. (2001). Bayesian nonparametric testing of constant versus nondecreasing hazard rates. In System and Bayesian Reliability: Essays in Honor of Professor Richard E. Barlow on His 70th Birthday (eds: Y Hayakawa, T Irony and M Xie), 391–406, World Scientific.
- [20] Ho, M.-W. (2002). Bayesian inference for models with monotone densities and hazard rates. Unpublished Ph.D. thesis. Department of Information and Systems Management. Hong Kong University of Science and Technology.
- [21] Ho, M.-W. (2005a). Bayes estimation of a monotone density. Preprint.
- [22] Ho, M.-W. (2005b). A Bayes method for an asymmetric unimodal density with mode at zero. Preprint.
- [23] HO, M.-W. AND LO, A. Y. (2001). Bayesian nonparametric estimation of a monotone hazard rate. In System and Bayesian Reliability: Essays in Honor of Professor Richard E. Barlow on His 70th Birthday (eds: Y Hayakawa, T Irony and M Xie), 301–314, World Scientific.
- [24] HUANG, J. AND WELLNER, J. A. (1995). Estimation of a monotone density or a monotone hazard under random censoring. *Scand. J. Statist.*, 22, 3–33.
- [25] IBRAHIM, J. G., CHEN, M.-H. AND MACEACHERN, S. N. (1999). Bayesian variable selection for proportional hazards models. *Canadian J. Statist.*, 37, 701–717.
- [26] ISHWARAN, H. AND JAMES, L. F. (2001). Gibbs sampling methods for stick-breaking priors. J. Amer. Statist. Assoc., 96, 161–173.
- [27] ISHWARAN, H. AND JAMES, L. F. (2003). Generalized weighted Chinese restaurant processes for species sampling mixture models. *Statistica Sinica*, 13, 1211–1235.
- [28] ISHWARAN, H. AND JAMES, L. F. (2004). Computational methods for multiplicative intensity models using weighted gamma processes: proportional hazards, marked point processes and panel count data. J. Amer. Statist. Assoc., 99, 175–190.
- [29] JAMES, L. F. (2002). Poisson process partition calculus with applications to exchangeable models and Bayesian nonparametrics, arXiv:math.PR/0205093, 2002. Available at http://arxiv.org/abs/math.PR/0205093.
- [30] JAMES, L. F. (2003). Bayesian calculus for gamma processes with applications to semiparametric intensity models. Sankhyā, 65, 196–223.
- [31] JAMES, L. F. (2004). Bayesian Poisson process partition calculus with an application to Bayesian Lévy moving averages. Forthcoming in Ann. Statist..

- [32] KALBFLEISCH, J. D. (1978). Non-parametric Bayesian analysis of survival time data. J. Royal Statist. Soc., Series B, 40, 214–221.
- [33] KINGMAN, J. F. C. (1967). Completely random measures. Pacific Journal of Mathematics, 21, 59–78.
- [34] KINGMAN, J. F. C. (1993). Poisson Processes. Oxford University Press, Oxford.
- [35] Lo, A. Y. (1982). Bayesian nonparametric statistical inference for Poisson point process, Z. Wahrsch. Verw. Gebiete, 59, 55–66.
- [36] LO, A. Y. (1984). On a class of Bayesian nonparametric estimates: I. Density estimation. Ann. Statist., 12, 351–357.
- [37] LO, A. Y., BRUNNER, L. J. AND CHAN, A. T. (1996). Weighted Chinese restaurant processes and Bayesian mixture models, research report. ISMT Department, HKUST, Hong Kong, & Revision I (1998)
- [38] LO, A. Y. AND WENG, C. S. (1989). On a class of Bayesian nonparametric estimates: II. Hazard rate estimates. Ann. Instit. Statist. Math., 41, 227–245.
- [39] LO, S. H. AND PHADIA, E. (1992). On estimation of a survival function in reliability theory based on censored data. Preprint, Columbia University.
- [40] MACEACHERN, S. N. (1994). Estimating normal means with a conjugate style Dirichlet process prior. Commun. Statist.-Simula., 23, 727-741.
- [41] MYKYTYN, S. W. AND SANTNER, T. J. (1981). Maximum likelihood estimation of the survival function based on censored data under hazard rate assumptions. *Commun. Statist.-Theor. Meth.*, A11, 1369–1387.
- [42] PADGETT, W. J. AND WEI, L. J. (1980). Maximum likelihood estimation of a distribution function with increasing failure rate based on censored observations. *Biometrika*, 67, 470–474.
- [43] PITMAN, J. AND YOR, M. (1997). The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator. Ann. Probab., 25, 855–900.
- [44] PRAKASA RAO, B. L. S. (1970). Estimation of distributions with monotone failure rate. Ann. Math. Statist., 41, 507–519.
- [45] PROPP, J. AND WILSON, D. (1996). Exact sampling with coupled Markov chains and applications to statistical mechanics. *Random Structure and Algorithm*, 9, 223–252.
- [46] TIERNEY, L. (1994). Markov chains for exploring posterior distributions. Ann. Statist., 22, 1701–1762.
- [47] VILLALOBOS, M. AND WAHBA, G. (1987). Inequality-constrained multivariate smoothing splines with application to the estimation of posterior probabilities. J. Amer. Statist. Assoc., 82, 239–248.

22

MAN-WAI HO DEPARTMENT OF STATISTICS AND APPLIED PROBABILITY NATIONAL UNIVERSITY OF SINGAPORE 6 SCIENCE DRIVE 2 SINGAPORE 117546 REPUBLIC OF SINGAPORE E-mail: stahmw@nus.edu.sg