

# Isomorphic groupoid $C^*$ -algebras associated with different Haar systems

Mădălina Roxana Buneci\*  
University Constantin Brâncuși, Târgu-Jiu

## Abstract

We shall consider a locally compact groupoid endowed with a Haar system  $\nu$  and having proper orbit space. We shall associated to each appropriated cross section  $\sigma : G^{(0)} \rightarrow G^F$  for  $d_F : G^F \rightarrow G^{(0)}$  (where  $F$  is a Borel subset of  $G^{(0)}$  meeting each orbit exactly once) a  $C^*$ -algebra  $M_\sigma^*(G, \nu)$ . We shall prove that the  $C^*$ -algebras  $M_\sigma^*(G, \nu_i)$   $i = 1, 2$  associated with different Haar systems are  $*$ -isomorphic

AMS 2000 Subject Classification: 22A22, 43A22, 43A65, 46L99.

Key Words: locally compact groupoid,  $C^*$ -algebra,  $*$ -isomorphism.

## 1 Introduction

The  $C^*$ -algebra of a locally compact groupoid was introduced by J. Renault in [8]. The construction extends the case of a group: the space of continuous functions with compact support on groupoid is made into a  $*$ -algebra and endowed with the smallest  $C^*$ -norm making its representations continuous. In order to define the convolution on groupoid one needs to assume the existence of a Haar system which is an analogue of Haar measure on a group. Unlike the case for groups, Haar systems need not to be unique. A result of Paul Muhly, Jean Renault and Dana Williams establishes that the  $C^*$ -algebras of  $G$  associated with two Haar systems are strongly Morita equivalent (Theorem 2.8/p. 10 [3]). If the groupoid is transitive they have proved that if  $G$  is transitive then the  $C^*$ -algebra of  $G$  is isomorphic to  $C^*(H) \otimes \mathcal{K}(L^2(\mu))$ , where  $H$  is the isotropy group  $G_u^u$  at any unit  $u \in G^{(0)}$ ,  $\mu$  is an essentially unique measure on  $G^{(0)}$ ,  $C^*(H)$  denotes the group  $C^*$ -algebra of  $H$ , and  $\mathcal{K}(L^2(\mu))$  denotes the compact operators on  $L^2(\mu)$  (Theorem 3.1/p. 16 [3]). Therefore the  $C^*$ -algebras of a *transitive* groupoid  $G$  associated with two Haar systems are  $*$ -isomorphic.

In [7] Arlan Ramsay and Martin E. Walter have associated to a locally compact groupoid  $G$  a  $C^*$ -algebra denoted  $M^*(G, \nu)$ . They have considered the universal representation  $\omega$  of  $C^*(G, \nu)$  -the usual  $C^*$ -algebra associated to

---

\*This work was partly supported by the MEC-CNCSIS grant At127/2004 and by the Postdoctoral Training Program HPRN-CT-2002-0277.

a Haar system  $\nu = \{\nu^u, u \in G^{(0)}\}$  (constructed as in [8]). Since every cyclic representation of  $C^*(G, \nu)$  is the integrated form of a representation of  $G$ , it follows that  $\omega$  can be also regarded as a representation of  $\mathcal{B}_c(G)$ , the space of compactly supported Borel bounded function on  $G$ . Arlan Ramsay and Martin E. Walter have used the notation  $M^*(G, \nu)$  for the operator norm closure of  $\omega(\mathcal{B}_c(G))$ . Since  $\omega$  is an  $*$ -isomorphism on  $C^*(G, \nu)$ , we can regard  $C^*(G, \nu)$  as a subalgebra of  $M^*(G, \nu)$ .

We shall assume that the orbit space of the groupoid  $G$  is proper and we shall choose a Borel subset  $F$  of  $G^{(0)}$  meeting each orbit exactly once and such that  $F \cap [K]$  has a compact closure for each compact subset  $K$  of  $G^{(0)}$ . For each appropriated cross section  $\sigma : G^{(0)} \rightarrow G^F$  for  $d_F : G^F \rightarrow G^{(0)}$ ,  $d_F(x) = d(x)$ , we shall construct a  $C^*$ -algebra  $M_\sigma^*(G, \nu)$  which can be viewed as a subalgebra of  $M^*(G, \nu)$ . If  $\nu_1 = \{\nu_1^u, u \in G^{(0)}\}$  and  $\nu_2 = \{\nu_2^u, u \in G^{(0)}\}$  are two Haar systems on  $G$ , we shall prove that the  $C^*$ -algebras  $M_\sigma^*(G, \nu_1)$  and  $M_\sigma^*(G, \nu_2)$  are  $*$ -isomorphic.

For a transitive (or more generally, a locally transitive) groupoid  $G$  we shall prove that the  $C^*$ -algebras  $C^*(G, \nu)$ ,  $M^*(G, \nu)$  and  $M_\sigma^*(G, \nu)$  coincide.

For a principal proper groupoid  $G$ , we shall prove that

$$C^*(G, \nu) \subset M_\sigma^*(G, \nu) \subset M^*(G, \nu).$$

Let  $\pi : G^{(0)} \rightarrow G^{(0)}/G$  be the quotient map and let  $\nu_i = \{\varepsilon_u \times \mu_i^{\pi(u)}, u \in G^{(0)}\}$ ,  $i = 1, 2$  be two Haar systems on the principal proper groupoid  $G$ . We shall also prove that if the Hilbert bundles determined by the systems of measures  $\{\mu_i^u\}_u$  have continuous bases in the sense of Definition 24, then  $*$ -isomorphism between  $M_\sigma^*(G, \nu_1)$  and  $M_\sigma^*(G, \nu_2)$  can be restricted to a  $*$ -isomorphism between  $C^*(G, \nu_1)$  and  $C^*(G, \nu_2)$ .

For establishing notation, we include some definitions that can be found in several places (e.g. [8], [4]). A groupoid is a set  $G$  endowed with a product map

$$(x, y) \rightarrow xy \quad [ : G^{(2)} \rightarrow G ]$$

where  $G^{(2)}$  is a subset of  $G \times G$  called the set of composable pairs, and an inverse map

$$x \rightarrow x^{-1} \quad [ : G \rightarrow G ]$$

such that the following conditions hold:

(1) If  $(x, y) \in G^{(2)}$  and  $(y, z) \in G^{(2)}$ , then  $(xy, z) \in G^{(2)}$ ,  $(x, yz) \in G^{(2)}$  and  $(xy)z = x(yz)$ .

(2)  $(x^{-1})^{-1} = x$  for all  $x \in G$ .

(3) For all  $x \in G$ ,  $(x, x^{-1}) \in G^{(2)}$ , and if  $(z, x) \in G^{(2)}$ , then  $(zx)x^{-1} = z$ .

(4) For all  $x \in G$ ,  $(x^{-1}, x) \in G^{(2)}$ , and if  $(x, y) \in G^{(2)}$ , then  $x^{-1}(xy) = y$ .

The maps  $r$  and  $d$  on  $G$ , defined by the formulae  $r(x) = xx^{-1}$  and  $d(x) = x^{-1}x$ , are called the range and the source maps. It follows easily from the definition that they have a common image called the unit space of  $G$ , which is denoted  $G^{(0)}$ . Its elements are units in the sense that  $xd(x) = r(x)x = x$ .

Units will usually be denoted by letters as  $u, v, w$  while arbitrary elements will be denoted by  $x, y, z$ . It is useful to note that a pair  $(x, y)$  lies in  $G^{(2)}$  precisely when  $d(x) = r(y)$ , and that the cancellation laws hold (e.g.  $xy = xz$  iff  $y = z$ ). The fibres of the range and the source maps are denoted  $G^u = r^{-1}(\{u\})$  and  $G_v = d^{-1}(\{v\})$ , respectively. More generally, given the subsets  $A, B \subset G^{(0)}$ , we define  $G^A = r^{-1}(A)$ ,  $G_B = d^{-1}(B)$  and  $G_B^A = r^{-1}(A) \cap d^{-1}(B)$ . The reduction of  $G$  to  $A \subset G^{(0)}$  is  $G|_A = G_A^A$ . The relation  $u \sim v$  iff  $G_v^u \neq \emptyset$  is an equivalence relation on  $G^{(0)}$ . Its equivalence classes are called orbits and the orbit of a unit  $u$  is denoted  $[u]$ . A groupoid is called transitive iff it has a single orbit. The quotient space for this equivalence relation is called the orbit space of  $G$  and denoted  $G^{(0)}/G$ . We denote by  $\pi : G^{(0)} \rightarrow G^{(0)}/G, \pi(u) = \dot{u}$  the quotient map. A subset of  $G^{(0)}$  is said saturated if it contains the orbits of its elements. For any subset  $A$  of  $G^{(0)}$ , we denote by  $[A]$  the union of the orbits  $[u]$  for all  $u \in A$ .

A topological groupoid consists of a groupoid  $G$  and a topology compatible with the groupoid structure. This means that:

- (1)  $x \rightarrow x^{-1} [ : G \rightarrow G ]$  is continuous.
- (2)  $(x, y) [ : G^{(2)} \rightarrow G ]$  is continuous where  $G^{(2)}$  has the induced topology from  $G \times G$ .

We are exclusively concerned with topological groupoids which are second countable, locally compact Hausdorff. It was shown in [6] that measured groupoids may be assumed to have locally compact topologies, with no loss in generality.

If  $X$  is a locally compact space,  $C_c(X)$  denotes the space of complex-valued continuous functions with compact support. The Borel sets of a topological space are taken to be the  $\sigma$ -algebra generated by the open sets. The space of compactly supported bounded Borel function on  $X$  is denoted by  $\mathcal{B}_c(X)$ .

For a locally compact groupoid  $G$ , we denote by

$$G' = \{x \in G : r(x) = d(x)\}$$

the isotropy group bundle of  $G$ . It is closed in  $G$ .

Let  $G$  be a locally compact second countable groupoid equipped with a Haar system, i.e. a family of positive Radon measures on  $G$ ,  $\{\nu^u, u \in G^{(0)}\}$ , such that

- 1) For all  $u \in G^{(0)}$ ,  $\text{supp}(\nu^u) = G^u$ .
- 2) For all  $f \in C_c(G)$ ,

$$u \rightarrow \int f(x) d\nu^u(x) [ : G^{(0)} \rightarrow \mathbf{C} ]$$

is continuous.

- 3) For all  $f \in C_c(G)$  and all  $x \in G$ ,

$$\int f(y) d\nu^{r(x)}(y) = \int f(xy) d\nu^{d(x)}(y)$$

As a consequence of the existence of continuous Haar systems,  $r, d : G \rightarrow G^{(0)}$  are open maps ([10]). Therefore, in this paper we shall always assume that  $r : G \rightarrow G^{(0)}$  is an open map

If  $\mu$  is a measure on  $G^{(0)}$ , then the measure  $\nu = \int \nu^u d\mu(u)$ , defined by

$$\int f(y) d\nu(y) = \int \left( \int f(y) d\nu^u(y) \right) d\mu(u), \quad f \geq 0 \text{ Borel}$$

is called the measure on  $G$  induced by  $\mu$ . The image of  $\nu$  by the inverse map  $x \rightarrow x^{-1}$  is denoted  $\nu^{-1}$ .  $\mu$  is said quasi-invariant if its induced measure  $\nu$  is equivalent to its inverse  $\nu^{-1}$ . A measure belonging to the class of a quasi-invariant measure is also quasi-invariant. We say that the class is invariant.

If  $\mu$  is a quasi-invariant measure on  $G^{(0)}$  and  $\nu$  is the measure induced on  $G$ , then the Radon-Nikodym derivative  $\Delta = \frac{d\nu}{d\nu^{-1}}$  is called the modular function of  $\mu$ .

In order to define the  $C^*$ -algebra of a groupoid the space of continuous functions with compact support on groupoid is made into a  $*$ -algebra and endowed with the smallest  $C^*$ -norm making its representations continuous. For  $f, g \in C_c(G)$  the convolution is defined by:

$$f * g(x) = \int f(xy) g(y^{-1}) d\nu^{d(x)}(y)$$

and the involution by

$$f^*(x) = \overline{f(x^{-1})}.$$

Under these operations,  $C_c(G)$  becomes a topological  $*$ -algebra.

A representation of  $C_c(G)$  is a  $*$ -homomorphism from  $C_c(G)$  into  $\mathcal{B}(H)$ , for some Hilbert space  $H$ , that is continuous with respect to the inductive limit topology on  $C_c(G)$  and the weak operator topology on  $\mathcal{B}(H)$ . The full  $C^*$ -algebra  $C^*(G)$  is defined as the completion of the involutive algebra  $C_c(G)$  with respect to the full  $C^*$ -norm

$$\|f\| = \sup \|L(f)\|$$

where  $L$  runs over all non-degenerate representation of  $C_c(G)$  which are continuous for the inductive limit topology.

Every representation  $(\mu, G^{(0)} * \mathcal{H}, L)$  (see Definition 3.20/p.68 [4]) of  $G$  can be integrated into a representation, still denoted by  $L$ , of  $C_c(G)$ . The relation between the two representation is:

$$\langle L(f) \xi_1, \xi_2 \rangle = \int f(x) \langle L(x) \xi_1(d(x)), \xi_2(r(x)) \rangle \Delta^{-\frac{1}{2}}(x) d\nu^u(x) d\mu(u)$$

where  $f \in C_c(G)$ ,  $\xi_1, \xi_2 \in \int_{G^{(0)}}^{\oplus} \mathcal{H}(u) d\mu(u)$ .

Conversely, every non-degenerate  $*$ -representation of  $C_c(G)$  is obtained in this fashion (see [8] or [4]).

## 2 The decomposition of a Haar system over the principal groupoid

First we present some results on the structure of the Haar systems, as developed by J. Renault in Section 1 of [9] and also by A. Ramsay and M.E. Walter in

Section 2 of [7].

In Section 1 of [9] Jean Renault constructs a Borel Haar system for  $G'$ . One way to do this is to choose a function  $F_0$  continuous with conditionally support which is nonnegative and equal to 1 at each  $u \in G^{(0)}$ . Then for each  $u \in G^{(0)}$  choose a left Haar measure  $\beta_u^u$  on  $G_u^u$  so the integral of  $F_0$  with respect to  $\beta_u^u$  is 1.

Renault defines  $\beta_v^u = x\beta_v^v$  if  $x \in G_v^u$  (where  $x\beta_v^v(f) = \int f(xy) d\beta_v^v(y)$  as usual). If  $z$  is another element in  $G_v^u$ , then  $x^{-1}z \in G_v^v$ , and since  $\beta_v^v$  is a left Haar measure on  $G_v^v$ , it follows that  $\beta_v^u$  is independent of the choice of  $x$ . If  $K$  is a compact subset of  $G$ , then  $\sup_{u,v} \beta_v^u(K) < \infty$ . Renault also defines a 1-cocycle  $\delta$  on  $G$  such that for every  $u \in G^{(0)}$ ,  $\delta|_{G_u^u}$  is the modular function for  $\beta_u^u$ .  $\delta$  and  $\delta^{-1} = 1/\delta$  are bounded on compact sets in  $G$ .

Let

$$R = (r, d)(G) = \{(r(x), d(x)), x \in G\}$$

be the graph of the equivalence relation induced on  $G^{(0)}$ . This  $R$  is the image of  $G$  under the homomorphism  $(r, d)$ , so it is a  $\sigma$ -compact groupoid. With this apparatus in place, Renault describes a decomposition of the Haar system  $\{\nu^u, u \in G^{(0)}\}$  for  $G$  over the equivalence relation  $R$  (the principal groupoid associated to  $G$ ). He proves that there is a unique Borel Haar system  $\alpha$  for  $R$  with the property that

$$\nu^u = \int \beta_t^s d\alpha^u(s, t) \quad \text{for all } u \in G^{(0)}.$$

In Section 2 [7] A. Ramsay and M.E. Walter prove that

$$\sup_u \alpha^u((r, d)(K)) < \infty, \text{ for all compact } K \subset G$$

For each  $u \in G^{(0)}$  the measure  $\alpha^u$  is concentrated on  $\{u\} \times [u]$ . Therefore there is a measure  $\mu^u$  concentrated on  $[u]$  such that  $\alpha^u = \varepsilon_u \times \mu^u$ , where  $\varepsilon_u$  is the unit point mass at  $u$ . Since  $\{\alpha^u, u \in G^{(0)}\}$  is a Haar system, we have  $\mu^u = \mu^v$  for all  $(u, v) \in R$ , and the function

$$u \rightarrow \int f(s) \mu^u(s)$$

is Borel for all  $f \geq 0$  Borel on  $G^{(0)}$ . For each  $u$  the measure  $\mu^u$  is quasi-invariant (Section 2 [7]). Therefore  $\mu^u$  is equivalent to  $d_*(v^u)$  (Lemma 4.5/p. 277 [5]).

If  $\eta$  is a quasi-invariant measure for  $\{\nu^u, u \in G^{(0)}\}$ , then  $\eta$  is a quasi-invariant measure for  $\{\alpha^u, u \in G^{(0)}\}$ . Also if  $\Delta_R$  is the modular function associated to  $\{\alpha^u, u \in G^{(0)}\}$  and  $\eta$ , then  $\Delta = \delta\Delta_R \circ (r, d)$  can serve as the modular function associated to  $\{\nu^u, u \in G^{(0)}\}$  and  $\eta$ .

Since  $\mu^u = \mu^v$  for all  $(u, v) \in R$ , the system of measures  $\{\mu^u\}_u$  may be indexed on the elements of the orbit space  $G^{(0)}/G$ .

**Definition 1** We shall call the pair of the system of measures

$$\left( \{\beta_v^u\}_{(u,v) \in R}, \{\mu^{\dot{u}}\}_{\dot{u} \in G^{(0)}/G} \right)$$

(described above) the decomposition of the Haar system  $\{\nu^u, u \in G^{(0)}\}$  over the principal groupoid associated to  $G$ . Also we shall call  $\delta$  the 1-cocycle associated to the decomposition.

**Remark 2** Let us note that the system of measures  $\{\beta_v^u\}$  and the 1-cocycle in the preceding definition do not depend on the Haar system.

**Lemma 3** Let  $G$  be a locally compact second countable groupoid with the bundle map  $r|_{G'}$  of  $G'$  open. Let  $\{\nu^u, u \in G^{(0)}\}$  be a Haar system on  $G$  and  $(\{\beta_v^u\}, \{\mu^{\dot{u}}\})$  its decomposition over the principal groupoid associated to  $G$ . Then for each  $f \in C_c(G)$  the function

$$x \rightarrow \int f(y) d\beta_{d(x)}^{r(x)}(y)$$

is continuous on  $G$ .

**Proof.** By Lemma 1.3/p. 6 [9], for each  $f \in C_c(G)$  the function  $u \rightarrow \int f(y) d\beta_u^u(y)$  is continuous.

Let  $x \in G$  and  $(x_i)_i$  be a sequence in  $G$  converging to  $x$ . Let  $f \in C_c(G)$  and let  $g$  be a continuous extension on  $G$  of  $y \rightarrow f(xy) [ : G^{d(x)} \rightarrow \mathbf{C}]$ . Let  $K$  be the compact set

$$\left( \{x, x_i, i = 1, 2, \dots\}^{-1} \text{supp}(f) \cup \text{supp}(g) \right) \cap r^{-1}(\{d(x), d(x_i), i = 1, 2, \dots\}).$$

We have

$$\begin{aligned} & \left| \int f(y) d\beta_{d(x)}^{r(x)}(y) - \int f(y) d\beta_{d(x_i)}^{r(x_i)}(y) \right| \\ &= \left| \int f(xy) d\beta_{d(x)}^{d(x)}(y) - \int f(x_i y) d\beta_{d(x_i)}^{d(x_i)}(y) \right| \\ &= \left| \int g(y) d\beta_{d(x)}^{d(x)}(y) - \int f(x_i y) d\beta_{d(x_i)}^{d(x_i)}(y) \right| \\ &\leq \left| \int g(y) d\beta_{d(x)}^{d(x)}(y) - \int g(y) d\beta_{d(x_i)}^{d(x_i)}(y) \right| + \\ &\quad + \left| \int g(y) d\beta_{d(x_i)}^{d(x_i)}(y) - \int f(x_i y) d\beta_{d(x_i)}^{d(x_i)}(y) \right| \\ &\leq \left| \int g(y) d\beta_{d(x)}^{d(x)}(y) - \int g(y) d\beta_{d(x_i)}^{d(x_i)}(y) \right| + \\ &\quad + \sup_{y \in G_{d(x_i)}^{d(x_i)}} |g(y) - f(x_i y)| \beta_{d(x_i)}^{d(x_i)}(K) \end{aligned}$$

A compactness argument shows that  $\sup_{y \in G_{d(x_i)}^{d(x_i)}} |g(y) - f(x_i y)|$  converges to 0. Also  $\left| \int g(y) d\beta_{d(x)}^{d(x)}(y) - \int g(y) d\beta_{d(x_i)}^{d(x_i)}(y) \right|$  converges to 0 because the function  $u \rightarrow \int f(y) d\beta_u^u(y)$  is continuous. Hence

$$\left| \int f(y) d\beta_{d(x)}^{r(x)}(y) - \int f(y) d\beta_{d(x_i)}^{r(x_i)}(y) \right|$$

converges to 0. ■

**Definition 4** A locally compact groupoid  $G$  is proper if the map  $(r, d) : G \rightarrow G^{(0)} \times G^{(0)}$  is proper (i.e. the inverse map of each compact subset of  $G^{(0)} \times G^{(0)}$  is compact). (Definition 2.1.9/p. 37 [1]).

Throughout this paper we shall assume that  $G$  is a second countable locally compact groupoid for which the orbit space is Hausdorff and the map

$$(r, d) : G \rightarrow R, (r, d)(x) = (r(x), d(x))$$

is open, where  $R$  is endowed with the product topology induced from  $G^{(0)} \times G^{(0)}$ . Therefore  $R$  will be a locally compact groupoid. The fact that  $R$  is a closed subset of  $G^{(0)} \times G^{(0)}$  and that it is endowed with the product topology is equivalent to the fact  $R$  is a proper groupoid.

Throughout this paper by a groupoid with proper orbit space we shall mean a groupoid  $G$  for which the orbit space is Hausdorff and the map

$$(r, d) : G \rightarrow R, (r, d)(x) = (r(x), d(x))$$

is open, where  $R$  is endowed with the product topology induced from  $G^{(0)} \times G^{(0)}$ .

**Proposition 5** Let  $G$  be a second countable locally compact groupoid with proper orbit space. Let  $\{\nu^u, u \in G^{(0)}\}$  be a Haar system on  $G$  and  $(\{\beta_v^u\}, \{\mu^u\})$  its decomposition over the principal groupoid associated to  $G$ . Then for each  $g \in C_c(G^{(0)})$ , the map

$$u \rightarrow \int g(v) d\mu^{\pi(u)}(v)$$

is continuous.

**Proof.** Let  $g \in C_c(G^{(0)})$  and  $u_0 \in G^{(0)}$ . Let  $K_1$  be a compact neighborhood of  $u_0$  and  $K_2$  be the support of  $g$ . Since  $G$  is locally compact and  $(r, d)$  is open from  $G$  to  $(r, d)(G)$ , there is a compact subset  $K$  of  $G$  such that  $(r, d)(K)$  contains  $(K_1 \times K_2) \cap (r, d)(G)$ . Let  $F_1 \in C_c(G)$  be a nonnegative function equal to 1 on a compact neighborhood  $U$  of  $K$ . Let  $F_2 \in C_c(G)$  be a function which extends to  $G$  the function  $x \rightarrow F_1(x) / \int F_1(y) d\beta_{d(x)}^{r(x)}(y)$ ,  $x \in U$ . We have  $\int F_2(y) d\beta_v^u(y) = 1$  for all  $(u, v) \in (r, d)(K)$ . Since for all  $u \in K_1$ ,

$$\begin{aligned} \int g(v) d\mu^{\pi(u)}(v) &= \int g(v) \int F_2(y) d\beta_v^u(y) d\mu^{\pi(u)}(v) \\ &= \int g(d(y)) F_2(y) d\nu^u(y), \end{aligned}$$

it follows that  $u \rightarrow \int g(v) d\mu^{\pi(u)}(v)$  is continuous at  $u_0$ . ■

**Remark 6** Let  $G$  be a locally compact second countable groupoid with proper orbit space. Let  $\{\nu^u, u \in G^{(0)}\}$  be a Haar system on  $G$  and  $(\{\beta_v^u\}, \{\mu^u\})$  be its decomposition over the associated principal groupoid. If  $\mu$  is a quasi-invariant probability measure for the Haar system, then  $\mu_1 = \int \mu^{\pi(u)} d\mu(u)$  is a Radon

measure which is equivalent to  $\mu$ . Indeed, let  $f \geq 0$  Borel on  $G^{(0)}$  such that  $\mu(f) = 0$ . Since  $\mu$  is quasi-invariant, it follows that for  $\mu$  a.a.  $u$   $\nu^u(f \circ d) = 0$ , and since  $\mu^{\pi(u)}$  is equivalent to  $d_*(\nu^u)$ , it results  $\mu^{\pi(u)}(f) = 0$  for  $\mu$  a.a.  $u$ . Conversely, if  $\mu_1(f) = 0$ , then  $\mu^{\pi(u)}(f) = 0$  for  $\mu$  a.a.  $u$ , and therefore  $\nu^u(f \circ d) = 0$ . Thus the quasi-invariance of  $\mu$  implies  $\mu(f) = 0$ . Thus each Radon quasi-invariant measure is equivalent to a Radon measure of the form  $\int \mu^u d\tilde{\mu}(u)$ , where  $\tilde{\mu}$  is a probability measure on the orbit space  $G/G^{(0)}$ .

### 3 A $C^*$ -algebra associated to a locally compact groupoid with proper orbit space

Let  $G$  be a locally compact second countable groupoid with proper orbit space. Let

$$\pi : G^{(0)} \rightarrow G^{(0)}/G$$

be the quotient map. Since the quotient space is proper,  $G^{(0)}/G$  is Hausdorff. Let us assume that the range map  $r$  is open. As a consequence, the map  $\pi$  is open. Applying Lemma 1.1 [2] to the locally compact second countable spaces  $G^{(0)}$  and  $G^{(0)}/G$  and to the continuous open surjection  $\pi : G^{(0)} \rightarrow G^{(0)}/G$ , it follows that there is a Borel set  $F$  in  $G^{(0)}$  such that:

1.  $F$  contains exactly one element in each orbit  $[u] = \pi^{-1}(\pi(u))$ .
2. For each compact subset  $K$  of  $G^{(0)}$ ,  $F \cap [K] = F \cap \pi^{-1}(\pi(K))$  has a compact closure.

For each unit  $u$  let us define  $e(u) = F \cap [u]$  ( $e(u)$  is the unique element in the orbit of  $u$  contained in  $F$ ). For each Borel subset  $B$  of  $G^{(0)}$ ,  $\pi$  is continuous and one-to-one on  $B \cap F$  and hence  $\pi(B \cap F)$  is Borel in  $G^{(0)}/G$ . Therefore the map  $e : G^{(0)} \rightarrow G^{(0)}$  is Borel (for each Borel subset  $B$  of  $G^{(0)}$ ,  $e^{-1}(B) = [B \cap F] = \pi^{-1}(\pi(B \cap F))$  is Borel in  $G^{(0)}$ ). Also for each compact subset  $K$  of  $G^{(0)}$ ,  $e(K)$  has a compact closure because  $e(K) \subset F \cap [K]$ .

Since the orbit space  $G^{(0)}/G$  is proper the map

$$(r, d) : G \rightarrow R, (r, d)(x) = (r(x), d(x))$$

is open and  $R$  is closed in  $G^{(0)} \times G^{(0)}$ . Applying Lemma 1.1 [2] to the locally compact second countable spaces  $G$  and  $R$  and to the continuous open surjection  $(r, d) : G \rightarrow R$ , it follows that there is a *regular cross section*  $\sigma_0 : R \rightarrow G$ . This means that  $\sigma_0$  is Borel,  $(r, d)(\sigma_0(u, v)) = (u, v)$  for all  $(u, v) \in R$ , and  $\sigma_0(K)$  is relatively compact in  $G$  for each compact subset  $K$  of  $R$ .

Let us define  $\sigma : G^{(0)} \rightarrow G^F$  by  $\sigma(u) = \sigma_0(e(u), u)$  for all  $u$ . It is easy to note that  $\sigma$  is a cross section for  $d : G^F \rightarrow G^{(0)}$  and  $\sigma(K)$  is relatively compact in  $G$  for all compact  $K \subset G^{(0)}$ .

Replacing  $\sigma$  by

$$v \rightarrow \sigma(e(v))^{-1} \sigma(v)$$



we may assume that  $\sigma(e(v)) = e(v)$  for all  $v$ . Let us define  $q : G \rightarrow G_F^F$  by

$$q(x) = \sigma(r(x)) x \sigma(d(x))^{-1}, \quad x \in G.$$

Let  $\nu = \{\nu^u : u \in G^{(0)}\}$  be a Haar system on  $G$  and let  $(\{\beta_v^u\}, \{\mu^{\dot{u}}\})$  be its decompositions over the principal groupoid. Let  $\delta$  be the 1-cocycle associated to the decomposition.

Let us denote by  $\mathcal{B}_\sigma(G)$  the linear span of the functions of the form

$$x \rightarrow g_1(r(x)) g(q(x)) g_2(d(x))$$

where  $g_1, g_2$  are compactly supported Borel bounded function on  $G^{(0)}$  and  $g$  is a Borel bounded function on  $G_F^F$  such that if  $S$  is the support of  $g$ , then the closure of  $\sigma(K_1)^{-1} S \sigma(K_2)$  is compact in  $G$  for all compact subsets  $K_1, K_2$  of  $G^{(0)}$ .  $\mathcal{B}_\sigma(G)$  is a subspace of  $\mathcal{B}_c(G)$ , the space of compactly supported Borel bounded function on  $G$ .

If  $f_1, f_2 \in \mathcal{B}_\sigma(G)$  are defined by

$$\begin{aligned} f_1(x) &= g_1(r(x)) g(q(x)) g_2(d(x)) \\ f_2(x) &= h_1(r(x)) h(q(x)) h_2(d(x)) \end{aligned}$$

then

$$\begin{aligned} f_1 * f_2(x) &= g * h(q(x)) g_1(r(x)) h_2(d(x)) \langle g_2, \overline{h_1} \rangle_{\pi(r(x))} \\ f_1^*(x) &= \overline{g_2(r(x)) g(q(x)^{-1}) g_1(d(x))} \end{aligned}$$

Thus  $\mathcal{B}_\sigma(G)$  is closed under convolution and involution.

Let  $\omega$  be the universal representation of  $C^*(G, \nu)$  the usual  $C^*$ -algebra associated to a Haar system  $\nu = \{\nu^u, u \in G^{(0)}\}$  (constructed as in [8]). Since every cyclic representation of  $C^*(G, \nu)$  is the integrated form of a representation of  $G$ , it follows that  $\omega$  can be also regarded as a representation of  $\mathcal{B}_c(G)$ , the space of compactly supported Borel bounded function on  $G$ . Arlan Ramsay and Martin E. Walter have used the notation  $M^*(G, \nu)$  for the operator norm closure of  $\omega(\mathcal{B}_c(G))$ . Since  $\omega$  is an  $*$ -isomorphism on  $C^*(G, \nu)$ , we can regard  $C^*(G, \nu)$  as a subalgebra of  $M^*(G, \nu)$ .

**Definition 7** We denote by  $M_\sigma^*(G, \nu)$  the operator norm closure of  $\omega(\mathcal{B}_\sigma(G))$ .

**Lemma 8** Let  $\{\mu_1^{\dot{u}}\}_u$  and  $\{\mu_2^{\dot{u}}\}_u$  be two systems of measures on  $G^{(0)}$  satisfying:

1.  $\text{supp}(\mu_i^{\dot{u}}) = [u]$  for all  $\dot{u}, i = 1, 2$
2. For all compactly supported Borel bounded function  $f$  on  $G^{(0)}$  the function

$$u \rightarrow \int f(v) \mu_i^{\pi(u)}(v)$$

is bounded and Borel.

Then there is a family  $\{U_{\dot{u}}\}_{\dot{u}}$  of unitary operators with the following properties:

1.  $U_{\dot{u}} : L^2(\mu_1^{\dot{u}}) \rightarrow L^2(\mu_2^{\dot{u}})$  is a unitary operator for each  $\dot{u} \in G^{(0)}/G$ .
2. For all Borel bounded function  $f$  on  $G^{(0)}$ ,

$$u \rightarrow U_{\pi(u)}(f)$$

is a bounded Borel function with compact support.

3. For all Borel bounded function  $f$  on  $G^{(0)}$ ,

$$U_{\pi(u)}(\overline{f}) = \overline{U_{\pi(u)}(f)}$$

**Proof.** Using the same argument as in [6] (p. 323) we can construct a sequence  $f_1, f_2, \dots$  of real valued Borel bounded function on  $G^{(0)}$  such that  $\dim(L^2(\mu_1^{\dot{u}})) = \infty$  if and only if  $\|f_n\|_2$  for  $n = 1, 2, \dots$  and then  $\{f_1, f_2, \dots\}$  gives an ortonormate basis of  $L^2(\mu_1^{\dot{u}})$ , while  $\dim(L^2(\mu_1^{\dot{u}})) = k < \infty$  if and only if  $\|f_n\|_2 = 1$  for  $n \leq k$ , and  $\|f_n\|_2 = 0$  for  $n > k$  and then  $\{f_1, f_2, \dots, f_k\}$  gives an ortonormate basis of  $L^2(\mu_1^{\dot{u}})$ . Let  $g_1, g_2, \dots$  be a sequence with the same properties as  $f_1, f_2, \dots$  corresponding to  $\{\mu_2^{\dot{u}}\}_{\dot{u}}$ . Let us define  $U_{\dot{u}} : L^2(\mu_1^{\dot{u}}) \rightarrow L^2(\mu_2^{\dot{u}})$  by

$$U_{\dot{u}}(f_n) = g_n \text{ for all } n$$

Then the family  $\{U_{\dot{u}}\}_{\dot{u}}$  has the required properties. ■

**Theorem 9** Let  $G$  be a locally compact second countable groupoid with proper orbit space. Let  $\{\nu_i^u, u \in G^{(0)}\}$ ,  $i = 1, 2$  be two Haar systems on  $G$ . Let  $F$  be a Borel subset of  $G^{(0)}$  containing only one element  $e(u)$  in each orbit  $[u]$ . Let  $\sigma : G^{(0)} \rightarrow G^F$  be a cross section for  $d : G^F \rightarrow G^{(0)}$  with  $\sigma(e(v)) = e(v)$  for all  $v \in G^{(0)}$  and  $\sigma(K)$  relatively compact in  $G$  for all compact sets  $K \subset G^{(0)}$ . Then the  $C^*$ -algebras  $M_\sigma^*(G, \nu_1)$  and  $M_\sigma^*(G, \nu_2)$  are  $*$ -isomorphic.

**Proof.** Let  $(\{\beta_v^u\}, \{\mu_i^{\dot{u}}\})$  be the decompositions of the Haar systems over the principal groupoid. Let  $\delta$  be the 1-cocycle associated to the decompositions,  $i = 1, 2$ .

We shall denote by  $\langle \cdot, \cdot \rangle_{i, \dot{u}}$  the inner product of  $(L^2(G^{(0)}, \delta(\sigma(\cdot))\mu_i^{\dot{u}}))$ ,  $i = 1, 2$ .

Let us define  $q : G \rightarrow G_F^F$  by

$$q(x) = \sigma(r(x))x\sigma(d(x))^{-1}, x \in G.$$

We shall define a  $*$ -homomorphism  $\Phi$  from  $\mathcal{B}_\sigma(G)$  to  $\mathcal{B}_\sigma(G)$ . It suffices to define  $\Phi$  on the set of function on  $G$  of the form

$$x \rightarrow g_1(r(x))g(q(x))g_2(d(x))$$

Let  $\{U_{\dot{u}}\}_{\dot{u}}$  be the family of unitary operators with the properties stated in Lemma 8, associated to the systems of measures  $\{\delta(\sigma(\cdot))\mu_i^{\dot{u}}\}_{\dot{u}}$ ,  $i = 1, 2$ .

Let us define  $\Phi$  by

$$\Phi(f) = (x \rightarrow U_{\pi(r(x))}(g_1)(r(x))g(q(x))U_{\pi(d(x))}(g_2)(d(x)))$$

where  $f$  is defined by

$$f(x) = g_1(r(x))g(q(x))g_2(d(x))$$

If  $f_1$  and  $f_2$  are defined by

$$\begin{aligned} f_1(x) &= g_1(r(x))g(q(x))g_2(d(x)) \\ f_2(x) &= h_1(r(x))h(q(x))h_2(d(x)) \end{aligned}$$

then

$$f_1 * f_2(x) = g * h(q(x))g_1(r(x))h_2(d(x))\langle g_2, \overline{h_1} \rangle_{1, \pi(r(x))}$$

and consequently

$$\begin{aligned} &\Phi(f_1 * f_2) \\ &= g * h(q(x))U_{\pi(r(x))}(g_1)(r(x))U_{\pi(r(x))}(h_2)(d(x))\langle g_2, \overline{h_1} \rangle_{1, \pi(r(x))} \\ &= \Phi(f_1) * \Phi(f_2). \end{aligned}$$

Let  $\tilde{\eta}$  be a probability measure on  $G^{(0)}/G$  and  $\eta_i = \int \mu_i^{\dot{u}} d\tilde{\eta}(\dot{u})$ ,  $i = 1, 2$ . Let  $L_1$  be the integrated form of a representation  $(L, \mathcal{H} * G^{(0)}, \eta_1)$  and  $L_2$  be the integrated form of  $(L, \mathcal{H} * G^{(0)}, \eta_2)$ . Let  $B$  be the Borel function defined by:

$$B(u) = L(\sigma(u))$$

and  $W : \int_{G^{(0)}}^{\oplus} \mathcal{H}(u) d\eta_1(u) \rightarrow \int_{G^{(0)}}^{\oplus} \mathcal{H}(e(u)) d\eta_1(u)$  be defined by

$$W(\zeta) = (u \rightarrow B(u)(\zeta(u)))$$

Since every element of  $L^2(G^{(0)}, \delta(\sigma(\cdot))\mu_1^{\dot{w}}, \mathcal{H}(e(w)))$  is a limit of linear combinations of elements  $u \rightarrow a(u)\xi$  with  $a \in L^2(G^{(0)}, \delta(\sigma(\cdot))\mu_1^{\dot{w}})$  and  $\xi \in \mathcal{H}(e(w))$ , we can define a unitary operator

$$V_{\dot{w}} : L^2(G^{(0)}, \delta(\sigma(\cdot))\mu_1^{\dot{w}}, \mathcal{H}(e(w))) \rightarrow L^2(G^{(0)}, \delta(\sigma(\cdot))\mu_2^{\dot{w}}, \mathcal{H}(e(w)))$$

by

$$V_{\dot{w}}(u \rightarrow a(u)\xi) = U_{\dot{w}}(a)\xi$$

Let  $V : \int_{G^{(0)}}^{\oplus} \mathcal{H}(e(u)) d\eta_1(u) \rightarrow \int_{G^{(0)}}^{\oplus} \mathcal{H}(e(u)) d\eta_2(u)$  be defined by

$$V(\zeta) = (u \rightarrow V_{\dot{u}}(\zeta(u)))$$

If  $\zeta_1, \zeta_2 \in \int_{G^{(0)}}^{\oplus} \mathcal{H}(e(u)) d\eta_1(u)$  and  $f$  is of the form

$$f(x) = g_1(r(x))g(q(x))g_2(d(x)),$$

we have

$$\begin{aligned} & \langle WL_1(f) W^* \zeta_1, \zeta_2 \rangle \\ &= \int \int g(x) \delta(x)^{\frac{-1}{2}} \langle L(x) A_1(\dot{w}), B_1(\dot{w}) \rangle d\beta_{e(w)}^{e(w)}(x) d\tilde{\eta}(\dot{w}) \end{aligned}$$

where

$$\begin{aligned} A_1(\dot{w}) &= \int g_2(v) \zeta_1(v) \delta(\sigma(v))^{\frac{1}{2}} d\mu_1^{\dot{w}}(v) \\ B_1(\dot{w}) &= \int g_1(u) \zeta_2(u) \delta(\sigma(u))^{\frac{1}{2}} d\mu_1^{\dot{w}}(u) \end{aligned}$$

Moreover, if  $f$  is of the form  $f(x) = g_1(r(x))g(q(x))g_2(d(x))$  and  $\zeta_1, \zeta_2 \in \int_{G^{(0)}}^{\oplus} \mathcal{H}(e(u)) d\eta_2(u)$ , then

$$\begin{aligned} & \langle VWL_1(f) W^* V^* \zeta_1, \zeta_2 \rangle \\ &= \int \int g(x) \delta(x)^{\frac{-1}{2}} \langle L(x) A_2(\dot{w}), B_2(\dot{w}) \rangle d\beta_{e(u)}^{e(u)}(x) d\tilde{\eta}(\dot{w}) \\ &= \langle WL_2(\Phi(f)) W^* \zeta_1, \zeta_2 \rangle \end{aligned}$$

where

$$\begin{aligned} A_2(\dot{w}) &= \int g_2(v) V^* \zeta_1(v) \delta(\sigma(v))^{\frac{1}{2}} d\mu_1^{\dot{w}}(v) \\ &= \int U_{\dot{v}}(g_2)(v) \zeta_1(v) \delta(\sigma(v))^{\frac{1}{2}} d\mu_2^{\dot{w}}(v) \\ B_2(\dot{w}) &= \int g_1(v) V^* \zeta_2(v) \delta(\sigma(v))^{\frac{1}{2}} d\mu_1^{\dot{w}}(v) \\ &= \int U_{\dot{u}}(g_1)(u) \zeta_2(u) \delta(\sigma(u))^{\frac{1}{2}} d\mu_2^{\dot{w}}(u) \end{aligned}$$

Therefore  $\|L_1(f)\| = \|L_2(\Phi(f))\|$ . Consequently we can extend  $\Phi$  to a \*-homomorphism between the  $M_\sigma^*(G, \nu_1)$  and  $M_\sigma^*(G, \nu_2)$ . It is not hard to see that  $\Phi$  is in fact a \*-isomorphism:

$$\Phi^{-1}(f) = \left( x \rightarrow U_{\pi(r(x))}^*(g_1)(r(x))g(q(x))U_{\pi(d(x))}^*(g_2)(d(x)) \right)$$

for each  $f$  of the form

$$f(x) = g_1(r(x))g(q(x))g_2(d(x)).$$

■

## 4 The case of locally transitive groupoids

A locally compact transitive groupoid  $G$  is a groupoid for which all orbits  $[u]$  are open in  $G^{(0)}$ . We shall prove that if  $G$  is a locally compact second countable locally transitive groupoid endowed with a Haar system  $\nu$ , then

$$C^*(G, \nu) = M^*(G, \nu) = M_\sigma^*(G, \nu)$$

for any regular cross section  $\sigma$ .

**Notation 10** Let  $\{\nu^u, u \in G^{(0)}\}$  be a fixed Haar system on  $G$ . Let  $\mu$  be a quasi-invariant measure,  $\Delta$  its modular function,  $\nu_1$  be the measure induced by  $\mu$  on  $G$  and  $\nu_0 = \Delta^{-\frac{1}{2}}\nu_1$ . Let us denote by  $II_\mu(G)$  the set

$$\left\{ f \in L^1(G, \nu_0) : \|f\|_{II, \mu} < \infty \right\},$$

where  $\|f\|_{II, \mu}$  is defined by

$$\|f\|_{II, \mu} = \sup \left\{ \int |f(x) j(d(x)) k(r(x))| d\nu_0(x), \int |j|^2 d\mu = \int |k|^2 d\mu = 1 \right\}.$$

If  $\mu_1$  and  $\mu_2$  are two equivalent quasi-invariant measures, then  $\|f\|_{II, \mu_1} = \|f\|_{II, \mu_2}$ , because  $\|f\|_{II, \mu} = \|II_\mu(|f|)\|$  for each quasi-invariant measure  $\mu$ , where  $II_\mu$  is the one dimensional trivial representation on  $\mu$

Define  $\|f\|_{II}$  to be

$$\sup \left\{ \|f\|_{II, \mu} : \mu \text{ quasi-invariant Radon measure on } G^{(0)} \right\}$$

The supremum can be taken over the classes of quasi-invariant measure.

If  $\|\cdot\|$  is the full  $C^*$ -norm on  $C_c(G)$ , then

$$\|f\| \leq \|f\|_{II} \text{ for all } f.$$

(see [7])

**Lemma 11** Let  $G$  be a locally compact second countable groupoid with proper orbit space. Let  $\{\nu^u, u \in G^{(0)}\}$  be a Haar system on  $G$ ,  $(\{\beta_v^u\}, \{\mu^u\})$  its decomposition over the principal groupoid associated to  $G$  and  $\delta$  the associated 1-cocycle. If  $f$  is a universally measurable function on  $G$ , then

$$\|f\|_{II} \leq \sup_{\dot{w}} \left( \int \int \left( \int |f(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 d\mu^{\dot{w}}(v) d\mu^{\dot{w}}(u) \right)^{\frac{1}{2}}.$$

**Proof.** Each Radon quasi-invariant measure is equivalent with a Radon measure of the form  $\int \mu^{\dot{u}} d\tilde{\mu}(\dot{u})$ , where  $\tilde{\mu}$  is a probability measure on the orbit space  $G/G^{(0)}$ . Therefore for the computation of  $\|\cdot\|_{II}$  it is enough to consider only the quasi-invariant measures of the form  $\mu = \int \mu^{\dot{u}} d\tilde{\mu}(\dot{u})$ , where  $\tilde{\mu}$  is a probability measure on  $G^{(0)}/G$ . It is easy to see that the modular function of  $\int \mu^{\dot{u}} d\tilde{\mu}(\dot{u})$  is  $\Delta = \delta$ .

Let  $j, k \in L^2(G^{(0)}, \mu)$  with  $\int |j|^2 d\mu = \int |k|^2 d\mu = 1$ . We have

$$\begin{aligned}
& \int \int \int \int |f(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) |j(v)| |k(u)| d\mu^{\dot{v}}(v) d\mu^{\dot{u}}(u) d\tilde{\mu}(\dot{w}) \\
& \leq \int \left( \int \int \left( \int |f(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 d\mu^{\dot{v}}(v) d\mu^{\dot{u}}(u) \right)^{\frac{1}{2}} \\
& \quad \cdot \left( \int \int |j(v)|^2 |k(u)|^2 d\mu^{\dot{v}}(v) d\mu^{\dot{u}}(u) \right)^{\frac{1}{2}} d\tilde{\mu}(\dot{w}) \\
& = \sup_{\dot{w}} \left( \int \int \left( \int |f(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 d\mu^{\dot{v}}(v) d\mu^{\dot{u}}(u) \right)^{\frac{1}{2}} \\
& \quad \cdot \int \left( \int |j(v)|^2 d\mu^{\dot{v}}(v) \right)^{\frac{1}{2}} \left( \int |k(u)|^2 d\mu^{\dot{u}}(u) \right)^{\frac{1}{2}} d\tilde{\mu}(\dot{w}) \\
& \leq \sup_{\dot{w}} \left( \int \int \left( \int |f(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 d\mu^{\dot{v}}(v) d\mu^{\dot{u}}(u) \right)^{\frac{1}{2}}
\end{aligned}$$

Consequently,

$$\|f\|_{II} \leq \sup_{\dot{w}} \left( \int \int \left( \int |f(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 d\mu^{\dot{v}}(v) d\mu^{\dot{u}}(u) \right)^{\frac{1}{2}}$$

■

If  $G$  is locally transitive, each orbit  $[u]$  is open in  $G^{(0)}$ . Each measure  $\mu^{\dot{u}}$  is supported on  $[u]$ . Since  $([u])$  is a partition of  $G^{(0)}$  into open sets, it follows that there is a unique Radon measure  $m$  on  $G^{(0)}$  such that the restriction of  $m$  at  $C_c([u])$  is  $\mu^{\dot{u}}$  for each  $[u]$ .

**Corollary 12** *Let  $G$  be a locally compact second countable locally transitive groupoid endowed with a Haar system  $\nu = \{\nu^u, u \in G^{(0)}\}$ . Let  $f$  be a universally measurable function such that  $\|f\|_{II} < \infty$ .*

1. *If  $(f_n)_n$  is a uniformly bounded sequence of universally measurable functions supported on a compact set, and if  $(f_n)_n$  converges pointwise to  $f$ , then  $(f_n)_n$  converges to  $f$  in the norm of  $C^*(G, \nu)$ .*
2. *If  $(f_n)_n$  is an increasing sequence of universally measurable nonnegative functions on  $G$  that converges pointwise to  $f$ , then  $(f_n)_n$  converges to  $f$  in the norm of  $C^*(G, \nu)$ .*

**Proof.** Let  $(\{\beta_v^u\}, \{\mu^u\})$  be the decomposition of the Haar system over the principal groupoid associated to  $G$  and  $\delta$  the associated 1-cocycle. Let  $m$  be the unique measure such that restriction of  $m$  at  $C_c([u])$  is  $\mu^u$  for each  $[u]$ . Let  $(f_n)_n$  be a sequence of universally measurable functions supported on a compact set  $K$ . Let us write

$$M = \sup_{u,v} \beta_u^v(K^{-1}).$$

and let us assume that  $(f_n)_n$  converges pointwise to  $f$ . According to Lemma 11, we have

$$\|f - f_n\|_{II} \leq \sup_{\dot{w}} \left( \int \int \left( \int |f(x) - f_n(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 d\mu^{\dot{w}}(v) d\mu^{\dot{w}}(u) \right)^{\frac{1}{2}}.$$

Hence

$$\begin{aligned} \|f - f_n\|_{II} &\leq \sup_{\dot{w}} M \left( \int \int \left( \int |f(x) - f_n(x)|^2 d\beta_v^u(x) \right) d\mu^{\dot{w}}(v) d\mu^{\dot{w}}(u) \right)^{\frac{1}{2}} \\ &\leq M \left( \int \int \left( \int |f(x) - f_n(x)|^2 d\beta_u^v(x) \right) dm(v) dm(u) \right)^{\frac{1}{2}}. \end{aligned}$$

If  $\|\cdot\|$  denotes the  $C^*$ -norm, then

$$\lim_n \|f - f_n\| \leq \lim_n \|f - f_n\|_{II} = 0,$$

because

$$\int \int \left( \int |f(x) - f_n(x)|^2 d\beta_u^v(x) \right) dm(v) dm(u)$$

converges to zero, by the Dominated Convergence Theorem.

Let  $(f_n)_n$  be an increasing sequence of universally measurable nonnegative functions that converges pointwise to  $f$ . Since

$$\begin{aligned} \|f - f_n\|_{II} &\leq \sup_{\dot{w}} \left( \int \int \left( \int |f(x) - f_n(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 d\mu^{\dot{w}}(v) d\mu^{\dot{w}}(u) \right)^{\frac{1}{2}} \\ &\leq \left( \int \int \left( \int |f(x) - f_n(x)| \delta(x)^{-\frac{1}{2}} d\beta_v^u(x) \right)^2 dm(v) dm(u) \right)^{\frac{1}{2}} \end{aligned}$$

it follows that

$$\lim_n \|f - f_n\|_{II} = 0.$$

■

**Proposition 13** *Let  $G$  be a locally compact second countable locally transitive groupoid endowed with a Haar system  $\nu = \{\nu^u, u \in G^{(0)}\}$ . Then any function in  $\mathcal{B}_c(G)$ , the space of compactly supported Borel bounded function on  $G$ , can be viewed as an element of  $C^*(G, \nu)$ .*

**Proof.** Let  $(\{\beta_v^u\}, \{\mu^{\dot{u}}\})$  be the decomposition of the Haar system over the principal groupoid associated to  $G$ , and  $\delta$  the associated 1-cocycle. Let  $m$  be the unique measure such that restriction of  $m$  at  $C_c([u])$  is  $\mu^{\dot{u}}$  for each  $[u]$ . Let  $m$  be a dominant for the family  $\{\mu^{\dot{u}}\}$ . Let  $\nu_1$  be the measure on  $G$  define by

$$\int f(x) d\nu_1(x) = \left( \int \int \left( \int f(x) d\beta_u^v(x) \right) dm(v) dm(u) \right)$$

for all Borel nonnegative function  $f$ . If  $f \in \mathcal{B}_c(G)$ , then  $f$  is the limit in  $L^2(G, \nu_1)$  of a sequence,  $(f_n)_n$ , in  $C_c(G)$  that is supported on some compact set  $K$  supporting  $f$ . If we write

$$M = \sup_{u,v} \beta_u^v(K^{-1}),$$

we have

$$\begin{aligned} \|f - f_n\|_{II} &\leq \sup_{\dot{w}} M \left( \int \int \left( \int |f(x) - f_n(x)|^2 d\beta_v^u(x) \right) d\mu^{\dot{w}}(v) d\mu^{\dot{w}}(u) \right)^{\frac{1}{2}} \\ &\leq M \left( \int \int \left( \int |f(x) - f_n(x)|^2 d\beta_u^v(x) \right) dm(v) dm(u) \right)^{\frac{1}{2}}. \end{aligned}$$

If  $\|\cdot\|$  denotes the  $C^*$ -norm, then

$$\lim_n \|f - f_n\| \leq \lim_n \|f - f_n\|_{II} = 0.$$

Thus  $f$  can be viewed as an element in  $C^*(G, \nu)$ . ■

**Proposition 14** *If  $G$  is a locally compact second countable locally transitive groupoid endowed with a Haar system  $\{\nu^u, u \in G^{(0)}\}$  with bounded decomposition, then*

$$C^*(G, \nu) = M^*(G, \nu).$$

**Proof.** It follows from the Proposition 13. ■

**Remark 15** *Let  $G$  be locally compact locally transitive groupoid. Let  $F$  be a subset of  $G^{(0)}$  containing only one element  $e(u)$  in each orbit  $[u]$ . It is easy to see that  $F$  is a closed subset of  $G$  and that  $F$  is a discrete space. Let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section of  $d_F$ . Let us endow  $\bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  with the topology induced from  $G^{(0)} \times G_F^F \times G^{(0)}$ . The topology of  $\bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  is locally compact because  $\bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  is a closed subset of the locally compact space  $G^{(0)} \times G_F^F \times G^{(0)}$ . With the operations:*

$$\begin{aligned} (u, x, v)(v, y, w) &= (u, xy, w) \\ (u, x, v)^{-1} &= (v, x^{-1}, u) \end{aligned}$$



$\bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  becomes a groupoid. Let us define  $\phi : G \rightarrow \bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  by

$$\phi(x) = \left( r(x), \sigma(r(x)) x \sigma(d(x))^{-1}, d(x) \right)$$

and note that  $\phi$  is a Borel isomorphism which carries compact sets to relatively compact sets .

**Lemma 16** *Let  $G$  be locally compact second countable locally transitive groupoid. Let  $F$  be a subset of  $G^{(0)}$  containing only one element  $e(u)$  in each orbit  $[u]$ . Let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section of  $d_F$ . Then any compactly supported Borel bounded function on  $G$  is pointwise limit of a uniformly bounded sequence  $(f_n)_n$  of Borel functions supported on a compact set supporting  $f$ , having the property that each  $f_n$  is a linear combination of functions of the form*

$$x \rightarrow g_1(r(x)) g \left( \sigma(r(x)) x \sigma(d(x))^{-1} \right) g_2(d(x))$$

where  $g_1, g_2$  are compactly supported Borel bounded function on  $G^{(0)}$  and  $g$  is a compactly supported Borel bounded function on  $G_F^F$ .

**Proof.** Let us endow  $\bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  with the topology induced from  $G^{(0)} \times G_F^F \times G^{(0)}$  as in Remark.15. The topology of  $\bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  is locally compact. Any compactly supported Borel bounded function on  $G^{(0)} \times G_F^F \times G^{(0)}$  is pointwise limit of uniformly bounded sequences  $(f_n)_n$  of Borel functions supported on a compact set, such that each function  $f_n$  is a linear combination of functions of the form

$$(u, x, v) \rightarrow g_1(u) g(x) g_2(v)$$

where  $g_1, g_2$  are compactly supported Borel bounded function on  $G^{(0)}$  and  $g$  is a compactly supported Borel bounded function on  $G_F^F$  . Consequently, any compactly supported Borel bounded function on  $\bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  has the same property. Since  $\phi : G \rightarrow \bigcup_{[u]} [u] \times G_{e(u)}^{e(u)} \times [u]$  defined by

$$\phi(x) = \left( r(x), \sigma(r(x)) x \sigma(d(x))^{-1}, d(x) \right)$$

is a Borel isomorphism which carries compact sets to relatively compact sets, it follows that any compactly supported Borel bounded function on  $G$  can be represented as a pointwise limit of a uniformly bounded sequence  $(f_n)_n$  of Borel functions supported on a compact set supporting  $f$  , having the property that each  $f_n$  is a linear combination of functions of the form

$$x \rightarrow g_1(r(x)) g \left( \sigma(r(x)) x \sigma(d(x))^{-1} \right) g_2(d(x))$$

■

**Corollary 17** *Let  $G$  be locally compact second countable locally transitive groupoid. Let  $F$  be a subset of  $G^{(0)}$  containing only one element  $e(u)$  in each orbit  $[u]$ . Let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section of  $d_F$ . Then the linear span of the functions of the form*

$$x \rightarrow g_1(r(x)) g\left(\sigma(r(x)) x \sigma(d(x))^{-1}\right) g_2(d(x))$$

where  $g_1, g_2 \in \mathcal{B}_c(G^{(0)})$  and  $g \in \mathcal{B}_c(G_F^F)$ , is dense in the full  $C^*$ -algebra of  $G$ .

**Proof.** Let  $f$  be a function on  $G$ , defined by

$$f(x) = g_1(r(x)) g\left(\sigma(r(x)) x \sigma(d(x))^{-1}\right) g_2(d(x))$$

where  $g_1, g_2 \in \mathcal{B}_c(G^{(0)})$  and  $g \in \mathcal{B}_c(G_F^F)$ . Then  $f \in \mathcal{B}_c(G)$ , therefore it can be viewed as an element of the  $C^*(G, \nu)$  as we note in Proposition 13. Each  $f \in \mathcal{B}_c(G)$  (in particular in  $C_c(G)$ ) is the limit (pointwise and consequently in the  $C^*$ -norm according to Corollary 12) of a uniformly bounded sequence  $(f_n)_n$  of Borel functions supported on a compact set supporting  $f$ , having the property that each  $f_n$  is a linear combination of functions of the required form. ■

**Proposition 18** *Let  $G$  is a locally compact second countable locally transitive groupoid endowed with a Haar system  $\{\nu^u, u \in G^{(0)}\}$ . Let  $F$  be a subset of  $G^{(0)}$  containing only one element  $e(u)$  in each orbit  $[u]$ . Let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section of  $d_F$ . Then*

$$C^*(G, \nu) = M^*(G, \nu) = M_\sigma^*(G, \nu).$$

**Proof.** We have proved that  $C^*(G, \nu) = M^*(G, \nu)$ . From the preceding corollary, it follows that the linear span of the functions of the form

$$x \rightarrow g_1(r(x)) g\left(\sigma(r(x)) x \sigma(d(x))^{-1}\right) g_2(d(x))$$

where  $g_1, g_2 \in \mathcal{B}_c(G^{(0)})$  and  $g \in \mathcal{B}_c(G_F^F)$  is dense in  $C^*(G, \nu)$ . But this space is contained in  $\mathcal{B}_\sigma(G)$ . Therefore  $C^*(G, \nu) = M^*(G, \nu) = M_\sigma^*(G, \nu)$ . ■

## 5 The case of principal proper groupoids case

**Notation 19** *Let  $G$  be a locally compact second countable groupoid with proper orbit space. Let  $F$  be a Borel subset of  $G^{(0)}$  containing only one element  $e(u)$  in each orbit  $[u]$ . Let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section for  $d_F : G^F \rightarrow$*

$G^{(0)}$ ,  $d_F(x) = d(x)$  with  $\sigma(e(v)) = e(v)$  for all  $v \in G^{(0)}$ . Let  $q : G \rightarrow G_F^F$  be defined by

$$q(x) = \sigma(r(x))x\sigma(d(x))^{-1}$$

We shall endow  $G_F^F$  with the quotient topology induced by  $q$ . We shall denote by  $\mathbf{C}_\sigma(G)$  the linear span of the functions of the form

$$x \rightarrow g_1(r(x))g\left(\sigma(r(x))x\sigma(d(x))^{-1}\right)g_2(d(x))$$

where  $g_1, g_2 \in C_c(G^{(0)})$  and  $g \in C_c(G_F^F)$ .

**Proposition 20** *With the Notation 19, if the space of continuous functions with compact support on  $G_F^F$  (with the respect to the quotient topology induced by  $q$ ) separates the points of  $G_F^F$ , then  $\mathbf{C}_\sigma(G)$  is dense in  $C_c(G)$  (for the inductive limit topology). In particular, if the quotient topology induced by  $q$  on  $G_F^F$  is a locally compact (Hausdorff) topology, then  $\mathbf{C}_\sigma(G)$  is dense in  $C_c(G)$ .*

**Proof.** If the space of continuous functions with compact support on  $G_F^F$  (with the respect to the quotient topology induced by  $q$ ) separates the points of  $G_F^F$ , then  $\mathbf{C}_\sigma(G)$  separates the points of  $G$ . By Stone-Weierstrass Theorem, it follows that  $\mathbf{C}_\sigma(G)$  is dense in  $C_c(G)$  (for the inductive limit topology) ■

**Proposition 21** *Let  $G$  be a locally compact principal groupoid. If  $G$  is proper, then the quotient topology induced by  $q$  on  $G_F^F$  is a locally compact (Hausdorff) topology. Consequently,  $\mathbf{C}_\sigma(G)$  is dense in  $C_c(G)$  for the inductive limit topology (we use the Notation 19).*

**Proof.** Let  $\pi : G \rightarrow G^{(0)}/G$  be the canonical projection. Let us note that for a principal groupoid the condition

$$q(x) = q(y)$$

is equivalent with

$$\pi(r(x)) = \pi(r(y)).$$

First we shall prove that the topology on  $G_F^F$  is Hausdorff. Let  $(x_i)_i$  and  $(y_i)_i$  be two nets with  $q(x_i) = q(y_i)$  for every  $i$ . Let us suppose that  $(x_i)_i$  converges to  $x$  and  $(y_i)_i$  converges to  $y$ . Then

$$\lim \pi(r(x_i)) = \lim \pi(r(y_i)) = \pi(r(x)) = \pi(r(y))$$

Hence  $q(x) = q(y)$ , and therefore the topology on  $G_F^F$  is Hausdorff. We shall prove that  $q$  is open. If  $(z_i)_i$  is a net converging to  $q(x)$  in  $G_F^F$ , then  $\pi \circ r(z_i)$  converges to  $\pi \circ r(x)$ . Since

$$\pi \circ r : G \rightarrow G^{(0)}/G$$

is an open map, there is a net  $(x_i)_i$  converging to  $x$ , such that  $\pi \circ r(x_i) = \pi \circ r(z_i)$ , and consequently  $q(x_i) = q(z_i) = z_i$ . Hence  $q$  is an open map and the quotient topology induced by  $q$  on  $G_F^F$  is locally compact. ■

**Theorem 22** *Let  $G$  be a locally compact second countable groupoid with proper orbit space. Let  $F$  be a Borel subset of  $G^{(0)}$  meeting each orbit exactly once. Let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section for  $d : G^F \rightarrow G$ . Let us assume that the quotient topology induced by  $q$  on  $G_F^F$  is a locally compact (Hausdorff) topology. Let  $\{\nu^u, u \in G^{(0)}\}$  be a Haar system on  $G$ . Then*

$$C^*(G, \nu) \subset M_\sigma^*(G, \nu) \subset M^*(G, \nu).$$

**Proof.** From Proposition 20  $\mathbf{C}_\sigma(G)$  is dense in  $C_c(G)$  for the inductive limit topology and hence is dense in  $C^*(G, \nu)$ . Since  $\mathbf{C}_\sigma(G) \subset \mathcal{B}_\sigma(G)$ , it follows that  $C^*(G, \nu) \subset M_\sigma^*(G, \nu)$ . ■

**Corollary 23** *Let  $G$  be a locally compact second countable principal proper groupoid. Let  $F$  be a Borel subset of  $G^{(0)}$  meeting each orbit exactly once. Let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section for  $d : G^F \rightarrow G$ . Let  $\{\nu^u, u \in G^{(0)}\}$  be a Haar system on  $G$ . Then*

$$C^*(G, \nu) \subset M_\sigma^*(G, \nu) \subset M^*(G, \nu).$$

**Proof.** Applying Proposition 21, we obtain that the quotient topology induced by  $q$  on  $G_F^F$  is a locally compact (Hausdorff) topology. Therefore  $G$  satisfies the hypothesis of Theorem 22. ■

**Definition 24** *Let  $\{\mu^{\dot{u}}\}_{\dot{u}}$  be a system of measures on  $G^{(0)}$  satisfying:*

1.  $\text{supp}(\mu^{\dot{u}}) = [u]$  for all  $\dot{u}$ .
2. For all compactly supported continuous functions  $f$  on  $G^{(0)}$  the function

$$u \rightarrow \int f(v) \mu^{\pi(u)}(v)$$

*is continuous*

*We shall say that the Hilbert bundle determined by the system of measures  $\{\mu^{\dot{u}}\}_{\dot{u}}$  has a continuous basis if there is sequence  $f_1, f_2, \dots$  of real valued continuous functions on  $G^{(0)}$  such that  $\dim(L^2(\mu^{\dot{u}})) = \infty$  if and only if  $\|f_n\|_2$  for  $n = 1, 2, \dots$  and then  $\{f_1, f_2, \dots\}$  gives an ortonormate basis of  $L^2(\mu^{\dot{u}})$ , while  $\dim(L^2(\mu^{\dot{u}})) = k < \infty$  if and only if  $\|f_n\|_2 = 1$  for  $n \leq k$ , and  $\|f_n\|_2 = 0$  for  $n > k$  and then  $\{f_1, f_2, \dots, f_k\}$  gives an ortonormate basis of  $L^2(\mu^{\dot{u}})$ .*

**Remark 25** *Let  $\{\mu_1^{\dot{u}}\}_{\dot{u}}$  and  $\{\mu_2^{\dot{u}}\}_{\dot{u}}$  be two systems of measures on  $G^{(0)}$  satisfying:*

1.  $\text{supp}(\mu_i^{\dot{u}}) = [u]$  for all  $\dot{u}$ ,  $i = 1, 2$ .

2. For all compactly supported continuous functions  $f$  on  $G^{(0)}$  the function

$$u \rightarrow \int f(v) \mu_i^{\pi(u)}(v)$$

is continuous

Let us assume that the Hilbert bundles determined by the systems of measures  $\{\mu_i^{\dot{u}}\}_{\dot{u}}$  have continuous bases. Let  $f_1, f_2, \dots$  be a continuous basis for Hilbert bundle determined by  $\{\mu_1^{\dot{u}}\}_{\dot{u}}$  and let  $g_1, g_2, \dots$  be a continuous basis for Hilbert bundle determined by  $\{\mu_2^{\dot{u}}\}_{\dot{u}}$ . Let us define a unitary operator  $U_{\dot{u}} : L^2(\mu_1^{\dot{u}}) \rightarrow L^2(\mu_2^{\dot{u}})$  by

$$U_{\dot{u}}(f_n) = g_n \text{ for all } n$$

Then the family  $\{U_{\dot{u}}\}_{\dot{u}}$  has the following properties:

1. For all Borel bounded function  $f$  on  $G^{(0)}$ ,

$$u \rightarrow U_{\pi(u)}(f)$$

is a bounded Borel function with compact support.

2. For all Borel bounded function  $f$  on  $G^{(0)}$ ,

$$U_{\pi(u)}(\overline{f}) = \overline{U_{\pi(u)}(f)}$$

3. For all compactly supported continuous functions  $f$  on  $G^{(0)}$  there is a sequence  $(h_n)_n$  of compactly supported continuous functions on  $G^{(0)}$  such that

$$\sup_{\dot{u}} \int |U_{\dot{u}}(f) - h_n|^2 d\mu_2^{\dot{u}} \rightarrow 0 \text{ (} n \rightarrow \infty \text{)}$$

Indeed, we can define

$$h_n(v) = \sum_{k=1}^n g_k(v) \int f(u) f_k(u) d\mu_1^{\pi(v)}(u).$$

**Remark 26** Let  $G$  be a locally compact second countable groupoid with proper orbit space. Let  $F$  be a Borel subset of  $G^{(0)}$  containing only one element  $e(u)$  in each orbit  $[u]$ . Let us assume that  $F \cap [K]$  has a compact closure for each compact subset  $K$  of  $G^{(0)}$ , and let  $\sigma : G^{(0)} \rightarrow G^F$  be a regular cross section for  $d_F : G^F \rightarrow G^{(0)}$ . Let us endow  $G_F^F$  with the quotient topology induced by  $q : G \rightarrow G_F^F$

$$q(x) = \sigma(r(x)) x \sigma(d(x))^{-1}, \quad x \in G$$

If  $g \in C_c(G_F^F)$  and  $g_1, g_2$  are two functions on  $G^{(0)}$  with the property that there

is two sequences  $(h_n^1)_n$  and  $(h_n^2)_n$  of compactly supported continuous functions on  $G^{(0)}$  such that

$$\sup_{\dot{u}} \int |g_i - h_n^i|^2 d\mu_{\dot{u}}^i \rightarrow 0 \quad (n \rightarrow \infty)$$

for  $i = 1, 2$ , then

$$x \xrightarrow{f} g_1(r(x)) g(\sigma(r(x)) x \sigma(d(x))^{-1}) g_2(d(x))$$

can be viewed as an element of  $C^*(G, \nu)$ . Indeed, it is easy to see that

$$\|f - (h_n^1 \circ r)(g \circ q)(h_n^2 \circ d)\|_{II} \rightarrow 0 \quad (n \rightarrow \infty).$$

**Proposition 27** *Let  $G$  be a locally compact second countable principal proper groupoid. Let  $\nu_i = \{\nu_i^u, u \in G^{(0)}\}$ ,  $i = 1, 2$  be two Haar system on  $G$  and  $(\{\beta_v^u\}, \{\mu_i^{\dot{u}}\})$  the corresponding decompositions over the principal groupoid. If the Hilbert bundles determined by the systems of measures  $\{\mu_i^{\dot{u}}\}_{\dot{u}}$  have continuous bases, then the  $C^*$ -algebras  $C^*(G, \nu_1)$  and  $C^*(G, \nu_2)$  are  $*$ -isomorphic.*

**Proof.** We use the Notation 19. From Proposition 20  $\mathbf{C}_\sigma(G)$  is dense in  $C_c(G)$  for the inductive limit topology and hence is dense in  $C^*(G, \nu_1)$ . We shall define a  $*$ -homomorphism  $\Phi$  from  $\mathbf{C}_\sigma(G)$  to  $C^*(G, \nu_2)$ . It suffices to define  $\Phi$  on the set of function on  $G$  of the form

$$x \rightarrow g_1(r(x)) g(q(x)) g_2(d(x))$$

where  $g_1, g_2 \in C_c(G^{(0)})$  and  $g \in C_c(G_F^F)$ . Let  $\{U_{\dot{u}}\}_{\dot{u}}$  be the family of unitary operators with the properties stated in Remark 25 associated to the systems of measures  $\{\mu_i^{\dot{u}}\}_{\dot{u}}$ ,  $i = 1, 2$ .

Let us define  $\Phi$  by

$$\Phi(f) = (x \rightarrow U_{\pi(r(x))}(g_1)(r(x)) g(q(x)) U_{\pi(d(x))}(g_2)(d(x)))$$

where  $f$  is defined by

$$f(x) = g_1(r(x)) g(q(x)) g_2(d(x))$$

with  $g_1, g_2 \in C_c(G^{(0)})$  and  $g \in C_c(G_F^F)$ .

We noted in Remark 26 that the functions of the form  $\Phi(f)$  can be viewed as elements of  $C^*(G, \nu_2)$ . With the same argument as in the proof of Theorem 9, it follows that  $\Phi$  can be extended to  $*$ -isomorphism between  $C^*(G, \nu_1)$  and  $C^*(G, \nu_2)$ . ■

## References

- [1] C. Anantharaman-Delaroche, J. Renault, *Amenable groupoids*, Monographie de L'Enseignement Mathematique No **36**, Geneve, 2000.
- [2] G. Mackey, *Induced representations of locally compact groups. I*, Ann. of Math., **55**(1952), 101-139.
- [3] P. Muhly, J. Renault and D. Williams, *Equivalence and isomorphism for groupoid  $C^*$ -algebras*, J. Operator Theory **17**(1987), 3-22.
- [4] P. Muhly, *Coordinates in operator algebra*, (Book in preparation).
- [5] A. Ramsay, *Virtual groups and groups actions*, Adv. in Math. **6**(1971), 253-322.
- [6] A. Ramsay, *Topologies on measured groupoids*, J. Funct. Anal. **47**(1982), 314-343.
- [7] A. Ramsay and M. E. Walter, *Fourier-Stieltjes Algebras of locally compact groupoids*, J. Funct. Anal. **148**(1997), 314-367.
- [8] J. Renault, *A groupoid approach to  $C^*$ -algebras*, Lecture Notes in Math., Springer-Verlag, **793**, 1980.
- [9] J. Renault, *The ideal structure of groupoid crossed product algebras*, J. Operator Theory, **25**(1991), 3-36.
- [10] J. Westman, *Nontransitive groupoid algebras*, Univ. of California at Irvine, 1967.

University Constantin Brâncuși of Târgu-Jiu  
Bulevardul Republicii, Nr. 1,  
210152 Târgu-Jiu , Gorj  
Romania  
e-mail: ada@utgjiu.ro