Navier-Stokes Equation on the Rectangle¹

Sérgio S. Rodrigues²

April 11, 2005

¹Supported by FCT (Portuguese Foundation for Science and Tecnology). ²SISSA-ISAS, via Beirut 2-4, Trieste 34014, Italy & University of Aveiro, Portugal; e-mail: srodrigs@sissa.it

Contents

2Existence, Continuity and Uniqueness.72.1The Spaces.72.1.1Recollection of Auxiliary Material on the Spaces $L^2(R)$ and $H^1(R)$ 2.1.2Some Properties of the Spaces V and H.82.1.3Poincaré Inequality. Equivalent Norms.92.2The Duals of V and H.92.3Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V.102.3.1Fourier Characterization of H and V.112.4The Operators A and B.132.4.1The Operator A.132.4.2The Operator B.162.5Classical and Weak Formulations.172.6Existence212.6.1Fourier Transform. Fractional Derivatives.212.7Uniqueness.292.8Continuity on Initial Data.312.9Some More Estimates for the Form b.332.10Strong Formulation.342.11Existence.352.12Uniqueness.372.13Continuity on Initial Data.37
2.1The Spaces.72.1.1Recollection of Auxiliary Material on the Spaces $L^2(R)$ and $H^1(R)$.2.1.2Some Properties of the Spaces V and H.82.1.3Poincaré Inequality. Equivalent Norms.92.2The Duals of V and H.92.3Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V.102.3.1Fourier Characterization of H and V.112.4The Operators A and B.132.4.1The Operator A.132.4.2The Operator B.162.5Classical and Weak Formulations.172.6Existence212.6.1Fourier Transform. Fractional Derivatives.212.6.2The Existence Theorem.212.7Uniqueness.292.8Continuity on Initial Data.332.9Some More Estimates for the Form b.332.10Strong Formulation.373Controllability of Galerkin Approximations.41
2.1.1Recollection of Auxiliary Material on the Spaces $L^2(R)$ and $H^1(R)$.2.1.2Some Properties of the Spaces V and H .82.1.3Poincaré Inequality. Equivalent Norms.92.2The Duals of V and H .92.3Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V .102.3.1Fourier Characterization of H and V .112.4The Operators A and B .132.4.1The Operator A .132.4.2The Operator B .162.5Classical and Weak Formulations.172.6Existence212.6.1Fourier Transform. Fractional Derivatives.212.6.2The Existence Theorem.292.8Continuity on Initial Data.312.9Some More Estimates for the Form b .332.10Strong Formulation.342.11Existence.372.13Continuity on Initial Data.373Controllability of Galerkin Approximations.41
2.1.2Some Properties of the Spaces V and H.82.1.3Poincaré Inequality. Equivalent Norms.92.2The Duals of V and H.92.3Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V.102.3.1Fourier Characterization of H and V.112.4The Operators A and B.132.4.1The Operator A.132.4.2The Operator B.162.5Classical and Weak Formulations.172.6Existence212.6.1Fourier Transform. Fractional Derivatives.212.6.2The Existence Theorem.212.7Uniqueness.292.8Continuity on Initial Data.312.9Some More Estimates for the Form b.332.10Strong Formulation.342.11Existence.372.13Continuity on Initial Data.373Controllability of Galerkin Approximations.41
2.1.3Poincaré Inequality. Equivalent Norms.92.2The Duals of V and H.92.3Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V.102.3.1Fourier Characterization of H and V.112.4The Operators A and B.132.4.1The Operator A.132.4.2The Operator B.162.5Classical and Weak Formulations.172.6Existence212.6.1Fourier Transform. Fractional Derivatives.212.6.2The Existence Theorem.212.7Uniqueness.292.8Continuity on Initial Data.332.10Strong Formulation.342.11Existence.352.12Uniqueness.372.13Controllability of Galerkin Approximations.41
2.2The Duals of V and H.92.3Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V.102.3.1Fourier Characterization of H and V.112.4The Operators A and B.132.4.1The Operator A.132.4.2The Operator B.162.5Classical and Weak Formulations.172.6Existence212.6.1Fourier Transform. Fractional Derivatives.212.6.2The Existence Theorem.212.7Uniqueness.292.8Continuity on Initial Data.332.10Strong Formulation.342.11Existence.352.12Uniqueness.372.13Continuity of Galerkin Approximations.41
2.3Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V .102.3.1Fourier Characterization of H and V .112.4The Operators A and B .132.4.1The Operator A .132.4.2The Operator B .162.5Classical and Weak Formulations.172.6Existence212.6.1Fourier Transform. Fractional Derivatives.212.6.2The Existence Theorem.212.6.2The Existence Theorem.212.7Uniqueness.292.8Continuity on Initial Data.332.10Strong Formulation.342.11Existence.352.12Uniqueness.373Controllability of Galerkin Approximations.41
2.3.1 Fourier Characterization of H and V. 11 2.4 The Operators A and B. 13 2.4.1 The Operator A. 13 2.4.2 The Operator B. 16 2.5 Classical and Weak Formulations. 17 2.6 Existence 21 2.6.1 Fourier Transform. Fractional Derivatives. 21 2.6.2 The Existence Theorem. 21 2.6.2 The Existence Theorem. 29 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Continuity on Initial Data. 37 3.13 Continuity on Initial Data. 37 3.14 Data. 37 3 Continuity on Initial Data. 37 3 Continuity on Initial Data. 37 3 Continuity on Initial Data. 37 3
2.4 The Operators A and B. 13 2.4.1 The Operator A. 13 2.4.2 The Operator B. 16 2.5 Classical and Weak Formulations. 17 2.6 Existence 17 2.6.1 Fourier Transform. Fractional Derivatives. 21 2.6.2 The Existence Theorem. 21 2.7 Uniqueness. 29 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.4.1 The Operator A. 13 2.4.2 The Operator B. 16 2.5 Classical and Weak Formulations. 17 2.6 Existence 17 2.6.1 Fourier Transform. Fractional Derivatives. 21 2.6.2 The Existence Theorem. 21 2.7 Uniqueness. 21 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Continuity on Initial Data. 37 3.1 Existence. 37 3.1 Existence. 37 3.1 Existence. 37 3.1 Strong Formulation. 37 3.13 Continuity on Initial Data. 37 3 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.4.2 The Operator B. 16 2.5 Classical and Weak Formulations. 17 2.6 Existence 21 2.6.1 Fourier Transform. Fractional Derivatives. 21 2.6.2 The Existence Theorem. 21 2.7 Uniqueness. 21 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3
2.5 Classical and Weak Formulations. 17 2.6 Existence 21 2.6.1 Fourier Transform. Fractional Derivatives. 21 2.6.2 The Existence Theorem. 21 2.7 Uniqueness. 21 2.7 Uniqueness. 29 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.6 Existence 21 2.6.1 Fourier Transform. Fractional Derivatives. 21 2.6.2 The Existence Theorem. 21 2.7 Uniqueness. 29 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.6.1 Fourier Transform. Fractional Derivatives. 21 2.6.2 The Existence Theorem. 21 2.7 Uniqueness. 29 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.6.2 The Existence Theorem. 21 2.7 Uniqueness. 29 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.7 Uniqueness. 29 2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.8 Continuity on Initial Data. 31 2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.9 Some More Estimates for the Form b. 33 2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 35 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41
2.10 Strong Formulation. 34 2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 37 37 38 37 39 Controllability of Galerkin Approximations. 41 41
2.11 Existence. 35 2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41 2.1 The ECE Decedure 41
2.12 Uniqueness. 37 2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41 2.1 The ECE Decedure 41
2.13 Continuity on Initial Data. 37 3 Controllability of Galerkin Approximations. 41 2.1 The ECE Decedure 41
3 Controllability of Galerkin Approximations. 41
2.1 The ECE Dressdume 41
\mathbf{a} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{c}
3.1.1 A Lemma from Linear Algebra
3.1.2 Factorization.
3.1.3 Convexification.
3.1.4 Extraction.
3.1.5 Iterating FCEs. 43
3.2 The Projection onto H
3.3 Spectral Method
3.4 The Infinite ODE System
3.4.1 Galerkin Approximations and G-Saturating Sets
3.5 Looking for Saturating Sets
3.6 Exact Controllability of Galerkin Approximations

		3.6.1 Lie Brackets. Full Lie Rank Property	3
		3.6.2 Zero Orbits and Zero Ideal)
4	Con	trollability in Observed Component. 73	3
	4.1	Controlled N-S Problem. Existence, Uniqueness and Continuity. 73	3
	4.2	Change of Variables	1
	4.3	Weak Case	1
	4.4	Existence	5
	4.5	Uniqueness. 7'	7
	4.6	Continuity 78	R
	4.7	Strong Case 8	ŝ
	1.1	471 Existence 84	1
		4.7.2 Uniqueness 86	т 3
		4.7.2 Continuity 86	3
	10	Continuity in Polaration Matric) 1
	4.0	A Demark on Finite System yargue Infinite System	L 2
	4.9	Calid Controllability in Observed Control System.) 2
	4.10	Solid Controllability in Observed Component)
		4.10.1 First Step. Proposition 4.10.2: N Big	(
		4.10.2 Second Step. Proposition 4.10.2: "Back-Induction" 10	L
	4.11	Proof of Lemma 4.10.9)
5	\mathbf{L}^2 -A	Approximate Controllability. 131	L

Chapter 1

Introduction

Following part of the work iniciated by A. Agrachev and A. Sarychev in [3] we study controllability, by means of low modes forcing, of incompressible 2D Navier-Stokes (NS) Equations on the two dimensional rectangle with tangent velocity on the boundary.

In the present paper we deal with the 2D NS system

$$u_t + (u \cdot \nabla)u + \nabla p = \nu \Delta u + F(x_1, x_2) + v(t, x_1, x_2)$$
(1.0.1)

- $\nabla \cdot u = 0 \quad \text{on} \quad R \tag{1.0.2}$
- $u \cdot \mathbf{n} = 0 \quad \text{on} \quad \partial R;$ (1.0.3)

$$\nabla^{\perp} \cdot u = 0 \quad \text{on} \quad \partial R \tag{1.0.4}$$

Where $R := \{(x_1, x_2) \in \mathbb{R}^2 \mid a_1 < x_1 < a_2; b_1 < x_2 < b_2\}$ and $\nabla^{\perp} := \begin{pmatrix} -\frac{\partial}{\partial x_2} \\ \frac{\partial}{\partial x_1} \end{pmatrix}$

and **n** is the unit normal to the boundary. In the equation (1.0.1) u is the velocity of the fluid "particle"; p is the pressure; the only nonlinear term of the equation $(u \cdot \nabla)u$ is called the *inertial term*; $\nu \Delta u$ is called the viscosity term, $\nu > 0$ is the coeficient of viscosity; F is an external force and; v is a control at our disposal. We are interested in the case where v is a degenerate forcing, i.e., v is a finite sum of harmonics $-v = \sum_{k \in \mathcal{K}^1} v_k(t) E_k$, where E_k are eigenfunctions of the Stokes operator. So the components $v_k(t)$, $k \in \mathcal{K}^1$, $t \in [0, T]$ are our controls which are measurable essentially bounded functions. We shall study Galerkin approximations, say, big enough to contain a set of modes we want to observe.

A natural way to study the NSE is to study its evolution on subspaces of Sobolev spaces; such subspaces depend on the boundary conditions.

We shall note by $L^2(R)$ the space of Lebesgue measurable square integrable real functions defined on R and by $\mathbf{L}^2(R)$ the product space $L^2(R)^2$. Similarly $H^1 := \{f \in L^2(R) \mid \frac{\partial f}{\partial x_j} \in L^2(R), j = 1, 2\}$ and, $\mathbf{H}^1(R) := H^1(R)^2$.

The most studied boundary conditions are the full Dirichlet (u = 0 on the boundary, also called no-slip) and periodic (u_1 and u_2 are periodic). For the full Dirichlet conditions the study is done in the spaces

$$H_D := \{ u \in \mathbf{L}^2(R) \mid \nabla \cdot u = 0 \& u \cdot \mathbf{n} = 0 \text{ on } \partial R \}$$
$$V_D := \{ u \in \mathbf{H}^1(R) \mid \nabla \cdot u = 0 \& u = 0 \text{ on } \partial R \}$$

and for the periodic boundary conditions the study is done in

$$H_P := \{ u \in \mathbf{L}^2_{per}(R) \mid \nabla \cdot u = 0 \}$$
$$V_P := \{ u \in \mathbf{H}^1_{per}(R) \mid \nabla \cdot u = 0 \}.$$

In the periodic case sometimes the special case of zero average space — $\int_R u \, dx = 0$ — is considered, in this case we have to take this new condition into account in the definition of the spaces H_P and V_P . For more information on the boundary conditions above see for example [15] or [14].

For the boundary conditions (1.0.3), (1.0.4) the spaces

$$H := \{ u \in \mathbf{L}^{2}(R) \mid \nabla \cdot u = 0 \& u \cdot \mathbf{n} = 0 \text{ on } \partial R \};$$

$$V := \text{ closure of } \mathcal{D}_{1}(R) \text{ on } \mathbf{H}^{1}(R); \qquad (1.0.5)$$

$$D(A) := \{ u \in \mathbf{H}^{2}(R) \mid \nabla \cdot u = 0 \& (u \cdot \mathbf{n} = 0 \land \nabla^{\perp} \cdot u = 0) \text{ on } \partial R \};$$

where $\mathcal{D}_1(R) := \{ u \in C^{\infty}(\overline{R}) \mid \nabla \cdot u = 0 \& (u \cdot \mathbf{n} = 0 \land \nabla^{\perp} \cdot u = 0) \text{ on } \partial R \},\$ are those where we shall consider the evolution of the NSE on.

In Chapter 2 we prove the existence of weak and strong solutions for (1.0.1)-(1.0.4), as well as its uniqueness and continuous dependence in the initial data. For the case of weak solutions we proceed as in [15] for the case of no-slip boundary conditions. To obtain strong solutions we just have to ask for some regularity on the initial data.

In Chapter 3, for strong solutions, we prove controllability, by means of low modes forcing, of Galerkin approximations of the infinite-dimensional system associated with (1.0.1)-(1.0.4).

In Chapter 4 we prove the so called *controllability in observed component*, again by means of low modes forcing. In other words we prove that the projection, onto any finite dimensional subspace (the space spanned by the modes we want to observe), of the attainable set from any point is surjective.

Finally, we end with Chapter 5 and with the prove of \mathbf{L}^2 -Approximate Controllability the is a straightforward corollary of some tools we have presented in Chapter 4.

The author is gratefull to A. Agrachev and A. Sarychev for the inspiring and helpfull discussions on the subject and, for the suggestions in the improvement of the text.

The author would like to thank FCT (Portuguese Foundation for Science and Tecnology) for financial support and, SISSA-ISAS (International School for Advanced Studies) for hospitality.

Chapter 2

Existence, Continuity and Uniqueness.

2.1 The Spaces.

2.1.1 Recollection of Auxiliary Material on the Spaces $L^2(R)$ and $H^1(R)$.

Recall that since $L^2(R)$ is a Hilbert space for the scalar product

$$(u, v)_1 := \int_R uv \, dx,$$
 (2.1.1)

then $\mathbf{L}^2(R)$ is a Hilbert space for the product topology and the scalar product is

$$(u, v) := \int_R u \cdot v \, dx. \tag{2.1.2}$$

We note that

$$(u, v) = \int_{R} u \cdot v \, dx = \sum_{i=1}^{2} \int_{R} u_{i} v_{i} \, dx = (u_{1}, v_{1})_{1} + (u_{2}, v_{2})_{1}.$$

The norms associated with the previous scalar products shall be represented by

$$|u|_1^2 := (u, u)_1 \tag{2.1.3}$$

$$|u|^2 := (u, u) \tag{2.1.4}$$

We note that

$$|u|^2 := |u_1|_1^2 + |u_2|_1^2$$

so, the norm $|\cdot|$ is the product norm $|\cdot|_1 \times |\cdot|_1$. Similarly, the Sobolev space $H^1(R)$ is a Hilbert space for the scalar product

$$((u, v))_{01} := (u, v)_1 + \sum_{i=1}^2 (\frac{\partial u}{\partial x_i}, \frac{\partial v}{\partial x_i})_1 = (u, v)_1 + (\nabla u, \nabla v).$$
(2.1.5)

then $\mathbf{H}^1(R)$ is a Hilbert space for the product topology and the scalar product is

$$((u, v))_0 := \sum_{j=1}^2 ((u_j, v_j))_{01} = (u, v) + \sum_{i=1}^2 (\nabla u_i, \nabla v_i)$$
(2.1.6)

The norms associated with the previous scalar products shall be represented by

$$||u||_{01}^2 := ((u, u))_{01}$$
(2.1.7)

$$||u||_0^2 := ((u, u))_0 \tag{2.1.8}$$

We note that

$$||u||_0^2 := ||u_1||_{01}^2 + ||u_2||_{01}^2$$

so, the norm $\|\cdot\|_0$ is the product norm $\|\cdot\|_{01} \times \|\cdot\|_{01}$.

2.1.2 Some Properties of the Spaces V and H.

Lemma 2.1.1. *H* coincides with the closure of $\mathcal{D}_1(R)$ in $\mathbf{L}^2(R)$.

Proof. It is well known from the study of the NSE with full Dirichlet boundary conditions that H is the closure of $\mathcal{D}(R)$ in $\mathbf{L}^2(R)$, where $\mathcal{D}(R)$ is the set of solenoidal (or divergence free) smooth functions with support in R. Since

$$\mathcal{D}(R) \subset \mathcal{D}_1(R) \subset H$$

we conclude that H coincides with the closure of $\mathcal{D}_1(R)$ in $\mathbf{L}^2(R)$.

We use the same argument, since $\mathcal{D}_1 \subset V \subset H$ to conclude that

Corollary 2.1.2. *H* is the closure of *V* in L^2 .¹

In the study of NSE some classical imbedding and compactness theorems are frequently used, we start by presenting some of them which we shall need (for our particular equation).

Proposition 2.1.3. For all $u \in H^1(R)$ and for all $q, 1 \leq q < \infty$ we have

$$||u||_{L^q(R)} \le C_1(q) ||u||_{01}$$

Corollary 2.1.4. For all $u \in \mathbf{H}^1(R)$ and for all $q, 1 \leq q < \infty$ we have

$$||u||_{\mathbf{L}^{q}(R)} \leq C_{1}(q)||u||_{0}$$

Proposition 2.1.5. The imbedding

$$H^1(R) \to L^2(R)$$
$$u \mapsto u$$

is compact.

Corollary 2.1.6. The imbedding

$$\mathbf{H}^{1}(R) \to \mathbf{L}^{2}(R)$$
$$u \mapsto u$$

is compact.

¹In other words V is dense in the closed space H

Corollary 2.1.7. The inclusion

$$\begin{aligned} i: V \to H \\ u \mapsto u \end{aligned} \tag{2.1.9}$$

is continuous and compact.

Proof. By Corollary 2.1.4, inclusion (2.1.9) is continuous and, by Corollary 2.1.6 it is compact. $\hfill \Box$

2.1.3 Poincaré Inequality. Equivalent Norms.

We have the following Lemma: 2

Lemma 2.1.8.

• For any seminorm p in $\mathbf{H}^1(R)$ satisfying " $(p(a) = 0 \land a \in \mathbb{R}^2) \Rightarrow a = 0$ " we have

$$|u| \le c \Big(\|\nabla u\|_{\mathbf{L}^2(R)^2} + p(u) \Big), \quad \forall u \in \mathbf{H}^1(R)$$

and,

• The seminorm

$$p(u) := \int_{\Gamma} |u \cdot \mathbf{n}| \, d\Gamma$$

satisfies the required condition.

Remark 1. The product space $\mathbf{L}^2(R)^2$ is a Hilbert space for the scalar product

$$(U, V) := ((U_1, V_1))_0 + ((U_2, V_2))_0, \quad U = \begin{pmatrix} U_1 \\ U_2 \end{pmatrix}, V = \begin{pmatrix} V_1 \\ V_2 \end{pmatrix} \in \mathbf{L}^2(R)^2$$

and, $\nabla \begin{pmatrix} V_1 \\ V_2 \end{pmatrix} := \begin{pmatrix} \nabla V_1 \\ \nabla V_2 \end{pmatrix} \quad \forall V \in \mathbf{L}^2(R)^2.$

A consequence of this lemma is a Poincaré-like inequality:

$$\forall u \in V[|u| \le c \|\nabla u\|_{\mathbf{L}^2(R)^2}].$$
(2.1.10)

Corollary 2.1.9. The norms $||u|| := ||\nabla u||_{\mathbf{L}^2(R)^2}$ and $||\cdot||_0$ are equivalent in $V.^3$

2.2 The Duals of V and H.

From now we shall consider the space V endowed with the norm $\|\cdot\|$. From Corollaries 2.1.2, 2.1.7 and 2.1.9 we obtain that the inclusion (2.1.9)(considering V endowed with $\|\cdot\|$) is dense continuous and compact.

Since H is an Hilbert Space due to the Riesz Representation Theorem we can identify H with its dual H'. In this way we arrive to the inclusions

$$V \subset H \equiv H' \subset V' \tag{2.2.1}$$

 $^{^2 \}mathrm{See}$ [13] subsections II.1.4 and III.2.2

³Note that $\|\cdot\|_0 = |\cdot| + \|\cdot\|$.

where both inclusions are linear, dense and continuous. Indeed we already know that the first inclusion has these properties. In the second inclusion we consider the map i', dual to the map i defined in (2.1.9),

$$\begin{aligned} i': H' \to V' \\ < i(v), h > = < v, i'(h) > \end{aligned}$$

that is linear and continuous — we consider V' endowed with the classical norm $||f|| := \sup_{\|v\|=1} |\langle f, v \rangle|$. The density of H' in V' is a corollary of the following Lemma, which one can find in [4], section (II.6),

Lemma 2.2.1. Let $G : D(G) \subset E \to F$ be a (nonbounded) closed operator with $\overline{D(G)} = E$. Then

(i)
$$N(G) = R(G')^{\perp};$$

(ii) $N(G') = R(G)^{\perp};$
(iii) $N(G)^{\perp} \supset \overline{R(G')};$
(iv) $N(G')^{\perp} = \overline{R(G)}.$

Where \overline{X} means closure of X; N(G) stays for the Kernel of G and R(G) for the image of G. Applying this Lemma to our inclusions the injectivity of the first inclusion implies the density of the second — (i). Moreover we also see that, since the first inclusion is dense, it follows the injectivity of the second (ii).

Remark 2. The domain D(i) of i is V. By the continuity of i we conclude its closedness. Recall that a operator $G : E \to F$ is said closed if for any sequence

$$(u_n) \text{ in } D(G) \text{ such that } \begin{cases} u_n \to u & \text{in } E\\ Au_n \to f & \text{in } F \end{cases} \text{ we have } \begin{cases} u \in D(G)\\ Au &= f \end{cases}$$

2.3 Fourier Series. A Basis in $\mathcal{D}_1(R)$ for H and V.

We start by noting that if in the case of the general rectangle $R := [a_1, a_2] \times [b_1, b_2]$ $a_1 < a_2, b_1 < b_2$ we make the change of variables

$$z_1 := x_1 - a_1, \quad z_2 := x_2 - b_1,$$

since the differential operators ∇ , Δ are invariant under tranlations, the Navier-Stokes Equation does not change in the variables (z_1, z_2) . To simplify the exposition from now we shall deal with the rectange

$$R := \{ (x_1, x_2) \in \mathbb{R}^2 \mid 0 \le x_1 \le a; 0 \le x_2 \le b \}.$$

First we note that under the condition $u \cdot \mathbf{n} = 0$ on ∂R , the condition $\nabla^{\perp} \cdot u = 0$ on ∂R equivals $\frac{\partial u_1}{\partial x_2} = \frac{\partial u_2}{\partial x_1} = 0$ on ∂R , because on the left and right faces we have $\frac{\partial u_1}{\partial x_2} = 0$ and, on the top and botton faces $\frac{\partial u_2}{\partial x_1} = 0$. It is well known that $\{\sin(\frac{n\pi x}{L}) \mid n \in \mathbb{N}_0\}$ is a basis for the functions in $L^2([0, L])$ vanishing on 0

and L and, $\{\cos(\frac{n\pi x}{L}) \mid n \in \mathbb{N}\}\$ is a basis for the functions in $L^2([0, L])$ with vanishing velocity on the boundary points 0 and L.⁴ So

$$\left\{\sin\left(\frac{n\pi x}{a}\right)\cos\left(\frac{m\pi x}{b}\right) \mid n \in \mathbb{N}_0, m \in \mathbb{N}\right\}$$

is a basis for $L^2(R)$ vanishing on the right and left faces of the rectangle and,

$$\left\{\cos\left(\frac{n\pi x}{a}\right)\sin\left(\frac{m\pi x}{b}\right) \mid n \in \mathbb{N}, \ m \in \mathbb{N}_0\right\}$$

is a basis for $L^2(R)$ vanishing on the botton and top faces of the rectangle.

Now consider a function $u \in H$, this function, being an element of $\mathbf{L}^{2}(R)$, can be written as

$$u = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} \sum_{\substack{n \in \mathbb{N}_0 \\ m \in \mathbb{N}}} u_{1k} \sin\left(\frac{n\pi x_1}{a}\right) \cos\left(\frac{m\pi x_2}{b}\right) \\ \sum_{\substack{m \in \mathbb{N}_0 \\ n \in \mathbb{N}}} u_{2k} \cos\left(\frac{n\pi x_1}{a}\right) \sin\left(\frac{m\pi x_2}{b}\right) \end{pmatrix}.$$

Since u is solenoidal — $\nabla \cdot u = 0$, we conclude that

$$u = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} \sum_{k \in \mathbb{N}_0^2} u_{1k} \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\ \sum_{k \in \mathbb{N}_0^2} u_{2k} \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix}.$$

and that for each $k \in \mathbb{N}_0^2$ we have

$$u_{1k}\frac{k_1\pi}{a} + u_{2k}\frac{k_2\pi}{b} = 0.$$

This means that $\frac{u_{1k}b}{-k_2\pi} = \frac{u_{2k}a}{k_1\pi} =: u_k$ and, we arrive to

$$u = \sum_{k \in \mathbb{N}_0^2} u_k W_k, \quad W_k := \begin{pmatrix} \frac{-k_2 \pi}{b} \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\ \frac{k_1 \pi}{a} \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix}.$$

We put

$$\mathcal{W} := \{ W_k \mid k \in \mathbb{N}_0^2 \}.$$
(2.3.1)

2.3.1 Fourier Characterization of *H* and *V*.

We start by computing the scalar product (W_k, W_z) between two elements of \mathcal{W} :

$$(W_k, W_z) = \int_R W_k \cdot W_z \, dx$$

= $\int_R \frac{k_2 z_2 \pi^2}{b^2} \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \sin\left(\frac{z_1 \pi x_1}{a}\right) \cos\left(\frac{z_2 \pi x_2}{b}\right) \, dx$
+ $\int_R \frac{k_1 z_1 \pi^2}{a^2} \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \cos\left(\frac{z_1 \pi x_1}{a}\right) \sin\left(\frac{z_2 \pi x_2}{b}\right) \, dx$
= $\begin{cases} 0 & \text{if } k \neq z \\ -\bar{k} \frac{ab}{4} & \text{if } k = z. \end{cases}$

⁴Here \mathbb{N} is the set of natural numbers and $\mathbb{N}_0 := \mathbb{N} \setminus \{0\}$

where $\bar{k} := -\pi^2 \left(\frac{k_1^2}{a^2} + \frac{k_2^2}{b^2}\right)$. Hence the square of the $\mathbf{L}^2(R)$ norm of each element in the orthogonal family \mathcal{W} equals $-\bar{k}\frac{ab}{4}$.

Given $u, v \in H$ the scalar product between them is then

$$(u, v) = \sum_{k \in \mathbb{N}_0^2} -\bar{k} \frac{ab}{4} u_k v_k,$$

where

$$u = \sum_{k \in \mathbb{N}_0^2} u_k W_k; \quad v = \sum_{k \in \mathbb{N}_0^2} v_k W_k.$$

The norm of u in H results

$$|u|^2 = \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} -\bar{k}u_k^2 \tag{2.3.2}$$

and $u \in H$ means that

$$\sum_{k\in\mathbb{N}_0^2} -\bar{k}u_k^2 < +\infty.$$

If $u, v \in V$ we want to compute the scalar product

$$((u, v)) = ((\nabla u_1, \nabla v_1)) + ((\nabla u_2, \nabla v_2)).$$

First we note that

$$\nabla u_1 = \left(\sum_{k \in \mathbb{N}_0^2} u_k \left(-\frac{k_2 \pi}{b} \frac{k_1 \pi}{a} \right) \cos\left(\frac{k_1 \pi x_1}{a} \right) \cos\left(\frac{k_2 \pi x_2}{b} \right) \\ \sum_{k \in \mathbb{N}_0^2} u_k \left(-\frac{k_2 \pi}{b} \frac{-k_2 \pi}{b} \right) \sin\left(\frac{k_1 \pi x_1}{a} \right) \sin\left(\frac{k_2 \pi x_2}{b} \right) \right),$$
$$\nabla u_2 = \left(\sum_{k \in \mathbb{N}_0^2} u_k \left(\frac{k_1 \pi}{a} \frac{-k_1 \pi}{a} \right) \sin\left(\frac{k_1 \pi x_1}{a} \right) \sin\left(\frac{k_2 \pi x_2}{b} \right) \\ \sum_{k \in \mathbb{N}_0^2} u_k \left(\frac{k_1 \pi}{a} \frac{k_2 \pi}{b} \right) \cos\left(\frac{k_1 \pi x_1}{a} \right) \cos\left(\frac{k_2 \pi x_2}{b} \right) \right)$$

and we obtain analogous expressions for ∇v_1 and ∇v_2 . Hence, doing the computations, we arrive to

$$((u, v)) = \sum_{k \in \mathbb{N}_0^2} \bar{k}^2 \frac{ab}{4} u_k v_k.$$

The norm of u in V results

$$||u||^2 = \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} \bar{k}^2 u_k^2 \tag{2.3.3}$$

and $u \in V$ means that

$$\sum_{k\in\mathbb{N}_0^2}\bar{k}^2u_k^2<+\infty.$$

From caracterizations (2.3.2) and (2.3.3) we can see easily, as referred in Corollary 2.1.9, that the norms $\|\cdot\|_0$ and $\|\cdot\|$ are equivalent in V. Indeed, $-\bar{k}$ goes

⁵This follows from the fact that the family of sines is orthogonal in $L^2([0, \pi])$, as well is the family of cosines.

to $+\infty$ when $\max\{k_1, k_2\}$ does⁶ so, there is $N \in \mathbb{N}_0$ and a constant $C \ge 1$ such that if $\max\{k_1, k_2\} > N$ we have $-\bar{k} \ge 1$ and,

$$\sum_{\max\{k_1, k_2\} \le N} -\bar{k} \le C \sum_{\max\{k_1, k_2\} \le N} \bar{k}^2.$$
(2.3.4)

Therefore $|u| \leq C ||u||$ and then $||\cdot||_0$ and $||\cdot||$ are equivalent.

When $u \in V$ we can not guarantee, in general, neither $\Delta u \in H$ nor $\Delta u \in \mathbf{L}^2$. Anyway if we compute $(\Delta u, v)$ for $u, v \in V$ we obtain

$$(\Delta u, v) = \sum_{k \in \mathbb{N}_0^2} \bar{k}^2 u_k^2 < +\infty = -((u, v)).$$
(2.3.5)

Hence the result of $(\Delta u, v)$ is finite and is properly defined even if $\Delta u \notin \mathbf{L}^2$ and so the operation $(\Delta u, v)$ seems to have no sense in itself. Below we will give a sense to it.

2.4 The Operators *A* and *B*.

2.4.1 The Operator A.

First we note that a consequence of the identifications (2.2.1), the scalar product in H of $f \in H$ and $u \in V$ is the same as the scalar product of f and u in the duality between V and V'

$$\forall f \in H \; \forall u \in V[\langle f, u \rangle = (f, u)]. \tag{2.4.1}$$

For each $u \in V$, the form

$$V \to \mathbb{R}$$
(2.4.2)
$$v \mapsto ((u, v))$$

is linear and continuous on V. Therefore there is an element of V' we shall denote by Au satisfying

$$\langle Au, v \rangle = ((u, v)), \quad \forall v \in V$$

$$(2.4.3)$$

and A is clearly linear and continuous; we can also see that A is an isomorphism between V and V'. Indeed the form a(u, v) := ((u, v)) is bilinear, continuous and coersive on $V \times V$ so, by Lax-Milgram Theorem we can conclude that A is an isomorphism between V and V'.

By equation (2.3.5) we see that

$$\langle Au, v \rangle = (-\Delta u, v) \quad \forall u, v \in V.$$
 (2.4.4)

⁶For $M := \max\{k_1, k_2\}$ and $m := \max\{a^2, b^2\}$, we have $-\bar{k} = \pi^2(\frac{k_1^2}{a^2} + \frac{k_2^2}{b^2}) \ge \pi^2 \frac{M^2}{m}$ that goes to ∞ when M does.

Further Properties of A.

As we have said above for $v \in V$, Av can be not in \mathbf{L}^2 . Now we define a subset D(A) of V, we shall call the domain of A, such that

$$A: D(A) \to H.$$

The domain of A is defined by

$$D(A) := \{ v \in \mathbf{H}^2 \mid v \in V \& \nabla^{\perp} \cdot v = 0 \text{ on } \Gamma \}$$

= $\{ v \in \mathbf{H}^2 \mid \nabla \cdot v = 0 \& (v \cdot \mathbf{n} = 0 \land \nabla^{\perp} \cdot v = 0 \text{ on } \Gamma) \}$
= closure of $\mathcal{D}_1(R)$ in \mathbf{H}^2 .

Where $\Gamma := \partial R$ and \mathbf{H}^2 is the Sobolev space $\mathbf{H}^2 = (W^{2,2})^2$ where

$$W^{2,2} := \{ u \in L^2 \mid \frac{\partial^{|\alpha|} u}{\partial x^{\alpha}} \in L^2, \quad \forall |\alpha| \le 2 \}$$

which is a Hilbert space for the scalar product

$$(u, v)_2 := \sum_{\substack{i \in \{1, 2\} \\ |\alpha| \le 2}} (\frac{\partial^{|\alpha|} u_i}{\partial x^{\alpha}}, \frac{\partial^{|\alpha|} v_i}{\partial x^{\alpha}})_1 = \sum_{|\alpha| \le 2} (\frac{\partial^{|\alpha|} u}{\partial x^{\alpha}}, \frac{\partial^{|\alpha|} v}{\partial x^{\alpha}}).$$

The norm corresponding to this scalar product is defined by

$$|u|_{2}^{2} = |u|^{2} + ||u||^{2} + |u|_{[2]}^{2},$$

with

$$u|_{[2]-}^{2} = \sum_{|\alpha|=2} |\frac{\partial^{|\alpha|} u}{\partial x^{\alpha}}|^{2} = \sum_{|\alpha|=2} \left(|\partial_{1}^{2} u|^{2} + |\partial_{2}^{2} u|^{2} + |\partial_{1,2}^{2} u|^{2} \right)^{-7}$$

We can see that the norm $|\cdot|_2$ is equivalent to that $|\cdot|_{2+}$ coming from the scalar product defined by

$$(u, v)_{2+} = (u, v)_2 + (\partial_{1,2}^2 u, \partial_{1,2}^2 v).$$

Indeed $|u|_2^2 \le |u|_{2+}^2 \le 2|u|_2^2$.

Now we look for a Fourier caracterization of the elements on D(A): For that we need to write down the expressions for the second order derivatives of u:

$$\partial_{1,2}^2 u = \sum_{k \in \mathbb{N}_0^2} u_k \begin{pmatrix} -\frac{k_2 \pi}{b} \left(-\frac{k_2 \pi}{b} \right) \frac{k_1 \pi}{a} \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \\ \frac{k_1 \pi}{a} \left(-\frac{k_1 \pi}{a} \right) \frac{k_2 \pi}{a} \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix};$$

$$\partial_1^2 u = \sum_{k \in \mathbb{N}_0^2} u_k \begin{pmatrix} \frac{k_2 \pi}{b} \left(\frac{k_1 \pi}{a}\right)^2 \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\ -\frac{k_1 \pi}{a} \left(\frac{k_1 \pi}{a}\right)^2 \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix};$$

$$\partial_2^2 u = \sum_{k \in \mathbb{N}_0^2} u_k \begin{pmatrix} \frac{k_2 \pi}{b} \left(\frac{k_2 \pi}{b}\right)^2 \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\ -\frac{k_1 \pi}{a} \left(\frac{k_2 \pi}{b}\right)^2 \cos\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix}.$$

$$^{7} \partial_j^2 \text{ stays for } \frac{\partial^2}{\partial x_j^2} \text{ and, } \partial_{1,2}^2 \text{ stays for } \frac{\partial^2}{\partial x_1^2}.$$

Computing $(u, v)_{2+}$ for $u, v \in D(A)$ we obtain

$$\begin{aligned} (u, v)_{2+} &= \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} -\bar{k}u_k v_k + \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} \bar{k}^2 u_k v_k \\ &+ \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} \pi^6 \bigg[2 \Big(\frac{k_2^2 k_1}{b^2 a} \Big)^2 + 2 \Big(\frac{k_1^2 k_2}{a^2 b} \Big)^2 + \Big(\frac{k_2 k_1^2}{b a^2} \Big)^2 \\ &+ \Big(\frac{k_2^2 k_1}{b^2 a} \Big)^2 + \Big(\frac{k_1^3}{a^3} \Big)^2 + \Big(\frac{k_2^3}{b^3} \Big)^2 \bigg] u_k v_k \\ &= \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} (-\bar{k} + \bar{k}^2 - \bar{k}^3) u_k v_k. \end{aligned}$$

Hence for an element u of D(A):

$$|u|_{2+}^2 = \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} (-\bar{k} + \bar{k}^2 - \bar{k}^3) u_k^2.$$

Using the constants C and N of the equation (2.3.4) and choosing C_1 such that

$$\sum_{\max\{k_1, k_2\} \le N} \bar{k}^2 \le C_1 \sum_{\max\{k_1, k_2\} \le N} -\bar{k}^3,$$
(2.4.5)

we conclude that

$$\sum_{k \in \mathbb{N}_0^2} -\bar{k}^3 \le \sum_{k \in \mathbb{N}_0^2} (-\bar{k} + \bar{k}^2 - \bar{k}^3) \le ((C+1)C_1 + 1) \sum_{k \in \mathbb{N}_0^2} -\bar{k}^3.$$

We define

$$(u, v)_{[2]} := (u, v)_{[2]-} + (\partial_{1,2}^2 u, \partial_{1,2}^2 v).$$

Then all the norms $|u|_{2+}$; $|u|_2$; $|u|_{[2]-}$ and $|u|_{[2]}$ are equivalent. Moreover for the last one we have the nice representations:

$$(u, v)_{[2]} = \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} -\bar{k}^3 u_k v_k;$$
$$|u|_{[2]}^2 = \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} -\bar{k}^3 u_k^2$$

Now we compute

$$abla (\nabla \cdot u) + \nabla^{\perp} (\nabla^{\perp} \cdot u), \quad u = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}:$$

$$\begin{aligned} \nabla(\nabla \cdot u) + \nabla^{\perp}(\nabla^{\perp} \cdot u) = &\nabla(\partial_1 u_1 + \partial_2 u_2) + \nabla^{\perp}(-\partial_2 u_1 + \partial_1 u_2) \\ &= \begin{pmatrix} \partial_1^2 u_1 + \partial_{1,2}^2 u_2 \\ \partial_{1,2}^2 u_1 + \partial_2^2 u_2 \end{pmatrix} + \begin{pmatrix} \partial_2^2 u_1 - \partial_{1,2}^2 u_2 \\ -\partial_{1,2}^2 u_1 + \partial_1^2 u_2 \end{pmatrix} \\ &= \begin{pmatrix} \partial_1^2 u_1 + \partial_2^2 u_1 \\ \partial_1^2 u_2 + \partial_2^2 u_2 \end{pmatrix} \end{aligned}$$

 \mathbf{so}

$$\Delta u = \nabla (\nabla \cdot u) + \nabla^{\perp} (\nabla^{\perp} \cdot u)$$
(2.4.6)

and, for u solenoidal we have $\Delta u = \nabla^{\perp} (\nabla^{\perp} \cdot u)$. Moreover for each $u \in D(A)$ we have

$$\nabla \cdot \Delta u = \nabla \cdot \nabla^{\perp} (\nabla^{\perp} \cdot u) = 0;$$

$$\mathbf{n} \cdot \Delta u = \mathbf{n} \cdot \nabla^{\perp} (\nabla^{\perp} \cdot u) = 0 \text{ on } \Gamma;$$

$$\Delta u \in H.$$
⁸
(2.4.7)

The operator A coincides with $-\Delta$ in D(A) because, due to (2.4.1) and to the definition of A we have

$$\langle Au, v \rangle = ((u, v)) = (-\Delta u, v) = \langle -\Delta u, v \rangle, \quad \forall v \in V.$$

2.4.2 The Operator B.

We define the form

$$b: (\mathbf{H}^{1}(R))^{3} \to \mathbb{R}$$
$$(u, v, w) \mapsto \sum_{i,j=1}^{2} \int_{R} u_{i}(\partial_{i}v_{j})w_{j} \, dx.$$
(2.4.8)

Lemma 2.4.1. The form b defined in (2.4.8) is trilinear and continuous on the product space $(\mathbf{H}^1(R))^3$.

Proof. The trilinearity is clear: $b(u, v, w) = ((u \cdot \nabla)v, w)$. For the continuity we start to compute $|\int_R u_i(\partial_i v_j)w_j dx|$: Since $u, v, w \in \mathbf{H}^1(R)$, by proposition 2.1.3, we have that

$$u_i, w_j \in L^4(R) \& \partial_i v_j \in L^2(R).$$

By Hölder Inequality we obtain that $u_i(\partial_i v_j)w_j \in L^1(R)$ and that

$$|\int_{R} u_{i}(\partial_{i}v_{j})w_{j} \, dx| \leq \int_{R} |u_{i}(\partial_{i}v_{j})w_{j}| \, dx \leq |u_{i}|_{L^{4}} |\partial_{i}v_{j}|_{L^{2}} |w_{j}|_{L^{4}}.$$

Therefore

$$|b(u, v, w)| \le |u|_{\mathbf{L}^4} |\nabla v|_{(\mathbf{L}^2)^2} |w|_{\mathbf{L}^4}$$

So, by Corollaries 2.1.4 and 2.1.9

$$|b(u, v, w)| \le C ||u|| ||v|| ||w||.$$

The form b, being continuous in $(\mathbf{H}^1(R))^3$, is continuous in V^3 and, for each pair $(u, v) \in V^2$ we define the operator $B(u, v) \in V'$ by

$$B(u, v): V \to \mathbb{R} \tag{2.4.9}$$

$$w \mapsto \langle B(u, v), w \rangle = b(u, v, w)$$
 (2.4.10)

and we set

$$B(u) := B(u, u) \in V' \quad \forall u \in V.$$

⁸Since $u \in \mathbf{H}^2$, $\overline{\Delta u \in \mathbf{L}^2}$.

2.5 Classical and Weak Formulations.

p

Classically the existence problem for (1.0.1)–(1.0.4), renaming $\tilde{F} := F + v$ amounts to finding a vector function

$$u: \overline{R} \times [0, T] \to \mathbb{R}^2$$

and a scalar function

$$: \overline{R} \times [0, T] \to \mathbb{R},$$

such that

$$\begin{split} u_t + (u \cdot \nabla)u + \nabla p &= \nu \Delta u + \tilde{F} & \text{in } R \times]0, T[; & (2.5.1) \\ \nabla \cdot u &= 0 & \text{in } R \times]0, T[; & (2.5.2) \\ u \cdot \mathbf{n} &= 0 & \text{on } \partial R \times]0, T[; & (2.5.3) \\ \nabla^{\perp} \cdot u &= 0 & \text{on } \partial R \times]0, T[; & (2.5.4) \\ u(x, 0) &= u_0(x) & \text{in } R. & (2.5.5) \end{split}$$

Where \tilde{F} and u_0 are given and defined in $R \times [0T]$ and R respectively. u_0 is the position at time 0 of the system.

Lemma 2.5.1. For $u, v \in V$ we have $(\Delta u, v) = -((u, v))$.

Proof. First we note that $(\Delta u, v) = -(A^{\frac{1}{2}}u, A^{\frac{1}{2}}v)$ where $A^{\frac{1}{2}}$ is the linear map defined by

$$A^{\frac{1}{2}}u = A^{\frac{1}{2}} \sum_{k \in \mathbb{N}_0^2} u_k W_k := \sum_{k \in \mathbb{N}_0^2} (-\bar{k})^{\frac{1}{2}} u_k W_k.$$

It is clear that $A^{\frac{1}{2}}$ maps V onto H continuously. Indeed for $u \in V$

$$|A^{\frac{1}{2}}u|^2 = \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} -\bar{k}(-\bar{k})u_k^2 = \|u\|^2 < \infty$$

and, the continuity follows from the linearity and from $|A^{\frac{1}{2}}(u-v)|^2 = ||u-v||^2$. Therefore it is enough to prove the Lemma for $u, v \in \mathcal{D}_1(R)$, by continuity it will be true for $u, v \in V$.

$$(\Delta u, v) = \sum_{i=1}^{2} \int_{R} \Delta u_{i} v_{i} \, dx = -\sum_{i=1}^{2} \int_{R} \nabla u_{i} \cdot \nabla v_{i} \, dx + \sum_{i=1}^{2} \int_{R} \nabla \cdot (\nabla u_{i} v_{i}) \, dx$$
$$= -((u, v)) + \sum_{i=1}^{2} \int_{\Gamma} v_{i} \nabla u_{i} \cdot \mathbf{n} \, d\Gamma.$$

Now we compute

$$\int_{\Gamma} v_1 \nabla u_1 \cdot \mathbf{n} \, d\Gamma = \int_{\Gamma} v_1 \partial_1 u_1 \mathbf{n}_1 \, d\Gamma + \int_{\Gamma} v_1 \partial_2 u_1 \mathbf{n}_2 \, d\Gamma.$$

The first term vanishes because v_1 vanishes where \mathbf{n}_1 does not ⁹ so we have

$$\int_{\Gamma} v_1 \nabla u_1 \cdot \mathbf{n} \, d\Gamma = 0 + \int_{f_b \cup f_t} v_1 \partial_2 u_1 \mathbf{n}_2 \, d\Gamma$$
$$= \int_{f_b \cup f_t} \partial_1 u_2 v_1 \mathbf{n}_2 \, d\Gamma = 0. \qquad [\text{by } \nabla^{\perp} \cdot u = 0 \text{ on } \Gamma],$$

 $^{{}^{9}\}mathbf{n}_{1}$ vanishes on the top and botton faces of the rectangle and v_{1} vanishes on the left and right ones.

where f_b , f_t and (below) f_l , f_r stay for the botton, top, left and right faces of R.

Analogously we conclude that

$$\int_{\Gamma} v_2 \nabla u_2 \cdot \mathbf{n} \, d\Gamma = \int_{f_l \cup f_r} v_2 \partial_2 u_1 \mathbf{n}_1 \, d\Gamma = 0.$$

Therefore

$$(\Delta u, v) = -((u, v))$$
 for $u, v \in \mathcal{D}_1(R)$.

г			
I			
I			
L			

Lemma 2.5.2. For $u, v \in V$ we have $(\Delta u, v) = -(\nabla^{\perp} \cdot u, \nabla^{\perp} \cdot v)$.

Proof. Again it is enough to prove the Lemma for $u, v \in \mathcal{D}_1(R)$. By (2.4.6)

$$\begin{split} \int_{R} \Delta u \cdot v \, dx &= \int_{R} \nabla^{\perp} (\nabla^{\perp} \cdot u) \cdot v \, dx \\ &= -\int_{R} (\nabla^{\perp} \cdot u) (\nabla^{\perp} \cdot v) \, dx + \int_{R} \nabla^{\perp} \cdot \left((\nabla^{\perp} \cdot u) v \right) dx \\ &= -\int_{R} (\nabla^{\perp} \cdot u) (\nabla^{\perp} \cdot v) \, dx + \int_{R} \nabla \cdot \left((\nabla^{\perp} \cdot u) v \right)^{\circ} dx \\ &= -\int_{R} (\nabla^{\perp} \cdot u) (\nabla^{\perp} \cdot v) \, dx + \int_{\Gamma} \left((\nabla^{\perp} \cdot u) v \right)^{\circ} \cdot \mathbf{n} \, d\Gamma \\ &= -\int_{R} (\nabla^{\perp} \cdot u) (\nabla^{\perp} \cdot v) \, dx. \end{split}$$

Where $\begin{pmatrix} u_1 \\ u_2 \end{pmatrix}^\circ := \begin{pmatrix} u_2 \\ -u_1 \end{pmatrix}$.

Corollary 2.5.3. The norms $u \mapsto ||u||$ and $u \mapsto |\nabla^{\perp} \cdot u|$ coincide in $\mathcal{D}_1(R)$ and then in V.

If u and p are classical solutions of $(2.5.1)-(2.5.5) - u \in C^2(\overline{R} \times [0, T])$, $p \in C^1(\overline{R} \times [0, T])$. Then clearly $u \in L^2(0, T, V)$ and if we fix $v \in \mathcal{D}_1(R)$, take the scalar product and use Lemma 2.5.1 obtain

$$\frac{d}{dt}(u, v) + \nu((u, v)) + b(u, u, v) = (\tilde{F}, v).$$
(2.5.6)

By continuity the previous expression holds for each $v \in V$.¹⁰ So is natural to define the following weak formulation of problem (2.5.1)–(2.5.5):

 $¹⁰v_k \to v \text{ in } \|\cdot\| \Rightarrow v_k \to v \text{ in } |\cdot|.$ (u, \cdot) continuous in $|\cdot|$ and $((u, \cdot))$ continuous in $\|\cdot\|$

Problem 2.5.1. Given

$$\tilde{F} \in L^2(0, T, V')$$
 (2.5.7)

&

$$u_0 \in H \tag{2.5.8}$$

to find

$$u \in L^2(0, T, V)$$
 (2.5.9)

satisfying (in the distribution sense)

$$\frac{d}{dt}(u, v) + \nu((u, v)) + b(u, u, v) = <\tilde{F}, v>, \quad \forall v \in V, (2.5.10)$$

and

$$u(0) = u_0. (2.5.11)$$

Remark 3. Note that (2.5.9) is not sufficient to give sense to (2.5.11). But we will show that if we have in addition (2.5.10) then u coincides almost everywhere with a continuous function giving a meaning to (2.5.11).

Now we present a proplem equivalent to Problem 2.5.1:

Problem 2.5.2. Given

$$\tilde{F} \in L^2(0, T, V'),$$
 (2.5.12)

&

$$u_0 \in H, \tag{2.5.13}$$

to find

$$u \in L^2(0, T, V), \qquad u' \in L^1(0, T, V')$$
 (2.5.14)

satisfying

$$u' + \nu Au + Bu = \tilde{F} \quad on \quad]0, T[, \qquad (2.5.15)$$

and

$$u(0) = u_0. (2.5.16)$$

To verify this equivalence we will need the following lemmas

Lemma 2.5.4. Let X be a Banach space with dual X' and let u and g be two functions belonging to $L^1(a, b, X)$. Then the following three conditions are equivalent

1. *u* is a.e. equal to a primitive function of *g*,

$$u(t) = \xi + \int_0^t g(s) \, ds, \quad \xi \in X, \ a.e. \ t \in [a, \ b];$$

2. For each test function $\phi \in \mathcal{D}(]a, b[)$,

$$\int_{a}^{b} u(t)\phi'(t) dt = -\int_{a}^{b} g(t)\phi(t) dt \qquad \left(\phi' = \frac{d\phi}{dt}\right);$$

3. For each $\eta \in X'$,

$$\frac{d}{dt} < u, \eta > = < g, \eta >$$

in the scalar distribution sense, on]a, b[

In particular if 1–3 is satisfied, u is a.e. equal to a continuous function $[a, b] \rightarrow X$.

The proof of Lemma 2.5.4 can be found in [15] section 3.1.

Lemma 2.5.5. If $u \in L^2(0, T, V)$, the function Bu defined by

$$< Bu(t), v >= b(u(t), u(t), v), \quad \forall v \in V, a.e. t \in [0, T],$$

belongs to $L^1(0, T, V')$.

Proof. For almost all t, Bu(t) is an element of V'. The measurability of the function

$$[0, T] \ni t \mapsto Bu(t) \in V',$$

is easy to check. Indeed it follows from the measurability of $t \mapsto u(t) \in V$ and from the continuity of $u \mapsto B(u) \in V'$. By the continuity and trilinearity of b on V we have

$$||Bw||_{V'} \le C||w||^2, \quad \forall w \in V, \quad C \in \mathbb{R}^+.$$
 (2.5.17)

Hence

$$\int_0^T \|Bu(t)\|_{V'} \, dt \le C \int_0^T \|u(t)\|^2 \, dt < +\infty.$$

Now let u satisfy both (2.5.9) and (2.5.10) then, due to the identity (2.4.1), to definition of A (given in (2.4.3)) and to the previous lemma we can write (2.5.10) as

$$\frac{d}{dt} < u, v \rangle = <\tilde{F} - \nu Au - Bu, v \rangle, \quad \forall v \in V.$$

By the linearity and continuity of $A: V \to V'$ we have that

$$Au \in L^2(0, T; V')$$
 (2.5.18)

so, $\tilde{F} - \nu Au - Bu \in L^1(0, T; V')$. Therefore, by Lemma 2.5.4¹¹ and

$$\begin{cases} u' \in L^1(0, T; V') \\ u' = f - \nu A u - B u, \end{cases}$$

u is a.e. equal to a continuous function $[0, T] \rightarrow V'$. Therefore (2.5.11) is meaningful and any solution of Problem 2.5.1 is a solution of Problem 2.5.2. Since any solution of Problem 2.5.2 is clearly a solution of Problem 2.5.1 we conclude that these two problems are equivalent.

¹¹Note that V, being a Hilbert space, is reflexive.

2.6 Existence

2.6.1 Fourier Transform. Fractional Derivatives.

We want to prove the existence of a solution for Problem 2.5.1, for that we will need another compactness theorem envolving fractional derivatives of a function. Given a function f from \mathbb{R} into a Hilbert space X_1 we denote its Fourier Transform by

$$\hat{f} = \int_{-\infty}^{+\infty} e^{-2i\pi t\tau} f(t) \, dt.$$

The derivative (in t) of order γ of f is the Fourier Transform of $(2i\pi\tau)^{\gamma}\hat{f}$ or

$$\widehat{D_t^{\gamma}f(\tau)} = (2\pi\tau)^{\gamma}\widehat{f}(\tau).$$

Now let X_0, X, X_1 be Hilbert spaces such that

$$X_0 \subset X \subset X_1$$

where, the injections are continuous and the first injection is in addiction compact. Given $\gamma > 0$, we define the space

$$\mathcal{H}^{\gamma}(\mathbb{R}, X_0, X_1) = \{ v \in L^2(\mathbb{R}, X_0) \mid D_t^{\gamma} v \in L^2(\mathbb{R}, X_1) \}.$$

 $\mathcal{H}^{\gamma}(\mathbb{R}, X_0, X_1)$ in a Hilbert space for the norm $\|\cdot\|_{\mathcal{H}^{\gamma}(\mathbb{R}, X_0, X_1)}$ defined by

$$\|v\|_{\mathcal{H}^{\gamma}(\mathbb{R}, X_{0}, X_{1})}^{2} = \|v\|_{L^{2}(\mathbb{R}, X_{0})}^{2} + \||\tau|^{\gamma} \hat{v}\|_{L^{2}(\mathbb{R}, X_{1})}.$$

For any set $K\subset \mathbb{R}$ we define the space

$$\mathcal{H}^{\gamma}_{K}(\mathbb{R}, X_{0}, X_{1}) = \{ u \in \mathcal{H}^{\gamma}(\mathbb{R}, X_{0}, X_{1}) \mid \operatorname{supp} u \subset K \}.$$

Now we can state the compactness theorem we will need:

Theorem 2.6.1. Let $X_0 \subset X \subset X_1$ be Hilbert spaces with both inclusions being continuous and the first one being also compact. Then for any bounded set K and any $\gamma > 0$ the injection of $\mathcal{H}_K^{\gamma}(\mathbb{R}, X_0, X_1)$ into $L^2(\mathbb{R}, X)$ is compact.

The proof can be found in [15] section 3.2.3.

2.6.2 The Existence Theorem.

In the proof of the Existence Theorem 2.6.4 we will need the following Lemma:

Lemma 2.6.2. Fixing the first variable in V, the form b defined in (2.4.8) results skew-symmetric in the last two variables, i.e.,

$$\forall u \in V \forall v, w \in \mathbf{H}^{1}[b(u, v, w) = -b(u, w, v)].$$

Due to the trilinearity of b the previous Lemma is equivalent to the following corollary

Corollary 2.6.3. Fixing the first variable in V, we have

$$\forall u \in V \forall v \in \mathbf{H}^1[b(u, v, v) = 0].$$

Proof. We prove the statement for $v \in C^{\infty}(\overline{R})$. Then by the continuity of b it holds for $v \in H^1$. For $u \in V$ and $v \in C^{\infty}(\overline{R})$ we have

$$\begin{split} b(u, v, v) &= \sum_{i,j=1}^{2} \int_{R} u_{i} \partial_{i} v_{j} v_{j} \, dx = \sum_{i,j=1}^{2} \int_{R} u_{i} \partial_{i} \frac{(v_{j})^{2}}{2} \, dx \\ &= \sum_{i,j=1}^{2} \int_{R} \partial_{i} \left(u_{i} \frac{(v_{j})^{2}}{2} \right) \, dx - \sum_{i,j=1}^{2} \int_{R} \partial_{i} u_{i} \frac{(v_{j})^{2}}{2} \, dx \\ &= \sum_{j=1}^{2} \int_{R} \nabla \cdot \left(u \frac{(v_{j})^{2}}{2} \right) \, dx - \sum_{j=1}^{2} \int_{R} (\nabla \cdot u) \frac{(v_{j})^{2}}{2} \, dx \\ &= \sum_{j=1}^{2} \int_{\Gamma} u \frac{(v_{j})^{2}}{2} \cdot \mathbf{n} \, d\Gamma - 0 = \sum_{j=1}^{2} \int_{\Gamma} \frac{(v_{j})^{2}}{2} \, u \cdot \mathbf{n} \, d\Gamma = 0. \end{split}$$

Theorem 2.6.4. Given \tilde{F} and u_0 satisfying (2.5.12) and (2.5.13). There is at least one function u satysfying (2.5.14)-(2.5.16).

Remark 4. The proof that follows is completely analogous to that for the case of full Dirichlet conditions that can be found in [15].

Proof. We start by defining, for each $m \in \mathbb{N}_0$, an approximate solution u^m of (2.5.10) as follows:

$$u^{m} := \sum_{\max\{i_{1}, i_{2}\} \le m} u^{m}_{i}(t)W_{i}$$

$$((u^{m})'(t), W_{j}) + \nu((u^{m}(t), W_{j})) + b(u^{m}(t), u^{m}(t), W_{j}) = <\tilde{F}(t), W_{j} >,$$

$$(2.6.2)$$

$$t \in [0, T], \quad \max\{j_1, j_2\} \le m,$$

 $u^m(0) = u_0^m.$ (2.6.3)

Where u_0^m is the orthogonal projection of u_0 onto $span\{W_i \mid \max\{i_1, i_2\} \leq m\}$. From (2.6.2) we obtain the nonlinear system of differential equations in the functions u_i^m , $\max\{i_1, i_2\} \leq m$:

$$\sum_{\substack{\max\{i_1, i_2\} \le m}} (u_i^m)'(t)(W_i, W_j) + \nu \sum_{\substack{\max\{i_1, i_2\} \le m}} u_i^m((W_i, W_j)) + \sum_{\substack{\max\{i_1, i_2\} \le m \\ \max\{l_1, l_2\} \le m}} u_i^m u_l^m b(W_i, W_l, W_j) = <\tilde{F}(t), W_j >, \quad (2.6.4)$$

that reduces to the ODE's system¹²

$$(u_{j}^{m})'(t)\left(-\bar{j}\frac{ab}{4}\right) + \nu u_{j}^{m}\bar{j}^{2}\frac{ab}{4} + \sum_{\substack{\max\{i_{1}, i_{2}\} \leq m \\ \max\{l_{1}, l_{2}\} \leq m}} u_{i}^{m}u_{l}^{m}b(W_{i}, W_{l}, W_{j}) = <\tilde{F}(t), W_{j} >;$$

$$\max\{j_{1}, j_{2}\} \leq m. \quad (2.6.5)$$

Now we note that (2.6.3) is the same as the *m* scalar conditions

$$u_j^m(0) =$$
the projection of u_0 onto $span\{W_j\} = u_{0j}$. ¹³ (2.6.6)

Therefore system (2.6.5) with initial condition (2.6.6) has a maximal solution defined in $[0, t_{max}]$ and, we will see that $t_{max} = T$. Indeed, if $t_{max} < T$ then $|u^m(t)|$ should tend to $+\infty$ as t goes to t_{max} . Below we show that $|u^m(t)|$ remains bounded in [0, T] so we have that the maximal solution is defined in [0, T].

A PRIORI ESTIMATES: We multiply (2.6.2) by u_j^m , $\max\{j_1, j_2\} \le m$ and add the equations obtained. Taking Corollary 2.6.3 into account, we arrive to

$$((u^m)'(t), u^m(t)) + \nu ||u^m(t)||^2 = \langle \tilde{F}(t), u^m(t) \rangle.$$

Hence

$$\frac{d}{dt}|u^{m}(t)|^{2} + 2\nu||u^{m}(t)||^{2} = 2 < \tilde{F}(t), \ u^{m}(t) >$$

$$\leq 2||\tilde{F}(t)||_{V'}||u^{m}(t)|| \leq \nu||u^{m}(t)||^{2} + \frac{1}{\nu}||\tilde{F}(t)||_{V'}^{2} \quad ^{14}$$

so,

$$\frac{d}{dt}|u^m(t)|^2 + \nu ||u^m(t)||^2 \le \frac{1}{\nu} ||\tilde{F}(t)||_{V'}^2.$$
(2.6.7)

In particular $\frac{d}{dt}|u^m(t)|^2 \leq \frac{1}{\nu} \|\tilde{F}(t)\|_{V'}^2$ and, integrating on [0, s] we obtain

$$|u^{m}(s)|^{2} \leq |u_{0}^{m}|^{2} + \frac{1}{\nu} \int_{0}^{s} \|\tilde{F}(t)\|_{V'}^{2} dt \leq |u_{0}|^{2} + \frac{1}{\nu} \int_{0}^{T} \|\tilde{F}(t)\|_{V'}^{2} dt.$$
(2.6.8)

Therefore

the sequence (u^m) remains in a bounded set of $L^{\infty}(0, T, H)$. (2.6.9)

Now we integrate (2.6.7) over [0, T] and obtain:

$$\begin{aligned} |u^m(T)|^2 + \nu \int_0^T \|u^m(t)\|^2 \, dt &\leq |u_0^m|^2 + \frac{1}{\nu} \int_0^T \|\tilde{F}(t)\|_{V'}^2 \, dt \\ &\leq |u_0|^2 + \frac{1}{\nu} \int_0^T \|\tilde{F}(t)\|_{V'}^2 \, dt. \end{aligned}$$

¹²In section 2.3 we have seen that the family \mathcal{W} is orthogonal in both H and V. ¹³If we write $u_0 = \sum_{i \in \mathbb{N}_0^2} u_{0i} W_i$. ¹⁴By expanded $(\sqrt{\nu}a - \frac{1}{\sqrt{\nu}}b)^2 \ge 0$ with $a = ||u^m(t)||$ and $b = ||\tilde{F}(t)||_{V'}$.

Therefore

the sequence
$$(u^m)$$
 remains in a bounded set of $L^2(0, T, V)$. (2.6.10)

Now we extend u^m to the entire real line putting

$$\tilde{u}^m(t) := \begin{cases} u^m(t) & \text{if } t \in [0, T] \\ 0 & \text{if } t \notin [0, T] \end{cases}$$

and the Fourier transform of \tilde{u}^m will be denoted by \hat{u}^m . APPLICATION OF THEOREM 2.6.1. We start by computing the integral

$$\int_{-\infty}^{+\infty} |\tau|^{2\gamma} |\hat{u}^m(\tau)|^2 \, d\tau.$$
 (2.6.11)

Equation (2.6.2) with u^m replaced by \tilde{u}^m results in

$$\frac{d}{dt}(\tilde{u}^m(t), W_j) = <\tilde{f}^m, W_j > +(u_0^m, W_j)\delta_0 - (u^m(T), W_j)\delta_T; \\ \max\{j_1, j_2\} \le m \quad (2.6.12)$$

where δ_0 , δ_T are the Dirac distributions at 0 and T and,

$$f^m = \tilde{F} - \nu A u^m - B u^m,$$

$$\tilde{f}^m(t) := \begin{cases} f^m(t) & \text{on}[0, T] \\ 0 & \text{outside}[0, T] \end{cases}.$$

Using the Fourier Transform (2.6.12) becomes

$$2\pi i\tau(\hat{u}^m(\tau), W_j) = \langle \hat{f}^m(\tau), W_j \rangle + (u_0^m, W_j) - (u^m(T), W_j) \exp(-2\pi i T\tau). \quad (2.6.13)$$

Multiplying (2.6.13) by $\hat{g}_i^m(\tau)$ and adding the obtained *m* equations we arrive to

$$2\pi i\tau |\hat{u}^m(\tau)|^2 = \langle \hat{f}^m(\tau), \, \hat{u}^m(\tau) \rangle + (u_0^m, \, \hat{u}^m(\tau)) - (u^m(T), \, \hat{u}^m(\tau)) \exp(-2\pi i T\tau). \quad (2.6.14)$$

By (2.5.17) we have

$$\|f^{m}(t)\|_{V'} \leq \|\tilde{F}(t)\|_{V'} + \nu \|u^{m}(t)\| + C\|u^{m}(t)\|^{2}; \text{ so}$$
$$\int_{0}^{T} \|f^{m}(t)\|_{V'} dt \leq \int_{0}^{T} \|\tilde{F}(t)\|_{V'} + \nu \|u^{m}(t)\| + C\|u^{m}(t)\|^{2} dt \qquad (2.6.15)$$

by (2.5.12) and (2.6.10) the integral (2.6.15) remains bounded. Hence ¹⁵

$$\sup_{\tau \in \mathbb{R}} \|\hat{f}^m(\tau)\|_{V'} \le \text{ const}, \quad \forall m \in \mathbb{N}_0.$$

¹⁵It is known that $|f(\tau)| \leq C \int_{\mathbb{R}} |f(t)| dt$. See for example [12].

By (2.6.8) both $|u^{m}(0)|$ and $|u^{m}(T)|$ are finite. Then by (2.6.14)

$$\tau ||\hat{u}^m(\tau)|^2 \le C_1 ||\hat{u}^m(\tau)|| + C_2 |\hat{u}^m(\tau)| \le D ||\hat{u}^m(\tau)||.$$
¹⁶ (2.6.16)

Fix $\gamma < \frac{1}{4}$ and define the real function $Q(x) := \frac{x^{2\gamma} + x}{1 + x}$, $x \in [0, +\infty[. Q \text{ is continuous and bounded},^{17}$ then we can find a constant $D_1 \in \mathbb{R}^+$ such that for all $\tau \in \mathbb{R}$:

$$\begin{aligned} Q(|\tau|) &\leq D_1 \\ \Leftrightarrow |\tau|^{2\gamma} &\leq D_1 \frac{1+|\tau|}{1+|\tau|^{1-2\gamma}} \end{aligned}$$

Therefore the integral (2.6.11) is bounded by

$$D_1 \int_{-\infty}^{+\infty} \frac{1+|\tau|}{1+|\tau|^{1-2\gamma}} |\hat{u}^m(\tau)|^2 d\tau$$

by (2.6.16)
$$\leq D_2 \int_{-\infty}^{+\infty} \frac{\|\hat{u}^m(\tau)\|}{1+|\tau|^{1-2\gamma}} d\tau + D_3 \int_{-\infty}^{+\infty} \|\hat{u}^m(\tau)\|^2 d\tau$$

The integral $D_3 \int_{-\infty}^{+\infty} \|\hat{u}^m(\tau)\|^2 d\tau$, by the Parseval Equality and (2.6.10), there is a constant D_4 such that

$$D_3 \int_{-\infty}^{+\infty} \|\hat{u}^m(\tau)\|^2 \, d\tau \le D_4, \qquad \forall m$$
 (2.6.17)

For the integral $D_2 \int_{-\infty}^{+\infty} \frac{\|\hat{u}^m(\tau)\|}{1+|\tau|^{1-2\gamma}} d\tau$ we apply Schwartz Inequality and Parseval Equality to obtain

$$D_2 \int_{-\infty}^{+\infty} \frac{\|\hat{u}^m(\tau)\|}{1+|\tau|^{1-2\gamma}} \, d\tau \le D_2 \left(\int_{-\infty}^{+\infty} \frac{1}{(1+|\tau|^{1-2\gamma})^2} \, d\tau \right)^{\frac{1}{2}} \left(\int_0^T \|u^m(t)\|^2 \, dt \right)^{\frac{1}{2}}$$
¹⁸

and, this product is finite and bounded as $m \to +\infty$, i.e., there is a constant D_5 such that

$$D_2 \int_{-\infty}^{+\infty} \frac{\|\hat{u}^m(\tau)\|}{1+|\tau|^{1-2\gamma}} d\tau \le D_5, \qquad \forall m.$$
(2.6.18)

By (2.6.17) and (2.6.18) we conclude that the integral (2.6.11) is finite:

$$\int_{-\infty}^{+\infty} |\tau|^{2\gamma} |\hat{u}^m(\tau)|^2 \, d\tau \le D_4 + D_5 =: E.$$

The finiteness of (2.6.11) with (2.6.10) implies that

the sequence (u^m) remains in a bounded set of $\mathcal{H}^{\gamma}(\mathbb{R}, V, H)$. (2.6.19)

THE LIMIT. We now need the following lemmas that can be found in [4] section III.6:

 $\begin{array}{l} \overset{1}{\overset{1}{\underset{1}{1}{1}}} \overset{1}{\overset{1}{\underset{1}{1}}} C_{1}, \ C_{2} \ \text{and} \ D \ \text{are constants.} \\ \overset{1}{\overset{1}{\underset{1}{1}}} \overset{1}{\overset{1}{\underset{1}{1}}} \overset{1}{\underset{1}{1}} \overset{1}{\underset{1}{1}} Q(0) = 0 \\ \overset{1}{\overset{1}{\underset{1}{1}}} \overset{1}{\underset{1}{1}} \overset{1}{\underset{1}{1}{1}} \overset{1}{\underset{1}{1}} \overset{1}{\underset{1}{$

Lemma 2.6.5. Let *E* be a reflexive Banach space and let (x_n) a bounded sequence in *E*. Then there is a subsequence $(x_{\sigma(n)})$ of (x_n) such that $x_{\sigma(n)} \rightharpoonup x$, for some $x \in E$.¹⁹

Lemma 2.6.6. Let E be a separable Banach space and let (f_n) a bounded sequence in E'. Then there is a subsequence $(f_{\sigma(n)})$ of (f_n) such that $f_{\sigma(n)} \rightharpoonup_* f$, for some $f \in E'$.

Lemma 2.6.6 and (2.6.9) implies the existence of a subsequence $(u^{\sigma(m)})$ of u^m and $u \in L^{\infty}(0, T, H)$ such that

$$u^{\sigma(m)} \rightharpoonup_* u$$
, in $L^{\infty}(0, T, H)$.²⁰

Analogously, Lemma 2.6.5 and (2.6.10) implies the existence of a subsequence $(u^{\alpha(\sigma(m))})$ of $u^{\sigma}(m)$ and $v \in L^2(0, T, V)$ such that

$$u^{\alpha(\sigma(m))} \rightharpoonup v, \text{ in } L^2(0, T, V).$$

The sequence (u^m) is in the space $\mathcal{H}^{\gamma}_{[0,T]}(\mathbb{R}, V, H)$ which injection in $L^2(\mathbb{R}, H)$ is compact due to Theorem 2.6.1.²¹ Then (2.6.19) implies the existence of a subsequence $\beta(\alpha(\sigma(m)))$ of $\alpha(\sigma(m))$ and $w \in L^2(0, T, H)$ satisfying

$$u^{\beta(\alpha(\sigma(m)))} \to w, \text{ in } L^2(0, T, H).$$

We put $\eta := \beta \circ \alpha \circ \sigma$ and we obtain

$$u = v = w \in L^2(0, T, H) \cap L^2(0, T, V) \cap L^\infty(0, T, H)$$

and

$$u^{\eta(m)} \rightharpoonup_* u, \text{ in } L^{\infty}(0, T, H);$$
 (2.6.20)

$$u^{\eta(m)} \rightharpoonup u, \text{ in } L^2(0, T, V);$$
 (2.6.21)

$$u^{\eta(m)} \to u, \text{ in } L^2(0, T, H).$$
 (2.6.22)

Indeed $L^{\infty}(0, T, H) = (L^1(0, T, H))'$ and $L^2(0, T, H) \subset L^1(0, T, H)$. So for each $f \in L^2(0, T, H)$:

$$\begin{cases} u^{\eta(m)}(f) & \to u(f) \\ u^{\eta(m)}(f) & \to v(f). \end{cases}$$

Since both $w, v \in L^2(0, T, H) = (L^2(0, T, H))'$, we can rewrite the previous expressions as

$$\forall f \in (L^2(0, T, H))' \begin{cases} f(u^{\eta(m)}) & \to f(u) \\ f(u^{\eta(m)}) & \to f(v). \end{cases}$$

Then $u = v \in L^2(0, T, H)$.

On the other side $(L^2(0, T, H))' \subseteq (L^2(0, T, V))'$ because a converging sequence in $L^2(0, T, V)$ converges in $L^2(0, T, H)$ too. So for each $f \in (L^2(0, T, H))'$ we have

$$\begin{cases} f(u^{\eta(m)}) & \to f(v) \\ f(u^{\eta(m)}) & \to f(w) \end{cases}$$

¹⁹We shall use the symbols: \rightarrow for weak convergence; \rightarrow_* for weak-star convergence and, \rightarrow for strong convergence.

 $^{^{20}}$ If wanted, without lack of generality we may assume $u^m \rightharpoonup_* u,$ in $L^\infty(0,\,T,\,H).$

²¹With $V = X_0$, & $H = X = X_1$.

2.6. EXISTENCE

But since both $w, v \in L^2(0, T, H)$ we have that $v = w.^{22}$ ENDING. To complete the proof we will need the following Lemma:

Lemma 2.6.7. If a sequence (u^n) satisfies $u^n \rightharpoonup u$ in $L^2(0, T, V)$ and $u^n \rightarrow u$ in $L^2(0, T, H)$, then for any vector function w with components in $C^1(\overline{R} \times [0, T])$,

$$\int_0^T b(u^n(t), \, u^n(t), \, w(t)) \, dt \to \int_0^T b(u(t), \, u(t), \, w(t)) \, dt.$$

Proof. From

$$b(u^{n}, u^{n}, w) - b(u, u, w) = b(u^{n} - u, u^{n}, w) + b(u, u^{n} - u, w)$$

we obtain

$$\begin{split} \left| \int_{0}^{T} b(u^{n}(t), u^{n}(t), w(t)) - b(u(t), u(t), w(t)) dt \right| \\ &\leq \int_{0}^{T} \left| b(u^{n}(t) - u(t), u^{n}(t), w(t)) \right| dt + \int_{0}^{T} \left| b(u(t), u^{n}(t) - u(t), w(t)) \right| dt \\ (\text{using}(2.7.1) \text{ below, and Lemma 2.6.2,}) \\ &\leq C \int_{0}^{T} \left[\left| u^{n}(t) - u(t) \right|^{\frac{1}{2}} \|u^{n}(t) - u(t)\|^{\frac{1}{2}} \|u^{n}(t)\|^{\frac{1}{2}} \|w\| \\ &+ |u(t)|^{\frac{1}{2}} \|u(t)\|^{\frac{1}{2}} \|u^{n}(t) - u(t)\|^{\frac{1}{2}} \|w\| \right] dt \\ &\leq C_{1} \left(\int^{T} \left| u^{n}(t) - u(t) \right| \|u^{n}(t) - u(t)\| dt \right)^{\frac{1}{2}} \|u^{n}(t)\|_{L^{2}(0, T, V)} \end{split}$$

$$+C_1 \|u(t)\|_{L^2(0,T,V)} \left(\int_0^T |u^n(t) - u(t)| \|u^n(t) - u(t)\| dt \right)^{\frac{1}{2}};$$

but u^n is bounded in $L^2(0, T, V)$, because $u^n \rightarrow u$ in $L^2(0, T, V)$, then

$$\begin{split} & \left| \int_0^T b(u^n(t), \, u^n(t), \, w(t)) - b(u(t), \, u(t), \, w(t)) \, dt \right| \\ \leq & C_2 \|u^n(t) - u(t)\|_{L^2(0, \, T, \, H)}^{\frac{1}{2}}. \end{split}$$

Hence $\left|\int_0^T b(u^n(t), u^n(t), w(t)) - b(u(t), u(t), w(t)) dt\right|$ goes to 0 as n goes to $+\infty$.

²²Note that since both $L^2(0, T, V)$ and $L^2(0, T, H)$ are both Hilbert spaces then each of them coincide with its dual. But we can not do these identifications here if we want to compare their duals, because the spaces H and V the norms are different. If we do the identifications we will obtain $(L^2(0, T, H))' \supseteq (L^2(0, T, V))'$ and that is not true.

Multiplying (2.6.2) by a function $\phi \in C^1([0, T])$ and, integrating we obtain:

$$\int_{0}^{T} ((u^{m})'(t), W_{j}\phi(t)) dt + \int_{0}^{T} \nu((u^{m}(t), W_{j}\phi(t))) dt$$

+
$$\int_{0}^{T} b(u^{m}(t), u^{m}(t), W_{j}\phi(t)) dt = \int_{0}^{T} \langle \tilde{F}(t), W_{j}\phi(t) \rangle dt$$

$$\iff -\int_{0}^{T} (u^{m}(t), W_{j}\phi'(t)) dt + \nu \int_{0}^{T} ((u^{m}(t), W_{j}\phi(t))) dt$$

+
$$\int_{0}^{T} b(u^{m}(t), u^{m}(t), W_{j}\phi(t)) dt = (u_{0}^{m}, W_{j})\phi(t)(0) - (u^{m}(T), W_{j})\phi(t)(T)$$

+
$$\int_{0}^{T} \langle \tilde{F}(t), W_{j}\phi(t) \rangle dt.$$

Due to Lemma 2.6.7 we can take the limit in the nonlinear term and, by (2.6.21)we can take the limite in the linear terms.²³ Hence

$$-\int_{0}^{T} (u(t), W_{j}\phi'(t)) dt + \nu \int_{0}^{T} ((u(t), W_{j}\phi(t))) dt + \int_{0}^{T} b(u(t), u(t), W_{j}\phi(t)) dt$$
$$= (u_{0}, W_{j})\phi(0) - (u(T), W_{j})\phi(T) + \int_{0}^{T} \langle \tilde{F}(t), W_{j}\phi(t) \rangle dt \quad (2.6.23)$$

Equation (2.6.23) being true for all W_j by linearity will be true for any finite combination of functions in \mathcal{W} and by continuity will be true for all $v \in V$. Then taking a test function $\phi \in \mathcal{D}(]0, T[)$ in (2.6.23) we conclude that

$$-\int_{0}^{T} (u(t), v\phi'(t)) dt + \nu \int_{0}^{T} ((u(t), v\phi(t))) dt + \int_{0}^{T} b(u(t), u(t), v\phi(t)) dt$$

=(u_0, v)\phi(0) - (u(T), v)\phi(T) + <\tilde{F}(t), v\phi(t) > dt; (2.6.24)
then

$$-\int_{0}^{T} (u(t), v\phi'(t)) dt + \nu \int_{0}^{T} ((u(t), v\phi(t))) dt + \int_{0}^{T} b(u(t), u(t), v\phi(t)) dt$$

= $\langle \tilde{F}(t), v\phi(t) \rangle dt$ (2.6.25)

what means that

Equation (2.5.10) is satisfied in the distribution sense.

Multiplying (2.5.10) by $\phi \in C^1([0, T])$ such that $\phi(0) = 1, \phi(T) = 0$ we obtain

$$-\int_{0}^{T} (u(t), v\phi'(t)) dt + \nu \int_{0}^{T} ((u(t), v\phi(t))) dt + \int_{0}^{T} b(u(t), u(t), v\phi(t)) dt$$
$$= (u(0), v) + \int_{0}^{T} \langle \tilde{F}(t), W_{j}\phi(t) \rangle dt.$$
(2.6.26)

²³Note that (2.6.21) implies that $u^{\eta(m)} \rightarrow u$ in $L^2(0, T, H)$.

The equation resulting from (2.6.24) with the same ϕ as in (2.6.26) is

$$-\int_{0}^{T} (u(t), v\phi'(t)) dt + \nu \int_{0}^{T} ((u(t), v\phi(t))) dt + \int_{0}^{T} b(u(t), u(t), v\phi(t)) dt$$

=(u₀, v)+ < $\tilde{F}(t)$, v $\phi(t)$ > dt (2.6.27)

From (2.6.26) and (2.6.27) we conclude that (2.5.11) is satisfied (and, we have finished the proof of theorem 2.6.4).

2.7 Uniqueness.

Lemma 2.7.1.

$$|b(u, v, w)| \le c|u|^{\frac{1}{2}} ||u||^{\frac{1}{2}} ||v|| |w|^{\frac{1}{2}} ||w||^{\frac{1}{2}}, \quad \forall u, v, w \in \mathbf{H}^{1}(R).$$
(2.7.1)

If u belongs to $L^2(0, T, V) \cap L^{\infty}(0, T, H)$, then Bu belongs to $L^2(0, T, V')$ and

$$||Bu||_{L^2(0,T,V')} \le c|u|_{L^{\infty}(0,T,H)} ||u|| L^2(0,T,V).$$
(2.7.2)

See [13] section III 3.2. for (2.7.1). Then (2.7.2) is a corollary of (2.7.1). Indeed

$$\begin{aligned} |b(u, u, w)| &= |b(u, w, u)| \le C |u|^{\frac{1}{2}} ||u||^{\frac{1}{2}} ||u||^{\frac{1}{2}} ||u||^{\frac{1}{2}} ||w|| \\ &= C |u| ||u|| ||w||, \quad w, u \in V. \end{aligned}$$
(2.7.3)

In the case $u \in L^2(0, T, V) \cap L^{\infty}(0, T, H)$ we obtain that

$$\int_{0}^{T} \|Bu(t)\|_{V'}^{2} dt \leq C \int_{0}^{T} \|u\|^{2} \|u\|^{2} dt$$
$$\leq C \|u\|_{L^{\infty}(0,T,H)}^{2} \int_{0}^{T} \|u(t)\|^{2} dt = C \|u\|_{L^{\infty}(0,T,H)}^{2} \|u\|_{L^{2}(0,T,V)}^{2} < +\infty$$

Theorem 2.7.2. The solution of Problems 2.5.1-2.5.2 given by Theorem 2.6.4 is unique. Moreover it is a.e. equal to a continuous function from [0, T] into H and,

$$u(t) \to u(t_1), \quad in \ H \quad as \ t \to t_1 \quad t_1 \in [0, \ T].$$
 (2.7.4)

In particular

$$\begin{split} & u(t) \to u_0, \quad in \ H \quad as \ t \to 0; \\ & u(t) \to u(T), \quad in \ H \quad as \ t \to T. \end{split}$$

For the proof we need the following lemma from [15]:

Lemma 2.7.3. Let V, H, V' three Hilbert spaces satisfying $V \subset H \subset V'$ with dense and continuous inclusions. If

$$u \in L^2(0, T, V) \& u' \in L^2(0, T, V')$$

then u is a.e. equal to a continuous function from [0, T] into H and

$$\frac{d}{dt}|u|^2 = 2 < u', u > \tag{2.7.5}$$

holds in the distribution sense on (0, T).

Remark 5. Note that (2.7.5) is meaningful because both

$$t \mapsto |u(t)|^2 \& t \mapsto \langle u'(t), u(t) \rangle$$

are integrable on [0, T].

Proof of Theorem 2.7.2. By (2.5.12), (2.5.15), (2.5.18) and (2.7.2) we have that $u' \in L^2(0, T, V')$. Then by (2.6.20), (2.6.21), and Lemma 2.7.3 we have that

$$u \in C([0, T], H).$$

Therefore (2.7.4) is satisfied.

To prove the uniqueness we consider two solutions u, v of the problems and put w := u - v. Then

$$w' = u' - v' = \tilde{F} - \nu Au - Bu - (\tilde{F} - \nu Av - Bv)$$
$$= -\nu Aw - Bu + Bv$$
and
$$w(0) = 0.$$

Now we take the scalar product with w in the duality between V and V' and obtain

$$< w', w > +\nu < Aw, w > = < Bv, w > - < Bu, w > -$$

We note that since $u, v \in L^2(0, T, V)$ & $u', v' \in L^2(0, T, V')$ then $w \in U'$ $L^2(0, T, V)$ & $w' \in L^2(0, T, V')$. Hence we can apply (2.7.5) and obtain

$$\begin{aligned} \frac{d}{dt}|w(t)|^2 + 2\nu||w(t)||^2 &= 2b(v(t), v(t) w(t)) - 2b(u(t), u(t), w(t)) \\ &= -2b(w(t), v(t), w(t)) = 2b(w(t), w(t), v(t)). \end{aligned}$$

Hence using (2.7.3)

$$\frac{d}{dt} |w(t)|^2 + 2\nu ||w(t)||^2 \le 2(C|w(t)|||w(t)||||v(t)||)$$
$$\le 2\nu ||w(t)||^2 + \frac{C^2}{2\nu} |w(t)|^2 ||v(t)||^2 = 25$$

Threfore

$$\frac{d}{dt}|w(t)|^2 \le \frac{C^2}{2\nu}|w(t)|^2||v(t)||^2.$$

Since $t \mapsto \frac{d}{dt}|w(t)|^2 = 2 < w'(t), w(t) >, t \mapsto |w(t)|^2 = < w(t), w(t) >$ and $t \mapsto ||v(t)||^2 = < v(t), v(t) >$ are integrable, by Gronwall Inequality (see (2.7.6)) below)

$$|w(t)|^{2} \leq |w(0)|^{2} \exp\left(\frac{C^{2}}{2\nu} \int_{0}^{t} ||v(\tau)||^{2} d\tau\right)$$
$$\leq |w(0)|^{2} \exp\left(\frac{C^{2}}{2\nu} \int_{0}^{T} ||v(\tau)||^{2} d\tau\right) = 0, \qquad t \in [0, T]$$

 ${}^{24}b(w,\,v,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,v,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,u,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,u,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,v,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ -b(v,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ b(u,\,w,\,w) \ + \ b(u,\,w,\,w) \ = \ b(u,\,w,\,w) \ + \ b(u,\,w) \ + \ b(u,\,w,\,w) \ + \ b(u,\,w) \ + \$ $\begin{array}{l} -b(v,v,w) + b(u,u,w). \\ {}^{25}\text{By expanded } (\sqrt{2\nu a} - \frac{C}{\sqrt{2\nu}}b)^2 \ge 0 \text{ with } a = \|w(t)\| \text{ and } b = |w(t)| \|v(t)\|. \end{array}$

(using the fact that w(0) = 0). Hence

u = v

concluding the uniqueness of the solution.

Lemma 2.7.4 (Gronwall Inequality). Let $g, h, y, \frac{dy}{dt}$ be locally integrable functions sattisfying

$$\frac{dy}{dt} \le gy + h \quad for \ t \ge t_0. \tag{2.7.6}$$

Then

$$y(t) \le y(t_0) \exp\left(\int_{t_0}^t g(\tau) \, d\tau\right) + \int_{t_0}^t h(s) \exp\left(-\int_t^s g(\tau) \, d\tau\right) \, ds, \quad t \ge t_0.$$

[The proof can be found in [13]].

2.8 Continuity on Initial Data.

Theorem 2.8.1. The map

$$\mathbb{S}: H \times L^2(0, T, V') \times]0, +\infty [\to C([0, T], H)$$
$$(u_0, \tilde{F}, \nu) \mapsto u$$

is continuous. Here $u \in C([0, T], H)$ is the unique solution of Problems 2.5.1-2.5.2 (see Theorem 2.7.2).

Proof. Fix a triple $(u_0, \tilde{F}, \nu) \in H \times L^2(0, T, V') \times]0, +\infty[$ and consider the solution of Theorem 2.7.2 induced by this triple. Then u satisfies

$$u' + \nu Au + Bu = F, \qquad u(0) = u_0.$$

Now consider another triple $(v_0, G, \eta) \in H \times L^2(0, T, V') \times]0, +\infty[$. The solution associated with this triple satisfies

$$v' + \eta Av + Bv = G, \qquad v(0) = v_0.$$

We put

$$w := v - u$$

and see that w satisfies the equation

$$w' = G - F - \eta Aw + (\nu - \eta)Au - Bv + Bu.$$

Taking the scalar product with w we obtain

$$< w', w > = < G - \tilde{F}, w > -\eta ||w||^{2} + (\nu - \eta)((u, w)) + b(w, w, u)$$
$$\frac{d}{dt}|w|^{2} \le 2||G - \tilde{F}||_{V'}||w|| - 2\eta ||w||^{2} + 2|\nu - \eta||u|||w|| + 2C|w|||w|||u||.$$

Expanding

$$\begin{split} & \left(\sqrt{\frac{\eta}{3}} \|w\| - \sqrt{\frac{3}{\eta}} \|G - \tilde{F}\|_{V'}\right)^2 \ge 0; \\ & \left(\sqrt{\frac{\eta}{3}} \|w\| - C\sqrt{\frac{3}{\eta}} \|w\| \|u\|\right)^2 \ge 0; \\ & \left(\sqrt{\frac{\eta}{3}} \|w\| - |\nu - \eta|\sqrt{\frac{3}{\eta}} \|u\|\right)^2 \ge 0; \end{split}$$

we obtain

$$\frac{d}{dt}|w|^2 + \eta \|w\|^2 \le \frac{3}{\eta} \|G - \tilde{F}\|_{V'}^2 + |\nu - \eta|^2 \frac{3}{\eta} \|u\|^2 + C^2 \frac{3}{\eta} |w|^2 \|u\|^2.$$
(2.8.1)

By (2.7.6) we have

$$\begin{split} &|w(t)|^{2} \\ \leq |w(0)|^{2} \exp\left(C^{2} \frac{3}{\eta} \int_{0}^{t} ||u(\tau)||^{2} d\tau\right) \\ &+ \int_{0}^{t} \left(\frac{3}{\eta} ||G(s) - \tilde{F}(s)||_{V'}^{2} + |\nu - \eta|^{2} \frac{3}{\eta} ||u(s)||^{2}\right) \exp\left(-\frac{3C^{2}}{\eta} \int_{t}^{s} ||u(\tau)||^{2} d\tau\right) ds \\ \leq \exp\left(C^{2} \frac{3}{\eta} \int_{0}^{T} ||u(\tau)||^{2} d\tau\right) \left(|w(0)|^{2} \\ &+ \int_{0}^{T} \left(\frac{3}{\eta} ||G(s) - \tilde{F}(s)||_{V'}^{2} + |\nu - \eta|^{2} \frac{3}{\eta} ||u(s)||^{2}\right) ds \right). \end{split}$$

Now if $|\nu - \eta| < \frac{\nu}{2}$, i.e., $\eta \in]\frac{\nu}{2}, \frac{3\nu}{2}[$, we have

$$\begin{split} |w(t)|^{2} &\leq \exp\left(C^{2}\frac{6}{\nu}\int_{0}^{T}\|u(\tau)\|^{2}\,d\tau\right)\left(|w(0)|^{2}\right.\\ &+\int_{0}^{T}\left(\frac{6}{\nu}\|G(s)-\tilde{F}(s)\|_{V'}^{2}+|\nu-\eta|^{2}\frac{6}{\nu}\|u(s)\|^{2}\right)ds\bigg). \end{split}$$

Fix $\varepsilon > 0$. Now if we put $E_0 := \exp\left(C^2 \frac{6}{\nu} \int_0^T \|u(\tau)\|^2 d\tau\right)$ and choose the triple (v_0, G, η)

$$\begin{split} |v_0 - u_0| &= |w(0)| < \frac{\varepsilon}{\sqrt{3}} E_0^{-\frac{1}{2}} =: \alpha_v; \\ \|G - \tilde{F}\|_{L^2(0, T, V')} &= \left(\int_0^T \|G(s) - \tilde{F}(s)\|_{V'}^2 \, ds\right)^{\frac{1}{2}} < \frac{\varepsilon\sqrt{\nu}}{3\sqrt{2}} E_0^{-\frac{1}{2}} =: \alpha_g; \\ |\nu - \eta| < \min\{\frac{\nu}{2}, \delta\} =: \alpha_\eta, \\ \text{where} \quad \delta &= \frac{\varepsilon\sqrt{\nu}}{3\sqrt{2}} \Big(\int_0^T \|u(s)\|^2 \, ds\Big)^{-\frac{1}{2}} E_0^{-\frac{1}{2}}, \quad ^{26} \end{split}$$

we obtain

$$|w(t)|^2 < \varepsilon^2 \Leftrightarrow |w(t)| < \varepsilon, \qquad t \in [0, T]$$

²⁶Since $u \in L^2(0, T, V)$ we have that α_v and α_η are finite. We can also see that $\alpha_v, \alpha_g, \alpha_\eta$ depend only on the (fixed) triple (u_0, \tilde{F}, ν) .

and, we have the continuity of S.

Theorem 2.8.2. The map

$$\begin{split} \mathbb{S}_2: \ H \times L^2(0,\,T,\,V') \times]0, \ +\infty [\rightarrow L^2(0,\,T,\,V) \\ (u_0,\,\tilde{F},\,\nu) \mapsto u \end{split}$$

is continuous. Here $u \in L^2(0, T, V)$ is the unique solution of Problems 2.5.1-2.5.2 (see Theorem 2.7.2).

Proof. As in the proof of Theorem 2.8.1 we fix $\varepsilon > 0$ and a triple $(u_0, \tilde{F}, \nu) \in H \times L^2(0, T, V') \times]0, +\infty[$ and consider the solution of Theorem 2.7.2 induced by this triple. Now consider another triple $(v_0, G, \eta) \in H \times L^2(0, T, V') \times]0, +\infty[$ and the solution associated with this triple. Put

$$w := v - u$$

and taking the scalar product with w we arrive to (2.8.1). Integrating over [0, T] we obtain

$$\begin{aligned} \frac{1}{\eta} |w(T)|^2 &- \frac{1}{\eta} |w(0)|^2 + \int_0^T \|w(t)\|^2 \, dt \le \frac{3}{\eta^2} \|G - \tilde{F}\|_{L^2(0, T, V')}^2 \\ &+ |\nu - \eta|^2 \frac{3}{\eta^2} \|u\|_{L^2(0, T, V)}^2 + \frac{3C^2}{\eta^2} \|w\|_{C([0, T], H)} \|u\|_{L^2(0, T, V)}^2 \end{aligned}$$

and, if $|\nu - \eta| < \frac{\nu}{2}$ we have

$$\begin{split} \int_0^T \|w(t)\|^2 \, dt &\leq \frac{12}{\nu^2} \|G - \tilde{F}\|_{L^2(0,\,T,\,V')}^2 + |\nu - \eta|^2 \frac{12}{\nu^2} \|u\|_{L^2(0,\,T,\,V)}^2 \\ &+ \Big(\frac{12C^2}{\nu^2} \|u\|_{L^2(0,\,T,\,V)}^2 + \frac{4}{\nu}\Big) \|w\|_{C(0,\,T,\,H)}^2. \end{split}$$

Using Theorem 2.8.1, there is $\delta > 0$ such that if (v_0, G, η) satisfies

$$|v_0 - u_0| < \delta, \, \|G - \tilde{F}\|_{L^2(0, T, V')} < \delta, \, |\eta - \nu| < \delta,$$

then $||w||_{C(0,T,H)} < \left(\frac{\varepsilon^2}{3}\right)^{\frac{1}{2}} \left(\frac{12C^2}{\nu^2} ||u||_{L^2(0,T,V)}^2 + \frac{4}{\nu}\right)^{-\frac{1}{2}}$. Hence for

$$\delta_1 = \min \Big\{ \frac{\nu}{2}, \ \delta, \ \Big(\frac{\varepsilon^2}{3}\Big)^{\frac{1}{2}} \Big(\frac{12}{\nu^2}\Big)^{-\frac{1}{2}}, \ \Big(\frac{\varepsilon^2}{3}\Big)^{\frac{1}{2}} \Big(\frac{12}{\nu^2} \|u\|_{L^2(0, T, V)}^2\Big)^{-\frac{1}{2}} \Big\},$$

we have $||w||_{L^2(0,T,V)} < \varepsilon$. Therefore the map \mathbb{S}_2 is continuous.

2.9 Some More Estimates for the Form b.

We present here some estimates we shall need later. We have 27

$$|b(u, v, w)| \le CK.$$

33

 $^{^{27}}$ These estimates can be found in [13] section III.3.2.. They can be obtained by interpolation ([9]), generalized Sobolev inequalities ([10]) and by a theorem by S.Agmon ([1]). See [14] for indications how to obtain them.

where C is a constant and K is one of the following products

$$||u|| ||v|| ||w|| u, v, w \in \mathbf{H}^1(R), (2.9.1)$$

$$\begin{split} &|u|^{\frac{1}{2}} \|u\|^{\frac{1}{2}} \|v\|^{\frac{1}{2}} \|v\|^{\frac{1}{2}} \|w\| & u \in \mathbf{H}^{1}(R), \, v \in \mathbf{H}^{2}(R), \, w \in \mathbf{L}^{2}(R), \quad (2.9.2) \\ &|u|^{\frac{1}{2}} \|u\|^{\frac{1}{2}} \|v\| \|w\| & u \in \mathbf{H}^{2}(R), \, v \in \mathbf{H}^{1}(R), \, w \in \mathbf{L}^{2}(R), \quad (2.9.3) \\ &|u| \|v\| \|w\|^{\frac{1}{2}} \|w\|^{\frac{1}{2}} & u \in \mathbf{L}^{2}(R), \, v \in \mathbf{H}^{1}(R), \, w \in \mathbf{H}^{2}(R), \quad (2.9.4) \\ &|u|^{\frac{1}{2}} \|u\|^{\frac{1}{2}} \|v\| \|w\|^{\frac{1}{2}} \|w\|^{\frac{1}{2}} & u, \, v, \, w \in \mathbf{H}^{1}(R). \quad (2.9.5) \end{split}$$

Strong Formulation. 2.10

Sometimes we want more regularity for the solutions of problem (1.0.1)-(1.0.4). Instead of Problems (2.5.1)-(2.5.2) where we ask for weak solutions we consider the following equivalent problems where we look for solutions more regular than weak.

Problem 2.10.1. Given

$$\dot{F} \in L^2(0, T, H)$$
 (2.10.1)

$$u_0 \in V \tag{2.10.2}$$

to find

&

$$u \in L^{2}(0, T, D(A)) \cap L^{\infty}(0, T, V)$$
satisfying (in the distribution sense)
$$(2.10.3)$$

$$\frac{d}{dt}(u, v) + \nu((u, v)) + b(u, u, v) = (\tilde{F}, v), \quad \forall v \in V, \ (2.10.4)$$

and

$$u(0) = u_0. (2.10.5)$$

Problem 2.10.2. Given

$$\tilde{F} \in L^2(0, T, H),$$
 (2.10.6)

$$u_0 \in V, \tag{2.10.7}$$

to find

&

$$u \in L^{2}(0, T, D(A)) \cap L^{\infty}(0, T, V), and$$

$$u' \in L^{2}(0, T, H)$$
(2.10.8)

satisfying

$$u' + \nu Au + Bu = \tilde{F}$$
 on]0, T[, (2.10.9)

and

$$u(0) = u_0. (2.10.10)$$

The equivalence of these problems follows from

1. A solution of Problem 2.10.2 is a solution of Problem 2.10.1;

2. For a solution u of Problem 2.10.1 we have

$$|Bu(t)| \le C ||u(t)||^{\frac{3}{2}} ||u(t)||^{\frac{1}{2}} \quad \text{a.e. } t \text{ by } (2.9.2), \text{ so}$$
$$\int_0^T |Bu(t)|^4 \le C \int_0^T ||u(t)||^6 ||u(t)||^2_2 \le C_1 \int_0^T ||u(t)||^2_2 < \infty$$

Hence

- (a) $Bu \in L^4(0, T, H)$. Since $u \in L^2(0, T, D(A))$ and $\int_0^T |Au(t)|^2 = \int_0^T ||u(t)||_2^2 < \infty$ we have
- (b) $Au \in L^2(0, T, H)$.

From (2a) and (2b) we have that

$$f - \nu Au - Bu \in L^2(0, T, H) \subseteq L^2(0, T, V').$$

Since $u \in L^2(0, T, D(A)) \subseteq L^2(0, T, V)$, by Lemma (2.5.4) and (2.10.4), we have that

$$u' = f - \nu Au - Bu$$
 a.e. and $u \in C(0, T, V')$ "a.e.".

Hence

3. u is a solution of Problem 2.10.2.

Lemma 2.10.1. The inclusions

$$D(A) \subseteq V \subseteq H$$

are both dense continuous and compact.

Proof. We have already seen in Corollary 2.1.9 and beginning of section 2.2 that the inclusion $V \subseteq H$ has the required properties. For the first inclusion we have that V, respectively D(A), are the closure of $\mathcal{D}_1(R)$ in \mathbf{H}^1 , respectively in \mathbf{H}^2 . From the density, continuity and compactness of the inclusion $\mathbf{H}^2 \subseteq \mathbf{H}^1$ come the same properties for the inclusion $D(A) \subseteq V$.

2.11 Existence.

Theorem 2.11.1. Given \tilde{F} and u_0 satisfying (2.10.6) and (2.10.7). There is at least one function u satisfying (2.10.8)-(2.10.10).

Remark 6. The proof that follows is completely analogous to that for the case of weak solutions and for no-slip boundary conditions that can be found in [15].

Proof. Outlines: We define an approximate solution u^m , for each $m \in \mathbb{N}_0$ like in (2.6.1)–(2.6.3), and arrive to the estimate (2.6.7) and conclusions (2.6.9) and (2.6.10) exactly in the same way. For Theorem we need only some more estimates: If we multiply 2.6.2 by $\bar{j}u_j^m$ and add the obtained equations we arrive to

$$((u^{m}(t))', Au^{m}(t)) + \nu((u^{m}(t), Au^{m}(t))) + Bu^{m}(t)(Au^{m}(t)) = (F(t), Au^{m}(t)),$$

i.e.,

$$(((u^{m}(t))', u^{m}(t))) + \nu(u^{m}(t), u^{m}(t))_{[2]} + Bu^{m}(t)(Au^{m}(t)) = (\tilde{F}(t), Au^{m}(t))_{[2]}$$
or

$$\frac{1}{2} \frac{d}{dt} \|u^{m}(t)\|^{2} + \nu |u^{m}(t)|^{2}_{[2]} \leq |Bu^{m}(t)(Au^{m}(t))| + |\tilde{F}(t)||Au^{m}(t)|$$

$$(by (2.9.2)) \leq C|u^{m}(t)|^{\frac{1}{2}} \|u^{m}(t)\| \|u^{m}(t)|^{\frac{3}{2}}_{[2]} + |\tilde{F}(t)||u^{m}(t)|_{[2]}$$

$$(by Young Inequalities)^{28}$$

$$\leq \frac{\nu}{4}(|u^{m}(t)|^{\frac{3}{2}}_{[2]})^{\frac{4}{3}} + C_{\frac{\nu}{4},\frac{4}{3}}(C|u^{m}(t)|^{\frac{1}{2}} \|u^{m}(t)\|)^{4}$$

$$+ \frac{\nu}{4}|u^{m}(t)|^{2}_{[2]} + C_{\frac{\nu}{4},2}|\tilde{F}(t)|^{2}$$

$$\frac{d}{dt} \|u^m\|^2 + \nu |u^m|^2_{[2]} \le +\frac{3^3}{2\nu^3} C^2 |u^m(t)|^2 \|u^m(t)\|^4 + \frac{2}{\nu} |\tilde{F}(t)|^2.$$
(2.11.1)

 $+\frac{\nu}{4}|u^{m}(t)|^{2}_{[2]}+\frac{1}{\nu}|\tilde{F}(t)|^{2}.$

 $\leq \frac{\nu}{4}(|u^m(t)|_{[2]}^{\frac{3}{2}})^{\frac{4}{3}} + \frac{3^3}{4\nu^3}(C|u^m(t)|^{\frac{1}{2}}\|u^m(t)\|)^4$

From equation (2.11.1) and from Gronwall Inequality we can derive for $s \in [0, T]$ (using (2.6.9) and (2.6.10))

$$\begin{aligned} &\|u^{m}(s)\|^{2} \\ \leq \exp\left(\int_{0}^{T} \frac{3^{3}}{2\nu^{3}} C^{2} |u^{m}(t)|^{2} \|u^{m}(t)\|^{2} dt\right) \left(\|u_{0}^{m}\|^{2} + \int_{0}^{T} \frac{2}{\nu} |\tilde{F}(t)|^{2} dt\right) \\ \leq K_{1} \end{aligned}$$

for some constant K_1 (independent of m). Hence

the sequence (u^m) remains in a bounded set of $L^{\infty}(0, T, V)$. (2.11.2)

Now we integrate (2.11.1) over [0, T] and obtain:

$$\|u^{m}(T)\|^{2} - \|u^{m}(0)\|^{2} + \nu \int_{0}^{T} |u^{m}(t)|^{2}_{[2]} dt$$

$$\leq \int_{0}^{T} \frac{3^{3}}{2\nu^{3}} C^{2} |u^{m}(t)|^{2} \|u^{m}(t)\|^{4} + \frac{2}{\nu} |\tilde{F}(t)|^{2} dt \leq K_{3}$$

with K_3 being a constant (independent of m, using (2.6.9), (2.11.2) and (2.10.6)). Therefore

 $\frac{\text{the sequence } (u^m) \text{ remains in a bounded set of } L^2(0, T, D(A)). \quad (2.11.3)}{^{28}\text{Young Inequality: } ab \le \varepsilon a^p + C_{\varepsilon,p}^Y b^{p'}. \text{ Where } a, b > 0, 1 0, \frac{1}{p} + \frac{1}{p'} = 1 \text{ and } C_{\varepsilon,p}^Y = \frac{p-1}{(p^{p'})(\varepsilon^{\frac{1}{p-1}})}.$
By 2.11.2, 2.11.3 and Lemmas 2.6.5 and 2.6.6 we conclude the existence of a subsequence $u^{\sigma(m)}$ such that

$$u^{\sigma(m)} \rightharpoonup u \text{ in } L^2(0, T, D(A))$$
$$u^{\sigma(m)} \rightharpoonup_* u \text{ in } L^{\infty}(0, T, V)$$

This limit is a solution for problems 2.10.1-2.10.2.

2.12Uniqueness.

Lemma 2.12.1. The inclusions

$$D(A) \subseteq V \subseteq D(A)'$$

are both dense and continuous.

Proof. We have just seen in Lemma 2.10.1 that the inclusion $D(A) \subseteq V$ has the required properties. For the second inclusion we proceed as in the beginning of section 2.2: We identify V with V'^{29} The continuity of the second inclusion follows from the continuity of the first one. The density of the second inclusion follows from Lemma 2.2.1 and the injectivity of the first one.

Theorem 2.12.2. The solution of problems 2.10.1-2.10.2 is unique. Moreover it is a.e. equal to a continuous function from [0, T] into V.

Proof. A solution of problems 2.10.1-2.10.2 is a solution of problems 2.5.1-2.5.2 as well, which is unique by Theorem 2.7.2. By $u \in L^2(0, T, D(A))$,

$$u' \in L^2(0, T, H) \subseteq L^2(0, T, V') \subseteq L^2(0, T, D(A)')$$

and by lemmas 2.12.1 and 2.7.3 we conclude that $u \in C([0, T], V)$.

2.13Continuity on Initial Data.

Theorem 2.13.1. The map

$$\mathbb{S}_s: V \times L^2(0, T, H) \times]0, +\infty[\to C([0, T], V)$$
$$(u_0, \tilde{F}, \nu) \mapsto u$$

is continuous. Where $u \in C([0, T], V)$ is the unique solution of Problems 2.10.1-2.10.2 (see Theorem 2.12.2).

Proof. Fix $\varepsilon > 0$ and a triple $(u_0, \tilde{F}, \nu) \in V \times L^2(0, T, H) \times [0, +\infty]$ and, consider the solution of Theorem 2.12.2 induced by this triple. Then u satisfies

$$u' + \nu Au + Bu = F, \qquad u(0) = u_0.$$

Now consider another triple $(v_0, G, \eta) \in V \times L^2(0, T, H) \times]0, +\infty[$. The solution associated with this triple satisfies

 $v' + \eta A v + B v = G,$ $v(0) = v_0.$ ²⁹Note that here *H* is not present. We are doing only one identification.

We put

$$w := v - u$$

and see that w satisfies the equation

$$w' = G - \tilde{F} - \eta Aw + (\nu - \eta)Au - Bv + Bu$$

Taking the scalar product with Aw we obtain

$$\frac{d}{dt}||w||^2 + 2\eta|w|_{[2]}^2 \le 2(G - \tilde{F}, Aw) + 2|\nu - \eta|(u, w)_{[2]} + 2(Bu - Bv)(Aw).$$

Choosing η such that $|\nu - \eta| < \frac{\nu}{2}$:

$$\frac{d}{dt}\|w\|^2 + \nu|w|_{[2]}^2 \le 2(G - \tilde{F}, Aw) + 2|\nu - \eta|(u, w)_{[2]} + 2(Bu - Bv)(Aw). \quad (2.13.1)$$

The last term is the more complicated so we work it a bit:

$$(Bu - Bv)(Aw) = b(u, u, Aw) - b(v, v, Aw)$$

= $b(u - v, u, Aw) + b(v, u, Aw) - b(v, v, Aw)$
= $-b(w, u, Aw) - b(v, w, Aw)$
= $-b(w, u, Aw) - b(w, w, Aw) - b(u, w, Aw)$

By (2.9.2) we have

$$\begin{aligned} |b(w, u, Aw)| &\leq C |w|^{\frac{1}{2}} ||w||^{\frac{1}{2}} ||u||^{\frac{1}{2}} |u||^{\frac{1}{2}}_{[2]} |w|_{[2]} \\ &\leq C_1 ||w|| |u|_{[2]} |w|_{[2]} \end{aligned}$$

and, by (2.9.3)

$$\begin{aligned} |b(w, w, Aw) + b(u, w, Aw)| &\leq C|w|^{\frac{1}{2}} |w||_{[2]}^{\frac{1}{2}} ||w|| |w|_{[2]} + C|u|_{[2]}^{\frac{1}{2}} ||u||^{\frac{1}{2}} ||w|| |w|_{[2]} \\ &\leq C|w|^{\frac{1}{2}} ||w|| |w|_{[2]}^{\frac{3}{2}} + C|u|_{[2]}^{\frac{1}{2}} ||u||^{\frac{1}{2}} ||w|| |w|_{[2]} \end{aligned}$$

1

Hence from (2.13.1) we have

$$\begin{aligned} &\frac{d}{dt} \|w\|^2 + \nu |w|_{[2]}^2 \\ &\leq 2|G - \tilde{F}||w|_{[2]} + 2|\nu - \eta||u|_{[2]}|w|_{[2]} + 2C_1 \|w\||u|_{[2]}|w|_{[2]} \\ &+ 2C|w|^{\frac{1}{2}} \|w\||w|_{[2]}^{\frac{3}{2}} + 2C|u|_{[2]}^{\frac{1}{2}}|u|^{\frac{1}{2}} \|w\||w|_{[2]}. \end{aligned}$$

Applying Young inequalities with suitable exponents and constants we arrive to

$$\begin{split} & \frac{d}{dt} \|w\|^2 + \nu |w|_{[2]}^2 \\ & \leq \frac{\nu}{10} |w|_{[2]}^2 + \frac{10}{\nu} |G - \tilde{F}|^2 + \frac{\nu}{10} |w|_{[2]}^2 + \frac{10}{\nu} |\nu - \eta|^2 |u|_{[2]}^2 \\ & + \frac{\nu}{10} |w|_{[2]}^2 + \frac{10}{\nu} C_1^2 \|w\|^2 |u|_{[2]}^2 \\ & + \frac{\nu}{10} |w|_{[2]}^2 + \frac{(10 \cdot 3)^3}{4^4 \cdot \nu^3} (2C)^4 |w|^2 \|w\|^4 + \frac{\nu}{10} |w|_{[2]}^2 + \frac{10}{\nu} C^2 |u| |u|_{[2]} \|w\|^2. \end{split}$$

Then

$$\frac{d}{dt} \|w\|^{2} + \frac{\nu}{2} |w|_{[2]}^{2}
\leq \frac{10}{\nu} |G - \tilde{F}|^{2} + \frac{10}{\nu} |\nu - \eta|^{2} |u|_{[2]}^{2} + \frac{10}{\nu} C_{1}^{2} \|w\|^{2} |u|_{[2]}^{2}
+ D_{1} |w|^{2} \|w\|^{4} + \frac{10}{\nu} C^{2} |u| |u|_{[2]} \|w\|^{2},$$
(2.13.2)

where $D_1 = \left(\frac{C}{2}\right)^4 \left(\frac{30}{\nu}\right)^3$. By Gronwall Inequality (2.7.6) we have for $t \in [0, T]$

$$\begin{aligned} \|w(t)\|^{2} \\ \leq \exp\left(\int_{0}^{T} \frac{10}{\nu} C_{1}^{2} |u(s)|_{[2]}^{2} + D_{1} |w(s)|^{2} \|w(s)\|^{2} + \frac{10}{\nu} C^{2} |u(s)| |u(s)|_{[2]} ds\right) \\ \cdot \left(\|w(0)\|^{2} + \int_{0}^{T} \frac{10}{\nu} |G(s) - \tilde{F}(s)|^{2} + \frac{10}{\nu} |\nu - \eta|^{2} |u(s)|_{[2]}^{2} ds\right). \end{aligned}$$
(2.13.3)

By theorems 2.8.1 and 2.8.2 and from $u \in L^2(0, T, D(A)) \cap L^{\infty}(0, T, V)$ the argument of the exponential is bounded, say less than a constant E, if we choose the triple (v_0, G, η) such that both $|v_0 - u_0|$, $||G - \tilde{F}||_{L^2(0, T, V')}$ and $|\nu - \eta|$ are less than δ for some $\delta > 0$ sufficiently small. Now put

$$\delta_1 := \min\left\{\frac{\nu}{2}, \ \delta, \ \frac{\varepsilon}{\sqrt{3}} (\exp E)^{-\frac{1}{2}}, \ \frac{\varepsilon}{\sqrt{3}} \left(\frac{10}{\nu}\right)^{-\frac{1}{2}} (\exp E)^{-\frac{1}{2}}, \\ \frac{\varepsilon}{\sqrt{3}} \left(\frac{10}{\nu} \|u\|_{L^2(0, T, D(A)}^2\right)^{-\frac{1}{2}} (\exp E)^{-\frac{1}{2}}\right\}.$$

It is clear from (2.13.3) that if both $|v_0 - u_0|$, $||G - \tilde{F}||_{L^2(0, T, V')}$ and $|\nu - \eta|$ are less than δ_1 , we have ||w(t)|| less than ε and, we have the continuity of \mathbb{S}_s .

Theorem 2.13.2. The map

$$\mathbb{S}_{2s}: V \times L^2(0, T, H) \times]0, +\infty[\rightarrow L^2(0, T, D(A))$$
$$(u_0, \tilde{F}, \nu) \mapsto u$$

is continuous. Where $u \in L^2(0, T, D(A))$ is the unique solution of Problems 2.10.1-2.10.2 (see Theorem 2.12.2).

Proof. As in the proof of Theorem 2.13.1 we fix $\varepsilon > 0$ and a triple $(u_0, \tilde{F}, \nu) \in H \times L^2(0, T, V') \times]0, +\infty[$ and consider the solution of Theorem 2.12.2 induced by this triple. Now consider another triple $(v_0, G, \eta) \in H \times L^2(0, T, V') \times]0, +\infty[$ and the solution associated with this triple. Put

$$w := v - u$$

and taking the scalar product with Aw we arrive to (2.13.2). Integrating over [0, T] we obtain

$$\begin{split} \|w(T)\|^{2} - \|w(0)\|^{2} + \int_{0}^{T} \frac{\nu}{2} \|w\|_{[2]}^{2} \\ \leq \frac{10}{\nu} \|G - \tilde{F}\|_{L^{2}(0,T,H)}^{2} + \frac{10}{\nu} \|u\|_{L^{2}(0,T,D(A))}^{2} \|\nu - \eta\|^{2} \\ + \frac{10}{\nu} C_{1}^{2} \|u\|_{L^{2}(0,T,D(A))}^{2} \|w\|_{C([0,T],V)}^{2} + TD_{1} \|w\|_{C([0,T],H)}^{2} \|w\|_{C([0,T],V)}^{4} \\ + \frac{10}{\nu} C^{2} \|u\|_{C([0,T],H)}^{2} \|u\|_{L^{2}(0,T,D(A))}^{2} \|w\|_{C([0,T],V)}^{2}. \end{split}$$

$$(2.13.4)$$

By Theorem 2.13.1, for a small enough $\delta > 0$ we have that if both $|v_0 - u_0|$, $\|\tilde{F} - G\|_{L^2(0,T,H)}$, and $|\eta - \nu|$ are less than δ , then $\|w\|_{C([0,T],V)} < 1$. Thus from (2.13.4) we have

$$\begin{split} &\int_{0}^{T}\|w\|_{[2]}^{2}\\ \leq &\frac{20}{\nu^{2}}\|G-\tilde{F}\|_{L^{2}(0,\,T,\,H)}^{2}+\frac{20}{\nu^{2}}\|u\|_{L^{2}(0,\,T,\,D(A))}^{2}|\nu-\eta|^{2}\\ &+ \Big(\frac{4}{\nu}+\frac{20}{\nu^{2}}C_{1}^{2}\|u\|_{L^{2}(0,\,T,\,D(A))}^{2}+\frac{2}{\nu}TD_{1}\|w\|_{C([0,\,T],\,H)}^{2}\\ &+ \frac{20}{\nu^{2}}C^{2}\|u\|_{C([0,\,T],\,H)}^{2}\|u\|_{L^{2}(0,\,T,\,D(A))}^{2}\Big)\|w\|_{C([0,\,T],\,V)}^{2}\\ \leq &\frac{20}{\nu^{2}}\|G-\tilde{F}\|_{L^{2}(0,\,T,\,H)}^{2}+D_{2}|\nu-\eta|^{2}+D_{3}\|w\|_{C([0,\,T],\,V)}^{2} \end{split}$$

for some constants D_2 , D_3 and all triples (v_0, G, η) satisfying

$$|v_0 - u_0| < \delta, \|\tilde{F} - G\|_{L^2(0, T, H)} < \delta, |\eta - \nu| < \delta.$$

Then for some δ_1 smaller than δ we have that if both $|v_0 - u_0|$, $\|\tilde{F} - G\|_{L^2(0, T, H)}$, and $|\eta - \nu|$ are less than δ_1 , we obtain

$$\left(\int_0^T \|w\|_{[2]}^2\right)^{\frac{1}{2}} < \varepsilon.$$

Therefore the map \mathbb{S}_{2s} is continuous

 \Box

Chapter 3

Controllability of Galerkin Approximations.

3.1 The FCE Procedure.

In this section we present the FCE Procedure, i.e., a procedure of *Factorization+Convexification+Extraction*:

3.1.1 A Lemma from Linear Algebra.

A result from Linear Algebra we will need is the following:

Lemma 3.1.1. Fix a linear space X of dimension $N \ge n + m$. Given two families $V := \{v_i \mid i = 1, ..., n\}$ and $W := \{w_j \mid j = 1, ..., m\}$ satisfying: The vectors in V are linearly independent and the vectors in $\pi W = \{\pi w_j \mid w_j \in W\}$ are linearly independent as well, where π is the projection onto some space V^c transversal to span(V), $(X = span(V) \oplus V^c)$. Then the family $V \cup W$ is linearly independent.

3.1.2 Factorization.

Consider a control-affine system

$$\dot{q} = f(q) + \sum_{i=1}^{r} v_i(t)g_i(q) \quad q \in \mathbb{R}^n, \, v_i \in \mathbb{R}$$
(3.1.1)

where f, g_i are smooth vector fields and $[g_i, g_j] = 0$ i = 1, ..., r.

In [2] it is proven that if we decompose the flow of system (3.1.1) as

$$\overrightarrow{exp} \int_0^t (f + gv(\tau))d\tau := \overrightarrow{exp} \int_0^t (f + \sum_{i=1}^r v_i(t)g_i(\tau))d\tau \qquad (3.1.2)$$

$$\overrightarrow{exp} \int_0^t (Ad \, G_{v(\tau)}^\tau) f \, d\tau \circ G_{v(\tau)}^\tau =: \overrightarrow{exp} \int_0^t (e^{-gw(\tau)})_* f \, d\tau \circ G_{v(t)}^\tau \tag{3.1.3}$$

where $g := (g_1, g_2, \cdots, g_r), v := (v_1, v_2, \cdots, v_r)^T$ and $G_{v(t)}^t$ stays for the flow $\overrightarrow{exp} \int_0^t gv(\tau) d\tau = e^{gw(t)}, w(t) = \int_0^t v(\tau) d\tau$, then

$$\overline{\mathcal{A}_{q_0}(f+gv)(t)} = \overline{\mathcal{A}_{q_0}((e^{-gV})_*f)(t) \circ \{G_{v(t)}^t \mid v(t) \in \mathbb{R}^r\}} \quad 1$$
(3.1.4)

Here $\mathcal{A}_x(y)(t)$ stays for the attainable set at time t from x following the vector fields y.

Similarly if we rewrite system (3.1.1) as

$$\dot{q} = f(q) + \sum_{i=1}^{r} (v_i^1(t) + v_i^2(t))g_i(q) \quad q \in \mathbb{R}^n, \, v_i^j \in \mathbb{R}$$

we arrive to

$$\overline{\mathcal{A}_{q_0}(f+gv)(t)} = \overline{\mathcal{A}_{q_0}((e^{-gV^2})_*f_1)(t) \circ \{G_{v^2(t)}^t \mid v^2(t) \in \mathbb{R}^r\}}$$
(3.1.5)

where $f_1(q) := f(q) + \sum_{i=1}^r v_i^1(t)g_i(q)$ and where v^1 , v^2 and V^2 are independent. The system $\dot{q} = (e^{-gV^2})_* f_1(q)$ is called *factorized system*.

Lemma 3.1.2. With $(e^{-gV^2})_*f_1$ and $G_{v^2}^t$ as in equation (3.1.5) we have

$$\overline{\mathcal{A}_{q_0}((e^{-gV^2})_*f_1)(t)} \circ \{G_{v^2(t)}^t \mid v^2(t) \in \mathbb{R}^r\}$$
$$\supseteq \overline{\mathcal{A}_{q_0}((e^{-gV^2})_*f_1)(t)} \circ \{G_{v^2(t)}^t \mid v^2(t) \in \mathbb{R}^r\}$$

Proof. Let $x \in \overline{\mathcal{A}_{q_0}((e^{-gV^2})_*f_1)(t)} \circ \{G_{v^2(t)}^t \mid v^2(t) \in \mathbb{R}^r\}$. Then there are $y \in \overline{\mathcal{A}_{q_0}((e^{-gV^2})_*f_1)(t)}$ and a control $u(t) \in \mathbb{R}^r t \in [0, T]$ such that $x = y \circ G_{u(t)}^T$. Let $y_n \longrightarrow y$, $y_n \in \mathcal{A}_{q_0}((e^{-gV^2})_*f_1)(t)$. Hence $x_n = y_n \circ G_{u(t)}^T$ is a sequence on $\mathcal{A}_{q_0}((e^{-gV^2})_*f_1)(t) \circ G_{u(t)}^t$ that converges to x.

So system (3.1.1) is approximately controllable in time t if

$$\overline{\mathcal{A}_q((\mathbf{e}^{-gV^2})_*f_1)(t)} \circ \{G_{v^2(t)}^t \mid v^2(t) \in \mathbb{R}^r\} = \mathbb{R}^n \quad \forall q \in \mathbb{R}^n.$$
(3.1.6)

If g_i are constant vector fields $-g_i(q) = g_i$ $i = 1, \ldots, r \quad q \in \mathbb{R}^n$ — they commute and the systems $\dot{q} = (e^{-gX(t)})_* f_1(q)$ and $\dot{q} = f_1(q + gX(t))$ coincide. A corollary of this is

Corollary 3.1.3. System (3.1.1) (with g constant) is approximately controllable in time t if

$$\overline{\mathcal{A}_q(f_{1X})(t)} \circ \{ e^{gV^2(t)} \} = \mathbb{R}^n \quad \forall q \in \mathbb{R}^n.$$

Where

$$f_{1X}(q) := f_1(q + gX(t)) = f(q + gX(t)) + g(q + gX(t))v^1$$

= $f_X(q) + gv^1 \quad X(t) \in \mathbb{R}^r.$

In particular it is approximately controllable in time t if

$$\overline{\mathcal{A}_q(f_{1X})(t)} = \mathbb{R}^n \quad \forall q \in \mathbb{R}^n.$$

¹Here V is independent of v.

²Here, for more precision, we should write $\{e^{gV^2(t)} \mid V^2(t) \in \mathbb{R}^r\}$.

3.1.3 Convexification.

If for some constant vector $\gamma \in \mathbb{R}^n$, $f(q) + \gamma$ belong to the convex set $Conv\{f_X \mid X \in \mathbb{R}^r\}$, then for every $u^1 \in \mathbb{R}^r$

$$f(q) + \gamma + gu^{1} \in Conv\{f_{X} \mid X \in \mathbb{R}^{r}\} + span(g)$$
$$\subseteq Conv\{f_{X} + gv^{1} \mid X, v^{1} \in \mathbb{R}^{r}\}.^{3}$$

This means that we can follow any of the vector fields $f(q) + \gamma + gv^1$ without changing the closure of the attainable set at time t (recall that convexification does not change the closure of attainable set at time t – see [8]). In particular system (3.1.1) is approximately controllable at time t if

$$\overline{\mathcal{A}_q(f(q) + \gamma + gv^1)(t)} = \mathbb{R}^n \quad \forall q \in \mathbb{R}^n.$$

3.1.4 Extraction.

Let C be a cone and suppose that

$$f(q) + C \subseteq Conv\{f_X \mid X \in \mathbb{R}^r\}.$$

Then putting G := span(g)

$$f(q) + C + G \subseteq f(q) + Conv(C) + G \subseteq Conv\{f_X(q) + G \mid X \in \mathbb{R}^r\}$$

= $Conv\{f_{1X}(q) \mid X \in \mathbb{R}^r\}.$

Now from Conv(C) + G we extract the linear space

$$G^1 := (G + Conv(C)) \cap (G - Conv(C)).$$

We shall call the directions on G^1 "extracted" directions. Since clearly $G \subseteq G^1$ because $0 \in C$, those directions in G will be called "old" directions and, those in $G^1 \setminus G$ "new" directions.

Adding new directions does not change the closure of attainable sets so, we can say that system (3.1.1) is appoximately controllable in time t if the "bigger" system $\dot{q} = f(q) + g_1 v^1$ is, where $v^1 \in \mathbb{R}^{r_1}$, $r_1 (\geq r)$ is the dimension of G^1 and g_1 is a matrix whose r_1 columns are vectors spanning G^1 .

3.1.5 Iterating FCEs.

Iterating FCE Procedures we obtain an increasing sequence

$$G =: G^0 \subseteq G^1 \subseteq \ldots \subseteq G^j \subseteq \ldots$$

of subspaces of controlled directions without changing the closure of the attainable set at time t. Obviously if for some $p \in \mathbb{N}$ we have $G^p = \mathbb{R}^n$, then the controllability in time t is an immediate consequence of Corollary 3.1.3 (note that in such a case we can set for V(t) any vector from \mathbb{R}^n).

³Here span(g) means the span of the columns of g. Conv(A) stays for convexification of the set A.

3.2 The Projection onto *H*.

Let $\mathbf{L}^2(R)_t$ the subspace of $\mathbf{L}^2(R)$ defined by

$$\mathbf{L}^{2}(R)_{t} := \{ v \in \mathbf{L}^{2}(R) \mid v \cdot \mathbf{n} = 0 \text{ on } \Gamma \}.$$

Let $v \in \mathbf{L}^2(R)_t$. We can write v as

$$v = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} \sum_{k \in \mathbb{N}_0 \times \mathbb{N}} v_{1k} \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\ \sum_{k \in \mathbb{N} \times \mathbb{N}_0} v_{2k} \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix}.$$

Now we want to write v as a sum of an element $u \in H$ (a solenoidal element) and a gradient of a function $q \in L^2(R) - v = u + \nabla q$. We put

$$\begin{pmatrix}
\sum_{k \in \mathbb{N}_0 \times \mathbb{N}} v_{1k} \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\
\sum_{k \in \mathbb{N} \times \mathbb{N}_0} v_{2k} \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right)
\end{pmatrix} = u + \nabla q;$$

$$u := \sum_{k \in \mathbb{N}_0^2} u_k W_k; \qquad q := \sum_{k \in \mathbb{N}^2 \setminus \{(0,0)\}} q_k \cos\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right).$$
(3.2.1)

Then

$$\nabla q = \begin{pmatrix} \sum_{k \in \mathbb{N}^2} -\frac{k_1 \pi}{a} q_k \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\ \sum_{k \in \mathbb{N}^2} -\frac{k_2 \pi}{b} q_k \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix}.$$

For $k \in \mathbb{N}^2 \setminus \mathbb{N}_0^2$ we find

$$\begin{cases} -q_k \frac{k_1 \pi}{a} = v_{1k}, & \text{for} \quad k \in \mathbb{N}_0 \times \{0\} \\ -q_k \frac{k_2 \pi}{b} = v_{2k}, & \text{for} \quad k \in \{0\} \times \mathbb{N}_0 \end{cases}$$

or,

$$q_k = \begin{cases} -\frac{a}{k_1 \pi} v_{1k}, & k \in \mathbb{N}_0 \times \{0\} \\ -\frac{b}{k_2 \pi} v_{2k}, & k \in \{0\} \times \mathbb{N}_0 \end{cases}.$$

and, for $k \in \mathbb{N}_0^2$ we obtain

$$-\frac{k_1\pi}{a}q_k = v_{1k} + \frac{k_2\pi}{b}u_k \tag{3.2.2}$$

$$-\frac{k_2\pi}{b}q_k = v_{2k} - \frac{k_1\pi}{a}u_k.$$
 (3.2.3)

Multiplying (3.2.2) by $-\frac{k_2\pi}{b}$ and (3.2.3) by $\frac{k_1\pi}{a}$ and then adding the products we obtain

$$\frac{k_2\pi}{b}v_{1k} - \frac{k_1\pi}{a}v_{2k} = \bar{k}u_k,$$

obtaining in this way u from v. Similarly we can obtain

$$-\left(\frac{k_2\pi}{b}\right)^2 q_k - \left(\frac{k_1\pi}{a}\right)^2 q_k = \frac{k_1\pi}{a} v_{1k} + \frac{k_2\pi}{b} v_{2k}$$

 or

$$\bar{k}q_k = \frac{k_1\pi}{a}v_{1k} + \frac{k_2\pi}{b}v_{2k}$$

Therefore, given

$$v = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} \sum_{k \in \mathbb{N}_0 \times \mathbb{N}} v_{1k} \sin\left(\frac{k_1 \pi x_1}{a}\right) \cos\left(\frac{k_2 \pi x_2}{b}\right) \\ \sum_{k \in \mathbb{N} \times \mathbb{N}_0} v_{2k} \cos\left(\frac{k_1 \pi x_1}{a}\right) \sin\left(\frac{k_2 \pi x_2}{b}\right) \end{pmatrix} \in \mathbf{L}^2(R)_t$$

we can write it, in a unique way, as a sum $v = u + \nabla q$, where $u \in H$ and $q \in L^2(R)$. So the projection map onto the solenoidal (divergence free) space H

$$P^{\nabla}: \mathbf{L}^2(R)_t \to H$$
$$v \mapsto u$$

is well defined and we have

$$u := P^{\nabla} v = \sum_{k \in \mathbb{N}_0^2} \frac{1}{\overline{k}} \left(\frac{k_2 \pi}{b} v_{1k} - \frac{k_1 \pi}{a} v_{2k} \right)$$
(3.2.4)
$$q = q_1 + q_2$$

$$q_1 = -\sum_{k \in \mathbb{N}_0 \times \{0\}} v_{1k} \frac{a}{k_1 \pi} \cos\left(\frac{k_1 \pi x_1}{a}\right) - \sum_{k \in \{0\} \times \mathbb{N}_0} v_{2k} \frac{b}{k_2 \pi} \cos\left(\frac{k_2 \pi x_2}{b}\right)$$
(3.2.5)

$$q_{2} = \sum_{k \in \mathbb{N}_{0}^{2}} \frac{1}{\bar{k}} \left(v_{1k} \frac{k_{1}\pi}{a} + v_{2k} \frac{k_{2}\pi}{b} \right) \cos\left(\frac{k_{1}\pi x_{1}}{a}\right) \cos\left(\frac{k_{2}\pi x_{2}}{b}\right). \quad (3.2.6)$$

It remains to show that $u \in H$ and $q_1, q_2 \in L^2(R)$. That follows from the fact that both v_1 and v_2 are in L^2 :

$$|u|^{2} = \sum_{k \in \mathbb{N}_{0}^{2}} -\bar{k} \frac{1}{\bar{k}^{2}} \left(\frac{k_{2}\pi}{b} v_{1k} - \frac{k_{1}\pi}{a} v_{2k} \right)^{2} \frac{ab}{4}$$

$$\leq \sum_{k \in \mathbb{N}_{0}^{2}} \left(|v_{1k}| + |v_{2k}| \right)^{2} \frac{ab}{4} \leq \left(||v_{1k}||_{L^{2}} + ||v_{2k}||_{L^{2}} \right)^{2};$$

$$\begin{aligned} \|q_1\|_{L^2}^2 &\leq \sum_{k \in \mathbb{N}_0 \times \{0\}} \left(v_{1k} \frac{a}{k_1 \pi} \right)^2 \frac{a}{2} + \sum_{k \in \{0\} \times \mathbb{N}_0} \left(v_{2k} \frac{b}{k_2 \pi} \right)^2 \frac{b}{2} \\ &\leq \frac{a^2 + b^2}{\pi^2} \left(\sum_{k \in \mathbb{N}_0 \times \{0\}} \left(v_{1k}^2 \frac{a}{2} + \sum_{k \in \{0\} \times \mathbb{N}_0} v_{2k}^2 \frac{b}{2} \right) \\ &\leq \frac{a^2 + b^2}{\pi^2} \left(\|v_{1k}\|_{L^2}^2 + \|v_{2k}\|_{L^2}^2 \right); \end{aligned}$$

$$q_{2} = \sum_{k \in \mathbb{N}_{0}^{2}} \frac{1}{\bar{k}^{2}} \left(v_{1k} \frac{k_{1}\pi}{a} + v_{2k} \frac{k_{2}\pi}{b} \right)^{2} \frac{ab}{4}$$

$$\leq \sum_{k \in \mathbb{N}_{0}^{2}} -\frac{1}{\bar{k}} \left(|v_{1k}| + |v_{2k}| \right)^{2} \frac{ab}{4} \leq -\frac{1}{(1, 1)} \left(\|v_{1k}\|_{L^{2}} + \|v_{2k}\|_{L^{2}} \right)^{2}.$$

3.3 Spectral Method.

We want to use *Spectral Algorithm*, i.e., we want to study NS equation in coordinates corresponding to the basis of eigenfunctions W defined in (2.3.1).

We consider the equation

$$u_t + (u \cdot \nabla)u + \nabla p = \Delta u + F + v;$$

and write

$$u = \sum_{k \in \mathbb{N}_0^2} u_k W_k.$$

Then we compute $(u \cdot \nabla)u = \begin{pmatrix} u \cdot \nabla u_1 \\ u \cdot \nabla u_2 \end{pmatrix}$:

$$\nabla u_1 = \begin{pmatrix} \sum_{k \in \mathbb{N}_0^2} u_k(-\frac{k_2\pi}{b}) \frac{k_1\pi}{a} \cos\left(\frac{k_1\pi x_1}{a}\right) \cos\left(\frac{k_2\pi x_2}{b}\right) \\ \sum_{k \in \mathbb{N}_0^2} u_k(-\frac{k_2\pi}{b})(-\frac{k_2\pi}{b}) \sin\left(\frac{k_1\pi x_1}{a}\right) \sin\left(\frac{k_2\pi x_2}{b}\right) \end{pmatrix}$$
$$\nabla u_2 = \begin{pmatrix} \sum_{k \in \mathbb{N}_0^2} u_k \frac{k_1\pi}{a}(-\frac{k_1\pi}{a}) \sin\left(\frac{k_1\pi x_1}{a}\right) \sin\left(\frac{k_2\pi x_2}{b}\right) \\ \sum_{k \in \mathbb{N}_0^2} u_k \frac{k_1\pi}{a} \frac{k_2\pi}{b} \cos\left(\frac{k_1\pi x_1}{a}\right) \cos\left(\frac{k_2\pi x_2}{b}\right) \end{pmatrix}.$$

Then

$$\begin{split} u \cdot \nabla u_1 \\ &= \sum_{m,n \in \mathbb{N}_0^2} u_m u_n (-\frac{n_2 \pi}{b}) (-\frac{m_2 \pi}{b}) \frac{m_1 \pi}{a} \sin\left(\frac{n_1 \pi x_1}{a}\right) \cos\left(\frac{n_2 \pi x_2}{b}\right) \\ &\quad \times \cos\left(\frac{m_1 \pi x_1}{a}\right) \cos\left(\frac{m_2 \pi x_2}{b}\right) \\ &\quad + \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{n_1 \pi}{a} (-\frac{m_2 \pi}{b}) (-\frac{m_2 \pi}{b}) \cos\left(\frac{n_1 \pi x_1}{a}\right) \sin\left(\frac{n_2 \pi x_2}{b}\right) \\ &\quad \times \sin\left(\frac{m_1 \pi x_1}{a}\right) \sin\left(\frac{m_2 \pi x_2}{b}\right) \\ &= \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{b} m_2 \left[\sin\left(\frac{(n_1 + m_1)\pi x_1}{a}\right) + \sin\left(\frac{(n_1 - m_1)\pi x_1}{a}\right) \right] \\ &\quad \times \left[\cos\left(\frac{(n_2 + m_2)\pi x_2}{b}\right) + \cos\left(\frac{(n_2 - m_2)\pi x_2}{b}\right) \right] \\ &\quad + \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{b} n_2 \left[\sin\left(\frac{(n_1 + m_1)\pi x_1}{a}\right) + \sin\left(\frac{(n_1 - m_1)\pi x_1}{a}\right) \right] \\ &\quad \times \left[- \cos\left(\frac{(n_2 + m_2)\pi x_2}{b}\right) + \cos\left(\frac{(n_2 - m_2)\pi x_2}{b}\right) \right]. \end{split}$$

We then arrive to

$$\begin{aligned} u \cdot \nabla u_{1} \\ &= \sum_{m,n \in \mathbb{N}_{0}^{2}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m_{1} n_{2} \frac{\pi}{b} (m_{2} - n_{2}) \sin\left(\frac{(n_{1} + m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} + m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{m,n \in \mathbb{N}_{0}^{2}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m_{1} n_{2} \frac{\pi}{b} (m_{2} + n_{2}) \sin\left(\frac{(n_{1} + m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} - m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{m,n \in \mathbb{N}_{0}^{2}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m_{1} n_{2} \frac{\pi}{b} (m_{2} - n_{2}) \sin\left(\frac{(n_{1} - m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} + m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{m,n \in \mathbb{N}_{0}^{2}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m_{1} n_{2} \frac{\pi}{b} (m_{2} + n_{2}) \sin\left(\frac{(n_{1} - m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} - m_{2})\pi x_{2}}{b}\right). \end{aligned}$$

$$(3.3.1)$$

Similarly:

$$\begin{split} u \cdot \nabla u_2 \\ &= \sum_{m,n \in \mathbb{N}_0^2} u_m u_n (-\frac{n_2 \pi}{b}) \frac{m_1 \pi}{a} (-\frac{m_1 \pi}{a}) \sin\left(\frac{n_1 \pi x_1}{a}\right) \cos\left(\frac{n_2 \pi x_2}{b}\right) \\ &\quad \times \sin\left(\frac{m_1 \pi x_1}{a}\right) \sin\left(\frac{m_2 \pi x_2}{b}\right) \\ &+ \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{n_1 \pi}{a} \frac{m_1 \pi}{a} \frac{m_2 \pi}{b} \cos\left(\frac{n_1 \pi x_1}{a}\right) \sin\left(\frac{n_2 \pi x_2}{b}\right) \\ &\quad \times \cos\left(\frac{m_1 \pi x_1}{a}\right) \cos\left(\frac{m_2 \pi x_2}{b}\right) \\ &= \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{a} m_1 \left[-\cos\left(\frac{(n_1 + m_1)\pi x_1}{a}\right) + \cos\left(\frac{(n_1 - m_1)\pi x_1}{a}\right)\right] \\ &\quad \times \left[\sin\left(\frac{(n_2 + m_2)\pi x_2}{b}\right) - \sin\left(\frac{(n_2 - m_2)\pi x_2}{b}\right)\right] \\ &+ \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{a} n_1 \left[\cos\left(\frac{(n_1 + m_1)\pi x_1}{a}\right) + \cos\left(\frac{(n_1 - m_1)\pi x_1}{a}\right)\right] \\ &\quad \times \left[\sin\left(\frac{(n_2 + m_2)\pi x_2}{b}\right) - \cos\left(\frac{(n_2 - m_2)\pi x_2}{b}\right)\right] \end{split}$$

and then,

$$\begin{aligned} u \cdot \nabla u_2 \\ &= \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{a} (n_1 - m_1) \cos\left(\frac{(n_1 + m_1)\pi x_1}{a}\right) \sin\left(\frac{(n_2 + m_2)\pi x_2}{b}\right) \\ &+ \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{a} (m_1 - n_1) \cos\left(\frac{(n_1 + m_1)\pi x_1}{a}\right) \sin\left(\frac{(n_2 - m_2)\pi x_2}{b}\right) \\ &+ \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{a} (m_1 + n_1) \cos\left(\frac{(n_1 - m_1)\pi x_1}{a}\right) \sin\left(\frac{(n_2 + m_2)\pi x_2}{b}\right) \\ &+ \sum_{m,n \in \mathbb{N}_0^2} u_m u_n \frac{\pi^2}{4ab} m_1 n_2 \frac{\pi}{a} (-m_1 - n_1) \cos\left(\frac{(n_1 - m_1)\pi x_1}{a}\right) \sin\left(\frac{(n_2 - m_2)\pi x_2}{b}\right). \end{aligned}$$

$$(3.3.2)$$

Grouping the terms in sum (3.3.1) and (3.3.2) envolving the product $u_m u_n$ we obtain

$$\begin{aligned} u \cdot \nabla u_{1} \\ &= \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \wedge n \frac{\pi}{b} (m_{2} - n_{2}) \sin\left(\frac{(n_{1} + m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} + m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \vee n \frac{\pi}{b} (m_{2} + n_{2}) \sin\left(\frac{(n_{1} + m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} - m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \vee n \frac{\pi}{b} (m_{2} - n_{2}) \sin\left(\frac{(n_{1} - m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} + m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \wedge n \frac{\pi}{b} (m_{2} + n_{2}) \sin\left(\frac{(n_{1} - m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} - m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{n} u_{n} \frac{\pi^{2}}{4ab} m \wedge n \frac{\pi}{b} (m_{2} + n_{2}) \sin\left(\frac{(n_{1} - m_{1})\pi x_{1}}{a}\right) \cos\left(\frac{(n_{2} - m_{2})\pi x_{2}}{b}\right) \end{aligned}$$
(3.3.3)

and,

$$\begin{aligned} u \cdot \nabla u_{2} \\ &= \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \wedge n \frac{\pi}{a} (n_{1} - m_{1}) \cos\left(\frac{(n_{1} + m_{1})\pi x_{1}}{a}\right) \sin\left(\frac{(n_{2} + m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \vee n \frac{\pi}{a} (m_{1} - n_{1}) \cos\left(\frac{(n_{1} + m_{1})\pi x_{1}}{a}\right) \sin\left(\frac{(n_{2} - m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \vee n \frac{\pi}{a} (m_{1} + n_{1}) \cos\left(\frac{(n_{1} - m_{1})\pi x_{1}}{a}\right) \sin\left(\frac{(n_{2} + m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} \frac{\pi^{2}}{4ab} m \wedge n \frac{\pi}{a} (-m_{1} - n_{1}) \cos\left(\frac{(n_{1} - m_{1})\pi x_{1}}{a}\right) \sin\left(\frac{(n_{2} - m_{2})\pi x_{2}}{b}\right) \\ &+ \sum_{n \in \mathbb{N}_{0}^{2}} u_{n}^{2} \frac{\pi^{2}}{4ab} n_{1} n_{2} \frac{\pi}{a} (2n_{1}) \sin\left(\frac{2n_{2}\pi x_{2}}{b}\right). \end{aligned}$$
(3.3.4)

Where the order under the sum sign, in (3.3.3) and (3.3.4), is the lexycographical one and, by definition:

$$m \lor n := m_1 n_2 + n_1 m_2;$$
 $m \land n := m_1 n_2 - n_1 m_2.$

Put

$$C_{m,n}^{\vee} = \frac{\pi^2}{4ab} m \vee n \quad \& \quad C_{m,n}^{\wedge} = \frac{\pi^2}{4ab} m \wedge n.$$

We rewrite⁵

$$u \cdot \nabla u_{1}$$

$$= \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\wedge} \frac{\pi}{b} (m_{2} - n_{2}) \sin\left(\frac{|n_{1} + m_{1}|\pi x_{1}}{a}\right) \cos\left(\frac{|n_{2} + m_{2}|\pi x_{2}}{b}\right)$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\vee} \frac{\pi}{b} (m_{2} + n_{2}) \sin\left(\frac{|n_{1} + m_{1}|\pi x_{1}}{a}\right) \cos\left(\frac{|n_{2} - m_{2}|\pi x_{2}}{b}\right)$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\vee} \frac{\pi}{b} (m_{2} - n_{2}) \sin\left(\frac{|n_{1} - m_{1}|\pi x_{1}}{a}\right) \cos\left(\frac{|n_{2} + m_{2}|\pi x_{2}}{b}\right)$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\wedge} \frac{\pi}{b} (m_{2} + n_{2}) \sin\left(\frac{|n_{1} - m_{1}|\pi x_{1}}{a}\right) \cos\left(\frac{|n_{2} - m_{2}|\pi x_{2}}{b}\right)$$

$$+ \sum_{\substack{n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{n}^{2} \frac{\pi^{2}}{4ab} n_{1} n_{2} \frac{\pi}{b} (2n_{2}) \sin\left(\frac{2n_{1}\pi x_{1}}{a}\right); \qquad (3.3.5)$$

⁴This last sum cooresponds to the sum over the diagonal $\{(m, n) \in (N_0^2)^2 \mid m = n\}$. ⁵m < n implies $m_1 \le n_1$. and

$$\begin{split} & u \cdot \nabla u_{2} \\ &= \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\wedge} \frac{\pi}{a} (n_{1} - m_{1}) \cos\left(\frac{|n_{1} + m_{1}|\pi x_{1}}{a}\right) \sin\left(\frac{|n_{2} + m_{2}|\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\vee} \frac{\pi}{a} (m_{1} - n_{1}) \cos\left(\frac{|n_{1} + m_{1}|\pi x_{1}}{a}\right) \sin\left(\frac{|n_{2} - m_{2}|\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\vee} \frac{\pi}{a} (-m_{1} + n_{1}) \cos\left(\frac{|n_{1} + m_{1}|\pi x_{1}}{a}\right) \sin\left(\frac{|n_{2} - m_{2}|\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\vee} \frac{\pi}{a} (-m_{1} + n_{1}) \cos\left(\frac{|n_{1} - m_{1}|\pi x_{1}}{a}\right) \sin\left(\frac{|n_{2} + m_{2}|\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\wedge} \frac{\pi}{a} (-m_{1} - n_{1}) \cos\left(\frac{|n_{1} - m_{1}|\pi x_{1}}{a}\right) \sin\left(\frac{|n_{2} - m_{2}|\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\wedge} \frac{\pi}{a} (m_{1} + n_{1}) \cos\left(\frac{|n_{1} - m_{1}|\pi x_{1}}{a}\right) \sin\left(\frac{|n_{2} - m_{2}|\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{m} u_{n} C_{m,n}^{\wedge} \frac{\pi}{a} (m_{1} + n_{1}) \cos\left(\frac{|n_{1} - m_{1}|\pi x_{1}}{a}\right) \sin\left(\frac{|n_{2} - m_{2}|\pi x_{2}}{b}\right) \\ &+ \sum_{\substack{n,n \in \mathbb{N}_{0}^{2} \\ m < n}} u_{n} u_{n} C_{m,n}^{\wedge} \frac{\pi}{a} (2n_{1}) \sin\left(\frac{2n_{2}\pi x_{2}}{b}\right). \end{split}$$
(3.3.6)

Now we project $(u \cdot \nabla)u$ in H:⁶ First we put

$$n(--)m := (n_1 - m_1, n_2 - m_2); \qquad n(-+)m := (n_1 - m_1, n_2 + m_2); n(+-)m := (n_1 + m_1, n_2 - m_2); \qquad n(++)m := (n_1 + m_1, n_2 + m_2); (a, b)^+ := (|a|, |b|), \quad a, b \in \mathbb{Z}$$

and then the projection can be written as

 $^{^{6}\}text{See}$ section 3.2. $^{7}\mathbb{Z}$ represents the set of integer numbers.

$$P^{\nabla}[(u \cdot \nabla)u]$$

$$= \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(++)m)^{+}} D_{++}W_{(n(++)m)^{+}}}$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\vee}}{(n(+-)m)^{+}} D_{+-}^{1}W_{(n(+-)m)^{+}}$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\vee}}{(n(+-)m)^{+}} D_{+-}^{2}W_{(n(+-)m)^{+}}$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\vee}}{(n(-+)m)^{+}} D_{-+}W_{(n(-+)m)^{+}}$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} D_{--}^{1}W_{(n(--)m)^{+}}$$

$$+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} D_{--}^{2}W_{(n(--)m)^{+}}; \qquad (3.3.7)$$

where

$$D_{++} = \frac{(n(++)m)_2^+ \pi}{b} \frac{\pi}{b} (m_2 - n_2) - \frac{(n(++)m)_1^+ \pi}{a} \frac{\pi}{a} (n_1 - m_1)$$

$$D_{+-}^1 = \frac{(n(+-)m)_2^+ \pi}{b} \frac{\pi}{b} (m_2 + n_2) - \frac{(n(+-)m)_1^+ \pi}{a} \frac{\pi}{a} (m_1 - n_1)$$

$$D_{+-}^2 = \frac{(n(+-)m)_2^+ \pi}{b} \frac{\pi}{b} (m_2 + n_2) - \frac{(n(+-)m)_1^+ \pi}{a} \frac{\pi}{a} (n_1 - m_1)$$

$$D_{-+} = \frac{(n(-+)m)_2^+ \pi}{b} \frac{\pi}{b} (m_2 - n_2) - \frac{(n(-+)m)_1^+ \pi}{a} \frac{\pi}{a} (m_1 + n_1)$$

$$D_{--}^1 = \frac{(n(--)m)_2^+ \pi}{b} \frac{\pi}{b} (m_2 + n_2) - \frac{(n(--)m)_1^+ \pi}{a} \frac{\pi}{a} (-m_1 - n_1)$$

$$D_{--}^2 = \frac{(n(--)m)_2^+ \pi}{b} \frac{\pi}{b} (m_2 + n_2) - \frac{(n(--)m)_1^+ \pi}{a} \frac{\pi}{a} (m_1 + n_1).$$

Equation (3.3.7) can be reduced to

Hence

$$P^{\nabla}[(u \cdot \nabla)u] = \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(++)m)^{+}} (\bar{n} - \bar{m})W_{(n(++)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\vee}}{(n(+-)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(+-)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\vee}}{(n(-+)m)^{+}} (\bar{n} - \bar{m})W_{(n(-+)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - m)\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - m)\mathrm{sign}(n_{2} - m)\mathrm{sign}(n_{2} - m)\mathrm{sign}(n_{2} - m)\mathrm{sign}(n_{2} - m)\mathrm$$

or, what is the same

$$P^{\nabla}[(u \cdot \nabla)u] = \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(++)m)^{+}} (\bar{n} - \bar{m})W_{(n(++)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\vee}}{(n(+-)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{2} - m_{2})W_{(n(+-)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} \frac{u_{m}u_{n}C_{m,n}^{\vee}}{(n(-+)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{1} - m_{1})W_{(n(-+)m)^{+}} + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{(n(--)m)^{+}} (\bar{n} - \bar{m})\mathrm{sign}(n_{1} - m_{1})\mathrm{sign}(n_{2} - m_{2})W_{(n(--)m)^{+}}$$

$$(3.3.8)$$

3.4 The Infinite ODE System.

We are now able to write an infinite system of ODEs related with the N-S Equation (1.0.1)

$$u_t + (u \cdot \nabla)u + \nabla p = \nu \Delta u + F + v.$$

As we can seen in the strong formulation of the N-S Problem (Problem 2.10.1), at each time τ the last equality is an equality between elements of V'. However if we project this equality onto the solenoidal space H we obtain

$$P^{\nabla}\Big(u_t + (u \cdot \nabla)u + \nabla p\Big) = P^{\nabla}\Big(\nu\Delta u + F + v\Big)$$

and, by (2.4.7) and the fact that $\tau \in D(A)$ a.e., we obtain

$$u_t(\tau) = -P^{\nabla} B u(\tau) + \nu \Delta u(\tau) + F + v(\tau).$$
(3.4.1)

We suppose ${\cal F}$ is solenoidal, otherwise we just take its solenoidal part. From

$$u = \sum_{k \in \mathbb{N}_0^2} u_k W_k$$

we derive

$$u_t = \sum_{\substack{m,n \in \mathbb{N}_0^2 \\ m < n}} (u_k)_t W_k$$
$$\Delta u = \sum_{\substack{m,n \in \mathbb{N}_0^2 \\ m < n}} \bar{k} u_k$$

Therefore we may write (3.4.1) in the form of the infinite system of ODEs:

$$\begin{split} \dot{u}_{k} &= \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(++)m)^{+} = k}} -\frac{u_{m}u_{n}C_{m,n}^{\wedge}}{\bar{k}}(\bar{n}-\bar{m}) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(--)m)^{+} = k}} \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{\bar{k}}(\bar{n}-\bar{m}) \mathrm{sign}(n_{1}-m_{1}) \mathrm{sign}(n_{2}-m_{2}) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(-+)m)^{+} = k}} -\frac{u_{m}u_{n}C_{m,n}^{\vee}}{\bar{k}}(\bar{n}-\bar{m}) \mathrm{sign}n_{1}-m_{1} \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(+-)m)^{+} = k}} \frac{u_{m}u_{n}C_{m,n}^{\vee}}{\bar{k}}(\bar{n}-\bar{m}) \mathrm{sign}(n_{2}-m_{2}) \\ &+ \nu \bar{k}u_{k} + F_{k} + v_{k}. \end{split}$$
(3.4.2)

3.4.1 Galerkin Approximations and G-Saturating Sets.

Trying to make the writting simpler we introduce some notation referring to the coefficients appearing in system (3.4.2):

Definition 3.4.1.

$$C_{m,n}^{++} := -\frac{\pi^2}{4ab} \frac{m \wedge n}{(m(++)n)^+} (\bar{n} - \bar{m})$$

$$C_{m,n}^{--} := \frac{\pi^2}{4ab} \frac{m \wedge n}{(m(--)n)^+} (\bar{n} - \bar{m}) \operatorname{sign}(n_1 - m_1) \operatorname{sign}(n_2 - m_2)$$

$$C_{m,n}^{-+} := -\frac{\pi^2}{4ab} \frac{m \vee n}{(m(-+)n)^+} (\bar{n} - \bar{m}) \operatorname{sign}(n_1 - m_1)$$

$$C_{m,n}^{+-} := \frac{\pi^2}{4ab} \frac{m \vee n}{(m(+-)n)^+} (\bar{n} - \bar{m}) \operatorname{sign}(n_2 - m_2).$$

Let g be a set of (coordinates of) r linearly independent vectors in H. Write also g for the matrix whose columns are those r vectors. Rewriting system (3.4.2) with the directions in g as the controlled ones, we obtain

$$\dot{u} = f(u) + gv \qquad v \in \mathbb{R}^r$$

or, equivalently

$$\dot{u}_k = (f(u))_k + (gv)_k \qquad v \in \mathbb{R}^r, \ k \in \mathbb{N}_0^2$$

where

$$(f(u))_{k} := \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(++)m)^{+} = k}} u_{m}u_{n}C_{m,n}^{++} \\ + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(--)m)^{+} = k}} u_{m}u_{n}C_{m,n}^{--} \\ + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(-+)m)^{+} = k}} u_{m}u_{n}C_{m,n}^{-+} \\ + \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(+-)m)^{+} = k}} u_{m}u_{n}C_{m,n}^{+-} \\ + \nu \bar{k}u_{k} + F_{k} \\ = B_{k}(u, u) + \nu \bar{k}u_{k} + F_{k},$$

defining $B_k(u, u) := (f(u))_k - (\nu \bar{k} u_k + F_k)$. Let us apply a FCE Procedure to this infinite system. Applying factorization to this system we obtain,⁸ the factorized system

$$(f_{1X}(u))_k = (f_X(u))_k + gv^1 = (f(u))_k + B_k(u, gX) + B_k(gX, u) + \nu \bar{k}(gX)_k + B_k((gX), (gX)) + gv^1$$
(3.4.3)

Now we put

$$\begin{aligned} & (\mathcal{V}_0(gX))_k &:= (f(u))_k + gv^1; \\ & (\mathcal{V}_1(gX))_k &:= B_k(u, gX) + B_k(gX, u) + \nu \bar{k}(gX)_k; \\ & (\mathcal{V}_2(gX))_k &:= B_k((gX), (gX)); \end{aligned}$$

and we note that \mathcal{V}_0 , \mathcal{V}_1 , and \mathcal{V}_2 are respectively, independent, linear and bilinear on gX vector fields.

Now, ginen $X \in \mathbb{R}^r$ we have

$$\mathcal{V}_0(g(-X)) = \mathcal{V}_0(gX); \quad \mathcal{V}_1(g(-X)) = -\mathcal{V}_1(gX) \text{ and } \mathcal{V}_2(g(-X)) = \mathcal{V}_2(gX).$$

So,

$$f(u) + B(gX, gX) = \frac{1}{2} \Big(f_X(u) + f_{-X}(u) \Big) \in Conv\{ f_X(u) \mid X \in \mathbb{R}^r \}.$$

The set

$$\{B(gX, gX) \mid X \in \mathbb{R}^r\} = \{B(Y, Y) \mid Y \in span(g)\}$$

is a cone.

 8 See section 3.1.

Next, from

$$Conv(\{B(gX, gX) \mid X \in \mathbb{R}^r\} + G = \{B(Y, Y) \mid Y \in span(g)\}) + G,$$

where G := span(g), we extract the subspace

$$G^{1} := \left(G + Conv\{B(Y, Y) \mid Y \in G\}\right) \cap \left(G - Conv\{B(Y, Y) \mid Y \in G\}\right).$$

Now we present two definitions:

Definition 3.4.2. A finite set of vectors $g \subseteq D(A)^{-9}$ is said **Saturating** for system (3.4.2) if the sequence $(G^j)_{n \in \mathbb{N}}$ of subspaces of H defined recursively by

$$G^{0} := G = span(g);$$

$$G^{j+1} := \left(G^{j} + Conv\{B(Y, Y) \mid Y \in G^{j}\}\right)$$

$$\cap \left(G^{j} - Conv\{B(Y, Y) \mid Y \in G^{j}\}\right) \cap D(A); \quad ^{10}$$

satisfies

$$\overline{\bigcup_{i\in\mathbb{N}}G^i} = H.$$

We say that g is **G-Saturating** for system (3.4.2) if $span(g) = span\{W_k \mid k \in \mathcal{K}^1\}$ for some finite set of modes $\mathcal{K}^1 \in \mathcal{FP}(\mathbb{N}^2_0)$, and there exists a sequence of subspaces H^j such that $H^0 := span(g)$ and

$$H^{j+1} \subseteq \left(H^j + Conv\{B(Y, Y) \mid Y \in H^j\}\right) \cap \left(H^j - Conv\{B(Y, Y) \mid Y \in H^j\}\right);$$
$$\overline{\bigcup_{i \in \mathbb{N}} H^i} = H;$$

and, besides there exists a finite subset $\mathcal{K}^{j+2} \in \mathcal{FP}(\mathbb{N}_0^2)$ such that $H^{j+1} = span\{W_k \mid k \in \mathcal{K}^{j+2}\}.$

Remark 7. In the previuos definition the condition

$$(A) \qquad \overline{\bigcup_{i \in \mathbb{N}} D^i} = H$$

 $is \ equivalent \ to$

(B)
$$\forall x \in H [j \to \infty \text{ only } if |x - \Pi_j x| \to 0]$$

where D^i stays for either G^i or H^i . Indeed if (B) is satisfied and $x \in H$, we have $\prod_i x \in D^i$ and so, $(\prod_i x)_{i \in \mathbb{N}}$ is a sequence in $\bigcup_{i \in \mathbb{N}} D^i$ converging to x. Thus $\overline{\bigcup_{i \in \mathbb{N}} D^i} \supseteq H$.

Conversely if (A) is satisfied and $x \in H$, we can find $j_0 \in \mathbb{N}$ such that $|x^0 - G^{j_0}| < 1$ and, for each $n \in \mathbb{N}_0$ we can find $j_n \in \mathbb{N}$ and $x^n \in D^{j_n}$ such that $j_n > j_{n-1}$ and $|x-x^n| < \frac{1}{n+1}$, then also $|x-\Pi_{j_n}x| < \frac{1}{n+1}$. ¹¹ So $(|x-\Pi_{j_n}x|)_{n\in\mathbb{N}}$ (that is a subsequence of $(|x - \Pi_j x|)_{j\in\mathbb{N}}$) converges to zero. Since the sequence $(D^i)_{i\in\mathbb{N}}$ is increasing (in the inclusion sense), the sequence $(|x - \Pi_j x|)_{j\in\mathbb{N}}$ is decreasing so, it converges to zero (as does its subsequence $(|x - \Pi_{j_n}x|)_{n\in\mathbb{N}}$).

⁹Recall that $D(A) \subset H$ has been defined in (1.0.5).

 $^{^{10}{\}rm We}$ intersect with D(A) because, it is helpful for the study of controllability issues of the infinite system.

¹¹Because $|x - \prod_{j_n} x| = min\{|x - y| \mid y \in G^{j_n}\}.$

Definition 3.4.3. A Galerkin approximation of system (3.4.2) is the same system with the additional condition

$$k, n, m \in \mathcal{G} \in \mathcal{FP}(\mathbb{N}_0^2).$$

From now we shall look for the existence of a G-Saturating set \mathcal{K}^1 for system (3.4.2). The existence of such a set means that for any given finite set of modes $\mathcal{O} \subseteq \mathbb{N}^2_0$, we want to observe, there exists $p \in \mathbb{N}$ such that $\mathcal{O} \subseteq H^p$.

For j < p the construction of H^{j+1} depends only on vectors of $H^j \subseteq H^p$. Therefore from the \mathcal{K}^{p+1} -Galerkin approximation with \mathcal{K}^j $(1 \leq j \leq p)$ as set of controlled modes, i.e., with H^{j-1} as set of controlled directions, we can arrive by a FCE Procedure to the same Galerkin approximation with \mathcal{K}^{j+1} as set of controlled modes.

Iterating FCE procedures, after p steps (starting with \mathcal{K}^1 as set of controlled modes), we arrive to the \mathcal{K}^{p+1} -Galerkin approximation with \mathcal{K}^{p+1} as set of controlled modes and, the approximate controllability at time t of such system is an immediate consequence of Corollary 3.1.3. The approximate controllability at time t of the \mathcal{K}^{p+1} -Galerkin approximation with \mathcal{K}^1 as set of controlled modes follows from the fact that a FCE Procedure does not change closure of attainable set at time t.

Hence the controlling of the modes in \mathcal{K}^1 in the \mathcal{K}^{p+1} -Galerkin approximation we "observe" approximate controllability in the state space $H^p = span(\mathcal{K}^{p+1})$ and so also in the " \mathcal{O} -space".

3.5 Looking for Saturating Sets.

We still working in the rectangle $[0, a] \times [0, b]$. We set two elements $n, m \in \mathbb{N}_0^2$ such that m < n. We have that

$$\bar{n} - \bar{m} = 0 \iff \frac{m_1^2}{a^2} + \frac{m_2^2}{b^2} = \frac{n_1^2}{a^2} + \frac{n_2^2}{b^2}$$

and, in this equation $m_1 = n_1 \iff m_2 = n_2$. So since m < n we have that $\bar{n} - \bar{m} = 0$ implies $m_1 \neq n_1 \& m_2 \neq n_2$. Then

$$\bar{n} - \bar{m} = 0 \iff \frac{a^2}{b^2} = -\frac{m_1^2 - n_1^2}{m_2^2 - n_2^2} = \frac{(n_1 - m_1)(n_1 + m_1)}{(m_2 - n_2)(m_2 + n_2)}.$$
 (3.5.1)

Now since $m_1 < n_1$ which, from (3.5.1), implies $m_2 > n_2$ we obtain

$$\bar{n} - \bar{m} = 0 \& m < n \Rightarrow n_1 > m_1 \& n_2 < m_2.$$
 (3.5.2)

By (3.5.2) we obtain

Corollary 3.5.1. Under the condition $m < n \land (m_1 = n_1 \lor n_2 \ge m_2)$ we have

$$C_{m,n}^{++} = 0 \iff m \land n = 0 \iff C_{m,n}^{--} = 0$$

and

$$C_{m,n}^{-+} \neq 0 \neq C_{m,n}^{+-}$$

Proposition 3.5.2. The set $\mathcal{K}^1 := \{(n_1, n_2) \in \mathbb{N}^2 \mid n_1, n_2 \leq 3\} \setminus \{(3, 3)\}$ is G-Saturating for system (3.4.2).¹²

Proof. For any $j \in \mathbb{N}_0$ put

and

and,

$$\mathcal{K}^j := \{ (n_1, n_2) \in \mathbb{N}_0^2 \mid n_1, n_2 \le j+2 \} \setminus \{ (j+2, j+2) \}.$$

We shall prove that the directions in $H^{j-1} := span(\mathcal{K}^j)$ can be obtained by a suitable FCE Procedure from the directions in H^{j-2} $(j \ge 2)$.

For every pair $(n, m) \in \mathcal{K}^1$, $n \neq m$ and for every $\lambda \in \mathbb{R}^+$ we define the vectors $v_{n,m}^{\lambda}$, $w_{m,n}^{\lambda} \in \mathbb{R}^r$ defined by

$$\begin{cases} (v_{m,n}^{\lambda})_n = \lambda;\\ (v_{m,n}^{\lambda})_m = 1;\\ (v_{m,n}^{\lambda})_k = 0, \quad k \in \mathcal{K}^1 \setminus \{n, m\};\\ \begin{cases} (w_{m,n}^{\lambda})_n = \lambda;\\ (w_{m,n}^{\lambda})_m = -1;\\ (w_{m,n}^{\lambda})_k = 0, \quad k \in \mathcal{K}^1 \setminus \{n, m\}. \end{cases}$$

The vector fields

$$\frac{f_{v_{m,n}^{\lambda}}(u) + f_{-v_{m,n}^{\lambda}}(u)}{2} = f(u) + \lambda \gamma_{m,n}$$
$$\frac{f_{w_{m,n}^{\lambda}}(u) + f_{-w_{m,n}^{\lambda}}(u)}{2} = f(u) - \lambda \gamma_{m,n},$$

where

$$\gamma_{m,n} = C_{m,n}^{--}\partial_{(n(--)m)^+} + C_{m,n}^{-+}\partial_{(n(-+)m)^+} + C_{m,n}^{+-}\partial_{(n(+-)m)^+} + C_{m,n}^{++}\partial_{(n(++)m)^+},$$

belong to $Conv\{f_{\pm v_{m,n}^{\lambda}}, f_{\pm w_{m,n}^{\lambda}} \mid \lambda \in \mathbb{R}^{+}, m, n \in \mathcal{K}^{1}, n \neq m\}.$ Now we shall extract from $\{f_{X(t)} \mid X(t) \in \mathbb{R}^{r}\}$ a family as follows: First we define for each pair $(m, n) \in \mathcal{K}^1$, $n \neq m$ the vector $\delta_{m,n}$ in \mathbb{R}^N by:

$$\delta_{m,n} = C_{m,n}^{--} e_{(n(--)m)^+} + C_{m,n}^{-+} e_{(n(-+)m)^+} + C_{m,n}^{+-} e_{(n(+-)m)^+} + C_{m,n}^{++} e_{(n(++)m)^+}$$
(3.5.3)

Now we select the subfamily $F_{S_1} := \{\delta_{m,n} \mid (m,n) \in S_1 \subseteq (K^1)^2\}$ of this vectors, where

$$S_{1} = \{((1,2), (2,1)); ((1,1), (2,3)); ((1,2), (2,2)); \\ ((1,1), (3,2)); ((2,1), (2,2)); ((1,1), (1,3)); ((1,1), (3,1))\}.$$
(3.5.4)

 $^{^{12}\}mathrm{We}$ take (3,3) off only because a writting reason. We shall apply a induction procedure in the proof and without these "corners" it becomes simpler.

The vectors of this family are precisely

$$\begin{split} \delta_{(1,2),(2,1)} &= \frac{9\pi^2(b^2 - a^2)}{4ab(a^2 + b^2)}e_{(1,1)} + \frac{15\pi^2(a^2 - b^2)}{4ab(9a^2 + b^2)}e_{(1,3)} \\ &+ \frac{15\pi^2(a^2 - b^2)}{4ab(a^2 + 9b^2)}e_{(3,1)} + \frac{\pi^2(b^2 - a^2)}{4ab(a^2 + b^2)}e_{(3,3)} \\ \delta_{(1,1),(2,3)} &= \frac{\pi^2(3b^2 + 8a^2)}{4ab(4a^2 + b^2)}e_{(1,2)} - \frac{5\pi^2(8a^2 + 3b^2)}{4ab(16a^2 + b^2)}e_{(1,4)} \\ &+ \frac{5\pi^2(8a^2 + 3b^2)}{4ab(4a^2 + 9b^2)}e_{(3,2)} - \frac{\pi^2(3b^2 + 8a^2)}{4ab(16a^2 + 9b^2)}e_{(3,4)} \\ \delta_{(1,2),(2,2)} &= -\frac{9b\pi^2}{2a(16a^2 + b^2)}e_{(1,4)} + \frac{3b\pi^2}{2a(16a^2 + 9b^2)}e_{(3,4)} \\ \delta_{(1,1),(3,2)} &= -\frac{\pi^2(8b^2 + 3a^2)}{4ab(a^2 + 4b^2)}e_{(2,1)} - \frac{5\pi^2(3a^2 + 8b^2)}{4ab(9a^2 + 4b^2)}e_{(2,3)} \\ &+ \frac{5\pi^2(3a^2 + 8b^2)}{4ab(a^2 + 16b^2)}e_{(4,1)} + \frac{\pi^2(8b^2 + 3a^2)}{4ab(9a^2 + 16b^2)}e_{(4,3)} \\ \delta_{(2,1),(2,2)} &= -\frac{9a\pi^2}{2b(16b^2 + a^2)}e_{(4,1)} - \frac{3a\pi^2}{2b(16b^2 + 9a^2)}e_{(4,3)} \\ \delta_{(1,1),(1,3)} &= \frac{2a\pi^2}{b(b^2 + a^2)}e_{(2,2)} - \frac{a\pi^2}{b(b^2 + 4a^2)}e_{(2,4)} \\ \delta_{(1,1),(3,1)} &= -\frac{2b\pi^2}{a(b^2 + a^2)}e_{(2,2)} + \frac{b\pi^2}{a(a^2 + 4b^2)}e_{(4,2)}, \end{split}$$

Projecting the vectors in this subfamily on the space $span\{e_k \mid k \in \mathcal{K}^2 \setminus \mathcal{K}^1\}$ we obtain

$$\Pi_{1}\delta_{(1,2),(2,1)} = \frac{\pi^{2}(b^{2}-a^{2})}{4ab(a^{2}+b^{2})}e_{(3,3)}$$
(3.5.5)

$$\Pi_{1}\delta_{(1,1),(2,3)} = -\frac{5\pi^{2}(8a^{2}+3b^{2})}{4ab(16a^{2}+b^{2})}e_{(1,4)} - \frac{\pi^{2}(3b^{2}+8a^{2})}{4ab(16a^{2}+9b^{2})}e_{(3,4)}$$
(3.5.5)

$$\Pi_{1}\delta_{(1,2),(2,2)} = -\frac{9b\pi^{2}}{2a(16a^{2}+b^{2})}e_{(1,4)} + \frac{3b\pi^{2}}{2a(16a^{2}+9b^{2})}e_{(3,4)}$$
(1.1)

$$\Pi_{1}\delta_{(1,1),(3,2)} = \frac{5\pi^{2}(3a^{2}+8b^{2})}{4ab(a^{2}+16b^{2})}e_{(4,1)} + \frac{\pi^{2}(8b^{2}+3a^{2})}{4ab(9a^{2}+16b^{2})}e_{(4,3)}$$
(1.1)

$$\Pi_{1}\delta_{(2,1),(2,2)} = -\frac{9a\pi^{2}}{2b(16b^{2}+a^{2})}e_{(4,1)} - \frac{3a\pi^{2}}{2b(16b^{2}+9a^{2})}e_{(4,3)}$$
(1.1)

$$\Pi_{1}\delta_{(1,1),(1,3)} = -\frac{a\pi^{2}}{b(b^{2}+4a^{2})}e_{(2,4)}$$
(1.1)

$$\Pi_{1}\delta_{(1,1),(3,1)} = \frac{b\pi^{2}}{a(a^{2}+4b^{2})}e_{(4,2)}.$$

Now we consider the case $a \neq b$, i. e., if our rectangle R is not a square (the case of the square will be considered below in Remark 9). If $a \neq b$ these projections are linearly independent so, by Lemma 3.1.1 the 15 vectors of the family $\{e_k \mid k \in \mathcal{K}^1\} \cup F_{S_1}$ are linearly independent in $span(\mathcal{K}^2) = H^1$ and then they span $span\{e_k \mid k \in \mathcal{K}^2\} \subset H$.

Now we extract the linear space

$$\tilde{H}^{1} := \left(H^{0} + Conv\{\lambda\delta_{m,n} \mid (m, n) \in S_{1}, \lambda \in \mathbb{R}\}\right)$$
$$\cap \left(H^{0} - Conv\{\lambda\delta_{m,n} \mid (m, n) \in S_{1}, \lambda \in \mathbb{R}\}\right)$$

from the space

$$\left(H^0 + Conv\{B(Y, Y) \mid Y \in H^0\}\right) \cap \left(H^0 - Conv\{B(Y, Y) \mid Y \in H^0\}\right).$$

Since $Conv\{\lambda\delta_{m,n} \mid (m, n) \in S_1, \lambda \in \mathbb{R}\}\$ coincides with $span\{\lambda\delta_{m,n} \mid (m, n) \in S_1, \lambda \in \mathbb{R}\}\$ $S_1, \lambda \in \mathbb{R}\}$, we have that

$$\tilde{H}^1 = span(\mathcal{K}^1 \cup \{\delta_{m,n} \mid (m, n) \in S_1\}) = span(\mathcal{K}^2) = H^1.$$

The following proposition completes the proof.

Proposition 3.5.3. From the directions in $H^j = span(\mathcal{K}^{j+1}), (j \ge 1)^{13}$ we can obtain the directions in $H^{j+1} = span(\mathcal{K}^{j+2})$ by a FCE Procedure.

Proof. We consider two cases "j even" and "j odd".

• j even: In this case

$$\mathcal{K}^{j+1} := \{ (n_1, n_2) \in \mathbb{N}_0^2 \mid n_1, n_2 \le j+3 \} \setminus \{ (j+3, j+3) \}.$$

can be written as

$$\mathcal{K}^{j+1} := \{ (n_1, n_2) \in \mathbb{N}_0^2 \mid n_1, n_2 \le 2p+1 \} \setminus \{ (2p+1, 2p+1) \},\$$

setting $p = \frac{j+2}{2}$. Then $p \ge 2$. As we did before in the case " $j = 0 \rightarrow j = 1$ ", we extract a subfamily $F_{S_{j+1}} := \{\delta_{m,n} \mid (m, n) \in S_{j+1} \subseteq (K^{j+1})^2\}$ of $\{B(Y, Y) \mid Y \in H^j\}$ where now the "selection" is

$$\begin{split} S_{j+1} &= \{((1,2),(2p,2p-1))\} \\ &\cup \{((1,1),(2z,2p+1)) \mid z=1,\ldots,p\} \cup \{((1,p+1)(2,p+1))\} \\ &\cup \{((1,1),(2p+1,2z)) \mid z=1,\ldots,p\} \cup \{((p+1,1),(p+1,2))\} \\ &\cup \{((s,1),(s,2p+1)) \mid s=1,\ldots,p\} \\ &\cup \{((1,s),(2p+1,s)) \mid s=1,\ldots,p\}. \end{split}$$

If we write explicitely the vectors of ${\cal F}_{S_{j+1}}$ we obtain quite long expressions, for example we have that $C^{--}_{(1,2),(2p,2p-1)}$ equals

$$-\frac{\left(a^2(-3+2p)+b^2(-1+2p)\right)(\pi+2p\pi)^2\operatorname{sign}(-3+2p)\operatorname{sign}(-1+2p)}{4ab(b^2|1-2p|^2+a^2|3-2p|^2)};$$

¹³The case j = 0 was already seen in the proof of proposition 3.5.2.

so, here we will not write those vectores explicitely. Anyway, those vectors are

And, projecting them onto the space $span\{e_k \mid k \in \mathcal{K}^{j+2} \setminus \mathcal{K}^{j+1}\}$ we arrive to the family $\prod_{j+1} F_{S_{j+1}}$ whose elements are

$$\Pi_{j+1}\delta_{(1,2),(2p,2p-1)} = C_{(1,2),(2p,2p-1)}^{++} e_{(2p+1,2p+1)};$$

$$\Pi_{j+1}\delta_{(1,1),(2z,2p+1)} = C_{(1,1),(2z,2p+1)}^{-+} e_{(2z-1,2(p+1))} + C_{(1,1),(2z,2p+1)}^{++} e_{(2z+1,2(p+1))}$$

$$z = 1, \dots, p;$$
(3.5.6)

$$\Pi_{j+1}\delta_{(1,p+1),(2,p+1)} = C^{-+}_{(1,p+1),(2,p+1)}e_{(1,2(p+1))} + C^{++}_{(1,p+1),(2,p+1)}e_{(3,2(p+1))};$$
(3.5.7)
$$\Pi_{j+1}\delta_{j+1}\delta_{j+1} = C^{+-}_{j+1}\delta_{j+$$

$$\Pi_{j+1}\delta_{(1,1),(2p+1,2z)} = C_{(1,1),(2p+1,2z)}^{+-}e_{(2(p+1),2z-1)} + C_{(1,1),(2p+1,2z)}^{++}e_{(2(p+1),2z+1)}$$
$$z = 1, \dots, p; \qquad (3.5.8)$$

 $\Pi_{j+1}\delta_{(p+1,1),(p+1,2)} = C^{+-}_{(p+1,1),(p+1,2)}e_{(2(p+1),1)} + C^{++}_{(p+1,1),(p+1,2)}e_{(2(p+1),3)};$ (3.5.9)

$$\Pi_{j+1}\delta_{(s,1),(s,2p+1)} = C^{++}_{(s,1),(s,2p+1)}e_{(2s,2(p+1))}, \qquad s = 1, \dots, p;$$
(3.5.10)

$$\Pi_{j+1}\delta_{(1,s),(2p+1,s)} = C_{(1,s),(2p+1,s)}^{++} e_{(2(p+1),2s)}, \qquad s = 1, \dots, p.$$
(3.5.11)

No one of the coefficients appearing in these expressions vanishes because all pairs (m, n) satisfy $m < n \land (m_1 = n_1 \lor n_2 \ge m_2)$ and because no one of the

following expressions vanish

$$\begin{array}{l} (1,2)\wedge(2p,2p-1)=-1-2p;\\ (1,1)\wedge(2z,2p+1)=1+2p-2z,\quad z=1,\ldots,\,p;\\ (1,p+1)\wedge(2,p+1)=-(1+p);\\ (1,1)\wedge(2p+1,2z)=-1-2p+2z,\quad z=1,\ldots,\,p;\\ (p+1,1)\wedge(p+1,2)=1+p;\\ (s,1)\wedge(s,2p+1)=2ps,\quad s=1,\ldots,\,p;\\ (1,s)\wedge(2p+1,s)=-2ps,\quad s=1,\ldots,\,p. \end{array}$$

Hence we can see that these vectors are linearly independent. Indeed it suffices to prove that:

- The vectors $\Pi_{j+1}\delta_{(1,1),(2,2p+1)}$, (z = 1 in (3.5.6)) and $\Pi_{j+1}\delta_{(1,p+1),(2,p+1)}$ are linearly independent; and
- The vectors $\Pi_{j+1}\delta_{(1,1),(2p+1,2)}$, (z = 1 in (3.5.8)) and $\Pi_{j+1}\delta_{(p+1,1),(p+1,2)}$ are linearly independent;

But that comes from

$$\frac{C_{(1,1),(2,2p+1)}^{-+}}{C_{(1,p+1),(2,p+1)}^{-+}} = \frac{(3+2p)(3b^2+4a^2p(1+p))}{9b^2(1+p)}$$

$$\neq -\frac{(-1+2p)(3b^2+4a^2p(1+p))}{3b^2(1+p)} = \frac{C_{(1,1),(2,2p+1)}^{++}}{C_{(1,p+1),(2,p+1)}^{++}};$$

and

.

$$\frac{C_{(1,1),(2p+1,2z)}^{+-}}{C_{(p+1,1),(p+1,2)}^{+-}} = \frac{(3+2p)(3a^2+4b^2p(1+p))}{9a^2(1+p)}$$

$$\neq -\frac{(-1+2p)(3a^2+4b^2p(1+p))}{3a^2(1+p)} = \frac{C_{(1,1),(2,2p+1)}^{++}}{C_{(1,p+1),(2,p+1)}^{++}}$$

By Lemma 3.1.1 the $(2(p+1))^2 - 1$ of $F_{S_{j+1}} \cup \{e_k \mid k \in \mathcal{K}^{j+1}\}$ are linearly independent and then they span

$$span\{e_k \mid 1 \le k_1, k_2 \le 2(p+1)\} \setminus \{(2(p+1), 2(p+1))\}$$

= $span\{e_k \mid k \in \mathcal{K}^{j+2}\}.$

Since with $\delta_{m,n}$ also $\lambda \delta_{m,n}$ belongs to $\{B(Y, Y) \mid Y \in H^j\}$ for every $\lambda \in \mathbb{R}$, we can select the linear space \tilde{H}^{j+1} defined by

$$\tilde{H}^{j+1} := \left(H^j + Conv\{\lambda \delta_{m,n} \mid (m, n) \in S_{j+1}, \lambda \in \mathbb{R} \} \right)$$
$$\cap \left(H^j - Conv\{\lambda \delta_{m,n} \mid (m, n) \in S_{j+1}, \lambda \in \mathbb{R} \} \right)$$

from the space

$$\left(H^{j} + Conv\{B(Y, Y) \mid Y \in H^{j}\}\right) \cap \left(H^{j} - Conv\{B(Y, Y) \mid Y \in H^{j}\}\right).$$

Since $Conv\{\lambda\delta_{m,n} \mid (m, n) \in S_{j+1}, \lambda \in \mathbb{R}\}\)$ coincides with $span\{\lambda\delta_{m,n} \mid \lambda\delta_{m,n} \mid 0 \leq N\}$ $(m, n) \in S_{j+1}, \lambda \in \mathbb{R}\}$, we have that

$$\tilde{H}^{j+1} = span(\mathcal{K}^{j+1} \cup \{\delta_{m,n} \mid (m, n) \in S_{j+1}\}) = span(\mathcal{K}^{j+2}) = H^{j+1}.$$

Now we study the other case

• j odd: In this case

$$\mathcal{K}^{j+1} := \{ (n_1, n_2) \in \mathbb{N}_0^2 \mid n_1, n_2 \le j+3 \} \setminus \{ (j+3, j+3) \}.$$

can be written as

$$\mathcal{K}^{j+1} := \{ (n_1, n_2) \in \mathbb{N}_0^2 \mid n_1, n_2 \le 2p \} \setminus \{ (2p, 2p) \},\$$

setting $p = \frac{j+3}{2}$. Then $p \ge 2$. We extract the subfamily $F_{S_{j+1}} := \{\delta_{m,n} \mid (m, n) \in S_{j+1} \subseteq (K^{j+1})^2\}$ of $\{B(Y, Y) \mid Y \in H^j\}$ where now the "selection" is

$$\begin{split} S_{j+1} &= \{ ((1,2p-1),(2p-1,1)) \} \\ &\cup \{ ((1,1),(2z-1,2p)) \mid z=2,\ldots,p \} \cup \{ ((1,p)(3,p+1)) \} \\ &\cup \{ ((1,1),(2p,2z-1)) \mid z=2,\ldots,p \} \cup \{ ((p,1),(p+1,3)) \} \\ &\cup \{ ((1,1),(2s,2p)) \mid s=1,\ldots,p-1 \} \cup \{ ((1,p),(2,p+1)) \} \\ &\cup \{ ((1,1),(2p,2s)) \mid s=1,\ldots,p-1 \} \cup \{ ((p,1),(p+1,2)) \}. \end{split}$$

Those vectors are

And, projecting them onto the space $span\{e_k \mid k \in \mathcal{K}^{j+2} \setminus \mathcal{K}^{j+1}\}$ we arrive to the family $\prod_{j+1} F_{S_{j+1}}$ which elements are

$$\begin{split} \Pi_{j+1}\delta_{(1,2p-1),(2p-1,1)} = C_{(1,2p-1),(2p-1,1)}^{++}e_{(2p,2p)} \\ \Pi_{j+1}\delta_{(1,1),(2z-1,2p)} = C_{(1,1),(2z-1,2p)}^{-+}e_{(2(z-1),2p+1)} + C_{(1,1),(2z-1,2p)}^{++}e_{(2z,2p+1)} \\ z = 2, \dots, p & (3.5.12) \\ \Pi_{j+1}\delta_{(1,p)(3,p+1)} = C_{(1,p)(3,p+1)}^{+-}e_{(2,2p+1)} + C_{(1,p)(3,p+1)}^{++}e_{(4,2p+1)} \\ \Pi_{j+1}\delta_{(1,1),(2p,2z-1)} = C_{(1,1),(2p,2z-1)}^{+-}e_{(2p+1,2(z-1))} + C_{(1,1),(2p,2z-1)}^{++}e_{(2p+1,2z)} \\ z = 2, \dots, p & (3.5.13) \\ \Pi_{j+1}\delta_{(p,1),(p+1,3)} = C_{(p,1),(p+1,3)}^{+-}e_{(2p+1,2)} + C_{(p,1),(p+1,3)}^{++}e_{(2p+1,4)} \\ \Pi_{j+1}\delta_{(1,1),(2s,2p)} = C_{(1,1),(2s,2p)}^{-+}e_{(2s-1,2p+1)} + C_{(1,1),(2s,2p)}^{++}e_{(2s+1,2p+1)} \\ s = 1, \dots, p-1 & (3.5.14) \\ \Pi_{j+1}\delta_{(1,1),(2p,2s)} = C_{(1,1),(2p,2s)}^{-+-}e_{(2p+1,2s-1)} + C_{(1,1),(2p,2s)}^{++}e_{(2p+1,2s+1)} \\ s = 1, \dots, p-1 & (3.5.15) \\ \Pi_{j+1}\delta_{(p,1),(p+1,2)} = C_{(p,1),(p+1,2)}^{+--}e_{(2p+1,1)} + C_{(p,1),(p+1,2)}^{++-}e_{(2p+1,3)}; & (3.5.16) \\ \end{split}$$

No one of the coefficients appearing in these expressions vanishes because all pairs (m, n) satisfy $m < n \land (m_1 = n_1 \lor n_2 \ge m_2)$ and because no one of the

following expressions vanish

$$\begin{split} (1,2p-1)\wedge(2p-1,1) &= 1-(2p-1)^2;\\ (1,1)\wedge(2z-1,2p) &= 1+2(p-z), \quad z=2,\,\ldots,\,p;\\ (1,p)\wedge(3,p+1) &= 1-2p;\\ (1,1)\wedge(2p,2z-1) &= -1-2(p-z), \quad z=2,\,\ldots,\,p;\\ (p,1)\wedge(p+1,3) &= 2p-1;\\ (1,1)\wedge(2s,2p) &= 2(p-s), \quad s=1,\,\ldots,\,p-1;\\ (1,p)\wedge(2,p+1) &= 1-p;\\ (1,1)\wedge(2p,2s) &= 2(s-p), \quad s=1,\,\ldots,\,p-1\\ (p,1)\wedge(p+1,2) &= p-1. \end{split}$$

Hence we can see that these vectors are linearly independent. To see this is enough to see that:

- The vectors $\Pi_{j+1}\delta_{(1,1),(3,2p)}$, (z = 2 in (3.5.12)) and $\Pi_{j+1}\delta_{(1,p)(3,p+1)}$ are linearly independent;
- The vectors $\Pi_{j+1}\delta_{(1,1),(2p,3)}$, (z = 2 in (3.5.13)) and $\Pi_{j+1}\delta_{(p,1),(p+1,3)}$ are linearly independent;
- The vectors $\Pi_{j+1}\delta_{(1,1),(2,2p)}$, (s = 1 in (3.5.14)) and $\Pi_{j+1}\delta_{(1,p)(2,p+1)}$ are linearly independent;
- The vectors $\Pi_{j+1}\delta_{(1,1),(2p,2)}$, (s = 1 in (3.5.15)) and $\Pi_{j+1}\delta_{(p,1),(p+1,2)}$ are linearly independent;

But that comes from

$$\frac{C_{(1,1),(3,2p)}^{-+}}{C_{(1,p)(3,p+1)}^{-+}} = \frac{(3+2p)(8b^2+a^2(4p^2-1))}{(1+4p)(8b^2+a^2(2p+1))} \\
\neq -\frac{(2p-3)(8b^2+a^2(4p^2-1))}{(2p-1)(8b^2+a^2(2p+1))} = \frac{C_{(1,1),(3,2p)}^{++}}{C_{(1,p)(3,p+1)}^{++}}; \\
\frac{C_{(1,1),(2p,3)}^{+-}}{C_{(p,1),(p+1,3)}^{+-}} = \frac{(3+2p)(8a^2+b^2(4p^2-1))}{(1+4p)(8a^2+b^2(2p+1))} \\
\neq -\frac{(2p-3)(8a^2+b^2(4p^2-1))}{(2p-1)(8a^2+b^2(2p+1))} = \frac{C_{(1,1),(2p,3)}^{++}}{C_{(p,1)(p+1,3)}^{++}};$$

$$\frac{C_{(1,1),(2,2p)}^{-+}}{C_{(1,p)(2,p+1)}^{-+}} = \frac{2(1+p)(3b^2+a^2(4p^2-1))}{(3p+1)(3b^2+a^2(2p+1))}$$

$$\neq -\frac{2(3b^2+a^2(4p^2-1))}{3b^2+a^2(2p+1)} = \frac{C_{(1,1),(2,2p)}^{++}}{C_{(1,p)(2,p+1)}^{++}};$$

and

$$\frac{C_{(1,1),(2p,2)}^{+-}}{C_{(p,1),(p+1,2)}^{+-}} = \frac{2(1+p)(3a^2+b^2(4p^2-1))}{(3p+1)(3a^2+b^2(2p+1))}$$
$$\neq -\frac{2(3a^2+b^2(4p^2-1))}{3b^2+a^2(2p+1)} = \frac{C_{(1,1),(2p,2)}^{++}}{C_{(p,1)(p+1,2)}^{++}}.$$

By Lemma 3.1.1 the $(2p+1)^2 - 1$ of $F_{S_{j+1}} \cup \{e_k \mid k \in \mathcal{K}^{j+1}\}$ are linearly independent and then they span

$$span\{e_k \mid 1 \le k_1, k_2 \le 2p+1\} \setminus \{(2p+1, 2p+1)\} = span\{e_k \mid k \in \mathcal{K}^{j+2}\}$$

Again, since with $\delta_{m,n}$ also $\lambda \delta_{m,n}$ belongs to $\{B(Y, Y) \mid Y \in H^j\}$ for every $\lambda \in \mathbb{R}$, we can select the linear space \tilde{H}^{j+1} defined by

$$\tilde{H}^{j+1} := \left(H^j + Conv\{\lambda \delta_{m,n} \mid (m, n) \in S_{j+1}, \lambda \in \mathbb{R} \} \right)$$
$$\cap \left(H^j - Conv\{\lambda \delta_{m,n} \mid (m, n) \in S_{j+1}, \lambda \in \mathbb{R} \} \right)$$

from the space

$$\Big(H^j + Conv\{B(Y, Y) \mid Y \in H^j\}\Big) \cap \Big(H^j - Conv\{B(Y, Y) \mid Y \in H^j\}\Big).$$

Since $Conv\{\lambda\delta_{m,n} \mid (m, n) \in S_{j+1}, \lambda \in \mathbb{R}\}\)$ coincides with $span\{\lambda\delta_{m,n} \mid (m, n) \in S_{j+1}, \lambda \in \mathbb{R}\}\)$, we have that

$$\tilde{H}^{j+1} = span(\mathcal{K}^{j+1} \cup \{\delta_{m,n} \mid (m, n) \in S_{j+1}\}) = span(\mathcal{K}^{j+2}) = H^{j+1}.$$

Remark 8. Starting with $span(\mathcal{K}^2)$, by Proposition 3.5.3, we can obtain the directions on $span(\mathcal{K}^j)$, $(j \ge 2)$ iterating FCE procedures. Since that Proposition is valid for any rectangle (including the square) we can say that \mathcal{K}^2 is G-Saturating for all rectangles.

Remark 9. We can show that \mathcal{K}^1 is saturating for the square too and so complete the proof of Proposition 3.5.3. The only step that is not clear in the square, is how to arrive to $\operatorname{span}(\mathcal{K}^2)$ (or to a superspace of its) from $\operatorname{span}(\mathcal{K}^1)$. In the case of the square the extracted family in the end of proof of Proposition 3.5.2 (Case j even: j = 0 in our present case) does not lead to a family of linearly independent vectors when projected in $\operatorname{span}\{e_k \mid k \notin \mathcal{K}^1\}$ because $\Pi\delta_{(1,2)2,1)} = 0$. Then we proceed as follows: First we select the family $F_{S_1} \setminus \{\delta_{(1,2),(2,1)}\}$. Projecting this family onto $\operatorname{span}(\mathcal{K}^2 \setminus (\{(3,3)\} \cup \mathcal{K}^1))$ we obtain a family of linearly independent vectors. So "adding" the vectors in $\{e_k \mid k \in \mathcal{K}^1\}$ to those F_{S_1} we obtain a family of $4^2 - 1 - 1$ linearly independent vectors spanning the space $span\{e_k \mid k \in \mathcal{K}^2 \setminus \{(3,3)\}\}$. So proceeding as before we reach the directions $span(\mathcal{K}^2 \setminus \{(3,3)\})$. Then we select the family

$$\begin{split} \delta_{(1,1),(2,4)} &= \frac{3\pi^2(5a^2+b^2)}{2ab(9a^2+b^2)}e_{(1,3)} - \frac{9\pi^2(5a^2+b^2)}{2ab(25a^2-b^2)}e_{(1,5)} \\ &+ \frac{\pi^2(5a^2+b^2)}{2ab(a^2+b^2)}e_{(3,3)} - \frac{3\pi^2(5a^2+b^2)}{2ab(25a^2+9b^2)}e_{(3,5)} \\ \delta_{(1,2),(2,3)} &= -\frac{\pi^2(5a^2+3b^2)}{4ab(a^2+b^2)}e_{(1,1)} - \frac{7\pi^2(5a^2+3b^2)}{4ab(25a^2+b^2)}e_{(1,5)} \\ &+ \frac{7\pi^2(5a^2+3b^2)}{4ab(a^2+9b^2)}e_{(3,1)} + \frac{\pi^2(5a^2+3b^2)}{4ab(25a^2+9b^2)}e_{(3,5)} \\ \delta_{(1,4),(2,1)} &= \frac{21\pi^2(b^2-5a^2)}{4ab(9a^2+b^2)}e_{(1,3)} + \frac{27\pi^2(5a^2-b^2)}{4ab(25a^2+b^2)}e_{(1,5)} \\ &+ \frac{3\pi^2(5a^2-b^2)}{4ab(a^2+b^2)}e_{(3,3)} + \frac{21\pi^2(b^2-5a^2)}{4ab(25a^2+9b^2)}e_{(3,5)}. \end{split}$$

Projecting in span $\{e_k \mid k \in \mathcal{K}^3 \setminus (\mathcal{K}^2 \setminus \{(3,3)\})\}$ we obtain $0\pi^2(5a^2 + b^2)$

$$\begin{split} \Pi \delta_{(1,1),(2,4)} &= -\frac{9\pi^2 (5a^2 + b^2)}{2ab(25a^2 - b^2)} e_{(1,5)} \\ &+ \frac{\pi^2 (5a^2 + b^2)}{2ab(a^2 + b^2)} e_{(3,3)} - \frac{3\pi^2 (5a^2 + b^2)}{2ab(25a^2 + 9b^2)} e_{(3,5)} \\ \Pi \delta_{(1,2),(2,3)} &= -\frac{7\pi^2 (5a^2 + 3b^2)}{4ab(25a^2 + b^2)} e_{(1,5)} + \frac{\pi^2 (5a^2 + 3b^2)}{4ab(25a^2 + 9b^2)} e_{(3,5)} \\ \Pi \delta_{(1,4),(2,1)} &= \frac{27\pi^2 (5a^2 - b^2)}{4ab(25a^2 + b^2)} e_{(1,5)} \\ &+ \frac{3\pi^2 (5a^2 - b^2)}{4ab(a^2 + b^2)} e_{(3,3)} + \frac{21\pi^2 (b^2 - 5a^2)}{4ab(25a^2 + 9b^2)} e_{(3,5)}. \end{split}$$

Since we are working in the square a = b and no one of the coefficients appearing in the last expressions vanish. We compute

$$Det \begin{pmatrix} C_{(1,1),(2,4)}^{+-} & C_{(1,1),(2,4)}^{+-} & C_{(1,1),(2,4)}^{++} \\ C_{(1,2),(2,3)}^{-+} & 0 & C_{(1,2),(2,3)}^{++} \\ C_{(1,4),(2,1)}^{+-} & C_{(1,4),(2,1)}^{+-} & C_{(1,4),(2,1)}^{+-} \end{pmatrix}$$
$$= -\frac{15\pi^2 (125a^6 + 75a^4b^2 - 5a^2b^4 - 32b^6)}{16a^3b^3(a^2 + b^2)(25a^2 + b^2)(25a^2 + 9b^2)}$$
$$(since \ a = b)$$
$$= -\frac{2880a^6}{28288a^{12}} = -\frac{45}{442a^6} \neq 0.$$

Hence the vectors are linearly independent. If we joint to the three $\delta_{m,n}$ vectors the vectors in $\{e_k \mid k \in \mathcal{K}^2 \setminus \{(3,3)\}\}$ we obtain a family of $4^2 + 1$ linearly independent vectors spanning the space $\{e_k \mid k \in \mathcal{K}^2 \cup \{(1,5), (3,3), (3,5)\}\}$. We arrive in this way to $\operatorname{span}(\mathcal{K}^2 \cup \{(1,5), (3,3), (3,5)\}) \supset \operatorname{span}(\mathcal{K}^2)$. Since \mathcal{K}^2 is G-Saturating for the square so is $\mathcal{K}^2 \cup \{(1,5), (3,3), (3,5)\}$. Hence \mathcal{K}^1 is saturating for the square.

3.6 Exact Controllability of Galerkin Approximations.

Write the \mathcal{K}^N -Galerkin approximation of the NSE system (3.4.2), with \mathcal{K}^1 as set of excited modes, in the concise form

$$N: \begin{cases} \dot{u}_k = \mathcal{B}_k(u) + \nu \Delta u_k + F_k + v_k & k \in \mathcal{K}^1 \\ \dot{u}_k = \mathcal{B}_k(u) + \nu \Delta u_k + F_k & k \in \mathcal{K}^N \setminus \mathcal{K}^1 \\ u \in \mathbb{R}^{\kappa_N}. \end{cases}$$
(3.6.1)

In [16] E. Weinam and J. Mattingly proved the Full Lie Rank Property for the 2D NSE with periodic conditions and for some class of few low modes controls. Similarly we prove that our equation also is "full Lie rank", i.e., Lie brackets at each point span the ambient space \mathbb{R}^{κ_N} .

Before we have proved that for all $N \in \mathbb{N}_0$ and all t > 0 the system [(3.6.1).N] is time-t approximately controllable:

$$\forall u \in \mathbb{R}^{\kappa_N} \quad \overline{\mathcal{A}_u(\mathcal{F}_N)(t)} = \mathbb{R}^{\kappa_N}$$

where \mathcal{F}_N is the family of vector fields of system [(3.6.1).N], i.e.,

$$\mathcal{F}_N = \{ \mathcal{B}(\cdot) + \nu \Delta(\cdot) + F^{\kappa_N} + v \mid v \in \mathbb{R}^{\kappa_1} \}.$$

Next we prove the (exact) controllability of system [(3.6.1).N]. For that we need to compute some Lie brackets.

3.6.1 Lie Brackets. Full Lie Rank Property.

From \mathcal{F}_N , we set the vector fields

$$V^{0} := \mathcal{B} + \nu \Delta + F^{\kappa_{N}}, \quad X^{i} := V^{0} + \frac{\partial}{\partial u_{i}}$$

where F^{κ_N} is the projection of F onto \mathbb{R}_N^{κ} and $\kappa_N = \#\mathcal{K}^N$ and, compute

$$V^i := [X^i, V^0] = [\frac{\partial}{\partial u_i}, V^0] = \frac{\partial V^0}{\partial u} \frac{\partial}{\partial u_i} = \frac{\partial V^0}{\partial u_i},$$

 \mathbf{SO}

$$V_k^i := \left(\sum_{\substack{k=(n++i)^+\\i< n}} u_n C_{i,n}^{++} + \sum_{\substack{k=(n+-i)^+\\i< n}} u_n C_{i,n}^{+-} + \sum_{\substack{k=(n--i)^+\\i< n}} u_n C_{i,n}^{--} \right) + \delta^{[k,i]} \nu \bar{k},$$

where $\delta^{[k,i]}$ is the Kronecker delta function: $\delta^{[k,i]} = \begin{cases} 1 & \text{if } k=i \\ 0 & \text{if } k \neq i \end{cases}$.

$$V^{j,i} := [X^j, V^i] = [X^j, [X^i, V^0]] = \frac{\partial V^i}{\partial u_j},$$

 $\mathrm{so},$

$$V^{j,i} = \gamma_{i,j}, \quad j > i.$$

At each given point $\bar{u} \in \mathbb{R}^{\kappa_N}$ the family of brackets¹⁴

$$\mathcal{H}^{1} := \{ V^{0} \pm \frac{1}{2} \frac{\partial}{\partial u_{n}}, \, \alpha \gamma_{i,j} \mid n \in \mathbb{R}^{\kappa_{1}}, \, (i, j) \in S_{1} \}$$

span a superspace of \mathbb{R}^{κ_2} . Indeed, $\frac{\partial}{\partial u_n} = \left(V^0 + \frac{1}{2}\frac{\partial}{\partial u_n}\right) - \left(V^0 - \frac{1}{2}\frac{\partial}{\partial u_n}\right)$.¹⁵ Now we have a technical difficulty, the vector fields have coefficients with

Now we have a technical difficulty, the vector fields have coefficients with not so nice expressions and, if we compute Lie brackets envolving them we will obtain even more complicated expressions. To avoid these expressions we prove by (finite) induction on $i \in \{1, ..., \kappa_N\}$ that each constant vector field $\frac{\partial}{\partial u_n}$, $n \in \mathcal{K}^i$ is a linear combination of brackets:

- For i = 1 we take the family \mathcal{H}^1 .
- Inductive Step: The induction hypothesis is:

"There is a family of brackets $\mathcal{H}^{p-1} = \{W^j \mid j = 1, \ldots, M_{p-1}\}$ such that every constant vector field $\frac{\partial}{\partial u_i}$, $i \in \mathcal{K}^{p-1}$ can be written as a linear combination of its elements:

$$\frac{\partial}{\partial u_i} = \sum_{j=1}^{M_{p-1}} \alpha_i^j W^j, \quad \alpha_i^j \in \mathbb{R};$$

Then for all $i \in \mathcal{K}^{p-1}$

$$V^{i} := \left[\frac{\partial}{\partial u_{i}}, V^{0}\right] \in span\{[W_{j}, V^{0}] \mid j = 1, \dots, M_{p-1}\}$$

and, for each $i, n \in \mathcal{K}^{p-1}$:

$$V^{n,i} := \left[\frac{\partial}{\partial u_n}, V^i\right] = \gamma_{i,n} \in span\{[W^k, [W_j, V^0]] \mid k, j = 1, \dots, M_{p-1}\}.$$

Since the vectors in $\{\frac{\partial}{\partial u_k}, \gamma_{i,n} \mid k \in \mathcal{K}^{p-1}, (i, n) \in S_{p-1}\}$ span \mathbb{R}^{κ_p} and can be written as a linear combination of brackets, then also the vector fields $\frac{\partial}{\partial u_i}$, $i \in \mathcal{K}^p \setminus \mathcal{K}^{p-1}$ are linear combinations of brackets. The wanted family is $\mathcal{H}^{p-1} \cup \{[W^k, [W_j, V^0]] \mid k, j = 1, \dots, M_{p-1}\}.$

Therefore, for all $N \in \mathbb{N}_0$, system [(3.6.1).N] is a full-rank bracket generating system. From that and from its approximate controllability¹⁶ we conclude its controllability. Unfortunately for fixed time the bracket generating property is not sufficient to conclude controllability from approximate controllability. To achieve controllability at time t we shall need some lemmas which proofs can be found in [8].

¹⁴Including the elements of \mathcal{F}_N we consider brackets of "length" 0.

¹⁵Recall that S_1 has been chosen so that $\{e_n, \delta_{i,j} \mid n \in \mathcal{K}^1, (i, j) \in S_1\}$ span \mathbb{R}^{κ_2} .

¹⁶Approximate controllability at time t implies, trivialy approximate controllability.

3.6.2 Zero Orbits and Zero Ideal.

Definition 3.6.1. A zero-time orbit N_{0u} through u of a family of vector fields \mathcal{F} is the set

$$N_{0u} := \{ u \circ e^{t_1 V_1} \circ \ldots \circ e^{t_p V_p} \mid p \in \mathbb{N}_0, V_i \in \mathcal{F}, t_i \in \mathbb{R}, \sum_{i=1}^p t_i = 0 \}$$

Definition 3.6.2. The **derived algebra** of \mathcal{F} , denoted $\mathcal{D}_{er}(\mathcal{F})$, is the set of all linear combinations of iterated brackets¹⁷

The zero-time ideal, denoted $\mathcal{I}(\mathcal{F})$, is the span of elements in $\mathcal{D}_{er}(\mathcal{F})$ and differences of the form X - Y with X and Y in \mathcal{F} .

Lemma 3.6.1. Let \mathcal{F} be any family of analytic vector fields on an analytic manifold M. Let N be an orbit of \mathcal{F} and, N_0 be a zero orbit of \mathcal{F} contained in N. Then we have the following:

- Each connected component of N_0 is an orbit of $\mathcal{I}(\mathcal{F})$;
- For each u ∈ N₀, the tangent space of N₀ at u is equal to the evaluation of I(F) at u;
- The dimension of \$\mathcal{I}_u(\mathcal{F})\$ is constant as u varies on N. It is equal either to dim(Lie_u(\mathcal{F})) 1 or to dim(Lie_u(\mathcal{F}));
- $dim(Lie_u(\mathcal{F})) = dim(\mathcal{I}_u(\mathcal{F}))$ iff $X(u) \in \mathcal{I}_u(\mathcal{F})$ for some $X \in \mathcal{F}$.

Lemma 3.6.2. Suppose that \mathcal{F} is a family of vector fields on M such that both \mathcal{F} and its zero-time ideal $\mathcal{I}(\mathcal{F})$ are Lie-determined (the evaluation of Lie brackets at each point span the tangent space to the orbit). In addiction, assume that \mathcal{F} contains a complete vector field. Then

- $\mathcal{A}_u(\mathcal{F})(t)$ is a connected subset of some zero orbit N_{0z} through some element $z \in M$.
- $\mathcal{A}_u(\mathcal{F})(t)$ has a nonempty interior in the manifold topology of the zeroorbit where it is contained. Moreover, the set of interior points is dense in $\mathcal{A}_u(\mathcal{F})(t)$.

Coming back to our system [(3.6.1).N], by Lemma 3.6.1 and due to the fact that $V^0(0) = 0 \in \mathcal{I}(\mathcal{F})$, we have

$$\dim(Lie_0(\mathcal{F}_N)) = \dim(\mathcal{I}_0(\mathcal{F}_N)) = \kappa_N;$$

which means that the zero-time orbit N_0 through 0 has dimension κ_N and, since that dimension is constant in all points in the unique orbit \mathbb{R}^{κ_N} of the system, we conclude that N_0 is a union of connected components of dimension κ_N . Since the dimension of that components is κ_N their topology coincide with that of \mathbb{R}^{κ_N} and, from the fact that the zero-time orbits form a partition of \mathbb{R}^{κ_N} we conclude that \mathbb{R}^{κ_N} is a union of connected open sets. Therefore there is only one zero-orbit, it is the whole state space \mathbb{R}^{κ_N} .

¹⁷Brackets of "length" ≥ 1 , considering the elements of \mathcal{F} brackets of length 0.

By Lemma 3.6.2, and by the fact that V^0 is a complete vector field which follows from the estimate $|u(s)| \leq |u(0)| + \frac{s}{\nu} ||F||_{V'}^2$ (see estimate (2.6.8) with $\tilde{F} = F$), the interior $int \mathcal{A}_u(\mathcal{F}_N)(t)$ of the attainable set from u at time t is dense in $\mathcal{A}_u(\mathcal{F})(t)$, where the interior and density are relative to the topology of \mathbb{R}^{κ_N} because that is the topology of the zero-orbit. Hence we arrive to the equality

$$\overline{int\mathcal{A}_u(\mathcal{F}_N)(t)} = \overline{\mathcal{A}_u(\mathcal{F}_N)(t)} = \mathbb{R}^{\kappa_N}$$

for all t > 0.

Now we can prove the controllability at time t of system [(3.6.1).N]: Let u, z be two elements in \mathbb{R}^{κ_N} . Since the intersection of two open dense sets stills open and dense, we may take a point

$$w \in int\mathcal{A}_u(\mathcal{F}_N)(t/2) \cap int\mathcal{A}_z(-F_N)(t/2).$$

[Note that the family $-\mathcal{F}_N := \{-V \mid V \in \mathcal{F}\}$ satisfies the requirements of lemmas 3.6.1 and 3.6.2.]

Then we can write

$$w = u \circ e^{t_1 V_1} \circ \dots \circ e^{t_n V_n}, \qquad V_i \in \mathcal{F}_N, \, t_i \ge 0, \, \sum_{i=1}^n t_i = \frac{t}{2};$$
$$w = z \circ e^{-s_1 W_1} \circ \dots \circ e^{-s_m W_m}, \qquad W_i \in \mathcal{F}_N, \, s_i \ge 0, \, \sum_{i=1}^m t_i = \frac{t}{2};$$

So, z is reachable from u in time t:

$$z = u \circ e^{t_1 V_1} \circ \cdots \circ e^{t_n V_n} \circ e^{s_m W_m} \circ \cdots \circ e^{s_1 W_1}$$

72CHAPTER 3. CONTROLLABILITY OF GALERKIN APPROXIMATIONS.
Chapter 4

Controllability in Observed Component.

4.1 Controlled N-S Problem. Existence, Uniqueness and Continuity.

In chapter 2 we have presented the classical, weak and strong formulations of (2.5.1)-(2.5.5). So for the controlled version with F + v in the place of \tilde{F} we have existence, unicity and continuity in the data $(u_0, F + v, \nu)$, because our control is an essencially bounded function, so that F+v will belong where \tilde{F} did: $(L^2(0, T, V') \text{ or } L^2(0, T, H))$. If we consider the initial data as (u_0, \tilde{F}, v, ν) , by Theorems 2.6.4, 2.8.1 and 2.8.2 we easily conclude that (considering again the external force depending on time):

Corollary 4.1.1. Given

 $(u_0, \tilde{F}, v, \nu) \in H \times L^2(0, T, V') \times L^\infty(0, T, V') \times]0, +\infty[,$

there is at least one weak solution $u \in C([0, T], H)$ for Problem 2.5.1 with $\tilde{F} + v$ in the place of \tilde{F} .

and,

Corollary 4.1.2. The maps

$$\mathbb{S}: H \times L^2(0, T, V') \times L^\infty(0, T, V') \times]0, +\infty[\to C([0, T], H)$$
$$(u_0, \tilde{F}, v, \nu) \mapsto u$$

and

$$\mathbb{S}_{2}: H \times L^{2}(0, T, V') \times L^{\infty}(0, T, V') \times]0, +\infty[\to L^{2}(0, T, V) \\ (u_{0}, \tilde{F}, v, \nu) \mapsto u$$

 $are\ continuous.$

Similarly, by Theorems 2.11.1, 2.13.1 and 2.13.2 we can easily see that

Corollary 4.1.3. Given

$$(u_0, \tilde{F}, v, \nu) \in V \times L^2(0, T, H) \times L^\infty(0, T, H) \times]0, +\infty[,$$

there is at least one weak solution $u \in C([0, T], V)$ for Problem 2.10.1 with $\tilde{F} + v$ in the place of \tilde{F} .

and,

Corollary 4.1.4. The maps

$$\mathbb{S}_s: V \times L^2(0, T, V') \times L^\infty(0, T, H) \times]0, +\infty[\to C([0, T], V)$$
$$(u_0, \tilde{F}, v, \nu) \mapsto u$$

and

$$\mathbb{S}_{2s}: V \times L^{2}(0, T, V') \times L^{\infty}(0, T, H) \times]0, +\infty[\to L^{2}(0, T, D(A))$$
$$(u_{0}, \tilde{F}, v, \nu) \mapsto u$$

are continuous.

4.2 Change of Variables.

If we make the change of variables

$$u = y + \mathbb{I}v$$

where \mathbb{I} is the primitive operator — $[\mathbb{I}v](t) = \int_0^t v(\tau) d\tau$, from

$$u' = -\nu Au - Bu + F + v$$

we arrive to the equation

$$y' = -\nu A(y + \mathbb{I}v) - B(y + \mathbb{I}v) + \tilde{F}.$$

Note that the function v appears only implicitly in the last equation. Now we forget that $\mathbb{I}v$ is a primitive of an essentially bounded function and replace it by P in the equation. Since v is a low modes forcing it takes value in a finite-dimensional space and, $\mathbb{I}v$ being a primitive we have $\mathbb{I}v \in C([0, T], D(A))$. But we take P in the larger space $L^4(0, T, D(A))$.

4.3 Weak Case.

We want to study the following equivalent problems 4.3.1 and 4.3.2.

Problem 4.3.1. Given

$$\tilde{F} \in L^2(0, T, V'), \quad P \in L^4(0, T, D(A))$$

&
(4.3.1)

$$y_0 \in H \tag{4.3.2}$$

$$y \in L^2(0, T, V) \tag{4.3.3}$$

satisfying (in the distribution sense)

$$\forall v \in V : \\ \frac{d}{dt}(y, v) + \nu((y + P, v)) + b(y + P, y + P, v) = <\tilde{F}, v >,$$
 (4.3.4) and

$$y(0) = y_0. (4.3.5)$$

Problem 4.3.2. Given

$$\tilde{F} \in L^2(0, T, V'), \quad P \in L^4(0, T, D(A))$$
&
(4.3.6)

$$y_0 \in H,$$
 (4.3.7)
to find

$$y \in L^2(0, T, V), \qquad y' \in L^1(0, T, V')$$
 (4.3.8)
satisfying

$$y' + \nu A(y+P) + B(y+P) = \tilde{F}$$
 on]0, T[, (4.3.9)

and

$$y(0) = y_0. (4.3.10)$$

Remark 10. The equivalence of these problems can be shown the way we proved the equivalence of problems 2.5.1 and 2.5.2 (see chapter 2). This equivalence follows from

$$\begin{split} \bullet \ &\tilde{F} \in L^2(0, \, T, \, V'); \\ \bullet \ & A(y+P) \in L^2(0, \, T, \, V'). \\ & Indeed \ \|A(y+P)\|_{V'} \leq \|y+P\| \ and \ y+P \in L^2(0, \, T, \, V); \\ \bullet \ & B(y+P) \in L^1(0, \, T, \, V'). \\ & Indeed \ \|B(y+P)\|_{V'} \leq C\|y+P\|^2 \ and \ y+P \in L^2(0, \, T, \, V) \end{split}$$

So that $\tilde{F} - \nu A(y+P) - B(y+P) \in L^1(0, T, V')$.

4.4 Existence.

We have the Theorem

Theorem 4.4.1. Given \tilde{F} , P and y_0 satisfying (4.3.6) and (4.3.7). There is at least one function y satisfying (4.3.8)-(4.3.10).

The proof is analogous to that of Theorem 2.6.4. Basically it differs only in some estimates we compute now: Following the proof of Theorem 2.6.4 presented in chapter 2, with the suitable adaptations to problems 4.3.1-4.3.2, we define an approximate solution

$$y^m = \sum_{\max\{i_1, i_2\} \le m} y_i^m(t) W_i$$

for each $m \in \mathbb{N}_0$ and arrive to the equation

$$<(y^{m})', y^{m} > +\nu < A(y^{m} + P^{m}), y^{m} >$$

+ $< B(y^{m} + P^{m}), y^{m} > = <\tilde{F}, y^{m} > .$ ¹ (4.4.1)

We note that

• $< A(y^m + P^m), y^m >= ||y^m||^2 + ((P^m, y^m))$ • $b(y^m + P^m, y^m + P^m, y^m)$ $= b(y^m + P^m, y^m + P^m, y^m + P^m) - b(y^m + P^m, y^m + P^m, P^m)$ $= -b(y^m + P^m, y^m, P^m) - b(y^m + P^m, P^m, P^m)$ $= -b(y^m + P^m, y^m, P^m) = -b(y^m, y^m, P^m) - b(P^m, y^m, P^m).$

Hence from (4.4.1) we have

$$\frac{d}{dt}|y^{m}|^{2} + 2\nu||y^{m}||^{2} = -2\nu((P^{m}, y^{m})) -b(y^{m}, P^{m}, y^{m}) + b(P^{m}, y^{m}, P^{m}) + \langle \tilde{F}, y^{m} \rangle;$$

hence

$$\frac{d}{dt}|y^{m}|^{2} + 2\nu||y^{m}||^{2} \le 2\nu||P||||y^{m}|| + C|y^{m}|||y^{m}|||P|| + C||P||^{2}||y^{m}|| + ||\tilde{F}||_{V'}||y^{m}|| = C||P||^{2}||y^{m}|| + ||\tilde{F}||_{V'}||y^{m}|| = C||P||^{2}||y^{m}|| + C||P||^{2}||y^{m}|| + ||\tilde{F}||_{V'}||y^{m}|| = C||P||^{2}||y^{m}|| + C||P||^{2}||y^{m}||||||^{2}||y^{m}|||y^{m}||||||^{2}||y^{m}||||||^{2}||y^{m}||||||$$

thus

$$\frac{d}{dt}|y^{m}|^{2} + \nu \|y^{m}\|^{2} \le 4\nu \|P\|^{2} + \frac{C^{2}}{\nu}|y^{m}|^{2}\|P\|^{2} + \frac{C^{2}}{\nu}\|P\|^{4} + \frac{1}{\nu}\|\tilde{F}\|_{V'}^{2}.$$
 (4.4.2)

From equation (4.4.2) and from Gronwall Inequality we can derive the estimates (4.4.3) and (4.4.5) below:

$$|y^{m}(s)|^{2} \leq \exp \int_{0}^{T} \frac{C^{2}}{\nu} ||P(t)||^{2} dt \left(|y_{0}|^{2} + \int_{0}^{T} 4\left(\nu ||P(t)||^{2} + \frac{C^{2}}{\nu} ||P(t)||^{4} + \frac{1}{\nu} ||\tilde{F}(t)||^{2}\right) dt\right).$$
(4.4.3)

Since $P \in L^4(0, T, D(A)) \subset L^4(0, T, V) \subset L^2(0, T, V)$ and $\tilde{F} \in L^2(0, T, V)$, (4.4.3) shows that

$$y^m$$
 remains in a bounded set of $L^{\infty}(0, T, H)$. (4.4.4)

¹Where P^m is the projection of P onto $span\{W_i \mid \max\{i_1, i_2\} \le m\}$.

From

$$|y^{m}(T)|^{2} - |y^{m}(0)|^{2} + \nu \int_{0}^{T} ||y^{m}(t)||^{2} dt$$

$$\leq \int_{0}^{T} \left(4\nu ||P(t)||^{2} + \frac{C^{2}}{\nu} ||P(t)||^{4} + \frac{1}{\nu} ||\tilde{F}(t)||^{2}\right) dt + \int_{0}^{T} \frac{C^{2}}{\nu} |y^{m}(t)|^{2} ||P(t)||^{2} dt.$$
(4.4.5)

By (4.4.4) the last integral is finite. Then (4.4.5) shows that

$$y^m$$
 remains in a bounded set of $L^2(0, T, V)$. (4.4.6)

The rest of the proof is completely analogous.

Remark 11. As we can see in the estimates (4.4.3) and (4.4.5), considering $P \in L^4(0, T, V)$ is sufficient to guarantee existence of weak solutions.

4.5 Uniqueness.

Theorem 4.5.1. The solution of Problems 4.3.1-4.3.2 given by Theorem 4.4.1 is unique. Moreover it is a.e. equal to a continuous function from [0, T] into H.

The continuity follows from

•
$$A(y+P) \in L^2(0, T, V')$$

• $\tilde{F} \in L^2(0, T, V')$

• $B(y+P) \in L^2(0, T, V').$

Indeed these expressions imply

$$y \in L^2(0, T, V) \& y' \in L^2(0, T, V').$$

The first two expressions we already know to be true. The last one follows from

$$\begin{split} \|B(y+P)\| &\leq C|y| \|P\| + C\|P\|^2; \\ P &\in L^4(0,\,T,\,D(A)) \subset L^4(0,\,T,\,V); \quad y \in L^\infty(0,\,T,\,H). \end{split}$$

To conclude the uniqueness we consider two solutions y and z of problems (4.3.1)-(4.3.2). The difference w := y - z satisfies

$$w' = -\nu Aw - B(y+P) + B(z+P)$$

from which we derive

$$\begin{aligned} \frac{d}{dt} |w|^2 + 2\nu ||w||^2 &\leq 2C |w| ||w|| ||z + P|| \\ \frac{d}{dt} |w|^2 &\leq \frac{C^2}{2\nu} |w|^2 ||z + P||^2 \\ |w(s)|^2 &\leq |w(0)|^2 \exp \int_0^T \frac{C^2}{2\nu} ||z(t) + P(t)||^2 \, dt = 0. \end{aligned}$$

4.6 Continuity.

Theorem 4.6.1. The map

is continuous. Where y is the unique solution of problems (4.3.1)-(4.3.2) corresponding to the data (y_0, \tilde{F}, P, ν) .

Proof. The proof is analogous to that of Theorem 4.1.2. We fix a quadruple

 $(y_0, \tilde{F}, P, \nu) \in H \times L^2(0, T, V') \times L^4(0, T, D(A)) \times]0, +\infty[$

and $\varepsilon > 0$. Then consider another quadruple

$$(z_0, G, Q, \eta) \in H \times L^2(0, T, V') \times L^4(0, T, D(A)) \times [0, +\infty[$$

Put

$$y := \mathbb{Y}(y_0, \tilde{F}, P, \nu) \& z := \mathbb{Y}(z_0, G, Q, \eta)$$

so that

$$y' + \nu A(y+P) + B(y+P) = \tilde{F} \& z' + \eta A(z+Q) + B(z+Q) = G.$$

Putting w := z - y we obtain

$$w' = G - \tilde{F} - \eta A w - \eta A (Q - P) + (\nu - \eta) A (y + P) - B(z + Q) + B(y + P).$$

Taking the scalar product with w we obtain

$$< w', w > = < G - \tilde{F}, w > -\eta ||w||^{2} - \eta((Q - P, w)) + (\nu - \eta)((y + P, w)) + b(y + P, y + P, w) - b(z + Q, z + Q, w).$$
(4.6.1)

Now we estimate the last term of the last equality:

$$\begin{split} b(y+P, \ y+P, \ w) &- b(z+Q, \ z+Q, \ w) \\ &= b(y, \ y, \ w) + b(y, \ P, \ w) + b(P, \ y, \ w) + b(P, \ P, \ w) \\ &- b(z, \ z, \ w) - b(z, \ Q, \ w) - b(Q, \ z, \ w) - b(Q, \ Q, \ w) \\ &= b(y, \ y, \ w) - b(z, \ z, \ w) + b(y, \ P, \ w) - b(z, \ Q, \ w) \\ &+ b(P, \ y, \ w) - b(Q, \ y, \ w) + b(P, \ P, \ w) - b(Q, \ Q, \ w). \end{split}$$

We put

$$Z_{1} := b(y, y, w) - b(z, z, w)$$

$$Z_{2} := b(y, P, w) - b(z, Q, w)$$

$$Z_{3} := b(P, y, w) - b(Q, z, w)$$

$$Z_{4} := b(P, P, w) - b(Q, Q, w).$$

Then

$$|b(y+P, y+P, w) - b(z+Q, z+Q, w)| \le |Z_1| + |Z_2| + |Z_3| + |Z_4| \quad (4.6.2)$$

Since

$$\begin{aligned} |Z_1| &= |b(y, y, w) - b(z, z, w)| = |-b(w, y, w)| \\ &\leq C|w|||w|||y||, \quad \text{by (2.9.5)} \\ (4.6.3) \\ |Z_2| &= |b(y, P, w) - b(z, Q, w)| = |b(y, P, w) - b(y, Q, w) - b(w, Q, w)| \\ &= |b(y, P - Q, w) - b(w, Q, w)| \leq |b(y, w, P - Q)| + |b(w, Q, w)| \\ &\leq C|y|||w|||P - Q|^{\frac{1}{2}}|P - Q|^{\frac{1}{2}}_{[2]} + C|w|||w|||Q|| \quad \text{by (2.9.4) and (2.9.5),} \\ &\leq C_1|y|||w|||P - Q|_{[2]} + C|w|||w|||Q||, \quad \text{by } |\cdot| \leq ||\cdot||_2 \\ (4.6.4) \\ |Z_3| &= |b(P, y, w) - b(Q, z, w)| = |b(P, y, w) - b(Q, y, w) - b(Q, w, w)| \\ &= |-b(P - Q, w, y)| \leq C|P - Q|^{\frac{1}{2}}|P - Q|^{\frac{1}{2}}_{[2]}||w|||y| \quad \text{by (2.9.3)} \\ &\leq C_1|P - Q|_{[2]}||w|||y|, \\ |Z_4| &= |b(P, P, w) - b(Q, Q, w)| = |b(P - Q, P, w) + b(Q, P, w) \\ &- b(Q, Q, w)| \leq |b(P - Q, P, w)| + |b(Q, P - Q, w)| \\ &\leq C||P - Q|||P||||w|| + C||Q||||P - Q|||w|| \quad \text{by (2.9.1).} \end{aligned}$$

From (4.6.2), (4.6.3)-(4.6.7) and (4.6.1) we obtain

$$\begin{aligned} \frac{d}{dt} |w|^2 &\leq 2 \|G - \tilde{F}\|_{V'} \|w\| - 2\eta \|w\|^2 + 2\eta \|Q - P\| \|w\| + 2|\nu - \eta| \|y + P\| \|w\| \\ &+ 2C|w| \|w\| \|y\| + 2C_1 |y| \|w\| |P - Q|_{[2]} + 2C|w| \|w\| \|Q\| \\ &+ 2C_1 |P - Q|_{[2]} \|w\| |y| + 2C \|P - Q\| \|P\| \|w\| + 2C \|Q\| \|P - Q\| \|w\|; \end{aligned}$$

Hence

$$\begin{aligned} &\frac{d}{dt}|w|^2 + \eta \|w\|^2 \leq \frac{8}{\eta} \|G - \tilde{F}\|_{V'}^2 + 8\eta \|Q - P\|^2 + \frac{8}{\eta}|\nu - \eta|^2 \|y + P\|^2 \\ &+ \frac{8}{\eta}C^2|w|^2(\|y\|^2 + \|Q\|^2) + \frac{8}{\eta}4C_1^2|y|^2|P - Q|_{[2]}^2 + \frac{8}{\eta}C^2\|P - Q\|^2(\|P\|^2 + \|Q\|^2). \end{aligned}$$

Now if η satisfies

$$|\nu - \eta| < \frac{\nu}{2}$$

we have $\eta \in]\frac{\nu}{2}, \frac{3\nu}{2}[$. Hence, from the last equation we obtain

$$\frac{d}{dt}|w|^{2} + \frac{\nu}{2}||w||^{2} \le \frac{16}{\nu}||G - \tilde{F}||_{V'}^{2} + |Q - P|_{[2]}^{2}(12\nu C_{2} + \frac{16}{\nu}4C_{1}^{2}|y|^{2})
+ \frac{16}{\nu}|\nu - \eta|^{2}||y + P||^{2} + \frac{16}{\nu}C^{2}|w|^{2}(||y||^{2} + ||Q||^{2}) + \frac{16}{\nu}C^{2}||P - Q||^{2}(||P||^{2} + ||Q||^{2}).$$
(4.6.8)

By the Gronwall Inequality:

$$\begin{split} |w(s)|^{2} &\leq \exp\left(\int_{0}^{T} \frac{16}{\nu} C^{2} \left(\|y(t)\|^{2} + \|Q(t)\|^{2}\right) dt\right) \left(|w(0)|^{2} \\ &+ \int_{0}^{T} \left[\frac{16}{\nu} \|G(t) - \tilde{F}(t)\|_{V'}^{2} + |Q(t) - P(t)|_{[2]}^{2} (12\nu C_{2} + \frac{16}{\nu} 4C_{1}^{2}|y|^{2}) \\ &+ \frac{16}{\nu} |\nu - \eta|^{2} \|y(t) + P(t)\|^{2} + \frac{16}{\nu} C^{2} \|P(t) - Q(t)\|^{2} \left(\|P(t)\|^{2} + \|Q(t)\|^{2}\right) dt \right). \end{split}$$

$$(4.6.9)$$

Now we consider two cases P = 0 & $P \neq 0$.

If $P \neq 0$ and if Q satisfies

$$||P - Q||_{L^4(0,T,D(A))} < \frac{||P||_{L^4(0,T,D(A))}}{2}$$

and since $||y||_{L^{\infty}(0,T,H)} =: D < +\infty$ we have that

$$|w(s)|^{2} \leq \exp\left[C_{3}\left(\int_{0}^{T} ||y(t)||^{2} dt + \left(\int_{0}^{T} ||P(t)||^{4} dt\right)^{\frac{1}{2}}\right)\right] \left(|w(0)|^{2} + C_{4}\left[\int_{0}^{T} ||G(t) - \tilde{F}(t)||^{2}_{V'} dt + \left(\int_{0}^{T} |Q(t) - P(t)|^{4}_{[2]} dt\right)^{\frac{1}{2}} + |\nu - \eta|^{2} \int_{0}^{T} ||y(t) + P(t)||^{2} dt + \left(\int_{0}^{T} ||P - Q||^{4} dt\right)^{\frac{1}{2}} \left(\int_{0}^{T} ||P||^{4} dt\right)^{\frac{1}{2}}\right].$$

Thus

$$|w(t)|^{2} \leq C_{5} \bigg(|w(0)|^{2} + \|G - \tilde{F}\|_{L^{2}(0, T, V')}^{2} + \|Q - P\|_{L^{4}(0, T, D(A))}^{2} + |\nu - \eta|^{2} + \|Q - P\|_{L^{4}(0, T, D(A))}^{2} \bigg).$$

We rewrite the last equation as

$$|w(t)|^{2} \leq C_{6}^{2} \left(|w(0)|^{2} + \|G - \tilde{F}\|_{L^{2}(0, T, V')}^{2} + \|Q - P\|_{L^{4}(0, T, D(A))}^{2} + |\nu - \eta|^{2} \right).$$

Then (in the case $P \neq 0$), if the quadruple (z_0, G, Q, η) satisfies

$$\begin{aligned} |\nu - \eta| &< \min\{\frac{\nu}{2}, \frac{\varepsilon}{2C_6}\};\\ \|P - Q\|_{L^4(0, T, D(A))} &< \min\{\frac{\|P\|_{L^4(0, T, D(A))}}{2}, \frac{\varepsilon}{2C_6}\};\\ |z_0 - y_0| &= |w(0)| < \frac{\varepsilon}{2C_6}; \quad \|G - \tilde{F}\|_{L^2(0, T, V')} < \frac{\varepsilon}{2C_6}; \end{aligned}$$

we have that $|w(s)| < \varepsilon$. Therefore we have the continuity of \mathbb{Y} in all quadruples such that $P \neq 0$. Note that C_6 depends only in the fixed quadruple (y_0, \tilde{F}, P, ν) where we are studying the continuity.²

 $^{{}^{2}}C_{6}$ depends in R and T too, but T and R are fixed in the statement of the Theorem.

4.6. CONTINUITY.

In the case P = 0 we can start from a bound for Q, say that Q satisfies $||Q||_{L^4(0,T,D(A))} < C_7$ (so that the argument of the exponential is bounded by a constant depending only on C_7 and (y_0, \tilde{F}, P, ν) , but not on Q). Then from (4.6.9) we can conclude that

$$|w(t)|^{2} \leq C_{8} \left(|w(0)|^{2} + ||G - \tilde{F}||^{2}_{L^{2}(0, T, V')} + ||Q||^{2}_{L^{4}(0, T, D(A))} + |\nu - \eta|^{2} + ||Q||^{4}_{L^{4}(0, T, D(A))} \right)$$

If $||Q||_{L^4(0,T,D(A))} < 1$ we can rewrite the last equation obtaining

$$|w(t)|^{2} \leq C_{9}^{2} \left(|w(0)|^{2} + \|G - \tilde{F}\|_{L^{2}(0, T, V')}^{2} + \|Q\|_{L^{4}(0, T, D(A))}^{2} + |\nu - \eta|^{2} \right)$$

So in the case P = 0 we choose, for the quadruple (z_0, G, Q, η) , the bounds

$$\begin{aligned} |\nu - \eta| &< \min\{\frac{\nu}{2}, \frac{\varepsilon}{2C_9}\}; \quad \|Q\|_{L^4(0, T, D(A))} &< \min\{1, \frac{\varepsilon}{2C_9}\}; \\ |z_0 - y_0| &= |w(0)| < \frac{\varepsilon}{2C_9}; \quad \|G - \tilde{F}\|_{L^2(0, T, V')} < \frac{\varepsilon}{2C_9}; \end{aligned}$$

and $\mathbb {Y}$ is continuous in this case too.

Theorem 4.6.2. The map

$$\mathbb{Y}_2: H \times L^2(0, T, V') \times L^4(0, T, D(A)) \times]0, +\infty[\to L^2([0, T], V) \\ (y_0, \tilde{F}, P, \nu) \mapsto y$$

is continuous. Where y is the unique solution of problems (4.3.1)-(4.3.2) corresponding to the data (y_0, \tilde{F}, P, ν) .

Proof. We Fix $\varepsilon > 0$, $(y_0, \tilde{F}, P, \nu) \in H \times L^2(0, T, V') \times L^4(0, T, D(A)) \times]0, +\infty[$ and put $y := \mathbb{Y}_2(y_0, \tilde{F}, P, \nu).$

Let (z_0, G, Q, η) be another quadruple in the product $H \times L^2(0, T, V') \times L^4(0, T, D(A)) \times]0, +\infty[$. Put $z := \mathbb{Y}_2(z_0, G, Q, \eta)$ and w := z - y and like in the proof of Theorem 4.6.1 we arrive to (4.6.8):

$$\begin{aligned} &\frac{d}{dt}|w|^2 + \frac{\nu}{2}\|w\|^2 \leq \frac{16}{\nu}\|G - \tilde{F}\|_{V'}^2 + \frac{16}{\nu}\|Q - P\|_2^2(1 + 4C_1^2|y|^2) \\ &+ \frac{16}{\nu}|\nu - \eta|^2\|y + P\|^2 + \frac{16}{\nu}C^2|w|^2(\|y\|^2 + \|Q\|^2) + \frac{16}{\nu}C^2\|P - Q\|^2(\|P\|^2 + \|Q\|^2). \end{aligned}$$

Integrating over [0, T]

$$\begin{split} |w(T)|^{2} - |w(0)|^{2} + \int_{0}^{T} \frac{\nu}{2} ||w(t)||^{2} dt &\leq \int_{0}^{T} \left[\frac{16}{\nu} ||G(t) - \tilde{F}(t)||_{V'}^{2} \\ &+ \frac{16}{\nu} ||Q(t) - P(t)||_{2}^{2} (1 + 4C_{1}^{2} |y(t)|^{2}) + \frac{16}{\nu} |\nu - \eta|^{2} ||y(t) + P(t)||^{2} \\ &+ \frac{16}{\nu} C^{2} |w(t)|^{2} (||y(t)||^{2} + ||Q(t)||^{2}) + \frac{16}{\nu} C^{2} ||P(t) - Q(t)||^{2} (||P(t)||^{2} + ||Q(t)||^{2}) \right] dt. \end{split}$$

Looking at the last two terms we see that

$$\int_0^T \|y(t)\|^2 + \|Q(t)\|^2 dt = \int_0^T \|y(t)\|^2 dt + \int_0^T \|Q(t)\|^2 dt$$
$$\leq D_1 + D_2 \|Q\|_{L^4(0, T, D(A))}^2$$

and

$$\int_{0}^{T} \|P(t) - Q(t)\|^{2} \|P(t)\|^{2} dt + \int_{0}^{T} \|P(t) - Q(t)\|^{2} \|Q(t)\|^{2} dt$$

$$\leq D_{3} \|P - Q\|_{L^{4}(0, T, D(A))}^{2} \left(\|P\|_{L^{4}(0, T, D(A))}^{2} + \|Q\|_{L^{4}(0, T, D(A))}^{2} \right)$$

so, we arrive to

$$\int_{0}^{T} \|w(t)\|^{2} dt \leq C_{1} \|G - \tilde{F}\|_{L^{2}(0,T,V'}^{2} + C_{2} \|Q - P\|_{L^{4}(0,T,D(A)}^{2} + C_{3} |\nu - \eta|^{2} + C_{4} |w|_{C([0,T],H)}^{2} (1 + \|Q\|_{L^{4}(0,T,D(A)}^{2}) + C_{5} \|P - Q\|_{L^{4}(0,T,D(A)}^{2} (\|P\|_{L^{4}(0,T,D(A)}^{2} + \|Q\|_{L^{4}(0,T,D(A)}^{2}))$$

where the constants do not depend on the quadruple (z_0, G, Q, η) . If we choose Q close to P, say $||P - Q||_{L^4(0, T, D(A)} < 1$ we have

$$\begin{split} \int_0^T \|w(t)\|^2 \, dt &\leq C_1 \|G - \tilde{F}\|_{L^2(0, T, V'}^2 + C_2 \|Q - P\|_{L^4(0, T, D(A)}^2 \\ &+ C_3 |\nu - \eta|^2 + D_4 |w|_{C([0, T], H)}^2 \left(1 + \|P\|_{L^4(0, T, D(A)}\right)^2 \\ &+ D_5 \|P - Q\|_{L^4(0, T, D(A)}^2 \left(1 + \|P\|_{L^4(0, T, D(A)}\right)^2 \right) \end{split}$$

or

$$\int_0^T \|w(t)\|^2 dt \le C_1 \|G - \tilde{F}\|_{L^2(0, T, V')}^2 + D_6 \|Q - P\|_{L^4(0, T, D(A))}^2 + C_3 |\nu - \eta|^2 + D_7 |w|_{C([0, T], H)}^2$$

(where the constants do not depend on the quadruple (z_0, G, Q, η)). By Theorem 4.6.1 there is a $\delta > 0$ such that if both $|y_0 - z_0|$, $||G - \tilde{F}||_{L^2(0, T, V')}$, $||P - Q||_{L^4(0, T, D(A)}$ and $|\eta - \nu|$ are less than δ , then $|w|_{C([0, T], H)} < \sqrt{\frac{\varepsilon}{4D_7}}$. Hence for some δ_1 smaller that δ we have that if both $|y_0 - z_0|$, $||G - \tilde{F}||_{L^2(0, T, V')}$, $||P - Q||_{L^4(0, T, D(A)}$ and $|\eta - \nu|$ are less than δ_1 , then $\left(\int_0^T ||w(t)||^2 dt\right)^{\frac{1}{2}} < \varepsilon$. Therefore the map \mathbb{Y}_2 is continuous.

Remark 12. As said in Remark 11 considering $P \in L^2(0, T, V)$ is enough to guarantee the existence of weak solutions but, to have the continuity of \mathbb{Y} or \mathbb{Y}_2 we have to consider P varying in $L^2(0, T, D(A))$ -norm. Variation in $L^2(0, T, V)$ -norm seems to be not stronger enough. Problem 4.7.1. Given

4.7 Strong Case.

Consider the equivalent problems 4.7.1 and 4.7.2:

$$\tilde{F} \in L^2(0, T, H), \quad P \in L^4(0, T, D(A))$$
&
(4.7.1)

$$y_0 \in V \tag{4.7.2}$$

$$y \in L^2(0, T, D(A)) \cap L^{\infty}(0, T, V)$$
satisfying (in the distribution sense)
$$(4.7.3)$$

satisfying (in the distribution sets

$$\forall v \in V$$
:

$$v \in V$$
:

$$\frac{d}{dt}(y, v) + \nu((y+P, v)) + b(y+P, y+P, v) = (\tilde{F}, v), \qquad (4.7.4)$$

and

$$y(0) = y_0. (4.7.5)$$

Problem 4.7.2. Given

$$\tilde{F} \in L^2(0, T, H), \quad P \in L^4(0, T, D(A))$$

&
(4.7.6)

$$y_0 \in V, \tag{4.7.7}$$

to find

$$y \in L^2(0, T, D(A)) \cap L^{\infty}(0, T, V), \qquad y' \in L^2(0, T, H)$$
 (4.7.8)

$$y' + \nu A(y+P) + B(y+P) = \tilde{F}$$
 on]0, T[, (4.7.9)
and

$$y(0) = y_0.$$

The equivalence of these problems follows from

• A solution of Problem 4.7.2 is a solution of Problem 4.7.1;
• Given a solution of Problem 4.7.1

$$- \tilde{F} \in L^2(0, T, H);$$

 $- A(y + P) \in L^2(0, T, H).$
Indeed $||A(y + P)||^2 = |y + P|_{[2]}^2$ and $y + P \in L^2(0, T, D(A));$
 $- B(y + P) \in L^2(0, T, H).$
See Remark 13 below.

So that $\tilde{F} - \nu A(y+P) - B(y+P) \in L^2(0, T, H) \subseteq L^2(0, T, V')$. By 4.7.4 and $y \in L^2(0, T, V)$ we have $y' = \tilde{F} - \nu A(y+P) - B(y+P)$ a.e., and $y \in C([0, T], V')$ "a.e.".

Remark 13. By definition and trilinearity we have

$$\begin{split} B(y+P)(h) &= b(y+P,\,y+P,\,h) \\ &= b(y,\,y,\,h) + b(y,\,P,\,h) + b(P,\,y,\,h) + b(P,\,P,\,h) \end{split}$$

(4.7.10)

or

$$B(y+P) = B(y, y) + B(y, P) + B(P, y) + B(P, P).$$
(4.7.11)

From (2.9.2) we obtain

$$\begin{split} |B(y, y)| &\leq C |y|^{\frac{1}{2}} \|y\| |y|^{\frac{1}{2}}_{[2]}; \\ |B(y, P)| &\leq C |y|^{\frac{1}{2}} \|y\|^{\frac{1}{2}} \|P\|^{\frac{1}{2}} |P|^{\frac{1}{2}}_{[2]}; \\ |B(y, P)| &\leq C |P|^{\frac{1}{2}} \|P\|^{\frac{1}{2}} \|y\|^{\frac{1}{2}} |y|^{\frac{1}{2}}_{[2]}; \text{ and} \\ |B(P, P)| &\leq C |P|^{\frac{1}{2}} \|P\| |P|^{\frac{1}{2}}_{[2]}. \end{split}$$

Thus for some constant C_1

$$|B(y, y)|^{4} \leq C_{1} ||y||^{6} |y|_{[2]}^{2} \in L^{1}(0, T, \mathbb{R});$$

$$|B(y, P)|^{4} \leq C_{1} ||y||^{4} |P|_{[2]}^{4} \in L^{1}(0, T, \mathbb{R});$$

$$|B(y, P)|^{2} \leq C_{1} |P|_{[2]}^{2} ||y|| ||y|_{[2]} \in L^{1}(0, T, \mathbb{R}); and$$

$$|B(P, P)|^{2} \leq C_{1} |P|_{[2]}^{4} \in L^{1}(0, T, \mathbb{R}).$$

So, B(y, y) and B(y, P) are elements of $L^4(0, T, H)$ and, B(P, y) and B(P, P) are elements of $L^2(0, T, H)$ thus, by (4.7.11), B(y+P) is a sum of elements of $L^2(0, T, H)$.

4.7.1 Existence.

Theorem 4.7.1. Given \tilde{F} , P and u_0 satisfying (4.7.6) and (4.7.7). There is at least one function y satisfying (4.7.8)-(4.7.10).

Proof. As we have done for Theorem 4.4.1 starting from approximate solutions we arrive to the conclusions (4.4.4) and (4.4.6). An approximate solution $y^m := \sum_{\max\{i_1, i_2\} \le m} y_i^m(t) W_i$ satisfy

$$((y^m)', v) + \nu(A(y^m + P^m), v) + (B(y^m + P^m), v) = (\tilde{F}, v), \quad v \in V,$$

setting $y_i^m \bar{\imath} W_i$ for v and summing up we arrive to

$$((y^m)', Ay^m) + \nu(A(y^m + P^m), Ay^m) + (B(y^m + P^m), Ay^m) = (\tilde{F}, Ay^m).$$
(4.7.12)

We have that

$$\begin{split} ((y^m)',\,Ay^m) &= (((y^m)',\,y^m)) = \frac{1}{2}\frac{d}{dt}\|y^m\|^2;\\ (A(y^m+P^m),\,Ay^m) &= |y^m|^2_{[2]} + ((P^m,\,y^m)); \end{split}$$

and for the term with B we have

By (2.9.3) we have

$$\begin{split} |b(y^{m}, P^{m}, Ay^{m})| &\leq C|y^{m}|^{\frac{1}{2}}|y^{m}|^{\frac{1}{2}}_{[2]}||P|||y^{m}|_{[2]} = C|y^{m}|^{\frac{1}{2}}||P|||y^{m}|^{\frac{3}{2}}_{[2]} \\ |b(y^{m}, y^{m}, Ay^{m})| &\leq C|y^{m}|^{\frac{1}{2}}|y^{m}|^{\frac{1}{2}}_{[2]}||y^{m}|||y^{m}|_{[2]} = C|y^{m}|^{\frac{1}{2}}||y^{m}|||y^{m}|^{\frac{3}{2}}_{[2]} \\ |b(P^{m}, P^{m}, Ay^{m})| &\leq C|P|^{\frac{1}{2}}|P|^{\frac{1}{2}}_{[2]}||P|||y^{m}|_{[2]} \\ |b(P^{m}, y^{m}, Ay^{m})| &\leq C|P|^{\frac{1}{2}}|P|^{\frac{1}{2}}_{[2]}||y^{m}|||y^{m}|_{[2]}. \end{split}$$

From (4.7.12) we obtain

$$\begin{split} \frac{d}{dt} \|y^m\|^2 + 2\nu |y^m|^2_{[2]} &\leq 2\nu |P|_{[2]} |y^m|_{[2]} + 2|\tilde{F}||y^m|_{[2]}; \\ &+ 2C|y^m|^{\frac{1}{2}} \|P\||y^m|^{\frac{3}{2}}_{[2]} + 2C|y^m|^{\frac{1}{2}} \|y^m\||y^m|^{\frac{3}{2}}_{[2]} \\ &+ 2C|P|^{\frac{1}{2}} |P|^{\frac{1}{2}}_{[2]} \|P\||y^m|_{[2]} + 2C|P|^{\frac{1}{2}} |P|^{\frac{1}{2}}_{[2]} \|y^m\||y^m|_{[2]}; \end{split}$$

by Young inequalities

$$\frac{d}{dt} \|y^{m}\|^{2} + \nu |y^{m}|_{[2]}^{2} \leq C_{1} |P|_{[2]}^{2} + C_{1} |\tilde{F}|^{2};$$

$$+ C_{2} |y^{m}|^{2} \|P\|^{4} + C_{2} |y^{m}|^{2} \|y^{m}\|^{4}$$

$$+ C_{3} |P||P|_{[2]} \|P\|^{2} + C_{3} |P||P|_{[2]} \|y^{m}\|^{2}.$$
(4.7.13)

By Gronwall Inequality

$$||y^{m}(t)||^{2} \leq \exp\left(\int_{0}^{T} C_{2}|y^{m}(s)|^{2}||y^{m}(s)||^{2} + C_{4}|P(s)|^{2}_{[2]} ds\right) \left(||y(0)||^{2} + \int_{0}^{T} C_{1}|P(s)|^{2}_{[2]} + C_{1}|\tilde{F}(s)|^{2} + C_{2}|y^{m}(s)|^{2}||P(s)||^{4} + C_{5}|P(s)|^{4}_{[2]} ds\right).$$

By (4.4.4), (4.4.6), $P \in L^4(0, T, D(A))$ and $\tilde{F} \in L^2(0, T, H)$ we have

 $||y^m(t)||^2 \le K_1$ for some constant K_1 .

Hence

$$y^m$$
 remains in a bounded set of $L^{\infty}(0, T, V)$. (4.7.14)

Integrating (4.7.13) over [0, T] we obtain

$$||y^{m}(T)||^{2} - ||y^{m}(0)||^{2} + \nu \int_{0}^{T} |y^{m}(t)|^{2}_{[2]} dt \leq \int_{0}^{T} C_{1} |P(t)|^{2}_{[2]} + C_{1} |\tilde{F}(t)|^{2} dt + \int_{0}^{T} C_{2} |y^{m}|^{2} ||P||^{4} + C_{2} |y^{m}|^{2} ||y^{m}||^{4} + C_{5} |P|^{4}_{[2]} + C_{6} |P|^{2}_{[2]} ||y^{m}||^{2} dt.$$

Again, by (4.4.4), (4.4.6), (4.7.14), $P \in L^4(0, T, D(A))$ and $\tilde{F} \in L^2(0, T, H)$ we arrive to

$$\int_0^1 |y^m(t)|_{[2]}^2 < K_2 \qquad \text{for some constant } K_2.$$

Hence

$$y^m$$
 remains in a bounded set of $L^2(0, T, D(A))$. (4.7.15)

The rest of the proof is analogous to that of Corollary 4.1.1. $\hfill \Box$

Remark 14. Contrary to the case of weak solutios (see Remark 11), we can not guarantee from the previous sketch that considering $P \in L^2(0, T, V)$ is enough to guarantee the existence of a strong solution.

4.7.2 Uniqueness.

Theorem 4.7.2. The solution of Problems 4.7.1-4.7.2 given by Theorem 4.7.1 is unique. Moreover it is a.e. equal to a continuous function from [0, T] into V.

Proof. The uniqueness follows from Theorem 4.5.1 and from the fact that a solution of problems 4.7.1-4.7.2 is a solution of problems 4.3.1-4.3.2. By $y' \in L^2(0, T, H)$ we have $y' \in L^2(0, T, D(A)')$. Since $y \in L^2(0, T, D(A))$ and the inclusions

$$D(A) \subseteq V \subseteq D(A)'$$

are dense and continuous we have that $y \in C([0, T], V)$. [See Lemma 1.2 in [15] section III.1].

4.7.3 Continuity.

Theorem 4.7.3. The map

$$\begin{aligned} \mathbb{Y}_s: \, V \times L^2(0, \, T, \, H) \times L^4(0, \, T, \, D(A)) \times]0, \, +\infty[\to C([0, \, T], \, V) \\ (y_0, \, \tilde{F}, \, P, \, \nu) \mapsto y \end{aligned}$$

is continuous. Where y is the unique solution of problems (4.7.1)-(4.7.2) corresponding to the data (y_0, \tilde{F}, P, ν) .

Proof. The proof is analogous to that of Theorem 4.6.1. We fix a quadruple

 $(y_0, \tilde{F}, P, \nu) \in V \times L^2(0, T, H) \times L^4(0, T, D(A)) \times]0, +\infty[$

and $\varepsilon > 0$. Then consider another quadruple

$$(z_0, G, Q, \eta) \in V \times L^2(0, T, H) \times L^4(0, T, D(A)) \times]0, +\infty[.$$

 Put

$$y := \mathbb{Y}_s(y_0, F, P, \nu) \& z := \mathbb{Y}_s(z_0, G, Q, \eta)$$

Putting w := z - y we obtain

$$w' = G - \tilde{F} - \eta A w - \eta A (Q - P) + (\nu - \eta) A (y + P) - B(z + Q) + B(y + P).$$

Taking the scalar product with Aw we obtain

$$(w', Aw) = \langle G - \tilde{F}, Aw \rangle -\eta |w|_{[2]}^2 - \eta (Q - P, w)_{[2]} + (\nu - \eta)(y + P, w)_{[2]} + b(y + P, y + P, w) - b(z + Q, z + Q, w).$$

$$(4.7.16)$$

Now we estimate the last difference:

$$b(y + P, y + P, w) - b(z + Q, z + Q, w)$$

= $b(y, y, Aw) + b(y, P, Aw) + b(P, y, Aw) + b(P, P, Aw)$
 $- b(z, z, Aw) - b(z, Q, Aw) - b(Q, z, Aw) - b(Q, Q, Aw).$

We put

$$\begin{split} &Z_1 := b(y, y, Aw) - b(z, z, Aw) \\ &= b(y, y, Aw) - \left(b(w, z, Aw) + b(y, z, Aw)\right) \\ &= b(y, y, Aw) - \left(b(w, w, Aw) + b(w, y, Aw) + b(y, w, Aw) + b(y, y, Aw)\right) \\ &= -b(w, w, Aw) - b(w, y, Aw) - b(y, w, Aw); \\ &Z_2 := b(y, P, Aw) - b(z, Q, Aw) \\ &= b(y, P, Aw) - \left(b(w, Q, Aw) + b(y, Q, Aw)\right) \\ &= b(y, P, Aw) - \left(b(w, Q, Aw) + b(y, Q - P, Aw) + b(y, P, Aw)\right) \\ &= -b(w, Q, Aw) - b(y, Q - P, Aw); \\ &Z_3 := b(P, y, Aw) - b(Q, z, Aw) \\ &= b(P, y, Aw) - \left(b(Q, w, Aw) + b(Q, P, y, Aw)\right) \\ &= b(P, y, Aw) - \left(b(Q, w, Aw) + b(Q - P, y, Aw) + b(P, y, Aw)\right) \\ &= -b(Q, w, Aw) - b(Q - P, y, Aw); \\ &Z_4 := b(P, P, Aw) - b(Q, Q, Aw) \\ &= b(P - Q, P, Aw) + \left(b(Q, P - Q, Aw); \right) \\ &= b(P - Q, P, Aw) + b(Q, P - Q, Aw); \end{split}$$

We have

$$|b(y+P, y+P, w) - b(z+Q, z+Q, w)| \le |Z_1| + |Z_2| + |Z_3| + |Z_4| \quad (4.7.17)$$

and, by (2.9.3)

$$|Z_{1}| \leq C|w|^{\frac{1}{2}}|w|^{\frac{1}{2}}_{[2]}||w|||w|_{[2]} + C|w|^{\frac{1}{2}}|w|^{\frac{1}{2}}_{[2]}||y|||w|_{[2]} + C|y|^{\frac{1}{2}}|y|^{\frac{1}{2}}_{[2]}||w|||w|_{[2]},$$
(4.7.18)

$$|Z_2| \le C|w|^{\frac{1}{2}} |w|^{\frac{1}{2}}_{[2]} ||Q|| |w|_{[2]} + C|y|^{\frac{1}{2}} |y|^{\frac{1}{2}}_{[2]} ||P - Q|| |w|_{[2]},$$
(4.7.19)

$$|Z_3| \le C|Q|^{\frac{1}{2}} |Q|^{\frac{1}{2}}_{[2]} ||w|| |w|_{[2]} + C|P - Q|^{\frac{1}{2}} |P - Q|^{\frac{1}{2}}_{[2]} ||y|| |w|_{[2]}, \qquad (4.7.20)$$

$$|Z_4| \le C|P - Q|^{\frac{1}{2}}|P - Q|^{\frac{1}{2}}_{[2]}||P|||w|_{[2]} + C|Q|^{\frac{1}{2}}|Q|^{\frac{1}{2}}_{[2]}||P - Q|||w|_{[2]}.$$
 (4.7.21)

From (4.7.17) and (4.7.16) we obtain

$$\begin{aligned} \frac{d}{dt} \|w\|^2 &\leq 2|G - \tilde{F}||w|_{[2]} - 2\eta|w|_{[2]}^2 + 2\eta|Q - P|_{[2]}|w|_{[2]} \\ &+ 2|\nu - \eta||y + P|_{[2]}|w|_{[2]} + 2\sum_{i=1}^4 |Z_i|. \end{aligned}$$

If η satisfies $|\eta - \nu| < \frac{\nu}{2}$, i.e., $\frac{\nu}{2} < \eta < \frac{3\nu}{2}$ we obtain

$$\begin{aligned} \frac{d}{dt} \|w\|^2 + \nu \|w\|_{[2]}^2 &\leq 2|G - \tilde{F}| \|w\|_{[2]} + 3\nu |Q - P|_{[2]}|w|_{[2]} \\ &+ 2|\nu - \eta| \|y + P|_{[2]}|w|_{[2]} + 2\sum_{i=1}^4 |Z_i| \end{aligned}$$

so, applying Young inequalities, for appropriate constants (independent of the quadruple (z_0, G, Q, η)) we have

$$\frac{d}{dt} \|w\|^{2} + \frac{\nu}{2} |w|_{[2]}^{2} \leq C_{1} |G - \tilde{F}|^{2} + C_{2} |Q - P|_{[2]}^{2} + C_{3} |\nu - \eta|^{2} |y + P|_{[2]}^{2}
+ C_{4} \left(|w|^{2} \|w\|^{4} + |w|^{2} \|y\|^{4} + |y||y|_{[2]} \|w\|^{2} + |w|^{2} \|Q\|^{4} + |y||y|_{[2]} \|P - Q\|^{2}
+ |Q||Q|_{[2]} \|w\|^{2} + |P - Q||P - Q|_{[2]} \|y\|^{2} + |P - Q||P - Q|_{[2]} \|P\|^{2} + |Q||Q|_{[2]} \|P - Q\|^{2} \right).$$
(4.7.22)

By the Gronwall Inequality:

$$\begin{aligned} \|w(s)\|^{2} \\ \leq \exp\left(\int_{0}^{T} C_{4}(|w(t)|^{2} \|w(t)\|^{2} + |y(t)||y(t)|_{[2]} + |Q(t)||Q(t)|_{[2]}) dt\right) \left(\|w(0)\|^{2} \\ + C_{5} \int_{0}^{T} \left[|Q(t) - P(t)|_{[2]}^{2} \left(1 + |y(t)||y(t)|_{[2]} + \|y(t)\|^{2} \right) \\ + \|P(t)\|^{2} + |Q(t)||Q(t)|_{[2]}\right) + |G(t) - \tilde{F}(t)|^{2} + |\nu - \eta|^{2}|y(t) + P(t)|_{[2]}^{2} \\ + |w(t)|^{2} \|y(t)\|^{4} + |w(t)|^{2} \|Q(t)\|^{4} dt \right). \end{aligned}$$

$$(4.7.24)$$

Given a constant E>0, by Theorems 4.6.1, 4.6.2 and by the existence of a constant C such that 3

$$\begin{split} |a| \leq C \|a\| \& \|f\|_{L^2(0,\,T,\,V')} \leq C \|f\|_{L^2(0,\,T,\,H)} \\ \text{for all } a \in V,\, f \in L^2(0,\,T,\,H). \end{split}$$

there is $\delta > 0$ such that if both $\|y_0 - z_0\|$, $\|G - \tilde{F}\|_{L^2(0, T, H)}$, $\|P - Q\|_{L^4(0, T, D(A))}$ and $|\eta - \nu|$ are less than δ we have

$$\|w\|_{C([0,T],H)} < E$$
 and $\int_0^T \|w(t)\|^2 dt < E$.

Therefore, if we choose in addition $\delta < \frac{\nu}{2}$, we have

³The existence of such a constant comes from the continuity of the inclusions $V \subseteq H \subseteq V'$.

$$\begin{split} \|w(s)\|^{2} &\leq D_{1} \left[\|w(0)\|^{2} + \|G - \tilde{F}\|_{L^{2}(0, T, H)}^{2} \\ &+ \int_{0}^{T} |Q(t) - P(t)|_{[2]}^{2} \left(1 + |y(t)|_{[2]} + |P(t)|_{[2]}^{2} + |Q(t)|_{[2]}^{2} \right) dt \\ &+ |\nu - \eta|^{2} + \|w\|_{C([0, T], H)}^{2} (1 + \|Q\|_{L^{4}(0, T, D(A))}^{4}) \right] \end{split}$$

and, because

$$\begin{split} &\int_{0}^{T} |Q(t) - P(t)|_{[2]}^{2} \left(1 + |y(t)|_{[2]} + |P(t)|_{[2]}^{2} + |Q(t)|_{[2]}^{2} \right) dt \\ \leq & \|Q - P\|_{L^{4}(0,T,D(A))}^{2} \left(\|1\|_{L^{2}(0,T,\mathbb{R})} + \|y\|_{L^{2}(0,T,D(A))} \right) \\ & + \|P\|_{L^{4}(0,T,D(A))}^{2} + \|Q\|_{L^{4}(0,T,D(A))}^{2} \right) \\ \leq & K_{1} \|Q - P\|_{L^{4}(0,T,D(A))}^{2} \left(1 + \left(\delta + \|P\|_{L^{4}(0,T,D(A))} \right)^{2} \right) \\ \leq & K_{2} \|Q - P\|_{L^{4}(0,T,D(A))}^{2}. \end{split}$$

we have

$$\|w(s)\|^{2} \leq K_{3} \Big[\|w(0)\|^{2} + \|G - \tilde{F}\|^{2}_{L^{2}(0, T, H)} + \|Q - P\|^{2}_{L^{4}(0, T, D(A))} + |\nu - \eta|^{2} + \|w\|^{2}_{C([0, T], H)} \Big]$$

$$(4.7.25)$$

(with K_3 independent of the quadruple (z_0, G, Q, η)). Then for some δ_1 smaller than δ , and using Theorem 4.6.1, we have $||w(s)|| < \varepsilon$ if both $||y_0 - z_0||$, $||\tilde{F} - G||_{L^2(0, T, H)}$, $||P - Q||_{L^4(0, T, D(A))}$ and $|\eta - \nu|$ are less than δ_1 . Thus the map \mathbb{Y}_s is continuous.

Theorem 4.7.4. The map

$$\mathbb{Y}_{2s}: V \times L^2(0, T, H) \times L^4(0, T, D(A)) \times]0, +\infty[\to L^2([0, T], D(A)) (y_0, \tilde{F}, P, \nu) \mapsto y$$

is continuous. Where y is the unique solution of problems (4.7.1)-(4.7.2) corresponding to the data (y_0, \tilde{F}, P, ν) .

Proof. We fix $(y_0, \tilde{F}, P, \nu) \in V \times L^2(0, T, H) \times L^4(0, T, D(A)) \times]0, +\infty[, \varepsilon > 0$ and, put $y := \mathbb{Y}_{2s}(y_0, \tilde{F}, P, \nu)$.

Let (z_0, G, Q, η) be another quadruple in the product space $V \times L^2(0, T, H) \times L^4(0, T, D(A)) \times]0, +\infty[$. Put $z := \mathbb{Y}_{2s}(z_0, G, Q, \eta)$ and w := z - y and like in

the proof of Theorem 4.7.3 we arrive to (4.7.22):

$$\begin{aligned} \frac{d}{dt} \|w\|^2 &+ \frac{\nu}{2} |w|_{[2]}^2 \leq C_1 |G - \tilde{F}|^2 + C_2 |Q - P|_{[2]}^2 + C_3 |\nu - \eta|^2 |y + P|_{[2]}^2 \\ &+ C_4 \left(|w|^2 \|w\|^4 + |w|^2 \|y\|^4 + |y||y|_{[2]} \|w\|^2 + |w|^2 \|Q\|^4 \\ &+ |y||y|_{[2]} \|P - Q\|^2 + |Q||Q|_{[2]} \|w\|^2 + |P - Q||P - Q|_{[2]} \|y\|^2 \\ &+ |P - Q||P - Q|_{[2]} \|P\|^2 + |Q||Q|_{[2]} \|P - Q\|^2 \right). \end{aligned}$$

 $\qquad \text{ or, } \qquad$

$$\begin{aligned} \frac{d}{dt} \|w\|^2 + \frac{\nu}{2} |w|_{[2]}^2 &\leq C_1 |G - \tilde{F}|^2 + C_2 |Q - P|_{[2]}^2 + C_3 |\nu - \eta|^2 |y + P|_{[2]}^2 \\ &+ C_5 \left(\|w\|^6 + \|w\|^2 + |y|_{[2]}^2 \|w\|^2 + \|w\|^2 \|Q\|^4 + |Q|_{[2]}^2 \|w\|^2 \\ &+ |P - Q|_{[2]}^2 \left(1 + |y|_{[2]} + |P|_{[2]}^2 + |Q|_{[2]}^2 \right) \right) \end{aligned}$$

(where the constants are independent of the quadruple (z_0, G, Q, η)). Integrating over [0, T]

$$\begin{split} \int_{0}^{T} |w(t)|_{[2]}^{2} dt &\leq D \left[\|w\|_{C([0,T],V)}^{2} + \|\tilde{F} - G\|_{L^{2}(0,T,H)}^{2} + \|Q - P\|_{L^{4}(0,T,D(A))}^{2} \\ &+ |\nu - \eta|^{2} + \|w\|_{C([0,T],V)}^{2} \left(\|w\|_{C([0,T],V)}^{4} + \|y\|_{L^{2}(0,T,D(A))}^{2} \\ &+ \|Q\|_{L^{4}(0,T,D(A))}^{4} + \|Q\|_{L^{4}(0,T,D(A))}^{2} \right) \\ &+ \|P - Q\|_{L^{4}(0,T,D(A))}^{2} \left(1 + \|y\|_{L^{2}(0,T,D(A))} \\ &+ \|P\|_{L^{4}(0,T,D(A))}^{2} + \|Q\|_{L^{4}(0,T,D(A))}^{2} \right) \end{split}$$

By Theorem 4.7.3, given a constant E > 0, there is a $\delta > 0$ such that, if both $\|y_0 - z_0\|$, $\|P - Q\|_{L^4(0, T, D(A))}$, $\|\tilde{F} - G\|_{L^2(0, T, H)}$, and $|\eta - \nu|$ are less than δ we have

$$||w||_{C([0,T],V)} < E.$$

Thus,

$$\begin{split} \int_{0}^{T} |w(t)|_{[2]}^{2} dt &\leq D_{1} \left[\|\tilde{F} - G\|_{L^{2}(0,T,H)}^{2} + \|Q - P\|_{L^{4}(0,T,D(A))}^{2} + |\nu - \eta|^{2} \\ &+ \|w\|_{C([0,T],V)}^{2} \left(\|w\|_{C([0,T],V)}^{4} + \|P\|_{L^{2}(0,T,D(A))} \right) \\ &+ \left(\delta + \|P\|_{L^{4}(0,T,D(A))} \right)^{4} + \left(\delta + \|P\|_{L^{4}(0,T,D(A))} \right)^{2} \\ &+ \|P - Q\|_{L^{4}(0,T,D(A))}^{2} \left(1 + \|y\|_{L^{2}(0,T,D(A))} \right) \\ &+ \|P\|_{L^{4}(0,T,D(A))}^{2} + \left(\delta + \|P\|_{L^{4}(0,T,D(A))} \right)^{2} \right) \end{split}$$

or

$$\begin{split} \int_0^T |w(t)|_{[2]}^2 dt &\leq D_2 \Biggl[\|\tilde{F} - G\|_{L^2(0, T, H)}^2 + \|Q - P\|_{L^4(0, T, D(A))}^2 \\ &+ |\nu - \eta|^2 + \|w\|_{C([0, T], V)}^2 \Biggr] \end{split}$$

(with D_2 independent of the quadruple (z_0, G, Q, η)).

Hence for some δ_1 smaller than δ we have $\int_0^T |w|_{[2]}^2 < \varepsilon^2$ if both $||y_0 - z_0||$, $\|\tilde{F} - G\|_{L^2(0, T, H)}, \|P - Q\|_{L^4(0, T, D(A))}$ and $|\eta - \nu|$ are less than δ_1 . Thus the map \mathbb{Y}_{2s} is continuous.

4.8 Continuity in Relaxation Metric.

We begin with a definition:

Definition 4.8.1. The relaxation metric in $L^1([0, T], \mathbb{R}^d)$ is defined by the norm

$$\|g\|_{rx} := \max_{t_1, t_2 \in [0, T]} \left\| \int_{t_1}^{t_2} g(\tau) \, d\tau \right\|_{\mathbb{R}^d},\tag{4.8.1}$$

where $\|\cdot\|_{\mathbb{R}^d}$ is the l_1 norm in \mathbb{R}^d .

If nothing in contrary is stated we consider the spaces \mathbb{R}^d $(d \in \mathbb{N}_0)$ endowed with l_1 -norm $- \|x\|_{\mathbb{R}^d} = \|x\|_{l_1} := \sum_{i=1}^d |x_i|$.

Remark 15. It is easy to check that (4.8.1) is a semi-norm and, since functions in $L^1([0, T], \mathbb{R}^d)$ coinciding on a set of measure T are identified we can conclude that (4.8.1) is a norm.

Consider, also, the **w-relaxation metric** on $L^1([0, T], \mathbb{R}^d)$ defined by the norm

$$||g||_{wrx} := \max_{t \in [0,T]} \left\| \int_0^t g(\tau) \, d\tau \right\|_{\mathbb{R}^d}.$$
(4.8.2)

Remark 16. The identity map

$$\left(L^1([0, T], \mathbb{R}^d), \|\cdot\|_{rx}\right) \mapsto \left(L^1([0, T], \mathbb{R}^d), \|\cdot\|_{wrx}\right)$$

is continuous.

Now we fix a finite subset $\mathbb{F} \subset \mathbb{N}_0^2$, put $\mathcal{F} := \#\mathbb{F}$ and define the space $S_{\mathbb{F}} := span\{W_k \mid k \in \mathbb{F}\}$. From the fact that the dimension of $S_{\mathbb{F}}$ is finite, we conclude the equivalence of all norms defined in it. In particular the norms $\|\cdot\|_{\mathbb{R}^F}$ and $|\cdot|_{[2]}$ are equivalent on $S_{\mathbb{F}}$. From this equivalence we have the following:

Proposition 4.8.1. The map

$$\mathbb{I}: L^{\infty}_{wrx}([0, T], \mathbb{R}^{\mathcal{F}}) \to C([0, T], S_{\mathbb{F}})$$
$$(v_k(t))_{k \in \mathbb{F}} \mapsto \sum_{k \in \mathbb{F}} \mathbb{I}v_k(t)W_k$$

is continuous. Where $\mathbb{I}v_k(t) := \int_0^t v_k(\tau) d\tau$ and the subscript "wrx" means that we are considering relaxation metric on the set $L^{\infty}([0,T], \mathbb{R}^{\mathcal{F}})$.

Proof. The continuity of the map

$$\mathbb{I}: L^{\infty}_{wrx}([0, T], \mathbb{R}^{\mathcal{F}}) \to C([0, T], \mathbb{R}^{\mathcal{F}})$$
$$(v_k(t))_{k \in \mathbb{F}} \mapsto (\mathbb{I}v_k(t))_{k \in \mathbb{F}}$$

is trivial.

Recall that by definition, the map S of Corollary 4.1.2 gives us the weak solution, belonging to C([0, T], H), of the NSE for an initial data in $\Pi :=$ $H \times L^2(0, T, V') \times L^{\infty}([0, T], S_{\mathbb{F}}) \times \mathbb{R}^+$. Changing the topology on the third factor of the previous product to the w-relaxation one we arrive to the space $L^{\infty}_{wrx}([0, T], \mathbb{R}^F)$ and we define the function \mathbb{S}_{wrx} as the function defined in the product $\Pi_{wrx} := H \times L^2(0, T, V') \times L^{\infty}_{wrx}([0, T], \mathbb{R}^F) \times \mathbb{R}^+$ and taking the same values as S.

Proposition 4.8.2. The map \mathbb{S}_{wrx} is continuous.

Proof. As in the beginning of section 4.2 we put $\mathbb{Y}_{wrx} := \mathbb{S}_{wrx} - \mathbb{I}_{\circ}$. But here we want an equality between functions defined on Π_{wrx} so, we just put $\mathbb{I}_{\circ}(u_0, \tilde{F}, v, \nu) := \mathbb{I}v$.

By Proposition 4.8.1 and Theorem 4.6.1 the map

$$\begin{aligned} \mathbb{Y}_{wrx} &: \Pi_{wrx} \to C([0, T], H) \\ (u_0, \tilde{F}, v, \nu) \mapsto \mathbb{Y}(u_0, \tilde{F}, \mathbb{I}v, \nu) = \mathbb{Y} \circ \mathbb{I}^{\circ}(u_0, \tilde{F}, v, \nu) \end{aligned}$$

is continuous. Where $\mathbb{I}^{\circ}(u_0, \tilde{F}, v, \nu) := (u_0, \tilde{F}, \mathbb{I}v, \nu)$. By the equality $\mathbb{S}_{wrx} = \mathbb{Y}_{wrx} + \mathbb{I}_{\circ}$ we conclude the continuity of \mathbb{S}_{wrx} .

Analogously, using Proposition 4.8.1 and theorems 4.6.2, 4.7.3 and 4.7.4, we can prove the continuity on relaxation metric of the maps S_2 , S_s and S_{2s} arriving to the Proposition

Proposition 4.8.3. The maps \mathbb{S}_{wrx} , \mathbb{S}_{2wrx} , \mathbb{S}_{swrx} are all continuous.

By Remark 16 we obtain

Corollary 4.8.4. The maps S_{rx} , S_{2rx} , S_{srx} S_{2srx} are all continuous.⁴

Remark 17. If, instead of a subspace spanned by a finite number of eigenfunctions, we consider any (with either finite or infinite dimension) subspace $\mathbb{F} \subseteq D(A)$ and, instead of defining the relaxation metric using the l_1 -norm we define it on $L^1(0, T, \mathbb{F})$ using the norm induced by D(A):

$$\|g\|_{rx[2]} := \max_{t_1, t_2 \in [0, T]} \left\| \int_{t_1}^{t_2} g(\tau) \, d\tau \right\|_{[2]}$$

and, its corresponding weak form by

$$\|g\|_{wrx[2]} := \max_{t \in [0, T]} \left\| \int_0^t g(\tau) \, d\tau \right\|_{[2]}$$

we still to have the continuity of the corresponding maps $\mathbb{S}_{rx[2]}$, $\mathbb{S}_{2rx[2]}$, $\mathbb{S}_{srx[2]}$, $\mathbb{S}_{2srx[2]}$, *i.e.*, the solutions of the NSE vary continuously when the control varies continuously in rx[2]-metric.

The reason why we have considered a finite space is that we are interested in the case where the control in a space spanned by a finite number of eigenfunctions and, the reason to consider l_1 -metric in the definition of relaxation metric is that it will be convenient later.

4.9 A Remark on Finite System versus Infinite System.

From now we consider only strong solutions and, the external force F and the coefficient of viscosity ν are fixed.

Definition 4.9.1. Given T > 0, we say that the N-S system is **time**-T **approximately controllable in observed component** if for any $\tilde{\phi} \in V$ and any finite subset of modes $\mathcal{O} \subseteq \mathbb{N}_0^2$, the projection of the closure of the attainable set at time T from $\tilde{\phi} \in V$ onto span $\{W_k \mid k \in \mathcal{O}\}$ is surjective. It is **time**-T **controllable in observed component** if the projection of the attainable set at time T from $\tilde{\phi} \in V$ on span $\{W_k \mid k \in \mathcal{O}\}$ is surjective.

In this section we ask ourselves if can we prove approximate controllability in observed component similarly as we have proved approximated controllability of Galerkin approximations.

As we have seen in (3.4.2) from the N-S equation

$$u_t + \nu A u + P^{\nabla} B u = F + v^{-5}$$

⁴These "rx"-maps are defined similarly as the "wrx" ones, just considering the "rx"-topology in the factor of essentially bounded functions.

we can derive the infinite ODE system

$$\begin{split} \dot{u}_{k} &= \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(++)m)^{+} = k}} - \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{\bar{k}}(\bar{n} - \bar{m}) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(--)m)^{+} = k}} \frac{u_{m}u_{n}C_{m,n}^{\wedge}}{\bar{k}}(\bar{n} - \bar{m}) \mathrm{sign}(n_{1} - m_{1}) \mathrm{sign}(n_{2} - m_{2}) \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(-+)m)^{+} = k}} - \frac{u_{m}u_{n}C_{m,n}^{\vee}}{\bar{k}}(\bar{n} - \bar{m}) \mathrm{sign}n_{1} - m_{1} \\ &+ \sum_{\substack{m,n \in \mathbb{N}_{0}^{2} \\ m < n \\ (n(+-)m)^{+} = k}} \frac{u_{m}u_{n}C_{m,n}^{\vee}}{\bar{k}}(\bar{n} - \bar{m}) \mathrm{sign}(n_{2} - m_{2}) \\ &+ \nu \bar{k}u_{k} + F_{k} + v_{k}. \end{split}$$
(4.9.1)

Now consider the subset $\mathcal{K}^1 = \{(n_1, n_2) \in \mathbb{N}_0^2 \mid n_1, n_2 \leq 3\} \setminus \{(3, 3)\}$ as the set of forced modes. Put $\kappa_1 := \# \mathcal{K}^1$. If we put $y := u - \mathbb{I}v$ we can write u in the form

$$u = y + \mathbb{I}v \tag{4.9.2}$$

and y satisfies

$$y_t = -\nu A(y + \mathbb{I}v) - P^{\nabla} B(y + \mathbb{I}v) + F.$$

Writing system (4.9.1) (with \mathcal{K}^1 as set of controlled modes) in the form

$$\begin{cases} \dot{u}_k = \mathcal{B}_k(u) + \nu \bar{k} u_k + F_k + v_k & k \in \mathcal{K}^1 \\ \dot{u}_k = \mathcal{B}_k(u) + \nu \bar{k} u_k + F_k & k \notin \mathcal{K}^1 \end{cases}$$
(4.9.3)

in the new variable "y" the system becomes

$$\dot{y}_k = \mathcal{B}_k(y + \mathbb{I}v) + \nu \bar{k}(y_k + \mathbb{I}v_k) + F_k \qquad v_k = 0 \text{ if } k \notin \mathcal{K}^1.$$

$$(4.9.4)$$

For any time T > 0 the closure of the attainable set at time T of system (4.9.3)— $\underline{\mathcal{A}_{u_0}(u)(T)}$ — and the closure of the attainable set at time T of system (4.9.4)— $\overline{\mathcal{A}_{u_0}(y)(T)}$ —⁶ are related by

$$\overline{\mathcal{A}_{u_0}(u)(T)} = \overline{\mathcal{A}_{u_0}(y)(T) \times \mathbb{R}^{\kappa_1}} = \overline{\mathcal{A}_{u_0}(y)(T)} \times \mathbb{R}^{\kappa_1}.$$

Indeed the inclusion $\overline{\mathcal{A}_{u_0}(u)(T)} \subseteq \overline{\mathcal{A}_{u_0}(y)(T) \times \mathbb{R}^{\kappa_1}}$ follows from (4.9.2) and the reverse one follows from the density of the map

$$L^{\infty}([0, T], \mathbb{R}^{\kappa_1}) \ni v \mapsto (\mathbb{I}v, \mathbb{I}v(T)) \in L^4(0, T, \mathbb{R}^{\kappa_1}) \times \mathbb{R}^{\kappa_1}$$

 $^{^5\}text{Recall that}\ P^\nabla$ is the projection on the space of divergence free functions.

⁶Note that, by (4.9.2), at initial time we have $u_0 = y_0$.

and from the continuity of the map

$$L^4(0, T, \mathbb{R}^{\kappa_1}) \ni P \mapsto \mathbb{Y}_s(u_0, F, P, \nu)(T) \in V$$

for fixed u_0 that is a consequence of Theorem 4.7.3. Similarly if we rewrite the NS Equation as

$$u_t + \nu A u + P^{\nabla} B u = F + v^1 + v^2$$

we obtain

$$\overline{\mathcal{A}_{u_0}(u)(T)} = \overline{\mathcal{A}_{u_0}(y^1)(T)} \times \mathbb{R}^{\kappa_1}$$

where $\mathcal{A}_{u_0}(y^1)(T)$ stays for the attainable set at time T from u_0 of the system

$$y_t = -\nu A(y + \mathbb{I}v^2) - P^{\nabla} B(y + \mathbb{I}v^2) + F + v^1.$$

So Factorization works like in the finite dimensional case.

In the step of convexification a problem arises: In section 3.5 we obtained the new vector fields $\gamma_{m,n}$ from the vector fields $v_{m,n}^{\lambda}$ and $w_{m,n}^{\lambda}$ and from the expressions

$$\begin{aligned} \frac{f_{v_{m,n}^{\lambda}}(u)+f_{-v_{m,n}^{\lambda}}(u)}{2} &= f(u)+\lambda\gamma_{m,n}\\ & \text{and}\\ \frac{f_{w_{m,n}^{\lambda}}(u)+f_{-w_{m,n}^{\lambda}}(u)}{2} &= f(u)-\lambda\gamma_{m,n}, \end{aligned}$$

i.e., we have extracted a vector field from the convexification of the vector fields $f_{\pm v_{m,n}^{\lambda}}$ and $f_{\pm w_{m,n}^{\lambda}}$ and, in the finite dimensional case we know that we can convex without changing the closure of attainable set at time t. From Factorization we know that we can follow the vector fields $f_{\pm v_{m,n}^{\lambda}}$ and $f_{\pm w_{m,n}^{\lambda}}$ without changing closure of attainable set but, in the infinite dimensional case we do not know if we can convex these new vector fields without changing the closure of attainable set.

If using the vector fields $f(u) + \lambda \gamma_{m,n}$ and $f(u) - \lambda \gamma_{m,n}$ we would not change closure of attainable set then we could add these new directions to the old ones — $span(\mathcal{K}^1)$ — without changing the closure of attainable set.

It is known that any control $v_0 \in span\{\delta_{m,n}, e_k \mid (m,n) \in S_1, k \in \mathcal{K}^1\}$ ⁷ can be approximated in relaxation metric by controls taking values on $\{\alpha\delta_{m,n}, \alpha e_k \mid (m,n) \in S_1, k \in \mathcal{K}^1\}$ where $\alpha \in \mathbb{R}$ is a positive constant depending on v_0 (as we will see in Lemma 4.10.5 below). By the continuity of \mathbb{S}_{srx} we do not change the closure of attainable set using controls in $span\{\delta_{m,n}, e_k \mid (m,n) \in S_1, k \in \mathcal{K}^1\}$. We have just applied a step of Convexification. Note that this last procedure of Convexification is quite different from that after Factorization Procedure that is more complicated: For example setting m := (1, 1), n := (2, 1) and considering the vector field $f_{v_{m,n}^{\lambda}}(u) = f(u) + \lambda \gamma_{m,n} + \mathcal{V}_1(u, v_{m,n}^{\lambda})$ (see (3.4.4)), we see that the candidate to new control $-\lambda \gamma_{m,n} + \mathcal{V}_1(u, v_{m,n}^{\lambda})$, contrary to what happens in the last step of Convexification depends on u and does not take values on a compact subset, indeed if we put $u^r := rW_m$ $r \in \mathbb{R}$ we obtain

$$\lambda \gamma_{m,n} + \mathcal{V}_1(u^r, v_{m,n}^{\lambda})$$

= $\lambda \gamma_{m,n} - \lambda r \left[P^{\nabla} B(W_n, W_m) + P^{\nabla} B(W_m, W_n) \right] + r \nu \bar{m}$

⁷See (3.5.4) for the definition of S_1 .

Since $\lambda \gamma_{m,n} - \lambda r \left[P^{\nabla} B(W_m, W_n) + P^{\nabla} B(W_m, W_n) \right]$ takes values on

$$span\{W_{(1,2)}, W_{(3,2)}\}$$

we obtain $r\nu(\overline{(1, 1)})$ for the projection of $\lambda\gamma_{m,n} + \mathcal{V}_1(u, v_{m,n}^{\lambda})$ onto $span\{W_{(1,1)}\}$. So, the projection goes to ∞ as r does.

Without compactness we are not able to use the Approximation Lemma 4.10.5 below.

In the case of Galerkin approximations from approximate controllability we could derive (exact) controllability using the bracket generating property of the system but, in the infinite case we do not have a property like that so, we would not be able to conclude controllability in observed component immediately from approximate controllability (in the case we could somehow prove approximate controllability).

In the next Section, using some more tools, we prove the so called *solid controllability in observed component* for system (4.9.3) which implies controllability in observed component.

4.10 Solid Controllability in Observed Component

Definition 4.10.1. Let $\phi^0 : M^1 \to M^2$ be a continuous map between two finite dimensional C^0 -manifolds, $\Omega \subset M^1$ be an open subset with compact closure and, $S \subseteq M^2$ be any subset. We say that $\phi^0(\Omega)$ covers S solidly, if for some C^0 -neighborhood \mathcal{N} of $\phi^0 \mid_{\overline{\Omega}}$ there holds: $S \subseteq \phi(\Omega)$.

Let $\mathcal{O} \subset \mathbb{N}_0^2$ be the finite set of modes we want to observe and, $\Pi_{\mathcal{O}}$ be the projection map from V onto $span\{W_k \mid k \in \mathcal{O}\}$. Define, for each T > 0 and each finite subset $\mathbb{F} \subset \mathbb{N}_0^2$, the "end point" map

$$\mathbb{E}_T: V \times L^{\infty}([0, T], \mathbb{R}^{\#\mathbb{F}}) \to \mathcal{O}$$
$$(u_0, v) \mapsto \Pi_{\mathcal{O}} \circ \mathbb{S}_s(u_0, F, v, \nu)(T).$$

For any $N \in \mathbb{N}_0$ define, also, the system

$$N: \begin{cases} \dot{u}_k = \mathcal{B}_k(u) + \nu \bar{k} u_k + F_k + v_k; & k \in \mathcal{K}^N \\ \dot{u}_k = \mathcal{B}_k(u) + \nu \bar{k} u_k + F_k; & k \notin \mathcal{K}^N. \end{cases}$$
(4.10.1)

that is the same as system (4.9.3) with \mathcal{K}^N as the finite set of controlled modes.

Definition 4.10.2. We shall say that system [(4.10.1).N] is **time**-T solidly controllable in observed component if for any $u_0 \in V$ and R > 0 there exists a family

$$\mathcal{V}_{u_0,R} := \{ v_b \in L^{\infty}([0, T], \mathbb{R}^{\kappa_N}) \mid b \in B_{u_0,R} \}$$

such that $\mathbb{E}_T(u_0, B_{u_0,R}) := \mathbb{E}_T(u_0, \mathcal{V}_{u_0,R})$ covers $\overline{\mathcal{O}}_R(u_0^{\#\mathcal{O}})$ solidly. Where, by $y^{\#\mathcal{O}}$ we mean the projection of y onto $\mathbb{R}^{\#\mathcal{O}} = \mathcal{O}$; $B_{u_0,R}$ is an open relatively compact subset of a C^0 -manifold and; $\overline{\mathcal{O}}_R(y)$ is the closed ball

$$\{x \in \mathcal{O} \mid ||x - y||_{l_1} \le R\} := \{x \in \mathbb{R}^{\#\mathcal{O}} \mid ||x - y||_{l_1} \le R\}.$$

Below we will need also open balls we define them by

$$\mathcal{O}_R(y) = \{ x \in \mathcal{O} \mid \|x - y\|_{l_1} < R \} := \{ x \in \mathbb{R}^{\#\mathcal{O}} \mid \|x - y\|_{l_1} < R \}.$$

Proposition 4.10.1. System [(4.10.1).1] is time-T solidly controllable in observed component.

Remark 18. Proposition 4.10.1 implies controllability in observed component and, it follows from Proposition 4.10.2 (with N = 1) below. Indeed given R > 0and $u_0 \in V$, if $T \leq T^0$ it is included in the statement of Proposition 4.10.2 (with N = 1), otherwise if $T > T^0$ we apply any control $\bar{v} \in L^{\infty}([0, T], \mathbb{R}^{\kappa_1})$ (for example $\bar{v} = 0$ — no control) up to time $T - T^0$ arriving to some point $y \in V$. Put $\bar{R} := R + ||y^{\kappa_1} - u_0^{\kappa_1}||$. Then apply first part of Proposition 4.10.2 (with N = 1 and $T = T^0$) to the pair $(y, \bar{R}) \in V \times]0, +\infty[$. The family $\mathcal{V}_{y,\bar{R}} \circ \bar{v}$ will do.

Proposition 4.10.2.

- 1. For some $T^0 > 0$, every $0 < T \leq T^0$ and every $N \in \mathbb{N}_0$ the system [(4.10.1).N] is time-T solid controllable in observed component;
- 2. For each pair $(u_0, R) \in V \times [0, +\infty[$ the family

$$\mathcal{V}_{u_0,R} := \{ v_b \mid b \in B_{u_0,R} \}$$

can be chosen satisfying:

- The map $b \mapsto v_b$ is $(B, L^2(0, T, \mathbb{R}^{\kappa_N}))$ -continuous and;
- The controls $v_b(t)$ are uniformly (w.r.t. b and t) l_1 -bounded:

$$||v_b(t)||_{l_1} \leq A = A(T, R, u_0)$$

Since \mathcal{K}^1 is saturating we have $\mathcal{O} \subseteq \mathcal{K}^M$ from some $M \in \mathbb{N}_0$. Fix M with this property. We shall prove Proposition 4.10.2 in two steps. Prove it in the case $N \geq M$ and prove the "back-induction" step " it holds for N implies it holds for N - 1" $(N = 2, \ldots, M)$. These steps are the following subsections 4.10.1 and 4.10.2.

4.10.1 First Step. Proposition 4.10.2: N Big

In this subsection we shall prove that the statement of Proposition 4.10.2 holds for $N \ge M$.

Decompose $u \in V$ as $u = u^{\kappa_N} + U^{\kappa_N}$ where $u^{\kappa_N} := P^{\kappa_N} u$, i.e., u^{κ_N} is the projection of u onto \mathbb{R}^{κ_N} . So $U^{\kappa_N} = P^{-\kappa_N} u$ is the projection of u onto $(\mathbb{R}^{\kappa_N})_V^{\perp}$. Now write system [(4.10.1).N] as

$$u_t^{\kappa_N} = P^{\kappa_N} (-\nu A u - P^{\nabla} B u + F + v)$$
$$U_t^{\kappa_N} = P^{-\kappa_N} (-\nu A u - P^{\nabla} B u + F + v),$$

i.e.,

$$\begin{split} u_t^{\kappa_N} &= -\nu A u^{\kappa_N} - P^{\kappa_N} P^{\nabla} B u + P^{\kappa_N} F + v =: g(u) + v \\ U_t^{\kappa_N} &= -\nu A U^{\kappa_N} - P^{-\kappa_N} P^{\nabla} B u + P^{-\kappa_N} F =: G(u). \end{split}$$

Concisely we have

$$u_t^{\kappa_N} = g(u) + v, \qquad U_t^{\kappa_N} = G(u) \qquad t \in [0, T].$$
 (4.10.2)

Let (u_0, R) be an element of $V \times]0, +\infty[$. Fix $\gamma > 1$. Let T > 0 be a positive real number.

For each $p \in \gamma \mathcal{O}_R(0) \subseteq \mathbb{R}^{\kappa_N}$ define, on [0, T] the constant control

$$v_p(t) := T^{-1}p.$$

Since $\gamma > 1$ we have that $\overline{\mathcal{O}}_R(0) \subseteq \gamma \mathcal{O}_R(0)$ and, $\int_0^T v_p(t) dt = \int_0^T T^{-1}p dt = p$. The family $\mathcal{V}_p := \{v_p \mid p \in \gamma \mathcal{O}_R(0)\}$ is parametrized continuously in L^q -norm (q > 0), i.e., the map $p \mapsto v_p$ is $(l_1, L^q(0, T, \mathbb{R}^{\#\mathcal{O}}))$ -continuous. Indeed for q > 0:

$$||T^{-1}p_2 - T^{-1}p_1||_{L^q}^q = \int_0^T T^{-q} ||p_2 - p_1||_{l_1}^q = T^{1-q} ||p_2 - p_1||_{l_1}^q$$

Hence

$$||T^{-1}p_2 - T^{-1}p_1||_{L^q} = T^{\frac{1-q}{q}} ||p_2 - p_1||_{l_1}$$

For $q = \infty$: $||T^{-1}p_2 - T^{-1}p_1||_{L^{\infty}} = T^{-1}||p_2 - p_1||_{l_1}$. We can also see that $||v_p(t)||_{l_1} < T^{-1}\gamma R$. To prove that the family \mathcal{V}_p is

We can also see that $||v_p(t)||_{l_1} < T^{-1}\gamma R$. To prove that the family \mathcal{V}_p is the one we are looking for it remains to check that $\mathbb{E}_T(u_0, \mathcal{V}_p)$ covers $\overline{\mathcal{O}}_R(u_0^{\#\mathcal{O}})$ solidly.

By the (l_1, L^2) -continuity of $p \mapsto v_p$ and $(L^2, C([0, T], V))$ -continuity of $v \mapsto S_s(u_0, F, v, \nu)$ we conclude the (l_1, l_1) -continuity of

$$p \mapsto \Pi_{\mathcal{O}} \circ \mathbb{S}_s(u_0, F, v_p, \nu)(T) = \mathbb{E}_T(u_0, v_p).$$

Rescaling time: $t = T\xi$, $\xi \in [0, 1]$. From (4.10.2) we obtain the system

$$u_{\xi}^{\kappa_N} = T(g(u) + v)$$
 $U_{\xi}^{\kappa_N} = TG(u)$ $u(0) = u_0, \, \xi \in [0, \, 1].$ (4.10.3)

which solutions, for $v = v_p$, will be compared with those of the following system:

$$y_{\xi}^{\kappa_N} = p$$
 $Y_{\xi}^{\kappa_N} = 0;$ $y(0) = y_0, \ \xi \in [0, 1].$ (4.10.4)

Put z = u - y. Then z satisfies

$$z_{\xi}^{\kappa_N} = T(g(u) + v_p) - p$$
 $Z_{\xi}^{\kappa_N} = TG(u)$ $\xi \in [0, 1],$

i.e.,

$$\frac{1}{T} z_{\xi}^{\kappa_N} = -\nu A u^{\kappa_N} - P^{\kappa_N} P^{\nabla} B u + P^{\kappa_N} F$$
$$\frac{1}{T} Z_{\xi}^{\kappa_N} = -\nu A U^{\kappa_N} - P^{-\kappa_N} P^{\nabla} B u + P^{-\kappa_N} F.$$

which is equivalent to

$$\frac{1}{T}z_{\xi} = -\nu Au - P^{\nabla}Bu + F.$$

Multiplying by z we arrive to

$$\frac{1}{2T}\frac{d}{d\xi}|z|^2 \le -\nu((u,z)) + |(P^{\nabla}Bu,z)| + |F||z|$$
$$\le -\nu||z||^2 - \nu((y,z)) + |(Bu,z)| + C_1|F|||z||$$

and, since

$$b(u, u, z) = b(y + z, y + z, z) = b(y + z, y, z)$$

= $b(y, y, z) + b(z, y, z)$

we arrive to

$$\frac{1}{2T}\frac{d}{d\xi}|z|^2 \le -\nu ||z||^2 - \nu((y, z)) + C||y||^2 ||z|| + C|z|||z|||y|| + C_1|F|||z||$$

from which we obtain

$$\frac{1}{T}\frac{d}{d\xi}|z|^2 + \nu ||z||^2 \le C_2 ||y||^2 + C_2 ||y||^4 + C_2 |z|^2 ||y||^2 + C_2 |F|^2.$$

By Gronwall inequality:

$$|z(s)|^{2} \leq \exp\left\{TC_{2}\int_{0}^{1}\|y(\xi)\|^{2} d\xi\right\} \left(|z(0)|^{2} + TC_{2}\int_{0}^{1}\|y(\xi)\|^{2} + \|y(\xi)\|^{4} + |F|^{2} d\xi\right)$$

$$\leq \exp(T)D_{1}(|z(0)|^{2} + TD_{2})$$

where D_1 and D_2 depend only on γ , R and $||y_0||$. Indeed $y(\xi) = y_0 + p\xi$ and $||y_0 + p\xi|| \le ||y_0|| + C ||p\xi||_{l_1}$ and, $||p\xi||_{l_1} < \gamma R$. In particular we have (for fixed u_0, γ and R):

Corollary 4.10.3.

- 1. If $y_0 = u_0$ then $|u y| \le [T \exp(T)]^{\frac{1}{2}} K$
- 2. For "bounded" T and y_0 , say $T \leq T_1$ and $||y_0 u_0|| < \beta$ we have

$$|u-y| \le K (|u_0-y_0|^2+1)^{\frac{1}{2}}$$

with K independent of T (K depends only on γ , R and $||y_0||$).

Note that

$$\mathbb{E}_T(u_0, v_p) = \Pi_{\mathcal{O}} \circ \Phi^T(u_0, p)(1)$$

where $\Phi^T(u_0, p)$ is the solution of system (4.10.3). Represent by $\Phi^0(u_0, p)$ the solution of system (4.10.4). Then $\Pi_{\mathcal{O}} \circ \Phi^0(u_0, p)(1) = u_0^{\#\mathcal{O}} + p$ Now we compute

$$\begin{aligned} \|\Pi_{\mathcal{O}} \circ \Phi^{T}(u_{0}, p)(1) - \Pi_{\mathcal{O}} \circ \Phi^{0}(u_{0}, p)(1)\|_{l_{\infty}} \\ \leq \|\Pi_{\mathcal{O}} \circ \Phi^{T}(u_{0}, p)(1) - \Pi_{\mathcal{O}} \circ \Phi^{0}(u_{0}, p)(1)\|_{l_{1}} \\ \leq C \|\Phi^{T}(u_{0}, p)(1) - \Phi^{0}(u_{0}, p)(1)\|_{H} \leq C \|\Phi^{T}(u_{0}, p) - \Phi^{0}(u_{0}, p)\|_{C([0, 1], H)}. \end{aligned}$$

By item 1 of of corollary 4.10.3:

$$|\Pi_{\mathcal{O}} \circ \Phi^{T}(u_{0}, p)(1) - \Pi_{\mathcal{O}} \circ \Phi^{0}(u_{0}, p)(1)||_{l_{\infty}} \leq CK[T \exp(T)]^{\frac{1}{2}}$$

where K is independent of T and p. Thus

 $\|\Pi_{\mathcal{O}} \circ \Phi^{T}(u_{0}, \cdot)(1) - \Pi_{\mathcal{O}} \circ \Phi^{0}(u_{0}, \cdot)(1)\|_{C(\gamma \mathcal{O}_{R}(0), \mathbb{R}_{\infty}^{\kappa_{N}})} \leq CK[T \exp(T)]^{\frac{1}{2}}$

or, defining in $\gamma \mathcal{O}_R(0)$:

$$\mathbb{G}_T(p) := \mathbb{E}_T(u_0, p); \quad \mathbb{G}_0(p) := \Pi_\mathcal{O} \circ \Phi^0(u_0, p)(1);$$

$$\left\|\mathbb{G}_{T} - \mathbb{G}_{0}\right\|_{C(\gamma \mathcal{O}_{R}(0), \mathbb{R}_{\infty}^{\kappa_{N}})} \leq CK[T\exp(T)]^{\frac{1}{2}}.$$
(4.10.5)

Put T^0 = the unique solution of $T^0 \exp(T^0) = \left(\frac{(\gamma-1)R}{2\#\mathcal{O}CK}\right)^2$. Then

$$\forall T \in]0, T^0] \left[\left\| \mathbb{G}_T - \mathbb{G}_0 \right\|_{C(\gamma \mathcal{O}_R(0), \mathbb{R}_\infty^{\#\mathcal{O}})} \le \frac{R(\gamma - 1)}{2 \# \mathcal{O}} \right].$$
(4.10.6)

Note that the map \mathbb{G}_0 is just the restriction of a translation in $\mathbb{R}^{\#\mathcal{O}} - \mathbb{G}_0(p) = u_0^{\#\mathcal{O}} + p$ — restricted to $\gamma \mathcal{O}_R(0) \equiv \mathcal{O}_{\gamma R}(0)$. By the Degree Theory⁹ we have that

$$\begin{bmatrix} p \notin \partial(\mathbb{G}_0 \mathcal{O}_{\gamma R}(0)) \equiv \partial \mathcal{O}_{\gamma R}(u_0^{\#\mathcal{O}}) \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} deg(\mathbb{G}_0, \mathcal{O}_{\gamma R}(0), p) = deg(I_{\gamma R}, \mathcal{O}_{\gamma R}(0), p - u_0^{\#\mathcal{O}}) = 1 \end{bmatrix}. \quad (4.10.7)$$

Where $I_{\gamma R}$ is the identity function on $\mathcal{O}_{\gamma R}(0)$.

Yet by the Degree Theory we know that for every $p \notin \partial \mathcal{O}_{\gamma R}(u_0^{\#\mathcal{O}})$ and every continuous function $\psi : \mathcal{O}_{\gamma R}(0) \to \mathbb{R}^{\#\mathcal{O}}$ such that

$$\begin{bmatrix} \|\psi - \mathbb{G}_0\|_{C(\mathcal{O}_{\gamma R}(0), \mathbb{R}^{\#\mathcal{O}}_{\infty})} < \|p - \partial \mathcal{O}_{\gamma R}(u_0^{\#\mathcal{O}})\|_{l_{\infty}} \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} deg(\psi, \mathcal{O}_{\gamma R}(0), p) = deg(\mathbb{G}_0, \mathcal{O}_{\gamma R}(0), p) \end{bmatrix}^{10} \quad (4.10.8)$$

We claim that for $T \leq T^0$ we have that $\mathbb{G}_T(\mathcal{O}_{\gamma R}(0))$ covers $\overline{\mathcal{O}}_R(u_0^{\#\mathcal{O}})$ solidly. Indeed, given $T \leq T^0$ and a continuous function $\phi : \mathcal{O}_{\gamma R}(0) \to \mathbb{R}^{\#\mathcal{O}}$ such that $\|\phi - \mathbb{G}_T\|_{C(\mathcal{O}_{\gamma R}(0), \mathbb{R}^{\#\mathcal{O}}_{\infty})} < \frac{(\gamma - 1)R}{2\#\mathcal{O}}$ we have, using (4.10.6),

$$\begin{split} \|\phi - \mathbb{G}_0\|_{C(\mathcal{O}_{\gamma R}(0), \mathbb{R}_{\infty}^{\#\mathcal{O}})} &\leq \|\phi - \mathbb{G}_T\|_{C(\mathcal{O}_{\gamma R}(0), \mathbb{R}_{\infty}^{\#\mathcal{O}})} + \|\mathbb{G}_T - \mathbb{G}_0\|_{C(\mathcal{O}_{\gamma R}(0), \mathbb{R}_{\infty}^{\#\mathcal{O}})} \\ &< \frac{(\gamma - 1)R}{2\#\mathcal{O}} + \frac{(\gamma - 1)R}{2\#\mathcal{O}} = \frac{(\gamma - 1)R}{\#\mathcal{O}}. \end{split}$$

 $^8 Where the subscript "<math display="inline">\infty$ " means that we are considering the "max"-norm — $l_\infty.$ $^9 See,$ for example, [6].

¹⁰Where ||p, A|| means the distance from the element p to the set A, i.e., $||p, A|| := \inf\{||p - a|| \mid a \in A\}$.

For $w \in \overline{\mathcal{O}}_R(u_0^{\#\mathcal{O}})$ we have:

$$\|w - \partial \mathcal{O}_{\gamma R}(u_0^{\#\mathcal{O}})\|_{l_{\infty}} \ge \frac{1}{\#\mathcal{O}} \|w - \partial \mathcal{O}_{\gamma R}(u_0^{\#\mathcal{O}})\|_{l_1} \ge \frac{(\gamma - 1)R}{\#\mathcal{O}}$$

so,

$$\|\phi - \mathbb{G}_0\|_{C(\mathcal{O}_{\gamma R}(0), \mathbb{R}_{\infty}^{\#\mathcal{O}})}\| < \|w - \partial \mathcal{O}_{\gamma R}(u_0^{\#\mathcal{O}})\|_{l_{\infty}}.$$

By (4.10.7) and (4.10.8) we conclude that

$$deg(\phi, \mathcal{O}_{\gamma R}(0), w) = 1.$$

which means, in particular, that the equation $\phi(y) = w$ has a solution on $\mathcal{O}_{\gamma R}(0)$, i.e., $\phi(\mathcal{O}_{\gamma R}(0))$ covers $\overline{\mathcal{O}}_{R}(u_{0}^{\#\mathcal{O}})$.

4.10.2 Second Step. Proposition 4.10.2: "Back-Induction".

In this subsection we "imitate" a driving using controls on \mathbb{R}^{κ_N} by a driving using controls on $\mathbb{R}^{\kappa_{N-1}}$, $N = 2, \ldots, M$, M is fixed and satisfies $\mathcal{O} \subseteq \mathcal{K}^M$. Both drivings leading to the same projection onto $\mathbb{R}^{\kappa_{N-1}}$ at final time but, possibly going by paths with projections "far from each other" in the middle. The projection onto the orthogonal space $(\mathbb{R}^{\kappa_N})^{\perp}_V$ of the paths will be *H*-close to each other so, at time *T* the two drivings lead to points close in *H*-metric. Hence the end points of the projection onto the finite dimensional observed space \mathcal{O} are close. Solid controllability will follow from this closeness and (again) from a Degree Theory argument.

Such imitation is then the key for the prove that if the system [(4.10.1).N] is solid controllable in observed component then so is system [(4.10.1).N-1].

After we prove this " $N \rightarrow N - 1$ " step it will be clear, from the fact that [(4.10.1).M] is solid controllable in observed component (see subsection 4.10.1), that system (4.9.3) is solid controllable in observed component, we just note that the systems [(4.10.1).1] and (4.9.3) are the same system.

To prove the "back-induction" step " $N \rightarrow N-1$ " we shall need some lemmas: For the next Lemma we may consider again the case where the external force depend on time:

Lemma 4.10.4. Given:

- A finite subset $J \subset \mathbb{N}_0^2$; $\mathbb{J} := span\{W_k \mid k \in J\}$
- A function $q \in W^{1,\infty}([t_i, t_f], \mathbb{J})$, such that $q(t_i) = q_i$
- An element $Q_i \in \mathbb{J}_V^{\perp}$, the orthogonal space to \mathbb{J} in V.

Then there exists a control $v^J(q, Q_i) \in L^{\infty}([t_i, t_f], \mathbb{J})$ depending on q and Q_i such that the projection onto \mathbb{J} of the solution of the NSE

$$u_t = -\nu Au - P^{\nabla} Bu + \tilde{F} + v^J(q, Q_i), \qquad u(t_i) = q_i + Q_i$$

equals q on $[t_i, t_f]$.

Moreover the map $v^J : (q, Q_i) \mapsto v^J(q, Q_i)$ is $(W^{1,2} \times \mathbb{J}_V^{\perp}, L^2(t_i, t_f, \mathbb{J})$ -continuous.

Proof. Let q, q_i, Q_i and \mathbb{J} be like in the statement of the Lemma. Consider the (non controlled) NSE: $u_t = -\nu Au - P^{\nabla} Bu + \tilde{F}$ with initial condition $u(t_i) = q_i + Q_i =: u_i$ and split it into

$$P^{J}(u_{t}) = P^{J}(-\nu Au - P^{\nabla}Bu + \tilde{F})$$
$$P^{-J}(u_{t}) = P^{-J}(-\nu Au - P^{\nabla}Bu + \tilde{F})$$
$$u(t_{i}) = u_{i}$$

with the same initial condition, where P^J (resp. P^{-J}) is the projection $H \to \mathbb{J}$ (resp. $H \to \mathbb{J}_H^{\perp}$). If we put $u^J := P^J u$ and $U^J := P^{-J} u$ we arrive to the systems

$$\begin{cases} u_t^J = -\nu A u^J - P^J P^{\nabla} B u + P^J \tilde{F} \\ u^J(t_i) = q_i \end{cases}$$

$$\tag{4.10.9}$$

$$\begin{cases} U_t^J = -\nu A U^J - P^{-J} P^{\nabla} B u + P^{-J} \tilde{F} \\ U^J(t_i) = Q_i. \end{cases}$$
(4.10.10)

In the system (4.10.10), for each $k \in \mathbb{J}$ replace $u_k(t)$ by $q_k(t)$ arriving, in this way, to the (closed) system

$$\begin{cases} U_t^J = -\nu A U^J - P^{-J} P^{\nabla} B (U^J + q) + P^{-J} \tilde{F} \\ U^J(t_i) = Q_i. \end{cases}$$
(4.10.11)

We can prove existence and uniqueness of a strong solution for this system as we prove existence and uniqueness for the "full" equation. We indicate only how to find some estimates: Starting from approximate solutions

$$U^{J,L} := \sum_{\substack{k \in \mathcal{K}^{L} \setminus \mathbb{J} \\ \mathbb{J} \subseteq \mathcal{K}^{L}}} U^{J,L}_{k} W_{k}$$

$$(U^{J,L}_{t}, W_{k}) = -\nu(AU^{J,L}, W_{k}) - (P^{-J}P^{\nabla}B(U^{J,L} + q), W_{k}) + (P^{-J}\tilde{F}, W_{k})$$

$$= -\nu((U^{J,L}, W_{k})) - (B(U^{J,L} + q), W_{k}) + (\tilde{F}, W_{k}) \qquad (4.10.12)$$

$$\forall k \in \mathcal{K}^{L} \setminus \mathbb{J} \qquad (4.10.13)$$

$$U^{J,L}_{k}(t) = O^{\kappa_{L}-J} \qquad \text{projection of } O \quad \text{erts} \ \mathbb{D}^{\kappa_{L}-J}$$

 $U^{J,L}(t_i) = Q_i^{\kappa_L - J} =$ projection of Q_i onto $\mathbb{R}^{\kappa_L - J}$,

from which we obtain the ODE

$$-\bar{k}\frac{ab}{4}\dot{U}_{k}^{J,L} = -\nu\bar{k}^{2}\frac{ab}{4}U_{k}^{J,L} - \sum_{m,n\in\mathcal{K}^{L}}u_{m}^{J,L}u_{n}^{J,L}b(W_{m}, W_{n}, W_{k}) - \bar{k}\frac{ab}{4}\tilde{F}_{k}$$
$$U_{i}^{J,L} = Q_{i}^{\kappa_{L}-J}$$

that has a maximal solution defined on $[t_i, t_{max}]$. Now we compute some estimates that, in particular, imply $t_{max} = t_f$:

Multiplying, for each k, the equation (4.10.12) by $U_k^{J,L}$ and summing up we obtain:

$$\frac{1}{2}\frac{d}{dt}|U^{J,L}|^2 \leq -\nu \|U^{J,L}\|^2 + |(B(U^{J,L}+q), U^{J,L})| + |\tilde{F}||U^{J,L}|$$

and, after simplify the term with B, using some estimates from section 2.9 and Young inequalities, we obtain

$$\frac{d}{dt}|U^{J,L}|^2 + \nu \|U^{J,L}\|^2 \le C|U^{J,L}|^2 \|q\|^2 + C\|q\|^4 + C|\tilde{F}|^2$$

from which we can conclude that for every $s \in [t_i, t_f]$:

$$|U^{J,L}(s)|^{2} \leq \exp \int_{t_{i}}^{t_{f}} C ||q(t)||^{2} dt \left(|Q_{i}|^{2} + C \int_{t_{i}}^{t_{f}} ||q(t)||^{4} + |\tilde{F}(t)|^{2} dt \right)$$

$$\leq D_{1}(||Q_{i}||^{2} + 1)$$

where D_1 can be taken depending only in $||q||_{C([t_i, t_f], \mathbb{J})}$ ¹¹. So for a constant C_1 depending only in $||q||_{C([t_i, t_f], \mathbb{J})}$ and $||Q_i||$ we have

$$\|U^{J,L}\|_{L^{\infty}(t_i, t_f, \mathbb{J}_H^{\perp})} \le C_1.$$
(4.10.14)

We also have

$$\int_{t_i}^{t_f} \|U^{J,L}(t)\|^2 dt \le D_3 \|U^{J,L}\|_{L^{\infty}(t_i, t_f, \mathbb{J}_H^{\perp})}^2 \left(1 + \int_{t_i}^{t_f} \|q(t)\|^2 dt\right) + D_3 \int_{t_i}^{t_f} \|q(t)\|^4 + |\tilde{F}(t)|^2 dt.$$

Hence for a constant C_2 depending only in $\|q\|_{C([t_i, t_f], \mathbb{J})}$ and $\|Q_i\|$ we have

$$\|U^{J,L}\|_{L^2(t_i, t_f, \mathbb{J}_V^{\perp})} \le C_2. \tag{4.10.15}$$

From (4.10.14) and (4.10.15) we have that

$$\begin{cases} (U^{J,L})_L \text{ remains in a bounded subset of } L^{\infty}([t_i, t_f], H) \\ (U^{J,L})_L \text{ remains in a bounded subset of } L^2(t_i, t_f, V). \end{cases}$$
(4.10.16)

Analogously if we multiply, for each k, the equation (4.10.12) by $-\bar{k}U_k^{J,L}$ and summing up we obtain:

$$\frac{1}{2}\frac{d}{dt}\|U^{J,L}\|^2 \le -\nu|U^{J,L}|^2_{[2]} + |(B(U^{J,L}+q), AU^{J,L})| + |\tilde{F}||U^{J,L}|_{[2]}$$

and,

$$\frac{d}{dt}\|U^{J,L}\|^2 + \nu |U^{J,L}|^2_{[2]} \le C|U^{J,L}|^2 \|U^{J,L}\|^4 + C|q|^4_{[2]} + C|q|^2_{[2]} \|U^{J,L}\|^2 + C|\tilde{F}|^2$$

from which, using (4.10.14) and (4.10.15), we conclude that for some constants C_3 and C_4 depending only in $||q||_{C([t_i, t_f], \mathbb{J})}$ and $||Q_i||$:

$$\|U^{J,L}\|_{L^{\infty}(t_i, t_f, \mathbb{J}_V^{\perp})} \le C_3; \tag{4.10.17}$$

$$\|U^{J,L}\|_{L^2(t_i, t_f, \mathbb{J}_{D(A)}^{\perp})} \le C_4.$$
(4.10.18)

 $^{{}^{11}\}tilde{F}$ is fixed.

 $\begin{cases} (U^{J,L})_L \text{ remains in a bounded subset of } L^{\infty}([t_i, t_f], V) \\ (U^{J,L})_L \text{ remains in a bounded subset of } L^2(t_i, t_f, D(A)) \end{cases}$ (4.10.19)

The existence of a strong solution follows some classical compactness theorems. For the uniqueness we consider the difference w of two solutions $V^J = V^J(q, Q_i)$ and $U^J = (q, Q_i) - w := V^J - U^J$. Then from

$$w_t = -\nu Aw - P^{-J} P^{\nabla} B(V^J + q) + P^{-J} P^{\nabla} B(U^J + q),$$

multiplying by w, using some estimates from section 2.9 and appropriate inequalities, we obtain

$$\frac{d}{dt}|w|^2 + \nu ||w||^2 \le C|w|^2 ||U^J + q||^2.$$

So, $|w(s)|^2 \leq |w(t_i)|^2 \exp \int_{t_i}^{t_f} C ||U^J(t) + q(t)|| dt = 0$. Then a weak solution is unique and, so is the strong one.

We have just proved that the map $(q, Q_i) \mapsto U^J(q, Q_i)$ is well defined. We claim that it is $(L^4(t_i, t_f, \mathbb{J}) \times (\mathbb{J}_V^{\perp}), X)$ -continuous, where X is either $L^{\infty}([t_i, t_f], \mathbb{J}_V^{\perp})$ or $L^2(t_i, t_f, \mathbb{J}_{D(A)}^{\perp})$. To prove these continuities we proceed as usually: Fix a pair $(q, Q_i) \in (W^{1,\infty} \times \mathbb{J}_V^{\perp})$ and consider another one (p, P_i) in the same product space. Define $w := U^J(q, Q_i) - U^J(p, P_i)$. Then we obtain the equation for w:

$$\dot{w} = -\nu Aw - P^{-J}P^{\nabla}B(U^{J}(q, Q_{i}) + q) + P^{-J}P^{\nabla}B(U^{J}(p, P_{i}) + p).$$

To simplify the writing we put $Q := U^J(q, Q_i)$ and $P := U^J(p, P_i)$ so,

$$\dot{w} = -\nu A w - P^{-J} P^{\nabla} B(Q+q) + P^{-J} P^{\nabla} B(P+p)$$
(4.10.20)

and, multiplying by w:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |w|^2 &\leq -\nu ||w||^2 + |b(P, P, w) - b(Q, Q, w)| + |b(P, p, w) - b(Q, q, w)| \\ &+ |b(p, P, w) - b(q, Q, w)| + |b(p, p, w) - b(q, q, w)| \\ &\leq -\nu ||w||^2 + |b(w, w, Q)| + \left\{ |b(-w, p, w) + b(Q, p - q, w)| \right\} \\ &+ |b(p - q, Q, w)| + \left\{ |b(p - q, p, w) + b(q, p - q, w)| \right\} \end{aligned}$$

from which we obtain

$$\frac{d}{dt}|w|^{2} + \nu||w||^{2} \leq C|w|^{2}||Q||^{2} + C||Q||^{2}||p - q||^{2} + C|w|^{2}||p||^{2} + C(||p||^{2} + ||q||^{2})||p - q||^{2}.$$
(4.10.21)

Then by Gronwall Inequality

$$\begin{split} \|w\|_{C([t_i, t_f], \mathbb{J}_H^{\perp})}^2 &\leq \exp\left[C\int_{t_i}^{t_f} \|Q(t)\|^2 + \|p(t)\|^2 \, dt\right] \left(|w(t_i)|^2 \\ &+ C\int_{t_i}^{t_f} \|p - q\|^2 (\|Q(t)\|^2 + \|p(t)\|^2 + \|q(t)\|^2) \, dt\right) \end{split}$$

For $||p-q||_{L^4} \leq 1$ we obtain¹²

$$\|w\|_{C([t_i, t_f], \mathbb{J}_H^{\perp})}^2 \le C_1 \|Q_i - P_i\|^2 + C_1 \|p - q\|_{L^4}^2$$
(4.10.22)

and conclude that the map U^J is $(L^4 \times \mathbb{J}_V^{\perp}, C([t_i, t_f], \mathbb{J}_H^{\perp}))$ -continuous.

From (4.10.21) we obtain

$$\begin{split} \int_{t_i}^{t_f} \|w(t)\|^2 \, dt &\leq C_0 \|w\|_{C([t_i, t_f], \mathbb{J}_H^{\perp})}^2 \left(1 + \int_{t_i}^{t_f} \|Q(t)\|^2 + \|p(t)\|^2 \, dt\right) \\ &+ C_0 \|p - q\|_{L^4}^2 \left(\|Q\|_{L^4}^2 + \|p\|_{L^4}^2 + \|q\|_{L^4}^2\right) \\ &\leq C_2 \|w\|_{C([0, T], \mathbb{J}_H^{\perp})}^2 + C_2 \|p - q\|_{L^4}^2, \end{split}$$

for $||p-q||_{L^4} \leq 1$. Hence by (4.10.22) we arrive to

$$\left[\|p-q\|_{L^4} \le 1\right] \Rightarrow \left[\|w\|_{L^2(t_i, t_f, \mathbb{J}_V^\perp)}^2 \le C_3 \|Q_i - P_i\|^2 + C_3 \|p-q\|_{L^4}^2\right].$$
(4.10.23)

and conclude that the map U^J is $(L^4 \times \mathbb{J}_V^{\perp}, L^2(t_i, t_f, \mathbb{J}_V^{\perp}))$ -continuous. Now from equation (4.10.20), multiplying it by Aw, we obtain

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|w\|^2 + \nu |w|_{[2]}^2 \le |(B(P+p), Aw) - (B(Q+q), Aw)| \\ &\le |b(P, P, Aw) - b(Q, Q, Aw)| + |b(P, p, Aw) - b(Q, q, Aw)| \\ &+ |b(p, P, Aw) - b(q, Q, Aw)| + |b(p, p, Aw) - b(q, q, Aw)|. \end{aligned}$$

Note that

$$\begin{array}{ll} b(P,\,P,\,Aw) - b(Q,\,Q,\,Aw) &= b(w,\,w,\,Aw) - b(w,\,Q,\,Aw) - b(Q,\,w,\,Aw) \\ b(P,\,p,\,Aw) - b(Q,\,q,\,Aw) &= -b(w,\,p,\,Aw) + b(Q,\,p-q,\,Aw) \\ b(p,\,P,\,Aw) - b(q,\,Q,\,Aw) &= -b(p,\,w,\,Aw) + b(p-q,\,Q,\,Aw) \\ b(p,\,p,\,Aw) - b(q,\,q,\,Aw) &= b(p-q,\,p,\,Aw) + b(q,\,p-q,\,Aw). \end{array}$$

Hence from (4.10.24), from the estimates of section 2.9 and from the continuity of the inclusions $D(A) \mapsto V \mapsto H$ we arrive to

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w\|^2 + \nu \|w\|_{[2]}^2 &\leq C \Big(\|w\|^{\frac{1}{2}} \|w\| \|w\|_{[2]}^{\frac{3}{2}} + \|w\|^{\frac{1}{2}} \|Q\| \|w\|_{[2]}^{\frac{3}{2}} \Big) \\ &+ C \Big(\|w\| \|p\|_{[2]} \|w\|_{[2]} + \|Q\| \|p-q\|_{[2]} \|w\|_{[2]} \Big) + C \Big(\|p\|_{[2]} \|w\| \|w\|_{[2]} + \|p-q\|_{[2]} \|Q\| \|w\|_{[2]} \Big) \\ &+ C \Big(\|p-q\|_{[2]} \|p\| \|w\|_{[2]} + \|q\| \|p-q\|_{[2]} \|w\|_{[2]} \Big). \end{aligned}$$

By appropriate Young inequalities:

$$\frac{d}{dt} \|w\|^{2} + \nu \|w\|_{[2]}^{2} \leq D\left(\|w\|^{2} \|w\|^{4} + \|w\|^{2} \|Q\|^{4}\right) + D\left(\|w\|^{2} |p|_{[2]}^{2} + \|Q\|^{2} |p-q|_{[2]}^{2}\right) \\
+ D\left(\|p\|_{[2]}^{2} \|w\|^{2} + |p-q|_{[2]}^{2} \|Q\|^{2}\right) + D\left(\|p-q\|_{[2]}^{2} \|p\|^{2} + \|q\|^{2} |p-q|_{[2]}^{2}\right). \tag{4.10.25}$$

¹²Of course we could ask for $||p-q||_{L^4} \leq D$ for any D > 0. What we need is a first bound for $||p||_{L^4}$.

Thus

$$\begin{split} \|w\|_{C([t_i, t_f], \mathbb{J}_V^{\perp})}^2 &\leq \exp\Big[D\int_{t_i}^{t_f} |w(t)|^2 \|w(t)\|^2 + \|Q(t)\|^4 + |p(t)|_{[2]}^2 dt\Big] \Big(\|w(t_i)\|^2 \\ &+ D\int_{t_i}^{t_f} |p(t) - q(t)|_{[2]}^2 (\|Q(t)\|^2 + \|p(t)\|^2 + \|q(t)\|^2) dt\Big). \end{split}$$

Then using (4.10.22) and (4.10.23)

$$\begin{cases} \|p-q\|_{L^4} \leq 1\\ \|P_i - Q_i\|_{\mathbb{J}^{\perp}_V} \leq 1 \end{cases} \Rightarrow \Big[\|w\|^2_{C([t_i, t_f], \mathbb{J}^{\perp}_V)} \leq D_1 \|Q_i - P_i\|^2 + D_1 \|p-q\|^2_{L^4} \Big]. \end{cases}$$

$$(4.10.26)$$

and conclude that the map U^J is $(L^4 \times (\mathbb{J}_V^{\perp}), C([t_i, t_f], \mathbb{J}_V^{\perp}))$ -continuous. (4.10.26)

From (4.10.25) we can also obtain

$$\begin{split} & \int_{t_i}^{t_f} |w(t)|_{[2]}^2 dt \\ & \leq D_2 \|w\|_{C([0,T], \mathbb{J}_V^{\perp})}^2 \Big(1 + \int_{t_i}^{t_f} |w(t)|^2 \|w(t)\|^2 + \|Q(t)\|^4 + |p(t)|_{[2]}^2 dt \Big) \\ & + D_2 \|p - q\|_{L^4}^2 \Big(\|Q\|_{L^4}^2 + \|p\|_{L^4}^2 + \|q\|_{L^4}^2 \big). \end{split}$$

Hence by (4.10.22) and (4.10.23) we have

$$\begin{cases} \|p-q\|_{L^4} \leq 1 \\ \|P_i - Q_i\|_{\mathbb{J}_V^{\perp}} \leq 1 \\ \Rightarrow \left[\|w\|_{L^2(t_i, t_f, \mathbb{J}_{D(A)}^{\perp})}^2 \leq D_3 \|w\|_{C([t_i, t_f], \mathbb{J}_V^{\perp})}^2 + D_3 \|p-q\|_{L^4}^2 \right] \end{cases}$$

and, by (4.10.26) we obtain

$$\begin{cases} \|p-q\|_{L^4} \leq 1 \\ \|P_i - Q_i\|_{\mathbb{J}_V^\perp} \leq 1 \end{cases} \Rightarrow \left[\|w\|_{L^2(t_i, t_f, \mathbb{J}_{D(A)}^\perp)}^2 \leq D_4 \|Q_i - P_i\|^2 + D_4 \|p-q\|_{L^4}^2 \right] \\ (4.10.27)\end{cases}$$

and conclude that the map U^J is $(L^4 \times (\mathbb{J}_V^{\perp}), L^2(t_i, t_f, D(A)))$ -continuous.

Now we define another map on the product $W^{1,\infty}([t_i, t_f], \mathbb{J}) \times \mathbb{J}_V^{\perp}$ taking values on H:

$$\Gamma^J: (q, Q_i) \mapsto -\nu A(U^J(q, Q_i) + q) - P^{\nabla} B(U^J(q, Q_i) + q) + \tilde{F}$$

Fix $(q, Q_i) \in W^{1,2}([t_i, t_f], \mathbb{J}) \times \mathbb{J}_V^{\perp}$ and, consider another pair (p, P_i) in the same space. Again, as we have done before put $Q = U^J(q, Q_i)$ and $P = U^J(p, P_i)$.

We compute the norm of the difference

$$\begin{split} |\Gamma^{J}(q, Q_{i}) - \Gamma^{J}(p, P_{i})| \\ = & |\nu A(P + p) + P^{\nabla}B(P + p) - \nu A(Q + q) - P^{\nabla}B(Q + q)| \\ \leq & |\nu A(P + p - Q - q)| + |P^{\nabla}[B(P + p) - B(Q + q)]| \\ \leq & |\nu A(P + p - Q - q)| + |B(P + p) - B(Q + q)| \\ \leq & \nu |P - Q|_{[2]} + \nu |p - q|_{[2]} + |BP - BQ| \\ + & |B(P, p) - B(Q, q)| + |B(p, P) - B(q, Q)| + |Bp - Bq| \\ \leq & \nu |P - Q|_{[2]} + \nu |p - q|_{[2]} + C|P - Q|_{[2]}(||P|| + ||Q||) \\ + & C \Big(|P - Q|_{[2]} ||q|| + ||P|||p - q|_{[2]} \Big) \\ + & C \Big(||P|||p - q|_{[2]} + |P - Q|_{[2]} ||q|| \Big) \\ + & C (||P|||p - q|_{[2]} + ||P - Q|_{[2]} ||q|| \Big) \end{split}$$

Therefore

$$\begin{aligned} |\Gamma^{J}(q, Q_{i}) - \Gamma^{J}(p, P_{i})| \\ \leq \nu |P - Q|_{[2]} + \nu |p - q|_{[2]} + C_{1} |P - Q|_{[2]} \Big(||P|| + ||Q|| + ||q|| \Big) \\ + C_{1} |p - q|_{[2]} \Big(||P|| + ||p|| + ||q|| \Big) \\ \leq C_{0} |P - Q|_{[2]} \Big(1 + ||P|| + ||Q|| + ||q|| \Big) + C_{0} |p - q|_{[2]} \Big(1 + ||P|| + ||p|| + ||q|| \Big). \end{aligned}$$

For $||p-q||_{L^4} \le 1$, and $||P_i-Q_i|| \le 1$, using (4.10.26) we obtain $||P(t)-Q(t)||^2 \le 2D_1$ and then $||P(t)|| \le \sqrt{2D_1} + ||Q(t)||$. So we can arrive to

$$|\Gamma^{J}(q, Q_{i}) - \Gamma^{J}(p, P_{i})| \le C_{2}|P - Q|_{[2]} + C_{2}|p - q|_{[2]}\left(1 + ||p||\right)$$
(4.10.28)

Note that we have used the fact that $q \in W^{1,\infty}$ and so, ||q|| is bounded. But we can not "replace" ||p|| by a constant in the last member because, unlike as q, p is not fixed and we are considering that p varying in L^4 topology.

$$\begin{split} &\int_{t_i}^{t_f} |\Gamma^J(q, Q_i) - \Gamma^J(p, P_i)|^2 dt \\ \leq & 2C_2^2 \int_{t_i}^{t_f} |P - Q|_{[2]}^2 dt + 2C_2^2 \int_{t_i}^{t_f} |p - q|_{[2]}^2 \Big(1 + \|p\|\Big)^2 dt \\ \leq & 2C_2^2 \|P - Q\|_{L^2(t_i, t_f, D(A))}^2 + 2C_2^2 \int_{t_i}^{t_f} |p - q|_{[2]}^2 \Big(1 + \|p\|^2\Big) dt \\ \leq & C_4 \|P - Q\|_{L^2(t_i, t_f, D(A))}^2 + C_4 \|p - q\|_{L^4}^2 (1 + \|p\|_{L^4}^2). \end{split}$$

Therefore, using (4.10.27):

$$\begin{cases} \|p-q\|_{L^4} \leq 1 \\ \|P_i - Q_i\|_{\mathbb{J}_V^{\perp}} \leq 1 \\ \Rightarrow \|\Gamma^J(q, Q_i) - \Gamma^J(p, P_i)\|_{L^2(t_i, t_f, H)}^2 dt \leq C_5 \|P_i - Q_i\|^2 + C_5 \|p-q\|_{L^4}^2 \\ (4.10.29) \end{cases}$$

and we conclude the $(L^4 \times (\mathbb{J}_V^{\perp}), L^2(t_i, t_f, H))$ -continuity of Γ^J .

Now we can indicate which is the control v^J appearing in the statement of the proposition: In fact

$$v^J(q, Q_i) := \dot{q} - P^J \Gamma^J(q, Q_i)$$

satisfies the statement. Indeed its $(W^{1,2}([t_i, t_f], \mathbb{J}) \times (\mathbb{J}_V^{\perp}), L^2(t_i, t_f, \mathbb{J})$ -continuity follows from the $(L^4 \times (\mathbb{J}_V^{\perp}), L^2(t_i, t_f, H)$ -continuity of Γ^J and from the $(W^{1,2}([t_i, t_f], \mathbb{J}), L^2(t_i, t_f, \mathbb{J})$ -continuity of $q \mapsto \dot{q}$.

To prove that the projection of the solution of the system

$$u_t = -\nu A u - B u + \tilde{F} + v^J(q, Q_i), \qquad u_i = q(t_i) + Q_i$$
(4.10.30)

coincides with q we differentiate $q + U^J(q, Q_i)$ obtaining

$$\begin{split} [q + U^{J}(q, Q_{i})]_{t} &= \dot{q} - \nu A(U^{J}(q, Q_{i})) - P^{-J}P^{\nabla}B(U^{J}(q, Q_{i}) + q) + P^{-J}\tilde{F} \\ &= -\nu A(q + U^{J}(q, Q_{i})) - P^{\nabla}B(q + U^{J}(q, Q_{i})) + \tilde{F} \\ &+ \dot{q} - (-\nu Aq - P^{J}P^{\nabla}B(q + U^{J}(q, Q_{i})) + P^{J}\tilde{F}) \\ &= -\nu A(q + U^{J}(q, Q_{i})) - P^{\nabla}B(q + U^{J}(q, Q_{i})) + \tilde{F} + v^{J}(q, Q_{i})) \end{split}$$

showing that $q + U^{J}(q, Q_{i})$ is the (unique) solution of (4.10.30).

To finish the prove remains to verify that $v^J \in L^{\infty}([t_i, t_f], \mathbb{J})$. Since $q \in W^{1,\infty}([t_i, t_f], \mathbb{J})$ we have $\dot{q} \in L^{\infty}([t_i, t_f], \mathbb{J})$ and by

$$\|\Gamma^{J}(q, Q_{i})\|_{V'} \leq \nu \|Q + q\| + \|Q + q\|^{2} + \|\tilde{F}\|_{V'}$$

$$\leq C \Big(\|Q + q\| + \|Q + q\|^{2} + |\tilde{F}| \Big) \leq C_{1}.$$
(4.10.31)

Hence $\Gamma^J(q, Q_i) \in L^{\infty}([t_i, t_f], V')$ and then, $P^J \Gamma^J(q, Q_i) \in L^{\infty}([t_i, t_f], P^J V')$, i.e., $P^J \Gamma^J(q, Q_i) \in L^{\infty}([t_i, t_f], \mathbb{J})$.¹³ Moreover we can see that

$$\begin{aligned} \|v^{J}(q, Q_{i})\|_{L^{\infty}([t_{i}, t_{f}], \mathbb{J})} &\leq \|\dot{q}\|_{L^{\infty}([t_{i}, t_{f}], \mathbb{J})} + \|\Gamma^{J}(q, Q_{i})\|_{L^{\infty}([t_{i}, t_{f}], \mathbb{J})} \\ &\leq \|\dot{q}\|_{L^{\infty}([t_{i}, t_{f}], \mathbb{J})} + D_{1}\|\Gamma^{J}(q, Q_{i})\|_{L^{\infty}([t_{i}, t_{f}], V')} \end{aligned}$$

and, by (4.10.31), we obtain

$$\|v^{J}(q, Q_{i})\|_{L^{\infty}([t_{i}, t_{f}], \mathbb{J})} \le D_{2}$$
(4.10.32)

where D_2 depends only on the norms $\|\dot{q}\|_{L^{\infty}([t_i, t_f], \mathbb{J})}$, $\|q\|_{L^{\infty}([t_i, t_f], \mathbb{J})}$ and $\|Q\|_{L^{\infty}([t_i, t_f], \mathbb{J}_V^{\perp})}$. Then using (4.10.17) the constant D_2 can be chosen depending only on $\|q\|_{W^{1,\infty}([t_i, t_f], \mathbb{J})}$ and $\|Q_i\|$.

Definition 4.10.3. We call δ -metric the function defined on the product space $(L^{\infty}([0, T], \mathbb{R}^d))^2$ by

$$\delta(u, v) := meas\{t \in [0, T] \mid u(t) \neq v(t)\}.$$

¹³By the equivalence of V'-norm and l_1 -norm in J. One can check that V' coincides with the domain $D(A^{-\frac{1}{2}})$ of the operator $A^{-\frac{1}{2}}$ and its Fourier characterization is $V' = \{\sum_{k \in \mathbb{N}_0^2} u_k W_k \mid \sum_{k \in \mathbb{N}_0^2} u_k^2 < +\infty\}$. For $v \in V'$, $||v||_{V'} = \frac{ab}{4} \sum_{k \in \mathbb{N}_0^2} u_k^2$.
Remark 19. The δ -metric is the restriction, to the space of ordinary controls, of the strong metric defined on the space of relaxed controls (see [7]).

In [7] (Chapter 3) we can find the so called *Approximation Lemma* that says that a strongly continuous family of relaxed controls can be weakly approximated with arbitrary accuracy by a strongly continuous family of ordinary controls. At the end of that chapter we can find a "Remark on the Terminology" leading us to the following Lemma:

Lemma 4.10.5 (Approximation Lemma; [7]). Let $A \subseteq \mathbb{R}^d$ be the convexification of a finite set of points:

$$A := Conv\{p_1, p_2, \ldots, p_r\}$$

and, $\mathcal{V} := \{v(t, b) \in L^{\infty}([0, T], A) \mid b \in B\}$ be a L^1 -continuous family of A-valued functions. Then for each $\varepsilon > 0$ one can construct a family $\mathcal{V}^{\varepsilon} := \{v^{\varepsilon}(t, b) \in L^{\infty}([0, T], \{p_1, \ldots, p_r\}) \mid b \in B\}$ of $\{p_1, p_2, \ldots, p_r\}$ -valued functions such that

- $\mathcal{V}^{\varepsilon}$ is δ -continuous, i.e., $b \mapsto v^{\varepsilon}(\cdot, b)$ is (B, δ) -continuous;
- $\mathcal{V}^{\varepsilon} \in \text{-approximates, uniformly w.r.t. } b, the family <math>\mathcal{V}$ in relaxation metric, i.e., $\forall b \in B \quad \|v^{\varepsilon}(\cdot, b) - v(\cdot, b)\|_{rx} < \varepsilon$ and;
- The elements of V^ε are piecewise constant and the number of intervals of constancy is the same for all b ∈ B.

Remark 20. Note that δ -metric and L^q -metric $(1 \leq q < +\infty)$ are equivalent in the subset of piecewise constant functions taking values on a fixed finite set. Also, since the controls take values on a finite set, the δ -continuity of $\mathcal{V}^{\varepsilon}$ is equivalent to the continuity of the lengths of the intervals of constancy of the controls.

In Lemma 4.10.5 is said that the intervals of constancy can be taken the same for all $b \in B$ but, looking at [[7]; ch. 3] some of those intervals may degenerate to a single point. We claim that we can suppose non-degeneracy of the intervals,¹⁴. We may even suppose that there exists a lower bound θ^{ε} for the lengths of the intervals of constancy of the family $\mathcal{V}^{\varepsilon}$, i.e., for all $b \in B$ none $v^{\varepsilon}(\cdot, b)$ has an interval of constancy with length less than θ^{ε} .

Looking at [[7]; ch. 3] we see that the intervals of constancy are (or can be) constructed in the following way: First we subdivide the interval [0, T] into n^2 intervals — L_i , $i = 1, \ldots, n^2$ — with the same length — $\frac{T}{n^2}$. Then subdivide each one of these intervals L_i into r subintervals — L_{ij} $j = 1, \ldots, r$ — which lengths $length(L_{ij}) = \int_{L_i} v_j(\tau, b) d\tau$, $v(\cdot, b) = \sum_{j=1}^r v_j(\cdot, b) p_j$, $v_j(t, b) \in [0, 1]$, depend (continuously) on b and the interval L_{ij_1} lays on the left of L_{ij_2} if $j_1 < j_2$. To this partition is associated the piecewise constant control $v_{n^2}(\cdot, b)$ defined by:

 $t \in L_{ij} \Rightarrow v_{n^2}(t, b) = p_j;$ $i = 1, ..., n^2, j = 1, ..., r,$

i.e., in each interval L_i we use all the controls from $\{p_1, \ldots, p_r\}$ using p_{j_1} before p_{j_2} if $j_1 < j_2$. Note that as we have said before some control may be used for time zero.

¹⁴Note that it is not enough to eliminate the degenerate intervals because the number of intervals would not be the same for all $b \in B$.

Each $v^{\varepsilon}(\cdot, b) \in \mathcal{V}^{\varepsilon}$ have the form $v^{\varepsilon} = v_{n^2}$ for some *n* depending only on ε . We shall need the following strong result without non degeneracy of the intervals of constancy:

Corollary 4.10.6 (Approximation Corollary). Let $A \subseteq \mathbb{R}^d$ be the convexification of a finite set of points:

$$A := Conv\{p_1, p_2, \ldots, p_r\}$$

and, $\mathcal{V} := \{v(t, b) \in L^{\infty}([0, T], A) \mid b \in B\}$ be a L^1 -continuous family of Avalued functions. Then for each $\varepsilon > 0$ there is $\theta^{\varepsilon} > 0$ and a family $\mathcal{Z}^{\varepsilon} := \{z^{\varepsilon}(t, b) \in L^{\infty}([0, T], \{p_1, p_2, \dots, p_r\}) \mid b \in B\}$ of $\{p_1, p_2, \dots, p_r\}$ -valued functions such that

- $\mathcal{Z}^{\varepsilon}$ is δ -continuous;
- $\mathcal{Z}^{\varepsilon} \varepsilon$ -approximates, uniformly w.r.t. b, the family \mathcal{V} in relaxation metric, i.e., $\forall b \in B \quad \|z^{\varepsilon}(\cdot, b) - v(\cdot, b)\|_{rx} < \varepsilon;$
- The elements of Z^ε are piecewise constant and the number of intervals of constancy is the same for all b ∈ B and,
- For all b ∈ B all the intervals of constancy of z^ε(·, b) have a length not less than θ^ε > 0.

Before the proof consider the following Lemma:

Lemma 4.10.7. Given K > 0, $\gamma > 0$ and $L \in \mathbb{N}_0$. Define the sets

$$\mathcal{P}_0 := \{ (x_1, \dots, x_L) \in \mathbb{R}^L \mid x_i \ge 0, \sum_{i=1}^L x_i = K \}$$
$$\mathcal{P}_\theta := \{ (x_1, \dots, x_L) \in \mathbb{R}^L \mid x_i \ge \theta, \sum_{i=1}^L x_i = K \}$$

where $\theta > 0$. Choose $n \in \mathbb{N}_0$ such that $\frac{(L+1)K}{nL} < \gamma$ and put

$$\theta = \frac{K}{nL}.$$
 ¹⁵ (4.10.33)

Then the map $P_{0,\theta} = (P_{0,\theta}^1, \ldots, P_{0,\theta}^L)$, defined on \mathcal{P}_0 by:

$$P_{0,\theta}^{i}(x) := \left(1 - \frac{1}{n}\right)\left(x_{i} - \frac{K}{L}\right) + \frac{K}{L},$$

is continuous, take its values on \mathcal{P}_{θ} and, satisfies $|P_{0,\theta}^i(x) - x_i| < \gamma$.

The Approximation Corollary follows from the Approximation Lemma and from the last Lemma. Indeed let $\varepsilon > 0$ be a positive real number, put $D := \max_{i \in \{1, ..., r\}} \{ \|p_i\| \}$, take K = 1, L = r, $\gamma < \frac{\varepsilon}{2TDr}$, $n \in \mathbb{N}_0$ such that $\frac{r+1}{nr} < \gamma$

¹⁵So, θ depends on both K, L and γ .

and $\theta = \frac{1}{nr}$.

The continuity of $P_{0,\theta}$ implies the continuity of the map

$$v(t, b) = \sum_{j=1}^{r} v_j(t, b) p_j \mapsto z(t, b) = \sum_{j=1}^{r} P_{0,\theta}^j(v(t, b))$$

and then the family of the controls $z(\cdot, b) = \sum_{j=1}^{r} P_{0,\theta}^{j}(v(t, b))$ is parametrized in $L^{2}(0, T, Conv\{p_{1}, \ldots, p_{r}\})$ -norm.

The family $\mathcal{Z} \stackrel{\varepsilon}{\xrightarrow{2}}$ -approximates the family \mathcal{V} in relaxation metric. Indeed

$$\begin{aligned} \|z(\cdot, b) - v(\cdot, b)\|_{rx} \\ &\leq \int_0^T |z(\tau, b) - v(\tau, b)| \, d\tau = \int_0^T |\sum_{j=1}^r z_j(t, b) p_j - \sum_{j=1}^r v_j(t, b) p_j| \, d\tau \\ &\leq \int_0^T |\sum_{j=1}^r |z_j(t, b) - v_j(t, b)| D \, d\tau < T\gamma Dr < \frac{\varepsilon}{2} \end{aligned}$$

Now apply Approximation lemma to the family \mathcal{Z} and find a family $\mathcal{Z}^{\frac{\varepsilon}{2}}$ that $\frac{\varepsilon}{2}$ -approximates \mathcal{Z} in relaxation metric. Hence

$$\|z^{\frac{\varepsilon}{2}}(\cdot, b) - v(\cdot, b)\|_{rx} \le \|z^{\frac{\varepsilon}{2}}(\cdot, b) - z(\cdot, b)\|_{rx} + \|z(\cdot, b) - v(\cdot, b)\|_{rx} < \varepsilon.$$

Therefore the family $\mathcal{Z}^{\frac{\varepsilon}{2}} \varepsilon$ -approximates \mathcal{V} in relaxation metric.

From the Approximation Lemma the number of intervals of constancy of $z^{\frac{\varepsilon}{2}}(\cdot, b)$ is the same for all $b \in B$ and, the L_{ij} interval of $z^{\frac{\varepsilon}{2}}(\cdot, b)$ has length $\int_{L_i} z_j(\tau, b) d\tau > \frac{T}{n^2} \theta$ for that $n \in \mathbb{N}_0$ such that $z^{\frac{\varepsilon}{2}}(\cdot, b) = z_{n^2}(\cdot, b)$. Thus all intervals have a length not smaller than $\frac{T}{n^2}\theta > 0$ — a positive constant depending only in ε .

Proof of Lemma 4.10.7. The continuity of each $P_{0,\theta}^i$, and then of $P_{0,\theta}$, is clear. From

$$\sum_{i=1}^{L} P_{0,\theta}^{i}(x) = \sum_{i=1}^{L} \left[\left(1 - \frac{1}{n}\right) \left(x_{i} - \frac{K}{L}\right) + \frac{K}{L} \right] = K$$

and

$$P_{0,\theta}^i(x) = \left(1 - \frac{1}{n}\right)x_i + \frac{K}{nL} \ge \frac{K}{nL} = \theta$$

we conclude that $P_{0,\theta}$ takes its values on \mathcal{P}_{θ} . It remais to estimate $|P_{0,\theta}^i(x) - x_i|$:

$$\left|P_{0,\theta}^{i}(x) - x_{i}\right| = \left|-\frac{x_{i}}{n} + \frac{K}{nL}\right| \le \frac{K}{n} + \frac{K}{nL} \le \frac{(L+1)K}{nL} < \gamma.$$

Now, we are ready to start the proof of the induction step. Fix $u_0 \in V$. By "back-induction" hypothesis system [(4.10.1).N] is time-T solid controllable in observed component so, there is a family $\mathcal{V} := \{v(\cdot, b) \in L^{\infty}([0, T], \mathbb{R}^{\kappa_N}) \mid b \in B\}$ parametrized continuously in L^2 -metric such that

$$\mathbb{E}_T(u_0, \mathcal{V}) \text{ covers } \overline{\mathcal{O}}_R(u_0^{\#\mathcal{O}}) \text{ solidly.}$$
 (4.10.34)

$$\|v(t, b)\|_{l_1} \le \Xi_0 \quad \forall (t, b) \in [0, T] \times B.$$
 (4.10.35)

Note that, by (4.10.35) we obtain

$$\forall (t, b) \in [0, T] \times B \quad v(t, b) \in Conv\{\pm \Xi_0 e_k \mid k \in \mathcal{K}^N\}$$
¹⁶

Then there exists $\Xi > 0$ such that

$$\forall (t, b) \in [0, T] \times B v(t, b) \in Conv\{\pm \Xi e_k, \ \pm \Xi \delta_{m,n} \mid k \in \mathcal{K}^{N-1}, \ (m, n) \in S_{N-1}\} =: C_N \quad (4.10.36)$$

Now fix $\varepsilon > 0$ and ε -approximate the family \mathcal{V} taking values on C_N by a family $\mathcal{Z}^{\varepsilon} := \{z^{\varepsilon}(t, b) \in | b \in B\}$ taking values on $\{\pm \Xi e_k, \pm \Xi \delta_{m,n} | k \in \mathcal{K}^{N-1}, (m, n) \in S_{N-1}\}$ like in Corollary 4.10.6. ¹⁷ Choose a real number γ such that

$$0 < \gamma < \|u_0^{\#\mathcal{O}} - \partial \mathbb{E}_T(u_0, \mathcal{V})\|_{l_1} - R.$$

Remark 21. It is clear that $||u_0^{\#\mathcal{O}} - \partial \mathbb{E}_T(u_0, \mathcal{V})||_{l_1} \geq R$ because of (4.10.34). To see that the inequality is, in fact, strict we suppose not and then, there is $p \in \partial \mathbb{E}_T(u_0, \mathcal{V})$ such that $||u_0^{\#\mathcal{O}} - p||_{l_1} = R$. For the sequence of continuous functions ϕ_n defined in B by

$$\phi_n(b) := \left(1 - \frac{1}{n}\right) \left(\mathbb{E}_T(u_0, v(\cdot, b)) - u_0^{\#\mathcal{O}} \right) + u_0^{\#\mathcal{O}}$$

 $we\ have$

$$\begin{aligned} \|\phi_n - \mathbb{E}_T\|_{C(B,\mathbb{R}_{\infty}^{\#^{\mathcal{O}}})} &= \frac{1}{n} \|\mathbb{E}_T\|_{C(B,\mathbb{R}_{\infty}^{\#^{\mathcal{O}}})} + \frac{1}{n} \|u_0^{\#^{\mathcal{O}}}\|_{l_{\infty}} \to 0; \\ \phi_n(\mathcal{V}) &= \left(1 - \frac{1}{n}\right) \mathbb{E}_T(u_0,\mathcal{V}) + \frac{1}{n} u_0^{\#^{\mathcal{O}}} \\ &\left\| \left(1 - \frac{1}{n}\right) p + \frac{1}{n} u_0^{\#^{\mathcal{O}}} - u_0^{\#^{\mathcal{O}}} \right\|_{l_1} = \left(1 - \frac{1}{n}\right) \|p - u_0^{\#^{\mathcal{O}}}\|_{l_1} \le R. \end{aligned}$$

Therefore

$$\left(1-\frac{1}{n}\right)p+\frac{1}{n}u_0^{\#\mathcal{O}}\in\partial\phi_n(\mathcal{V})\cap\mathcal{O}_R(u_0^{\#}\mathcal{O})$$

and then, for all $n \in \mathbb{N}_0$, $\phi_n(\mathcal{V})$ does not cover $\overline{\mathcal{O}}_R(u_0^{\#\mathcal{O}})$ which contradicts (4.10.34).

Now note that for $x \in \overline{\mathcal{O}}_R(u_0^{\#\mathcal{O}})$ we obtain $\|u_0^{\#\mathcal{O}} - \partial \mathbb{E}_T(u_0, \mathcal{V})\|_{l_1} \leq R + \|x - \partial \mathbb{E}_T(u_0, \mathcal{V})\|_{l_1}$, i.e., $\gamma < \|x - \partial \mathbb{E}_T(u_0, \mathcal{V})\|_{l_1}$ and then

$$\frac{\gamma}{\#\mathcal{O}} < \|x - \partial \mathbb{E}_T(u_0, \mathcal{V})\|_{l_\infty}.$$
(4.10.37)

By the continuity of \mathbb{S}_{srx} there is $\bar{\varepsilon} > 0$ such that

$$\|v - w\|_{rx} < \bar{\varepsilon} \Rightarrow$$

$$\|\Pi_{\mathcal{O}} \circ \mathbb{S}_s(u_0, F, w, \nu)(T) - \Pi_{\mathcal{O}} \circ \mathbb{S}_s(u_0, F, v, \nu)(T)\|_{C(B, \mathbb{R}^{\#\mathcal{O}}_{\infty})} < \frac{\gamma}{2\#\mathcal{O}}.$$

¹⁶Note that the norm used in (4.10.35) is the l_1 one.

¹⁷Recall that S_{N-1} has been defined, in the proof of Proposition 3.5.3 for N > 2 and in the proof of Proposition 3.5.2 for N = 2, as the set of "extracted" pairs of modes associated with the new directions.

Fix $\varepsilon < \bar{\varepsilon}$ and put

$$\mathbb{E}_T^{\varepsilon}(u_0, z^{\varepsilon}(b)) := \Pi_{\mathcal{O}} \circ \mathbb{S}_s(u_0, F, z^{\varepsilon}(b), \nu)(T)$$

and consider ϕ defined on B such that

$$\|\phi(\cdot) - \mathbb{E}_T^{\varepsilon}(u_0, z^{\varepsilon}(\cdot))\|_{C(B, \mathbb{R}_{\infty}^{\#\mathcal{O}}} < \frac{\gamma}{2\#\mathcal{O}}$$

Hence

$$\|\phi - \mathbb{E}_T\|_{C(B,\mathbb{R}^{\#\mathcal{O}}_{\infty})} \le \|\phi - \mathbb{E}_T^{\varepsilon}\|_{C(B,\mathbb{R}^{\#\mathcal{O}}_{\infty})} + \|\mathbb{E}_T^{\varepsilon} - \mathbb{E}_T\|_{C(B,\mathbb{R}^{\#\mathcal{O}}_{\infty})} < \frac{\gamma}{\#\mathcal{O}}$$

and, by (4.10.37) we conclude

$$\forall x \in \overline{\mathcal{O}}_R(u_0^{\#}\mathcal{O}) \Big[deg(\phi, B, x) = deg(\mathbb{E}_T, B, x) \neq 0 \Big]$$

so, $\phi(B)$ covers $\overline{\mathcal{O}}_R(u_0^{\#}\mathcal{O})$. Therefore $E_T^{\varepsilon}(u_0, \mathcal{Z}^{\varepsilon})$ covers $\overline{\mathcal{O}}_R(u_0^{\#}\mathcal{O})$ solidly for all positive $\varepsilon < \overline{\varepsilon}$. Moreover, for all pairs (t, b) we have $||z^{\varepsilon}(t, b)||_{l_1} \leq \Xi$. Therefore for $\varepsilon < \overline{\varepsilon}$ system [(4.10.1).N] is is solid controllable by means of the family $\mathcal{Z}^{\varepsilon}$.

Now we fix $\varepsilon \in]0, \overline{\varepsilon}[$ and define $\mathcal{Z} := \mathcal{Z}^{\varepsilon}; \quad z(t, b) := z^{\varepsilon}(t, b).$ We shall compare the trajectories generated by a control $z(t, b) \in \mathcal{Z}$ with another generated by some "appropriated" control taking values on $\mathbb{R}^{\kappa_{N-1}}$.

To the control $z(\cdot, b)$ is associated a partition X(b) of [0, T] into m nondegenerated intervals of lengths $x_i \ge \theta > 0$:

$$X(b) = (x_1, x_2, \dots, x_m) \in \mathbb{R}^m, \ m \in \mathbb{N}_0, \ \sum_{i=1}^m x_i = T,$$

where m and θ are independent of the parameter b. We put

$$A(b) = \{(0 = \alpha_0, \alpha_1, \dots, \alpha_m = T)\}$$

for the end points of the intervals in X(b). So,

$$A(b) \in \mathcal{A}_{\theta} := \{ (\alpha_0, \alpha_1, \dots, \alpha_m) \in \mathbb{R}^{m+1} \mid \alpha_0 = 0, \ \alpha_m = T, \\ \alpha_i - \alpha_{i-1} \ge \theta, \sum_{i=1}^m (\alpha_i - \alpha_{i-1}) = T \}.$$

Let $w \in \mathbb{R}$, $w \ge 3$. Now in [0, T] we want to define a function $\phi_w(\cdot, b)$ associated to X(b) with the following properties

- $\phi_w(\cdot, b)$ vanishes at the points α_i , $i = 0, \ldots, m$;
- $\phi_w(\cdot, b) \in W^{1,\infty}([0, T], \mathbb{R})$ with

$$\|\phi_w(\cdot, b)\|_{C([0,T],\mathbb{R})} \le 1; \qquad \|\dot{\phi}_w(\cdot, b)\|_{L^{\infty}([0,T],\mathbb{R})} \le \frac{w(1+\theta)}{\theta}$$

• $\delta(\phi_w(t, b), \sin(wt)) \leq \frac{2T}{w}$ and;

• For fixed w, the map $A(b) \mapsto \phi_w(\cdot, b)$ is $(\mathcal{A}_{\theta}, W^{1,2}(0, T, \mathbb{R}))$ -continuous (where \mathcal{A}_{θ} is endowed with the topology induced by \mathbb{R}^{m+1}).

We proceed with the construction of $\phi_w(\cdot, b)$: For each $i \in \{1, \ldots, m\}$ put $\rho_i = \frac{x_i}{w}$. Then subdivide each interval $[\alpha_{i-1}, \alpha_i]$ into

$$[\alpha_{i-1}, \alpha_i] = [\alpha_{i-1}, \alpha_{i-1} + \rho_i[\cup [\alpha_{i-1} + \rho_i, \alpha_i - \rho_i]\cup]\alpha_i - \rho_i, \alpha_i].$$
¹⁸

For each $i = 1, \ldots, m$ put

$$\phi_w(t, b) = \begin{cases} \frac{\sin(w(\alpha_{i-1} + \rho_i))}{\rho_i}(t - \alpha_{i-1}) & \text{if} \quad t \in [\alpha_{i-1}, \alpha_{i-1} + \rho_i];\\ \sin(wt) & \text{if} \quad t \in [\alpha_{i-1} + \rho_i, \alpha_i - \rho_i];\\ \frac{\sin(w(\alpha_i - \rho_i))}{-\rho_i}(t - \alpha_i) & \text{if} \quad t \in [\alpha_i - \rho_i, \alpha_i]. \end{cases}$$

Then the graph of the restriction of $\phi_w(\cdot, b)$ to an interval $[\alpha_{i-1}, \alpha_i]$ is a concatenation of a straight line, a piece of the graph of sin(wt) and another straight line. From the construction is clear that $\phi_w(\cdot, b)$ vanishes at the points $\alpha_i, \quad i=0,\ldots,m \text{ and that } \phi_w(\cdot,b) \text{ is continuous with } \|\phi_w(\cdot,b)\|_{C([0,T],\mathbb{R})} \leq 1.$ In the subintervals $]\alpha_{i-1} + \rho_i, \ \alpha_i - \rho_i [$ we have $\dot{\phi}_w(t, b) = w \cos(wt) \operatorname{so}_i |\dot{\phi}_w(t, b)| \leq 1$ $w \leq \frac{w(1+\theta)}{\theta}$. In the subintervals $]\alpha_{i-1}, \alpha_{i-1} + \rho_i[$ and $]\alpha_i - \rho_i, \alpha_i[$ we have $|\dot{\phi}_w(t,b)| \leq \frac{1}{\rho_i} = \frac{w}{x_i} \leq \frac{w}{\theta} \leq \frac{w(1+\theta)}{\theta}$. Hence we have $\|\dot{\phi}_w(\cdot,b)\|_{L^{\infty}([0,T],\mathbb{R})} \leq \frac{w}{\theta}$ $\frac{w(1+\theta)}{\theta}.$ Therefore $\phi_w(\cdot,\,b)\in W^{1,\infty}([0,\,T],\,\mathbb{R})$ and

$$\|\phi_w(\cdot, b)\|_{W^{1,\infty}([0,T],\mathbb{R})} \le 1 + \frac{w(1+\theta)}{\theta}.$$

We see that $\phi_w(t, b)$ differs from $\sin(wt)$ only in the intervals $[\alpha_{i-1}, \alpha_{i-1} + \rho_i]$ and $\left[\alpha_{i}-\rho_{i}, \alpha_{i}\right]$ so,

$$\delta(\phi_w(t, b), \sin(wt)) = \sum_{i=1}^m 2\rho_i = \sum_{i=1}^m 2\frac{x_i}{w} \le \frac{2T}{w}$$

It remains to check the continuity property. For that fix $w \geq 3$ and $A \in \mathcal{A}_{\theta}$. Let $\gamma > 0$ and $B \in \mathcal{A}_{\theta}$ such that $||A - B||_{l_1} < \gamma$. Let $A = (\alpha_0, \alpha_1, \ldots, \alpha_m)$ and $B = (\beta_0, \beta_1, \dots, \beta_m)$ be the coordinates of A and B. Then we have $|\alpha_i - \beta_i| < \gamma$ for $i = 1, \ldots, m - 1$, $\alpha_0 = \beta_0 = 0$ and $\alpha_m = \beta_m = T$. For small $\gamma \quad (\gamma < \frac{\theta}{w})^{19}$, putting ϕ_A and ϕ_B for the functions ϕ_w associated with A and B, we have that ϕ_A and ϕ_B differ only in the following union of subintervals

$$[0, \min\{\rho_{1}^{A}, \rho_{1}^{B}\}] \cup [\min\{\rho_{1}^{A}, \rho_{1}^{B}\}, \max\{\rho_{1}^{A}, \rho_{1}^{B}\}]$$

$$\cup [\min\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\}, \max\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\}]$$

$$\cup [\max\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\}, \min\{\alpha_{i}, \beta_{i}\}]$$

$$\cup [\min\{\alpha_{i}, \beta_{i}\}, \max\{\alpha_{i}, \beta_{i}\}]$$

$$\cup [\max\{\alpha_{i}, \beta_{i}\}, \min\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\}]$$

$$\cup [\min\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\}, \max\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\}]$$

$$\cup [\min\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\}, \max\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\}]$$

$$\cup [\max\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\}, T].$$
(4.10.38)

¹⁸Note that, since $w \ge 3$ we have $\rho_i \le \frac{x_i}{3}$ and the subdivision is well defined. ¹⁹Note that for $\gamma < \frac{\theta}{w}$ we have $\gamma < \rho_i^A$ and $\gamma < \rho_i^B$ for all $i \in \{1, \ldots, m\}$.

Now we prove that $\|\phi_A - \phi_B\|_{C([0,T],\mathbb{R})}$ goes to zero as γ does. For that is enough to prove that $\|\phi_A - \phi_B\|_{C(I,\mathbb{R})}$ goes to zero as γ does for every interval I of the union (4.10.38).

• For $I = [0, \min\{\rho_1^A, \rho_1^B\}]$:

$$\|\phi_A - \phi_B\|_{C(I,\mathbb{R})} = \max_{t \in I} \{\frac{\sin(w\rho_1^A)}{\rho_1^A} t - \frac{\sin(w\rho_1^B)}{\rho_1^B} t\}$$
$$= \left|\frac{\sin(w\rho_1^A)}{\rho_1^A} \min\{\rho_1^A, \rho_1^B\} - \frac{\sin(w\rho_1^B)}{\rho_1^B} \min\{\rho_1^A, \rho_1^B\}\right|^{-20}$$

• For
$$I = [\min\{\rho_1^A, \rho_1^B\}, \max\{\rho_1^A, \rho_1^B\}]$$
:

$$\begin{aligned} \|\phi_A - \phi_B\|_{C(I,\mathbb{R})} &\leq (\max\{\rho_1^A, \rho_1^B\} - \min\{\rho_1^A, \rho_1^B\}) \|\phi_A - \phi_B\|_{C([0,T],\mathbb{R})} \\ &\leq 2|\rho_1^A - \rho_1^B| = 2\frac{1}{w}|x_1 - y_1| \leq 2\frac{\gamma}{w} \end{aligned}$$

• For i = 1, ..., m - 1 and

$$- \text{ For } I = [\min\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\}, \max\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\}]:$$

$$\|\phi_{A} - \phi_{B}\|_{C(I,\mathbb{R})}$$

$$\leq (\max\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\} - \min\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\})\|\phi_{A} - \phi_{B}\|_{C^{0}}$$

$$\leq 2|(\alpha_{i} - \rho_{i}^{A}) - (\beta_{i} - \rho_{i}^{B})| \leq 2|\alpha_{i} - \beta_{i}| + 2|\rho_{i}^{A} - \rho_{i}^{B}|$$

$$\leq 2\gamma + 2\frac{1}{w}|x_{i} - y_{i}| \leq 2\gamma + 2\frac{2\gamma}{w} = 2\gamma(1 + \frac{2}{w}).$$

- For $I = [\max\{\alpha_i - \rho_i^A, \beta_i - \rho_i^B\}, \min\{\alpha_i, \beta_i\}]$:

$$\begin{split} \|\phi_{A} - \phi_{B}\|_{C(I,\mathbb{R})} \\ &= \max_{t \in I} \left\{ \begin{array}{c} \left| \frac{\sin(w(\alpha_{i} - \rho_{i}^{A})}{-\rho_{i}^{A}}(t - \alpha_{i}) - \frac{\sin(w(\beta_{i} - \rho_{i}^{B}))}{-\rho_{i}^{B}}(t - \beta_{i}) \right| \right\} \\ &= \max \left\{ \begin{array}{c} \left| \frac{\sin(w(\alpha_{i} - \rho_{i}^{A})}{-\rho_{i}^{A}}(\max\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\} - \alpha_{i}) \right. \\ &\left. - \frac{\sin(w(\beta_{i} - \rho_{i}^{B}))}{-\rho_{i}^{B}}(\max\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\} - \beta_{i}) \right| , \\ \left. \frac{\left| \sin(w(\alpha_{i} - \rho_{i}^{A})}{-\rho_{i}^{A}}(\min\{\alpha_{i}, \beta_{i}\} - \alpha_{i}) \right. \\ &\left. - \frac{\sin(w(\beta_{i} - \rho_{i}^{B}))}{-\rho_{i}^{B}}(\min\{\alpha_{i}, \beta_{i}\} - \beta_{i}) \right| \right\} \end{split}$$

- For $I = [\min\{\alpha_i, \beta_i\}, \max\{\alpha_i, \beta_i\}]$:

 $\|\phi_A - \phi_B\|_{C(I,\mathbb{R})} \le (\max\{\alpha_i, \beta_i\} - \min\{\alpha_i, \beta_i\}) \|\phi_A - \phi_B\|_{C^0} \le 2\gamma.$

²⁰Note that the maximum is attained at one of the end points of the interval I, because the functions ϕ_A and ϕ_B are affine in I.

$$- \text{ For } I = [\max\{\alpha_{i}, \beta_{i}\}, \min\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\}]:$$

$$\|\phi_{A} - \phi_{B}\|_{C(I,\mathbb{R})}$$

$$= \max\left\{ \left| \frac{\sin(w(\alpha_{i} + \rho_{i+1}^{A})}{\rho_{i}^{A}}(t - \alpha_{i}) - \frac{\sin(w(\beta_{i} + \rho_{i+1}^{B}))}{\rho_{i}^{B}}(t - \beta_{i}) \right| \right\}$$

$$= \max\left\{ \left| \frac{\sin(w(\alpha_{i} + \rho_{i+1}^{A})}{\rho_{i}^{A}}(\max\{\alpha_{i}, \beta_{i}\} - \alpha_{i}) - \frac{\sin(w(\beta_{i} + \rho_{i+1}^{B}))}{\rho_{i}^{B}}(\max\{\alpha_{i}, \beta_{i}\} - \beta_{i}) \right|, \left| \frac{\sin(w(\alpha_{i} + \rho_{i+1}^{A})}{\rho_{i}^{A}}(\min\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\} - \alpha_{i}) - \frac{\sin(w(\beta_{i} + \rho_{i+1}^{B}))}{\rho_{i}^{B}}(\min\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\} - \beta_{i}) \right| \right\}$$

- For $I = [\min\{\alpha_i + \rho_{i+1}^A, \beta_i + \rho_{i+1}^B\}, \max\{\alpha_i + \rho_{i+1}^A, \beta_i + \rho_{i+1}^B\}]$:

$$\begin{aligned} \|\phi_{A} - \phi_{B}\|_{C(I,\mathbb{R})} \\ \leq & (\max\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\} - \min\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\}) \|\phi_{A} - \phi_{B}\|_{C^{0}} \\ \leq & 2|(\alpha_{i} + \rho_{i+1}^{A}) - (\beta_{i} + \rho_{i+1}^{B})| \leq 2|\alpha_{i} - \beta_{i}| + 2|\rho_{i+1}^{A} - \rho_{i+1}^{B}| \\ \leq & 2\gamma + 2\frac{1}{w}|x_{i+1} - y_{i+1}| \leq 2\gamma + 2\frac{2\gamma}{w} = 2\gamma(1 + \frac{2}{w}). \end{aligned}$$

For $I = [\min\{T - \rho_m^A, T - \rho_m^B\}, \max\{T - \rho_m^A, T - \rho_m^B\}]:$

$$\begin{aligned} &\|\phi_A - \phi_B\|_{C(I,\mathbb{R})} \\ \leq &(\max\{T - \rho_m^A, T - \rho_m^B\} - \min\{T - \rho_m^A, T - \rho_m^B\}) \|\phi_A - \phi_B\|_{C([0,T],\mathbb{R})} \\ \leq &2|\rho_m^A - \rho_m^B| = \frac{2}{w}|x_m - y_m| \le \frac{2\gamma}{w}. \end{aligned}$$

For $I = [\max\{T - \rho_m^A, T - \rho_m^B\}, T]$:

$$\begin{split} &\|\phi_{A} - \phi_{B}\|_{C(I,\mathbb{R})} \\ &= \max_{t \in I} \Big\{ \Big| \frac{\sin(w(T - \rho_{m}^{A})}{\rho_{m}^{A}}(t - T) - \frac{\sin(w(T - \rho_{m}^{B}))}{\rho_{i}^{B}}(t - T) \Big| \Big\} \\ &= \Big| \frac{\sin(w(T - \rho_{m}^{A})}{\rho_{m}^{A}}(\max\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\} - T) \\ &- \frac{\sin(w(T - \rho_{m}^{B}))}{\rho_{i}^{B}}(\max\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\} - T) \Big| \end{split}$$

Therefore since $\beta_i \to \alpha_i$ and $\rho_m^B \to \rho_m^A$ as $\gamma \to 0$ we have that for every interval I in the union (4.10.38)

 $\|\phi_A - \phi_B\|_{C(I,\mathbb{R})} \to 0 \quad \text{as} \quad \gamma \to 0.$

Hence

$$\|\phi_A - \phi_B\|_{C(0,T,\mathbb{R})} \to 0 \text{ as } B \to A.$$
 (4.10.39)

For the derivatives we have

$$\begin{split} & \int_{0}^{T} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ ^{21} &= \int_{0}^{\min\{\rho_{1}^{A}, \rho_{1}^{B}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt + \int_{\min\{\rho_{1}^{A}, \rho_{1}^{B}\}}^{\max\{\rho_{1}^{A}, \rho_{1}^{B}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ &+ \sum_{i=1}^{m-1} \left[\int_{\min\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\}}^{\max\{\alpha_{i} - \rho_{i}^{A}, \beta_{i} - \rho_{i}^{B}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ &+ \int_{\max\{\alpha_{i}, \beta_{i}\}}^{\min\{\alpha_{i}, \beta_{i}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ &+ \int_{\min\{\alpha_{i}, \beta_{i}\}}^{\max\{\alpha_{i}, \beta_{i}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ &+ \int_{\max\{\alpha_{i}, \beta_{i}\}}^{\max\{\alpha_{i}, \beta_{i}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ &+ \int_{\max\{\alpha_{i}, \beta_{i}\}}^{\max\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ &+ \int_{\min\{\alpha_{i} + \rho_{i+1}^{A}, \beta_{i} + \rho_{i+1}^{B}\}}^{\max\{\alpha_{i}, \beta_{i}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \\ &+ \int_{\min\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\}}^{\max\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt \tag{4.10.40} \\ &+ \int_{\max\{T - \rho_{m}^{A}, T - \rho_{m}^{B}\}}^{T} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt. \end{aligned}$$

Now we prove that each one of the intervals on the right-hand side of (4.10.41) goes to zero when γ does:

$$\int_{0}^{\min\{\rho_{1}^{A}, \, \rho_{1}^{B}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt$$

$$\leq \left[\frac{\sin(w\rho_{1}^{A})}{\rho_{1}^{A}} - \frac{\sin(w\rho_{1}^{B})}{\rho_{1}^{B}}\right]^{2} \min\{\rho_{1}^{A}, \, \rho_{1}^{B}\}.$$

$$\int_{\min\{\rho_1^A, \rho_1^B\}}^{\max\{\rho_1^A, \rho_1^B\}} |\dot{\phi}_A(t) - \dot{\phi}_B(t)|^2 dt$$

$$\leq (\max\{\rho_1^A, \rho_1^B\} - \min\{\rho_1^A, \rho_1^B\}) ||\dot{\phi}_A - \dot{\phi}_B||_{L^{\infty}(0, T, \mathbb{R})}^2$$

$$= |\rho_1^A - \rho_1^B| \left(2\frac{w(1+\theta)}{\theta}\right)^2 = \left(2\frac{w(1+\theta)}{\theta}\right)^2 \frac{1}{w} |x_1 - y_1| \leq 4w\gamma \left(\frac{1+\theta}{\theta}\right)^2,$$

²¹Recall that out of the intervals of (4.10.38) $\dot{\phi}_A$ and $\dot{\phi}_B$ coincide.

For
$$i = 1, ..., m - 1$$
:

$$\int_{\min\{\alpha_i - \rho_i^A, \beta_i - \rho_i^B\}}^{\max\{\alpha_i - \rho_i^A, \beta_i - \rho_i^B\}} |\dot{\phi}_A(t) - \dot{\phi}_B(t)|^2 dt$$

$$\leq (\max\{\alpha_i - \rho_i^A, \beta_i - \rho_i^B\} - \min\{\alpha_i - \rho_i^A, \beta_i - \rho_i^B\}) \|\dot{\phi}_A - \dot{\phi}_B\|_{L^{\infty}(0, T, \mathbb{R})}^2$$

$$= 4w^2 \left(\frac{1+\theta}{\theta}\right)^2 |(\alpha_i - \rho_i^A) - (\beta_i - \rho_i^B)| \leq 4w^2 \left(\frac{1+\theta}{\theta}\right)^2 (|\alpha_i - \beta_i| + |\rho_i^A - \rho_i^B|)$$

$$\leq 4w^2 \left(\frac{1+\theta}{\theta}\right)^2 (\gamma + \frac{1}{w}|x_i - y_i|) \leq 4w^2 \left(\frac{1+\theta}{\theta}\right)^2 \left(\gamma + \frac{2\gamma}{w}\right)$$

$$= 4\gamma w^2 \left(\frac{1+\theta}{\theta}\right)^2 \left(1 + \frac{2}{w}\right);$$

$$\int_{\max\{\alpha_i,\beta_i\}}^{\min\{\alpha_i,\beta_i\}} |\dot{\phi}_A(t) - \dot{\phi}_B(t)|^2 dt$$
$$= \left[\frac{\sin(w(\alpha_i - \rho_i^A))}{-\rho_i^A} - \frac{\sin(w(\beta_i - \rho_i^B))}{-\rho_i^B}\right]^2$$
$$\cdot (\min\{\alpha_i,\beta_i\} - \max\{\alpha_i - \rho_i^A,\beta_i - \rho_i^B\});$$

$$\int_{\min\{\alpha_i,\,\beta_i\}}^{\max\{\alpha_i,\,\beta_i\}} |\dot{\phi}_A(t) - \dot{\phi}_B(t)|^2 \, dt$$

$$\leq (\max\{\alpha_i,\,\beta_i\} - \min\{\alpha_i,\,\beta_i\}) \|\dot{\phi}_A - \dot{\phi}_B\|_{L^{\infty}(0,\,T,\,\mathbb{R})}^2 \leq 4w^2 \Big(\frac{1+\theta}{\theta}\Big)^2 \gamma;$$

$$\int_{\max\{\alpha_{i},\beta_{i}\}}^{\min\{\alpha_{i}+\rho_{i+1}^{A},\beta_{i}+\rho_{i+1}^{B}\}} |\dot{\phi}_{A}(t) - \dot{\phi}_{B}(t)|^{2} dt$$

$$= \left[\frac{\sin(w(\alpha_{i}+\rho_{i+1}^{A}))}{\rho_{i+1}^{A}} - \frac{\sin(w(\beta_{i}+\rho_{i+1}^{B}))}{\rho_{i+1}^{B}}\right]^{2}$$

$$\cdot (\min\{\alpha_{i}+\rho_{i+1}^{A},\beta_{i}+\rho_{i+1}^{B}\} - \max\{\alpha_{i},\beta_{i}\});$$

$$\int_{\min\{\alpha_{i}+\rho_{i+1}^{A},\,\beta_{i}+\rho_{i+1}^{B}\}}^{\max\{\alpha_{i}+\rho_{i+1}^{A},\,\beta_{i}+\rho_{i+1}^{B}\}} |\dot{\phi}_{A}(t)-\dot{\phi}_{B}(t)|^{2} dt$$

$$\leq (\max\{\alpha_{i}+\rho_{i+1}^{A},\,\beta_{i}+\rho_{i+1}^{B}\}-\min\{\alpha_{i}+\rho_{i+1}^{A},\,\beta_{i}+\rho_{i+1}^{B}\})|\dot{\phi}_{A}-\dot{\phi}_{B}||_{L^{\infty}(0,\,T,\,\mathbb{R})}^{2}$$

$$\leq 4w^{2} \left(\frac{1+\theta}{\theta}\right)^{2} |(\alpha_{i}+\rho_{i+1}^{A})-(\beta_{i}+\rho_{i+1}^{B})|\leq 4\gamma w^{2} \left(\frac{1+\theta}{\theta}\right)^{2} \left(1+\frac{2}{w}\right);$$

$$\int_{\min\{T-\rho_m^A, T-\rho_m^B\}}^{\max\{T-\rho_m^A, T-\rho_m^B\}} |\dot{\phi}_A(t) - \dot{\phi}_B(t)|^2 dt \\
\leq (\max\{T-\rho_m^A, T-\rho_m^B\} - \min\{T-\rho_m^A, T-\rho_m^B\}) ||\dot{\phi}_A - \dot{\phi}_B||_{L^{\infty}(0, T, \mathbb{R})}^2 \\
\leq 4w^2 \Big(\frac{1+\theta}{\theta}\Big)^2 |\rho_m^A - \rho_m^B| = 4w^2 \Big(\frac{1+\theta}{\theta}\Big)^2 \frac{1}{w} |x_m - y_m| \leq 4\gamma w \Big(\frac{1+\theta}{\theta}\Big)^2;$$

Finally for the last integral of (4.10.41) we have

$$\int_{\max\{T-\rho_m^A, T-\rho_m^B\}}^{T} |\dot{\phi}_A(t) - \dot{\phi}_B(t)|^2 dt$$
$$= \left[\frac{\sin(w(T-\rho_m^A))}{-\rho_m^A} - \frac{\sin(w(T-\rho_m^B))}{-\rho_m^B}\right]^2 (T - \max\{T-\rho_m^A, T-\rho_m^B\}).$$

Since $\beta_i \to \alpha_i$ and $\rho_m^B \to \rho_m^A$ as $\gamma \to 0$ we have that all the intervals on the right-hand side of (4.10.41) go to zero as γ does, i.e.,

$$\|\dot{\phi}_A - \dot{\phi}_B\|_{L^2(0,T,\mathbb{R})} \to 0 \text{ as } B \to A.$$
 (4.10.42)

Hence from (4.10.39) and (4.10.42) follows the $(\mathcal{A}_{\theta}, W^{1,2}(0, T, \mathbb{R}))$ -continuity of $A \mapsto \phi_A$.

Now from the $(B, \mathcal{A}_{\theta})$ -continuity of the map $b \mapsto A(b)$ (which is equivalent to the δ -continuity of the family \mathcal{Z}) and, from the $(\mathcal{A}_{\theta}, W^{1,2})$ -continuity of $A \mapsto \phi_A$ we have the following corollary

Corollary 4.10.8. The map $b \mapsto \phi_w(\cdot, b)$ is $(B, W^{1,2}(0, T, \mathbb{R}))$ -continuous.

Imitation.

Now we "imitate" the control $z(\cdot, b) \in \mathbb{Z}$ taking values in $\{\pm \Xi e_k, \pm \Xi \delta_{m,n} \mid k \in \mathcal{K}^{N-1}, (m, n) \in S_{N-1}\}$ by a control $z^w(\cdot, b)$ taking values in $\mathbb{R}^{\kappa_{N-1}}$.

Take the solution $u^{\infty}(\cdot, b)$ of the equation

$$u_t^{\infty}(\cdot, b) = -\nu A u^{\infty} - B u^{\infty} + F + z(\cdot, b), \quad u(0) = u_0$$

and, consider its projection onto $\mathbb{R}^{\kappa_{N-1}}$:

$$q^{\infty}(\cdot, b) = P^{\kappa_{N-1}} u^{\infty}(\cdot, b).$$

Let $\{0 = \alpha_0 < \alpha_1 < \cdots < \alpha_m = T\}$ be the end-points of the intervals of constancy of $z(\cdot, b)$. For $w \ge 3$ define the control $z^w(\cdot, b)$ by recursion in the following way:

• In the first interval of constancy $[\alpha_0, \alpha_1]$:

$$z^{w}(\cdot, b) := \begin{cases} z(\cdot, b) & \text{if } z(\cdot, b) \in \{\pm \Xi e_{k} \mid k \in \mathcal{K}^{N-1}\}; \\ v^{\kappa_{N-1}}(q_{1}^{\infty}(\cdot, b) + \sqrt{2\Xi}\phi^{w}(\cdot, b)(e_{m} \pm e_{n}), U_{0}) & ^{22} \\ \text{if } z(\cdot, b) \in \{\pm \Xi \delta_{m,n} \mid (m, n) \in S_{N-1}\}. \end{cases}$$

where U_0 is the projection of u_0 onto $\mathbb{J}_V^{\perp} = (\mathbb{R}^{\kappa_N})_V^{\perp}$ — the orthogonal space to $\mathbb{R}^{\kappa_{N-1}}$ in V and, $q_1^{\infty}(\cdot, b)$ is the restriction of $q^{\infty}(\cdot, b)$ to $[\alpha_0, \alpha_1]$;

• If the control $z^{w}(\cdot, b)$ is already defined in the first p-1 intervals of constancy (up to α_{p-1}), we define it in the p^{th} interval $[\alpha_{p-1}, \alpha_{p}]$ by:

$$z^{w}(\cdot, b) := \begin{cases} z(\cdot, b) & \text{if } z(\cdot, b) \in \{ \pm \Xi e_{k} \mid k \in \mathcal{K}^{N-1} \} \\ v^{\kappa_{N-1}}(q_{p}^{\infty}(\cdot, b) + \sqrt{2\Xi}\phi^{w}(\cdot, b)(e_{m} \pm e_{n}), U^{w}(\alpha_{p-1})) \\ & \text{if } z(\cdot, b) \in \{ \pm \Xi \delta_{m,n} \mid (m, n) \in S_{N-1} \}. \end{cases}$$

²²Here $v^{\kappa_{N-1}}$ is the control given by Lemma 4.10.4, for $J = \mathcal{K}^{N-1}$.

where U^w is the projection onto $(\mathbb{R}^{\kappa_N})^{\perp}_V$ of the solution of the equation

$$u_t^w(\cdot, b) = -\nu A u^w - P^{\nabla} B u^w + F + z^w(\cdot, b), \quad u(0) = u_0, \, t \in [0, \, \alpha_{p-1}]$$

and, $q_p^{\infty}(\cdot, b)$ is the restriction of $q^{\infty}(\cdot, b)$ to $[\alpha_{p-1}, \alpha_p]$.

We shall prove that at time T, $u^w(T)$ goes, uniformly w.r.t. b, to $u^\infty(T)$ in \mathbf{L}^2 -norm as w goes to ∞ , i.e.,

Lemma 4.10.9. For any $\varepsilon > 0$ there exists $w_{\varepsilon} \geq 3$ such that

$$\forall b \in B \forall w \ge w_{\varepsilon} \quad |u^w(T, b) - u^\infty(T, b)| < \varepsilon.$$

We claim that if the statement of the previous Lemma is true and, if we put $\gamma = \frac{1}{2} \left(\|\partial \mathbb{E}_T(u_0, \mathcal{Z}), u_0\|_{l_1} - R \right)$ then, $\mathbb{E}_T(u_0, \mathcal{Z}^{w_{\varepsilon}})$ with $\varepsilon = \frac{\gamma}{\#\mathcal{O}C}$ and C is a constant such that $\|x\|_{l_1} \leq C \|x\|_H$ $(x \in \mathbb{R}^{\#\mathcal{O}})$, covers $\overline{\mathcal{O}}_R(u_0)$ solidly. Indeed, let ϕ be a continuous function defined on the closure of B such that

$$\|\phi - \mathbb{E}_T(u_0, \mathcal{Z}^{w_{\varepsilon}})\|_{C(B, \mathbb{R}^{\#\mathcal{O}}_{\infty})} < \frac{\gamma}{\#\mathcal{O}},$$

then

$$\begin{aligned} \|\phi - \mathbb{E}_T(u_0, \mathcal{Z})\|_{C(B, \mathbb{R}^{\#\mathcal{O}}_{\infty})} \\ \leq \|\phi - \mathbb{E}_T(u_0, \mathcal{Z}^{w_{\varepsilon}})\|_{C(B, \mathbb{R}^{\#\mathcal{O}}_{\infty})} + \|\mathbb{E}_T(u_0, \mathcal{Z}^{w_{\varepsilon}}) - \mathbb{E}_T(u_0, \mathcal{Z})\|_{C(B, \mathbb{R}^{\#\mathcal{O}}_{\infty})} \\ < \frac{\gamma}{\#\mathcal{O}} + C\varepsilon < 2\frac{\gamma}{\#\mathcal{O}}. \end{aligned}$$

For $x \in \overline{\mathcal{O}}_R(u_0)$ we obtain

$$\|\phi - \mathbb{E}(u_0, \mathcal{Z})\|_{C(B, \mathbb{R}^{\#\mathcal{O}}_{\infty})} < \frac{1}{\#\mathcal{O}} \|\mathbb{E}_T(u_0, \mathcal{Z}), u_0\|_{l_1} - R \le \frac{1}{\#\mathcal{O}} \|\mathbb{E}_T(u_0, \mathcal{Z}), x\|_{l_1} \le \|\mathbb{E}_T(u_0, \mathcal{Z}), x\|_{l_{\infty}}.$$

Hence $deg(\phi, B, x) = deg(\mathbb{E}_T, B, x) \neq 0$ and so, $\phi(B)$ covers $\overline{\mathcal{O}}_R(u_0)$.

Therefore what remains to conclude the proof of the back-induction step and then, the solid controllability in Observed Component of system (4.9.3) is the proof of Lemma 4.10.9 that we present in the next section.

4.11 Proof of Lemma 4.10.9.

First we note that at the times α_i we have that the projections of $u^{\infty}(\alpha_i, b)$ and $u^w(\alpha_i, b)$ onto $\mathbb{R}^{\kappa_{N-1}}$ coincide. In fact, denoting the projection of $u^w(\cdot, b)$ by $q^w(\cdot, b)$ we have

$$q^{w}(\cdot, b) = q^{\infty}(\cdot, b) + \sum_{(m,n)\in S_{N-1}} \phi_{m,n}^{w}(\cdot, b)$$

where

$$\phi_{m,n}^{w}(t, b) := \begin{cases} 0 & \text{if } v(t, b) \in \{ \pm \Xi e_k \mid k \in \mathcal{K}^{N-1} \} \\ \sqrt{2\Xi} \phi^{w}(t, b)(e_m \pm e_n) & \text{if } z(t, b) \in \{ \pm \Xi \delta_{m,n} \mid (m, n) \in S_{N-1} \}. \end{cases}$$

Since $\phi^w(\cdot, b)$ vanishes at the points α_i also $\phi_{m,n}^w(t, b)$ does. Hence $q^\infty(\alpha_i, b) = q^w(\alpha_i, b)$ for all $i = 0, 1, \ldots, m$. In particular for i = m we obtain $q^\infty(T, b) = q^w(T, b)$ and so, we need only to compare the projections onto $(\mathbb{R}^{\kappa_{N-1}})_H^{\perp}$. For that we shall need some lemmas:

Lemma 4.11.1. The solution of the controlled NSE

$$u_t = -\nu A u - P^{\nabla} B u + F + v, \qquad u(0) = u_0, \quad t \in [0, T]$$

satisfies

$$||u_t||_{L^2(0, T, H)} \le C_u$$

where C_v is a constant depending only on the norm $\|v\|_{L^2(0,T,H)}$.²³

Moreover for a given family of controls $\{v_{\beta} \mid \beta \in \mathcal{B}\}$ bounded in $L^{2}(0, T, H)$ norm, i.e., if there is a constant K > 0 such that for all $\beta \in \mathcal{B} \|v_{\beta}\|_{L^{2}(0, T, H)} \leq K$, then we can find a constant C_{K} depending only on K such that

$$||u_t^\beta||_{L^2(0,T,H)} \le C_K$$

where u^{β} is the solution of

$$u_t^{\beta} = -\nu A u^{\beta} - P^{\nabla} B u^{\beta} + F + v_{\beta}, \qquad u(0) = u_0, \quad t \in [0, T].$$

Proof. Multiplying the equation by u_t we obtain

$$|u_t|^2 = -\frac{\nu}{2} \frac{d}{dt} ||u||^2 - (P^{\nabla} Bu, u_t) + (F + v, u_t)$$

$$\leq -\frac{\nu}{2} \frac{d}{dt} ||u||^2 + C ||u|| |u|_{[2]} |u_t| + |F||u_t| + |v||u_t|$$

 $\mathbf{so},$

$$\frac{1}{2}|u_t|^2 \le -\frac{\nu}{2}\frac{d}{dt}||u||^2 + C_1||u||^2|u|_{[2]}^2 + C_1|F|^2 + C_1|v|^2$$
$$\int_0^T |u_t(t)|^2 dt \le C_2||u||_{C([0,T],V)}^2 (1 + \int_0^T |u(t)|_{[2]}^2 dt)$$
(4.11.1)

$$+ 2C_1 T |F|^2 + 2C_1 \int_0^T |v|^2 dt.$$
(4.11.2)

Multiplying the equation by u and Au, analogously as we have obtained the "a priori" estimates in the proofs of existence of weak and strong solutions, we obtain the inequalities

$$\begin{aligned} \|u\|_{C([0,T],H)} &\leq |u_0|^2 + C_1 \|F + v\|_{L^2(0,T,V')}^2 \\ \|u\|_{L^2(0,T,V)} &\leq 2\|u\|_{C([0,T],H)} + C_1 \nu\|F + v\|_{L^2(0,T,V')}^2 \\ \|u\|_{C([0,T],V)} &\leq \exp(C_2 \|u\|_{C([0,T],H)} \|u\|_{L^2(0,T,V)}) \Big(\|u_0\|^2 \\ &+ C_2 \|F + v\|_{L^2(0,T,H)}^2 \Big) \\ \|u\|_{L^2(0,T,D(A))} &\leq 2\|u\|_{C([0,T],V)} + C_3 \|u\|_{C([0,T],H)}^2 \|u\|_{C([0,T],V)}^4 \\ &+ C_3 \|F + v\|_{L^2(0,T,H)}^2. \end{aligned}$$
(4.11.3)

²³Recall that we have fixed ν , F, T and u_0 . Otherwise the constant would depend on them.

from which, looking at (4.11.2), we conclude that $||u_t||_{L^2(0,T,H)} \leq C_v$ for some constant C_v depending only on $||v||_{L^2(0,T,H)}$.

The statement relative to the family v_{β} is also clearly true from the previous expressions:

In each one of the previous estimates if we "replace $||v_{\beta}||_{L^{2}(0, T, H)}$ by the bound K" we obtain bounds independent of β . (4.11.4)

Corollary 4.11.2. There is a constant $C_{\infty} > 0$ such that

$$\forall b \in B \quad \|u_t^{\infty}(\cdot, b)\|_{L^2(0, T, H)} \le C_{\infty}.$$

Proof. For all $t \in [0, T]$, $b \in B$ we have

$$\|v(t, b)\|_{l_1} \le \Xi \max \Big\{ \|x\|_{l_1} \mid x \in \{e_k, \, \delta_{m,n} \mid k \in \mathcal{K}^{N-1}, \, (m, n) \in S_{N-1} \} \Big\}$$

so, we have that for some constant K_2

$$||v(\cdot, b)||_{L^{\infty}(0, T, H)} \leq K_2.$$

The result follows by the last part of Lemma 4.11.1 and the continuity of the inclusion $L^{\infty}(0, T, H) \rightarrow L^{2}(0, T, H)$.

If $v(t, b) \in \{\pm \Xi e_k \mid k \in \mathcal{K}^{N-1}\}$ in the interval of constancy $[\alpha_{i-1}, \alpha_i]$ let us call this interval of the **first kind**. Otherwise, if $z(t, b) \in \{\pm \Xi \delta_{m,n} \mid (m, n) \in S_{N-1}\}$ on $[\alpha_{i-1}, \alpha_i]$ we call the interval of the **second kind**.

Now we note that for $u^w(\cdot, b)$ we can not find a bound for $||u_t^w||_{L^2(0, T, H)}$ independent of the parameter w because the projection $q^w(\cdot, b)$ in an interval of constancy $I = [\alpha_{i-1}, \alpha_i]$ of the second kind, say $v(t, b) = \delta_{m,n}$ on I, reads

$$q^{w}(t, b) = q^{\infty}(t, b) + \phi^{w}_{m,n}(t, b)$$

and,

$$\begin{aligned} &\|\phi_t^w\|_{L^2(0,T,\mathbb{R})}^2 \\ &\geq \int_{[\alpha_{i-1}+\frac{L}{w},\,\alpha_i-\frac{L}{w}]} |\phi_t^w(t)|^2 \, dt = w^2 \int_{[\alpha_{i-1}+\frac{L}{w},\,\alpha_i-\frac{L}{w}]} \cos^2(wt) \, dt \\ &= w^2 \bigg[\frac{1}{2} (L-2\frac{L}{w}) + \frac{1}{4w} \Big(\sin(2w(\alpha_i-\frac{L}{w})) - \sin(2w(\alpha_{i-1}+\frac{L}{w})) \Big) \bigg]. \end{aligned}$$

where L is the length of I. Since $w \ge 3$ (and L > 0) we obtain

$$\|\phi_t^w\|_{L^2(0,T,\mathbb{R})}^2 \ge w^2 \left[\frac{L}{6} + \frac{1}{4w} \left(\sin(2w(\alpha_i - \frac{L}{w})) - \sin(2w(\alpha_{i-1} + \frac{L}{w}))\right)\right].$$

Then we see that when w goes to ∞ also $\|\phi_t^w\|_{L^2(0,T,\mathbb{R})}$ does and then, so does $\|u_t^w\|_{L^2(0,T,H)}$.

If we consider the projection $U^w(\cdot, b)$ of $u^w(\cdot, b)$ onto $(\mathbb{R}^{\kappa_{N-1}})_H^{\perp}$, there holds that we can find a bound for $\|U_t^w(\cdot, b)\|_{L^2(0, T, (\mathbb{R}^{\kappa_{N-1}})_H^{\perp})}$ independent of both parameters b and w. If fact multiplying by $U_t^w(\cdot, b)$ the equation

$$U_t^w(\cdot, b) = -\nu A U^w(\cdot, b) - P^{-\kappa_{N-1}} P^{\nabla} B u^w(\cdot, b) + P^{-\kappa_{N-1}} F$$

that is satisfied by U^w we obtain

$$|U_t^w(t,b)|^2 \le -\frac{\nu}{2} \frac{d}{dt} ||U^w(t,b)||^2 + |F||U_t^w(t,b)| + C||U^w(t,b) + q^w(t,b)||U^w(t,b) + q^w(t,b)|_{[2]}|U_t^w(t,b)|.$$
(4.11.5)

After using appropriate Young Inequalities and integrating

$$\int_{0}^{T} |U_{t}^{w}(t,b)|^{2} dt
\leq C_{1} ||U^{w}(\cdot,b)||_{C([0,T],(\mathbb{R}^{\kappa_{N-1}})_{V}^{\perp})}^{2} + C_{1}|F|^{2}
+ C_{1} ||U^{w}(\cdot,b) + q^{w}(\cdot,b)||_{C([0,T],V)}^{2} ||U^{w}(\cdot,b) + q^{w}(\cdot,b)||_{L^{2}(0,T,D(A))}^{2}
\leq C ||U^{w}(\cdot,b)||_{C([0,T],(\mathbb{R}^{\kappa_{N}})_{V}^{\perp})}^{2} + T|F|^{2}
+ C \Big[||U^{w}(\cdot,b)||_{C([0,T],(\mathbb{R}^{\kappa_{N}})_{V}^{\perp})}^{2} + ||q^{w}(\cdot,b)||_{C([0,T],\mathbb{R}^{\kappa_{N-1}})}^{2} \Big]
\times \Big[||U^{w}(\cdot,b)||_{L^{2}(0,T,(\mathbb{R}^{\kappa_{N-1}})_{D(A)}^{\perp})}^{2} + ||q^{w}(\cdot,b)||_{L^{2}(0,T,\mathbb{R}^{\kappa_{N-1}})}^{2} \Big].
(4.11.6)$$

Multiplying the equation by $U^w(\cdot, b)$ and by $AU^w(\cdot, b)$ we obtain that $U^w(\cdot, b)$ satisfy the estimates (4.10.14), (4.10.15), (4.10.17) and (4.10.18) with $U^{J,L}$ replaced by $U^w(\cdot, b)$ and we can easily see that there is a constant $C_{w,b}$ depending only on $\|q^w(\cdot, b)\|_{C([0, T], \mathbb{R}^{\kappa_{N-1}})}$ such that

$$\|U^{w}(\cdot,b)\|_{C([0,T],(\mathbb{R}^{\kappa_{N}})_{V}^{\perp})} \leq C_{w,b} \& \|U^{w}(\cdot,b)\|_{L^{2}(0,T,(\mathbb{R}^{\kappa_{N-1}})_{D(A)}^{\perp})} \leq C_{w,b}.$$
(4.11.7)

Moreover the family $\{U_t^w(\cdot, b)\}$ is uniformly bounded, w.r.t. b and w, in the norm of $L^2(0, T, H)$ because, in the "a priori"-like estimates above and in (4.11.6) we can take bounds independent of the parameters depending only in the bound

$$\max_{(b,w)\in B\times[3,+\infty[}\{\|q^{w}(\cdot,b)\|_{C([0,T],\mathbb{R}^{\kappa_{N-1}})}\}$$

$$\leq \|q^{\infty}(\cdot,b)\|_{C([0,T],\mathbb{R}^{\kappa_{N-1}})} + \left\|\sum_{(m,n)\in S_{N-1}}\phi_{m,n}^{w}(\cdot,b)\right\|_{C([0,T],\mathbb{R}^{\kappa_{N-1}})} \leq C$$

where C does not depend neither on b nor on w. Indeed by (4.10.36) the family $v(\cdot, b)$ is uniformly bounded in $L^2(0, T, \mathbb{R}^{\kappa_{N-1}})$ -norm which implies that $q^{\infty}(\cdot, b)$ is uniformly bounded in $C([0, T], \mathbb{R}^{\kappa_{N-1}})$ -norm. Indeed replacing u by $u^{\infty}(\cdot, b)$ in the equations (4.11.3) we see that, since the family of controls $\{v(\cdot, b)\}$ is uniformly bounded in $L^2(0, T, H)$ norm, we can find a uniform bound D for the solutions $u^{\infty}(\cdot, b)$ in anyone of the norms $C([0, T], H), L^2(0, T, V), C([0, T], V)$ and $L^2(0, T, D(A))$. In particular

 $\begin{aligned} \|q^{\infty}(\cdot, b)\|_{C([0,T], \mathbb{R}^{\kappa_{N-1}})} &\leq D_1 \|u^{\infty}(\cdot, b)\|_{C([0,T], H)} \leq D_1 D. \\ \text{On the other hand } \|\sum_{(m,n)\in S_{N-1}} \phi_{m,n}^w(\cdot, b)\|_{C([0,T], \mathbb{R}^{\kappa_{N-1}})} \end{aligned}$ never exceeds the

value $2\sqrt{2\Xi}$. The we can the write:

Corollary 4.11.3. There is a constant C > 0 such that for all (w, b) in the product $[3, +\infty] \times B$:

$$\|U_t^w(\cdot, b)\|_{L^2(0, T, (R^{\kappa_{N-1}})_H^{\perp})} \le C.$$

Another Lemma we shall use is the following:

Lemma 4.11.4. Let $\{z(\cdot, \sigma) \in W^{1,2}([t_i, t_f], \mathbb{R}) \mid \sigma \in \Sigma\}$ be a uniformly bounded, w.r.t. σ , family, i.e.,

$$\exists C > 0 \, \forall \sigma \in \Sigma \quad \|z(\cdot, \sigma)\|_{W^{1,2}([t_i, t_f], \mathbb{R})} \leq C.$$

Then there is a constant D_1 depending only in C and in the length $t_f - t_i$ of the interval $[t_i, t_f]$ and so, independent of σ and w, such that

$$\|\sin(wt)z(t,\sigma)\|_{rx} \le D_1 w^{-1}, \quad and \quad \|\cos(wt)z(t,\sigma)\|_{rx} \le D_1 w^{-1}.$$

Proof. The Lemma follows by direct computation: Let s, r belong to $[t_i, t_f]$:

$$\begin{split} &\int_{s}^{r} \sin(wt)z(t,\,\sigma)\,dt \\ &= -w^{-1}\int_{s}^{r} -w\sin(wt)z(t,\,\sigma)\,dt = w^{-1}\int_{s}^{r} \left(\frac{d}{dt}\cos(wt)\right)z(t,\,\sigma)\,dt \\ &= -w^{-1}\left(\left[\cos(wt)z(t,\,\sigma)\right]_{s}^{r} - \int_{s}^{r}\cos(wt)\left(\frac{d}{dt}z(t,\,\sigma)\right)\,dt\right) \\ &\leq -w^{-1}\left[\cos(wr)z(r,\,\sigma) - \cos(ws)z(s,\,\sigma)\right] \\ &+ w^{-1}\left(\int_{s}^{r}\cos^{2}(wt)\,dt\right)^{\frac{1}{2}}\left(\int_{s}^{r} \|\frac{d}{dt}z(t,\,\sigma)\|^{2}\,dt\right)^{\frac{1}{2}} \\ &\leq 2w^{-1}\|z(t,\,\sigma)\|_{C([t_{i},\,t_{f}],\,\mathbb{R})} + w^{-1}|r-s|^{\frac{1}{2}}C. \end{split}$$

By the continuity of the embedding $W^{1,2} \to C^0$ there is a constant C_1 such that $||u||_{C^0} \leq C_1 ||u||_{W^{1,2}}$. Putting $C_2 = \max\{C, C_1C\}$ we obtain

 $\int_{s}^{r} \sin(wt) z(t, \sigma) \, dt \le w^{-1} C_2 \left(2 + (t_f - t_i)^{\frac{1}{2}} \right).$

Analogously we arrive to

or

$$\int_{s}^{t} \cos(wt) z(t, \sigma) \, dt \le w^{-1} C_2 \left(2 + (t_f - t_i)^{\frac{1}{2}} \right).$$

Hence

$$\forall \sigma \in \Sigma \,\forall s, \, r \in [t_i, \, t_f] \left\{ \int_s^r \sin(wt) z(t, \, \sigma) \, dt \le w^{-1} C_2 \left(2 + (t_f - t_i)^{\frac{1}{2}} \right) \\ \int_s^r \cos(wt) z(t, \, \sigma) \, dt \le w^{-1} C_2 \left(2 + (t_f - t_i)^{\frac{1}{2}} \right). \right.$$

Therefore

$$\forall \sigma \in \Sigma \begin{cases} \|\sin(wt)z(t,\,\sigma)\|_{rx} \le w^{-1}C_2\left(2 + (t_f - t_i)^{\frac{1}{2}}\right) \\ \|\cos(wt)z(t,\,\sigma)\|_{rx} \le w^{-1}C_2\left(2 + (t_f - t_i)^{\frac{1}{2}}\right). \end{cases}$$

Choose $D_1 = C_2 (2 + (t_f - t_i)^{\frac{1}{2}}).$

Corollary 4.11.5. Let $\{z(\cdot, \sigma) \in W^{1,2}([0, T], \mathbb{R}) \mid \sigma \in \Sigma\}$ be a uniformly bounded, w.r.t. σ , family:

$$\exists C > 0 \,\forall \sigma \in \Sigma \quad \|z(\cdot, \, \sigma)\|_{W^{1,2}([0, \, T], \,\mathbb{R})} \le C.$$

Then there is a constant D_2 depending only in C and so, independent of σ and of the parameter b of our controls, such that

$$\|\phi_w(\cdot, b)z(\cdot, \sigma)\|_{rx} \le D_2 w^{-1}.$$

Proof. Let s, r be in [0, T] and suppose without loss of generality that s < r. Put $D := \{t \in [0, T] \mid \phi_w(t, b) \neq \sin(wt)\}.$

$$\int_{s}^{T} \phi_{w}(t, b) z(t, \sigma) dt = \int_{D \cap [s, r]} \phi_{w}(t, b) z(t, \sigma) dt + \int_{[s, r] \setminus D} \sin(wt) z(t, \sigma) dt$$
$$\leq 2 \frac{T}{w} C_{1} C + m w^{-1} C_{2} \left(2 + T^{\frac{1}{2}}\right)$$
$$\leq w^{-1} C_{2} \left(2T + 2m + T^{\frac{1}{2}}m\right).$$
²⁴

Choose $D_2 = C_2 (2T + 2m + T^{\frac{1}{2}}m).$

Now we compare the projections $U^w(\cdot, b)$ and $U^\infty(\cdot, b)$ of respectively $u^w(\cdot, b)$ and $u^\infty(\cdot, b)$ onto $(\mathbb{R}^{\kappa_{N-1}})^{\perp}_H$. We claim that at time α_i , $(i = 1, \ldots, m)$ there holds

$$|U^w(\alpha_i, b) - U^\infty(\alpha_i, b)| \le \overline{C_i} w^{-1}$$
(4.11.8)

where $\overline{C_i}$ is a constant independent of the parameters w and b. In particular at time T there holds

$$|U^{w}(T, b) - U^{\infty}(T, b)| \le \overline{C_{m}} w^{-1}.$$
(4.11.9)

Note that (4.11.9) implies Lemma 4.10.9 because as we have seen at time T we have $q^w(T, b) = q^{\infty}(T, b)$. To prove (4.11.9) we shall compare $U^w(\cdot, b)$ and $U^{\infty}(\cdot, b)$ in each interval of constancy:

• In an interval of the first kind $U^w(\cdot,\,b)$ and $U^\infty(\cdot,\,b)$ satisfy the same equation

$$\begin{aligned} U_t^{\infty}(\cdot, b) &= -\nu A U^{\infty}(\cdot, b) - P^{-\kappa_{N-1}} P^{\nabla} \big(U^{\infty}(\cdot, b) + q^{\infty}(\cdot, b) \big) + P^{-\kappa_{N-1}} F; \\ U_t^{w}(\cdot, b) &= -\nu A U^{w}(\cdot, b) - P^{-\kappa_{N-1}} P^{\nabla} \big(U^{w}(\cdot, b) + q^{\infty}(\cdot, b) \big) + P^{-\kappa_{N-1}} F \end{aligned}$$

• In an interval of the second kind, say $v(\cdot, b) = \pm \Xi \delta_{m,n}$ $(m,n) \in S_{N-1}$, $U^w(\cdot, b)$ and $U^\infty(\cdot, b)$ satisfy the equations

$$\begin{aligned} U_t^{\infty}(\cdot, b) &= -\nu A U^{\infty}(\cdot, b) - P^{-\kappa_{N-1}} P^{\nabla} \big(U^{\infty}(\cdot, b) + q^{\infty}(\cdot, b) \big) + P^{-\kappa_{N-1}} F; \\ U_t^{w}(\cdot, b) &= -\nu A U^{w}(\cdot, b) - P^{-\kappa_{N-1}} P^{\nabla} \Big[U^{w}(\cdot, b) + q^{\infty}(\cdot, b) \\ &+ \sqrt{2\Xi} \phi^{w}(\cdot, b) (e_m \pm e_n) \Big] + P^{-\kappa_{N-1}} F \pm \Xi \delta_{m,n}. \end{aligned}$$

 $^{^{24} \}mathrm{Note}$ that m being the number of intervals of constancy, then $[s,\,r] \setminus D$ is a union of at most m intervals.

Define the difference

$$\eta^w(\cdot, b) := U^w(\cdot, b) - U^\infty(\cdot, b).$$

For an interval $[\alpha_{i-1}, \alpha_i]$ of second kind we find

$$\eta_t^w(s, b) = -\nu A \eta^w(\cdot, b) + P^{-\kappa_{N-1}} P^{\nabla} \Big[U^\infty(\cdot, b) + q^\infty(\cdot, b) \Big] - P^{-\kappa_{N-1}} P^{\nabla} \Big[U^w(\cdot, b) + q^\infty(\cdot, b) + \sqrt{2\Xi} \phi^w(\cdot, b) (e_m \pm e_n) \Big) \pm \Xi P^{-\kappa_{N-1}} \delta_{m,n}.$$

Multiplying by $\eta^w(\cdot, b)$ we obtain

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} |\eta_t^w(s, b)|^2 \\ &= -\nu \|\eta^w(s, b)\|^2 + \left[P^{-\kappa_{N-1}} P^{\nabla} (U^{\infty}(s, b) + q^{\infty}(s, b)), \, \eta^w(s, b) \right] \\ &- \left[P^{-\kappa_{N-1}} P^{\nabla} (U^w(s, b) + q^{\infty}(s, b) + \sqrt{2\Xi} \phi^w(s, b) (e_m \pm e_n)), \, \eta^w(s, b) \right] \\ &\pm \left(\Xi P^{-\kappa_{N-1}} \delta_{m,n}, \, \eta^w(s, b) \right) \end{aligned}$$

 $\mathrm{so},$

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} |\eta^{w}(s, b)|^{2} + \nu \|\eta^{w}(s, b)\|^{2} \\ \leq &- \left(P^{-\kappa_{N-1}} P^{\nabla} \left(U^{w}(s, b) + q^{\infty}(s, b) \right), \eta^{w}(s, b) \right) \\ &+ \left(P^{-\kappa_{N-1}} P^{\nabla} \left(U^{\infty}(s, b) + q^{\infty}(s, b) \right), \eta^{w}(s, b) \right) \\ &- \left(P^{-\kappa_{N-1}} P^{\nabla} \left(U^{w}(s, b) + q^{\infty}(s, b), \sqrt{2\Xi} \phi^{w}(s, b)(e_{m} \pm e_{n}) \right), \eta^{w}(s, b) \right) \\ &- \left(P^{-\kappa_{N-1}} P^{\nabla} \left(\sqrt{2\Xi} \phi^{w}(s, b)(e_{m} \pm e_{n}), U^{w}(s, b) + q^{\infty}(s, b) \right), \eta^{w}(s, b) \right) \\ &- \left(P^{-\kappa_{N-1}} P^{\nabla} \left(\sqrt{2\Xi} \phi^{w}(s, b)(e_{m} \pm e_{n}) \right), \eta^{w}(s, b) \right) \\ &+ \left(\Xi P^{-\kappa_{N-1}} \delta_{m,n}, \eta^{w}(s, b) \right) \\ &= A_{1}(s) + A_{2}(s) + A_{3}(s) + A_{4}(s) \end{aligned}$$
(4.11.10)

where we define

$$\begin{split} A_{1}(s) &:= -\left(P^{-\kappa_{N-1}}P^{\nabla}\big(U^{w}(s,\,b) + q^{\infty}(s,\,b),\,\eta^{w}(s,\,b)\big) \\ &+ \left(P^{-\kappa_{N-1}}P^{\nabla}\big(U^{\infty}(s,\,b) + q^{\infty}(s,\,b)\big),\,\eta^{w}(s,\,b)\big) \\ A_{2}(s) &:= \\ &- \left(P^{-\kappa_{N-1}}P^{\nabla}\big(U^{w}(s,\,b) + q^{\infty}(s,\,b),\,\sqrt{2\Xi}\phi^{w}(s,\,b)(e_{m} \pm e_{n})\big),\,\eta^{w}(s,\,b)\right) \\ A_{3}(s) &:= \\ &- \left(P^{-\kappa_{N-1}}P^{\nabla}\big(\sqrt{2\Xi}\phi^{w}(s,\,b)(e_{m} \pm e_{n}),\,U^{w}(s,\,b) + q^{\infty}(s,\,b)\big),\,\eta^{w}(s,\,b)\right) \\ A_{4}(s) &:= -\left(P^{-\kappa_{N-1}}P^{\nabla}\big(\sqrt{2\Xi}\phi^{w}(s,\,b)(e_{m} \pm e_{n})\big),\,\eta^{w}(s,\,b)\right) \\ &\pm \left(\Xi P^{-\kappa_{N-1}}\delta_{m,n},\,\eta^{w}(s,\,b)\right). \end{split}$$

To estimate $|\eta^w(z, b)|^2 \quad z \in [\alpha_{i-1}, \alpha_i]$ we will integrate (4.11.10). In the second member, for A_1 we have:

$$|A_1| \le C |\eta^w(z, b)| \|\eta^w(z, b)\| \|U^\infty(z, b) + q^\infty(z, b)\|$$
(4.11.11)

and by Young Inequality we arrive to

$$\frac{d}{dt} |\eta^w(z, b)|^2 + \nu ||\eta^w(z, b)||^2$$

$$\leq D |\eta^w(z, b)|^2 ||U^\infty(z, b) + q^\infty(z, b)||^2 + 2\sum_{r=2}^4 A_r(z)$$

and, by Gronwall Inequality we obtain

$$|\eta^{w}(z, b)|^{2} \leq |\eta^{w}(\alpha_{i-1}, b)|^{2} \exp\left(D \int_{\alpha_{i-1}}^{\alpha_{i}} \|U^{\infty}(s, b) + q^{\infty}(s, b)\|^{2} ds\right) + 2\sum_{r=2}^{4} \int_{\alpha_{i-1}}^{z} A_{r}(s) E(s) ds$$
(4.11.12)

with $E(s) := \exp \int_{z}^{s} -D \| u^{\infty}(t, b) \|^{2} dt$. For r = 2 we have

$$\int_{\alpha_{i-1}}^{z} A_{2}(s)E(s) ds = -\sqrt{2\Xi} \int_{\alpha_{i-1}}^{z} \phi^{w}(s, b) \Big(P^{-\kappa_{N-1}}P^{\nabla} \big(U^{w}(s, b) + q^{\infty}(s, b), e_{m} \big), \eta^{w}(s, b) \Big) E(s) ds = \sqrt{2\Xi} \int_{\alpha_{i-1}}^{z} \phi^{w}(s, b) \Big(P^{-\kappa_{N-1}}P^{\nabla} \big(U^{w}(s, b) + q^{\infty}(s, b), e_{n} \big), \eta^{w}(s, b) \Big) E(s) ds.$$

$$(4.11.13)$$

Now we estimate the derivative of the product

$$\begin{split} & \left(P^{-\kappa_{N-1}}P^{\nabla}\left(U^{w}(s,\,b)+q^{\infty}(s,\,b),\,e_{m}\right),\,\eta^{w}(s,\,b)\right)E(s):\\ & \frac{d}{ds}\left(P^{-\kappa_{N-1}}P^{\nabla}\left(U^{w}(s,\,b)+q^{\infty}(s,\,b),\,e_{m}\right),\,\eta^{w}(s,\,b)\right)\\ & = \left(P^{-\kappa_{N-1}}P^{\nabla}\left(\dot{U}^{w}(s,\,b)+\dot{q}^{\infty}(s,\,b),\,e_{m}\right),\,\eta^{w}(s,\,b)\right)\\ & + \left(P^{-\kappa_{N-1}}P^{\nabla}\left(U^{w}(s,\,b)+q^{\infty}(s,\,b),\,e_{m}\right),\,\dot{\eta}^{w}(s,\,b)\right)\\ & \leq C|\dot{U}^{w}(s,\,b)+\dot{q}^{\infty}(s,\,b)||e_{m}|_{[2]}||\eta^{w}(s,\,b)||\\ & + C||U^{w}(s,\,b)+q^{\infty}(s,\,b)||e_{m}|_{[2]}|\dot{\eta}^{w}(s,\,b)||\\ & \leq D|\dot{U}^{w}(s,\,b)+\dot{q}^{\infty}(s,\,b)|||\eta^{w}(s,\,b)|| + D||U^{w}(s,\,b)+q^{\infty}(s,\,b)||\dot{\eta}^{w}(s,\,b)| \end{split}$$

so,

$$\begin{split} &\int_{\alpha_{i-1}}^{z} |\frac{d}{ds} \Big(P^{-\kappa_{N-1}} P^{\nabla} \big(U^{w}(s,\,b) + q^{\infty}(s,\,b),\,e_{m} \big),\,\eta^{w}(s,\,b) \Big)|^{2} \,ds \\ &\leq D_{1} \int_{\alpha_{i-1}}^{\alpha_{i}} |\dot{U}^{w}(s,\,b) + \dot{q}^{\infty}(s,\,b)|^{2} \|\eta^{w}(s,\,b)\|^{2} \,ds \\ &+ D_{1} \int_{\alpha_{i-1}}^{\alpha_{i}} \|U^{w}(s,\,b) + q^{\infty}(s,\,b)\|^{2} |\dot{\eta}^{w}(s,\,b)|^{2} \,ds \end{split}$$

and, for
$$\left| \left(P^{-\kappa_{N-1}} P^{\nabla} (U^w(s, b) + q^{\infty}(s, b), e_m), \eta^w(s, b) \right) \right|$$
 we obtain the bound
 $\left| \left(P^{-\kappa_{N-1}} P^{\nabla} (U^w(s, b) + q^{\infty}(s, b), e_m), \eta^w(s, b) \right) \right|$
 $\leq C \| U^w(s, b) + q^{\infty}(s, b) \| \| e_m \| \| \eta^w(s, b) \| \leq D \| U^w(s, b) + q^{\infty}(s, b) \| \| \eta^w(s, b) \|.$

Hence, by (4.11.4) and (4.11.7) and Corollaries 4.11.2, 4.11.3 we conclude that the family

$$\left\{ \left(P^{-\kappa_{N-1}} P^{\nabla} (U^w(s, b) + q^{\infty}(s, b), e_m), \eta^w(s, b) \right) \right\}$$

is uniformly bounded, w.r.t. w and b, in $W^{1,2}([\alpha_{i-1}, \alpha_i])$ -norm.

For E(s) we find

$$|E(s)| \le \exp(TD \| u^{\infty}(\cdot, b) \|_{C([0, T], V)}) \le C_1$$
(4.11.14)

$$\left|\frac{d}{ds}E(s)\right| \le \|u^{\infty}(\cdot, b)\|_{C([0, T], V)}C_1 \le C_2 \tag{4.11.15}$$

where C_1 and C_2 do not depend neither on b nor on s. In particular E is uniformly bounded in $W^{1,\infty}([\alpha_{i-1}, \alpha_i], \mathbb{R})$ -norm and then also the family

$$\left\{ \left(P^{-\kappa_{N-1}} P^{\nabla} (U^w(s, b) + q^{\infty}(s, b), e_m), \eta^w(s, b) \right) E(s) \right\}$$

is uniformly bounded in $W^{1,2}([\alpha_{i-1}, \alpha_i], \mathbb{R})$ -norm.

Therefore by (4.11.13) and Corollary 4.11.5 there is a constant K_1 , independent of z and of the parameters w and b, such that

$$\left| \int_{\alpha_{i-1}}^{z} \phi^{w}(s, b) \Big(P^{-\kappa_{N-1}} P^{\nabla} \big(U^{w}(s, b) + q^{\infty}(s, b), e_{m} \big), \eta^{w}(s, b) \Big) E(s) \, ds \right|$$

$$\leq K_{1} w^{-1}.$$

If we replace m by n we obtain a similar estimate and then, we conclude that for some constant K_2 independent of z, w and b:

$$\left| \int_{\alpha_{i-1}}^{z} A_2(s) E(s) \, ds \right| \le K_2 w^{-1}. \tag{4.11.16}$$

Analogously we conclude that for some constant K_3 independent of z, w and b:

$$\left| \int_{\alpha_{i-1}}^{z} A_3(s) E(s) \, ds \right| \le K_3 w^{-1}. \tag{4.11.17}$$

Recalling equation (3.3.8) and supposing without loss of generality that m < n, we have

$$P^{\nabla}B\left[\sqrt{2\Xi}\phi^{w}(s,b)(e_{m}\pm e_{n})\right] = 2\Xi\left(\phi^{w}(s,b)\right)^{2}P^{\nabla}B\left[(e_{m}\pm e_{n})\right]$$
$$= \pm 2\Xi\left(\phi^{w}(s,b)\right)^{2}\delta_{m,n}.$$

Therefore

$$\begin{split} &\int_{\alpha_{i-1}}^{z} A_4(s) E(s) \, ds \\ &= \int_{\alpha_{i-1}}^{z} \left[\mp 2\Xi \left(\phi^w(s, b) \right)^2 \pm \Xi \right] \left(P^{-\kappa_{N-1}} \delta_{m,n}, \eta^w(s, b) \right) E(s) \, ds \\ &= \int_{\alpha_{i-1}}^{z} \left[\mp 2\Xi \left(\phi^w(s, b) \right)^2 \pm \Xi \right] \left(\delta_{m,n}, \eta^w(s, b) \right) E(s) \, ds \\ &= \mp \Xi \int_{\alpha_{i-1}}^{z} \left[2 \left(\phi^w(s, b) \right)^2 - 1 \right] \left(\delta_{m,n}, \eta^w(s, b) \right) E(s) \, ds \\ &\leq \mp \Xi \int_{I} \cos(2wt) \left(\delta_{m,n}, \eta^w(s, b) \right) E(s) \, ds \\ &\mp \Xi \int_{[\alpha_{i-1}, z] \setminus I} |(\delta_{m,n}, \eta^w(s, b))| |E(s)| \, ds. \end{split}$$

where $I = [\alpha_{i-1}, z] \cap [\alpha_{i-1} + \frac{L_i}{w}, \alpha_i - \frac{L_i}{w}]$ and $L_i = \alpha_i - \alpha_{i-1}$. By (4.11.14) we have that

$$\int_{[\alpha_{i-1},z]\setminus I} |(\delta_{m,n},\eta^w(s,b))| E(s)| ds$$

$$\leq C_1 |(\delta_{m,n},\eta^w(s,b))|_{C([0,T],\mathbb{R})} \frac{2L_i}{w} \leq w^{-1} K_0 ||\eta^w(s,b)\rangle ||_{C([0,T],(\mathbb{R}^{\kappa_{N-1}})_H^{\perp})}$$

where the constant K_0 can be taken independent of $(m, n) \in S_{N-1}$, because S_{N-1} is finite.

For the derivative of the product $(\delta_{m,n}, \eta^w(s, b))$ we find

$$\frac{d}{dt}(\delta_{m,n}, \, \eta^w(s, \, b)) = (\delta_{m,n}, \, \dot{\eta}^w(s, \, b)) \le K_1 |\dot{\eta}^w(s, \, b)|$$

so, by (4.11.4), (4.11.7) and corollaries 4.11.2 and 4.11.3 we conclude that for some constant C_2 independent of w and b: $\|(\delta_{m,n}, \eta^w(\cdot, b))\|_{W^{1,2}([0,T],\mathbb{R})} \leq C_2$. So by the uniform boudedness of $E(\cdot)$ in $W^{1,\infty}([0,T],\mathbb{R})$ we conclude the uniform boudedness of $(\delta_{m,n}, \eta^w(s, b))E(s)$ in $W^{1,2}([0,T],\mathbb{R})$.

By Lemma 4.11.4 there is a constant C_1 , independent of w and b, such that

$$\left|\int_{I}\cos(2wt)\Big(\delta_{m,n},\,\eta^{w}(s,\,b)\Big)E(s)\,ds\right|\leq C_{1}(2w)^{-1}.$$

Hence

$$\left| \int_{\alpha_{i-1}}^{z} A_4(s) E(s) \, ds \right| \le \frac{C_1}{2} w^{-1} + w^{-1} K_0 \| \eta^w(s, b) \|_{C([0, T], \, (\mathbb{R}^{\kappa_{N-1}})_H^{\perp})}$$

and, again by (4.11.4) and (4.11.7) we arrive to

$$\left| \int_{\alpha_{i-1}}^{z} A_4(s) E(s) \, ds \right| \le K_2 w^{-1}. \tag{4.11.18}$$

where K_2 is independent of b and w.

By equation (4.11.12) and estimates (4.11.16), (4.11.17) and (4.11.18) we obtain

$$\begin{aligned} &|\eta^w(z, b)|^2 \\ \leq &|\eta^w(\alpha_{i-1}, b)|^2 \exp\left(D\int_0^T \|u^\infty(s, b)\|^2 \, ds\right) + 2\sum_{r=2}^4 \int_{\alpha_{i-1}}^z A_r(s) E(s) \, ds \\ \leq &D_1 |\eta^w(\alpha_{i-1}, b)|^2 + D_2 w^{-1}. \end{aligned}$$

In an interval of the first kind we just have $A_2 = A_3 = A_4 = 0$ on (4.11.10) and then we obtain

$$|\eta^w(z, b)|^2 \le |\eta^w(\alpha_{i-1}, b)|^2 \exp\left(D\int_0^T \|U^\infty(s, b) + q^\infty(s, b)\|^2 \, ds\right) \le D_1 |\eta^w(\alpha_{i-1}, b)|^2.$$

Therefore in any interval of constancy (either of first or second kind) we have

$$|\eta^w(z, b)|^2 \le D_1 |\eta^w(\alpha_{i-1}, b)|^2 + D_2 w^{-1}$$
(4.11.19)

where D_1 and D_2 are independent of $z \in [\alpha_{i-1}, \alpha_i]$, of the parameters w and b and of $i \in \{1, 2, \ldots, m\}$, i.e., of the interval of constancy.

We prove (4.11.8) by induction on *i*:

• i = 1: At time α_1 , by (4.11.19), we have

$$|U^w(\alpha_1, b) - U^\infty(\alpha_1, b)|^2 \le D_2 w^{-1} \quad ^{25}$$

Choose $\overline{C_1} = D_2$.

• Induction Step: Suppose that at time α_{i-1} we have

$$|U^w(\alpha_{i-1}, b) - U^\infty(\alpha_{i-1}, b)|^2 \le \overline{C_{i-1}}w^{-1}.$$

Then by (4.11.19), we have

$$|U^{w}(\alpha_{i-1}, b) - U^{\infty}(\alpha_{i-1}, b)|^{2} \leq D_{1}(\overline{C_{i-1}}w^{-1})^{2} + D_{2}w^{-1} = \left[D_{1}\overline{C_{i-1}}^{2}w^{-1} + D_{2}\right]w^{-1} \leq \left[D_{1}\overline{C_{i-1}}^{2}\frac{1}{3} + D_{2}\right]w^{-1}.$$

Choose $\overline{C_i} = D_1 \overline{C_{i-1}}^2 \frac{1}{3} + D_2.$

Therefore (4.11.8) holds.

²⁵Remember that both u^{∞} and u^w start at u_0 at time 0.

Chapter 5

L^2 -Approximate Controllability.

The following Proposition says that for any T > 0, system (4.9.3) is time-T approximately controllable in L²-norm.

Proposition 5.0.6. For any $u_0 \in V$ and T > 0, the attainable set at time T from u_0 of system (4.9.3) is dense in H.

Proof. Fix $\varepsilon > 0$, $u_0 \in V$, $x_1 \in H$ and T > 0. We prove that it is possible to drive the system from u_0 to the ball $\{y \in H \mid |y - x_1| \leq \varepsilon\}$. For that, first we set $M \in \mathbb{N}_0$ such that $|x_1 - P^{\kappa_M} x_1| < \frac{\varepsilon}{2}$, where P^{κ_M} is the projection map onto \mathbb{R}^{κ_M} and, consider the system [(4.10.1).M]. As we have seen in the first step of the proof of Proposition 4.10.2 (section 4.10.1), there is a control v_M taking values on \mathbb{R}^{κ_M} and driving the system from u_0 to some point u_T^M such that $P^{\kappa_M} u_T^M = P^{\kappa_M} x_1$. Moreover by item 1 of Corollary 4.10.3 we have

$$|u_T^M - P^{\kappa_M} u_T^M| \le K [T \exp(T)]^{\frac{1}{2}}.$$

Now (see Lemma 4.10.9) we imitate v_M by another control v_{M-1} taking values in $\mathbb{R}^{\kappa_{M-1}}$ and driving the system to a point $u_T^{M-1} \in V$ such that

$$|u_T^{M-1} - u_T^M| \le \frac{\varepsilon}{2M}.$$

Repeating this procedure of imitation, at each step we find a control v_{i-1} taking values in $\mathbb{R}^{\kappa_{i-1}}$ and driving the system to a point $u_T^{i-1} \in V$ such that

$$|u_T^{i-1} - u_T^i| \le \frac{\varepsilon}{2M}.$$

Let $T \leq T_1$ where T_1 equals the unique solution of $T_1 \exp(T_1) = \left(\frac{\varepsilon}{2KM}\right)^2$. The control v^1 takes its values on \mathbb{R}^{κ_1} and drives the system to a point $u_T^1 \in V$ satisfying

$$|u_T^1 - x_1| \le \sum_{i=2}^M |u_T^{i-1} - u_T^i| + |u_T^M - P^{\kappa_M} u_T^M| + |P^{\kappa_M} u_T^M - x_1|.$$

Since $P^{\kappa_M} u_T^M = P^{\kappa_M} x_1$ we have

$$|u_T^1 - x_1| \le \frac{\varepsilon}{2M}(M-1) + K[T\exp(T)]^{\frac{1}{2}} + \frac{\varepsilon}{2} \le \varepsilon.$$

Hence the Proposition is proved for $T \leq T_1$. If $T > T_1$ we apply an arbitrary control up to time $T - T_1$ arriving to some point $y_0 \in V$ and then drive the system in time T_1 from y_0 to the ball $\{y \in H \mid |y - x_1| \leq \varepsilon\}$.

Bibliography

- S. Agmon, Lectures on Elliptic Boundary Value Problems, van Nostrand 1965.
- [2] A. A. Agrachev, Y. L. Sachkov Control Theory from the Geometric Viewpoint, Encyclopaedia of Mathematical Sciences, 87, Springer,2004
- [3] A. A. Agrachev, A. V. Sarychev, Navier-Stokes Equations: Controllability by Means of Low Modes Forcing, To appear on the Journal of Mathematical Fluid Mechanics.
- [4] H. Brezis, Analyse Fonctionnelle, Théorie et Applications, Masson, 1993.
- [5] C. Foias, O. Manley, R. Rosa, R. Temam, Navier-Stokes Equations and Turbulence, Encyclopedia of Mathematics and its Applications, Cambridge university Press, 2001.
- [6] I. Fonseca, W. Gangbo, Degree Theory in Analysis and Applications; Oxford Lectures Series in Mathematics and its Applications, Oxford University Press, 1995.
- [7] R.V. Gamkrelidze, *Principles of Optimal Control Theory*, Plenum Press, 1978.
- [8] V. Jurdjevic, *Geometric Control Theory*, Cambridge Studies in Advanced Mathematics 51, Cambridge University Press, 1997.
- [9] P.L. Lions, E. Magenes, Non-Homogeneous Boundary Value Problems and Applications, vol.I, Die Grundlehren der Mathematischen Wissenschaften in Einzeldarstellungen, band 181, Springer-Verlag, 1972.
- [10] J. Peetre, Espaces d'Interpolation et théorème de Soboleff, Annales de l'Institut Fourier, 279-317, 1978.
- [11] M. Renardy, R.C. Rogers, An Introduction to Partial Differential Equations, Texts in Applied Mathematics, 13, Springer-Verlag, 1993.
- [12] R.S. Strichartz, A Guide to Distribution Theory and Fourier Transformations, World Scientific, 2003.
- [13] R. Temam, Infinite-Dimensional dynamical Systems in Mechanics and Physics, 2nd ed., Applied Mathematical Sciences, 68, Springer, 1997.

- [14] R. Temam, Navier-Stokes Equations and Nonlinear Functional Analysis, 2nd Ed., CBMS-NSF Regional Conference Series in Applied Mathematics, Society for Industrial and Applied Mathematics, 1995.
- [15] R. Temam, Navier-Stokes Equations: Theory and Numerical Analysis, AMS Chelsea Publishing, 2001.
- [16] E. Weinam, J.C. Mattingly Ergodicity for the Navier-Stokes Equation with Degenerate Random Forcing: Finite Dimensional Approximation, Comm. on Pure and Applied Math., Vol. 54, 1386-1402, 2001.