On N-differential graded algebras

Mauricio Angel[∗] and Rafael Díaz[†]

September 12, 2018

Abstract

We introduce the concept of N-differential graded algebras (N-dga), and study the moduli space of deformations of the differential of an N-dga. We prove that it is controlled by what we call the (M,N)-Maurer-Cartan equation.

Introduction

The goal of this paper is to take the first step towards finding a generalization of Homological Mirror Symmetry (HMS) [\[11\]](#page-15-0) to the context of N-homological algebra [\[5\]](#page-14-0). In [\[7\]](#page-15-1) Fukaya introduced HMS as the equivalence of the deformation functor of the differential of a differential graded algebra associated with the holomorphic structure, with the deformation functor of an A_{∞} -algebra associated with the symplectic structure of a Calabi-Yau variety. This idea motivated us to define deformation functors of the differential of an N-differential graded algebra. An N-dga is a graded associative algebra A, provided with an operator $d : A \to A$ of degree 1 such that $d(ab) = d(a)b + (-1)^{\bar{a}}ad(b)$ and $d^N = 0$. A nilpotent differential graded algebra (Nil-dga) will be an N-dga for some integer $N \geq 2$. Theorem [10](#page-3-0) endows the category of Nil-differential graded algebras with a symmetric monoidal structure. We remark that such a monoidal structure cannot be constructed in a natural way for a fixed N (except for $N = 2$), not even using the q-deformed Leibniz rule, see [\[13\]](#page-15-2).

In Section 2 we consider deformations of a 2-dga into an N-dga. By deforming 2-dgas one is able to construct a plethora of examples of N-dgas. Roughly speaking Theorem [16](#page-6-0) tell us that a derivation of a 2-dga $d_A + e$ is an N-differential iff

$$
(d_{End}(e) + e^{2})^{\frac{N-1}{2}}(d_{A} + e) = 0 \quad \text{for N odd,}
$$

$$
(d_{End}(e) + e^{2})^{\frac{N}{2}} = 0 \quad \text{for N even.}
$$

[∗]Work partially supported by IVIC.

[†]Work partially supported by IVIC.

In Section 3 we introduce a general formalism for discrete quantum mechanics. We introduce these models since they turn out, in a totally unexpected way, to be relevant in the problem of deforming an M-differential into an N-differential with $N \geq M$. Section 4 contains our main result, Theorem [19](#page-11-0) which provides an explicit identity called the (M, N) -Maurer-Cartan equation that controls deformations of an M-complex into an Ncomplex. The construction of the (M, N) -Maurer-Cartan equation is based on an explicit description of coefficients c_k such that

$$
(d_A + e)^N = \sum_{k=0}^{N-1} c_k d_A^k,
$$

where c_k depends on d_A and e . In Section 5 we define a functional $cs_{2,2N}$ whose critical points are naturally determined by the $(2, 2N)$ -Maurer-Cartan equation.

In conclusion in this paper we introduce the moduli space of deformations of the differential of an N-dga and prove that it is controlled by a generalized Maurer-Cartan equation. We point out that our methods and ideas can be applied in a wide variety of contexts. Examples of N-dga's coming from differential geometry are developed in [\[1\]](#page-14-1). A q-analogue, for q a primitive N-th root of unity, of our main result Theorem [19](#page-11-0) is provided in [\[2\]](#page-14-2). In [\[3\]](#page-14-3) we state an N-generalized Deligne's principle and use the constructions of this paper to study A_{∞} -algebras of depth N.

1 N-differential graded algebras and modules

Throughout this paper we shall work with the abelian category of k-modules over a commutative ring **k** with unit. We will denote by $A^{\bullet} \mathbb{Z}$ -graded **k**-modules $\bigoplus_{i\in\mathbb{Z}} A^i$. We let $\bar{a} \in \mathbb{Z}$ denote the degree of the element $a \in A^{\bar{a}}$. The following definition is taken from [\[9\]](#page-15-3).

Definition 1 Let $N \geq 1$ an integer. An N-complex is a pair (A^{\bullet}, d) , where A^{\bullet} is a $\mathbb{Z}\text{-}\text{graded object and } d: A^{\bullet} \to A^{\bullet} \text{ is a morphism of degree 1 such that } d^N = 0.$

Clearly an N-complex is a P-complex for all $P \geq N$. If **k** is a field, then an Ncomplex (A^{\bullet}, d) is referred as an N-differential graded vector space (N-dgvect). An *N*-complex (A^{\bullet}, d) such that $d^{N-1} \neq 0$ is said to be a **proper** *N*-complex. Let (A^{\bullet}, d_A) be an M-complex and (B^{\bullet}, d_B) be an N-complex, a **morphism** $f : (A^{\bullet}, d_A) \to (B^{\bullet}, d_B)$ is a morphism $f : A^{\bullet} \to B^{\bullet}$ of **k**-modules such that $d_B f = f d_A$.

Lemma 2 Let (A^{\bullet}, d_A) be a proper M-complex, (B^{\bullet}, d_B) be a proper N-complex and $f:(A^{\bullet},d_A)\rightarrow (B^{\bullet},d_B)$ be a morphism, then (1) If $Ker(f)=0$, then $M\leq N$; (2) If $Im(f) = B^{\bullet}$, then $M \ge N$ and (3) If $Ker(f) = 0$ and $Im(f) = B^{\bullet}$, then $M = N$.

Proof. 1. Assume that $N < M$ and let $a \in A^{\bullet}$ then $f(d_A^N(a)) = d_B^N(f(a)) = 0$. This implies that $d_A^N(a) \in Ker(f) = 0$, and therefore $d_A^N(a) = 0$ which is in contradiction with the fact that (A^{\bullet}, d_A) is a proper M-complex. The proof of 2. is analogous to 1., 3. follows from 1. and $2.\blacklozenge$

Example 3 Consider $V = \mathbb{C} \langle e_1, e_2, e_3 \rangle$ the complex vector space generated by e_1, e_2, e_3 . We endow V with a Z-graduation declaring $\bar{e_1} = 0$, $\bar{e_2} = 1$ and $\bar{e_3} = 2$. Define the linear map $d: V \to V$ on generators by

$$
d(e_1) = e_2
$$
, $d(e_2) = e_3$, and $d(e_3) = 0$.

 (V, d) is a proper 3-complex.

Definition 4 Let (A^{\bullet}, d) be an N-complex, we say that an element $a \in A^i$ is **p-closed** if $d^p(a) = 0$ and is **p-exact** if there exists an element $b \in A^{i-N+p}$ such that $d^{N-p}(b) = a$, for $1 \leq p \leq N$ fixed. The **cohomology groups** of are the **k**-modules

$$
_{p}H^{i}(A) = \frac{Ker\{d^{p} : A^{i} \to A^{i+p}\}}{Im\{d^{N-p} : A^{i-N+p} \to A^{i}\}},
$$

where $i \in \mathbb{Z}, p = 1, 2, ..., N-1$. We set $_k H^*(A) = 0$ for $k \ge N$.

Notice that a 2-complex A^{\bullet} is just a complex in the usual sense and in this case p is necessarily equal to 1 and $_1H^i(A)$ agrees with $H^i(A)$ for all $i \in \mathbb{Z}$.

- **Definition 5** (a) Let $N \geq 1$ an integer. An N-differential graded algebra or Ndga over **k**, is a triple (A^{\bullet}, m, d) where $m : A^k \otimes A^l \rightarrow A^{k+l}$ and $d : A^k \rightarrow A^{k+l}$ are k-modules homomorphisms satisfying
	- 1) The pair (A^{\bullet}, m) is a graded associative algebra.
	- 2) For all a, $b \in A^{\bullet}$, d satisfies the graded Leibniz rule $d(ab) = d(a)b + (-1)^{\bar{a}}ad(b)$.
	- 3) $d^N = 0$, i.e., (A^{\bullet}, d) is an N-complex.
	- (b) A nilpotent differential graded algebra (Nil-dga) is an N-dga for some integer $N \geq 2$.

A 1-dga is a graded associative algebra. A 2-dga is a differential graded algebra.

Lemma 6 Let (A^{\bullet}, m, d) be an N-dga, then if a is p-closed and b is q-closed then ab is $(p+q-1)$ -closed.

Proof. The Lemma follows from the identity

$$
d^n(ab) = \sum_{i=0}^n \left\{ \begin{array}{c} n \\ i \end{array} \right\} _{\bar{a}} d^i(a)d^{n-i}(b),
$$

where $\begin{cases} n \\ 0 \end{cases}$ $\overline{0}$ \mathcal{L} \bar{a} $=(-1)^{\bar{a}}$, and for $j \geq 1$, $\begin{cases} n+1 \\ i \end{cases}$ j \mathcal{L} \bar{a} = $\int n$ $j-1$ \mathcal{L} \bar{a} $+(-1)^{\bar{a}+j}\begin{cases} n \end{cases}$ j \mathcal{L} \bar{a} . When $n = p + q - 1$, since $d^{i}(a) = 0$ for $i \geq p$, we only consider the case $i < p$, then

 $n-i = p+q-1-i > q-1$ and $d^{n-i}(b) = 0$, because $d^j(b) = 0$ for $j \geq q$. Thus so either $d^{i}(a) = 0$ or $d^{n-i}(b) = 0$ for all *i*, and we have *ab* is $(p+q-1)$ -closed. ◆

Definition 7 Let (A^{\bullet}, m_A, d_A) be an M-dga and (B^{\bullet}, m_B, d_B) be an N-dga. A mor**phism** $f: A^{\bullet} \to B^{\bullet}$ is a linear map such that $fm_A = m_B(f \otimes Id) + m_B(Id \otimes f)$ and $d_Bf = f d_A.$

A morphism $f: A^{\bullet} \to B^{\bullet}$ such that $f(A^i) \subset B^{i+k}$ is said to be a morphism of degree k. A pair of morphisms $f, g : A^{\bullet} \to B^{\bullet}$ of N-dga are homotopic, if there exist $h: A^{\bullet} \to B^{\bullet}$ of degree $N-1$ such that

$$
f - g = \sum_{i=0}^{N-1} d_B^{N-1-i} h d_A^i.
$$

We remark that if two morphisms $f, g : A^{\bullet} \to B^{\bullet}$ of Nil-dga are homotopic then they induce the same maps in cohomology.

Let (A^{\bullet}, m_A, d_A) and (B^{\bullet}, m_B, d_B) be an M-dga and an N-dga, respectively. Define $d_{A\otimes B} = d_A \otimes Id + Id \otimes d_B$, the identity

$$
d_{A\otimes B}^n(a\otimes b) = \sum_{k=0}^n (-1)^{\bar{a}(n-k)} d_A^k(a) \otimes d_B^{n-k}(b) \text{ implies,}
$$

Proposition 8 The triple $(A^{\bullet} \otimes B^{\bullet}, m_{A \otimes B}, d_{A \otimes B})$ is an $(M+N-1)$ -dga, where $m_{A \otimes B} =$ $m_A \otimes m_B$.

Example 9 Let (V, d) be the 3-complex of in Example [3.](#page-2-0) On the space $V \otimes V^*$ consider the base given by $E_{ij} = e_i \otimes e_j^*, i, j = 1, 2, 3,$ and define

$$
D(E_{ij}) = E_{(i+1)j} + (-1)^{i+j} E_{i(j-1)},
$$

by Proposition [8](#page-3-1) and since $D^4(E_{13}) \neq 0$, then $(V \otimes V^*, D)$ is a proper 5-dga.

Theorem 10 The category Nil-dgvect is a symmetric monoidal category. Nil-dga is the category of monoids in Nil-dgvect. Nil-dga inherits a symmetric monoidal structure from Nil-dgvetc.

Let V^{\bullet} be an N-dga. By Proposition [8,](#page-3-1) $(V^{\bullet})^{\otimes 2}$ is a $(2N-1)$ -dga, $(V^{\bullet})^{\otimes 3}$ is a $(3N-2)$ dga and in general $(V^{\bullet})^{\otimes k}$ is a $[k(N-1)+1]$ -dga.

Definition 11 Let (A^{\bullet}, m_A, d_A) be an N-dga and M^{\bullet} a graded **k**-module. Let $K \geq 2$ an integer. A K-differential graded module $(K-dgm)$ over (A^{\bullet}, m_A, d_A) , is a triple (M^{\bullet}, m_M, d_M) with $m_M : A^k \otimes M^l \to M^{k+l}$ and $d_M : M^k \to M^{k+l}$, **k**-modules morphisms satisfying the following properties

- 1. For all $a, b \in A^{\bullet}$ and $m \in M^{\bullet}$, $m_M(a, m_M(b, m)) = m_M(m_A(a, b), m)$, If no confusion arises, we denote $m_M(a, m)$ by am.
- 2. For all $a \in A^{\bullet}$ and $m \in M^{\bullet}$, $d_M(am) = d_A(a)m + (-1)^{\bar{a}}ad_M(m)$.
- 3. The pair (M^{\bullet}, d_M) is a K-complex, $d_M^K = 0$.

Let (M^{\bullet}, m_M, d_M) be a K-dgm and (N^{\bullet}, m_N, d_N) be an L-dgm both over an Ndga (A^{\bullet}, m_A, d_A) . A morphism $f : M^{\bullet} \to N^{\bullet}$ of degree k is a linear map such that $f(m_M(a, b)) = (-1)^{\bar{a}\bar{f}} m_N(a, f(b))$ and $d_M(f(b)) = f(d_N(b))$, for all $a \in A^{\bullet}$ and $b \in M^{\bullet}$. Now let (M^{\bullet}, m_M, d_M) be a K-dgm over an M-dga (A^{\bullet}, m_A, d_A) and (N^{\bullet}, m_N, d_N) and L-dgm over an N-dga (B^{\bullet}, m_B, d_B) . The triple $(M \otimes N, m_{N \otimes M}, d_{M \otimes N})$ turns out to be a $(K+L-1)$ -dgm over $(A \otimes B, m_{A \otimes B}, d_{A \otimes B})$, where $m_{M \otimes N}$ and $d_{M \otimes N}$ are defined as before.

Definition 12 The space of endomorphisms of degree k of M^{\bullet} is $End^k(M) =$ $\prod_{i\in\mathbb{Z}} Hom(M^i, M^{i+k}),$ this is, $End^k(M)$ consist of maps $f : M^{\bullet} \to M^{\bullet}$ of degree k which are linear in regard to the action of A^{\bullet} but which does not satisfy necessarily the relation $d_Mf = (-1)^{\bar{f}} f d_M$.

There are operators $\circ_M : End(M) \otimes M^{\bullet} \to M^{\bullet}$ and $\circ_E : End(M) \otimes End(M) \to$ $End(M)$. Similarly to Proposition [8,](#page-3-1) Proposition [13](#page-4-0) below provides the natural algebraic structure on $End(M)$,

Proposition 13 Define $d_{End}(f) := d_M(f) - (-1)^{\bar{f}} f(d_M)$, for $f \in End(M)$. The triple $(End(M), \circ_E, d_{End})$ is a $(2N-1)$ -dga, and $(M^{\bullet}, \circ_M, d_M)$ is an N-dgm over $(End(M), \circ_E, d_{End}).$

Proof. Associativity of \circ_E follows from the associativity morphisms composition. The Leibniz rule for d_{End} is a consequence of the Leibniz rule for d_M . From the definition of d_{End} we obtain the identity

$$
d_{End}^n(f) = \sum_{k=0}^n (-1)^{\bar{f}(n-k)} d_M^k \circ f \circ d_M^{n-k}
$$

which can be proved by induction and holds for all $n \geq 1$. Let $n = 2N - 1$ if $k < N$ then $N-1 < n-k$ and thus $d_M^{n-k} = 0$. Similarly if $n-k < N$ then $d_M^k = 0$.

2 Deformation theory of 2-dgas into N-dgas

Let k a field and consider the category $Artin$ of finite dimensional local k -algebras. If $\mathcal{R} \in Ob(\mathbf{Artin})$ with maximal ideal \mathcal{R}_+ then $\mathbf{k} \cong \mathcal{R}/\mathcal{R}_+$ $(\mathcal{R} = \mathbf{k}[[t]]$ and $\mathcal{R}_+ = t\mathbf{k}[[t]]$ are examples to keep in mind). Since $k \approx \mathcal{R}/\mathcal{R}_+$ then $\mathcal{R} \approx k \oplus \mathcal{R}_+$ as vector spaces. We study deformation theory using the formalism which considers deformations as functors from Artin algebras to Sets for later convenience.

Definition 14 Let A^{\bullet} be an M-dga, an N-deformation of A^{\bullet} over $\mathcal R$ is an N-dga $A^{\bullet}_{\mathcal{R}}$ over \mathcal{R} , with $N \geq M$, such that $A^{\bullet}_{\mathcal{R}}/\mathcal{R}_+ A^{\bullet}_{\mathcal{R}}$ is isomorphic to A^{\bullet} as N-dga. Two N-deformations $A^{\bullet}_{\mathcal{R}}$ and $B^{\bullet}_{\mathcal{R}}$ are said to be **isomorphic** if there exists an isomorphism $\Phi: A^{\bullet}_\mathcal{R} \to B^{\bullet}_\mathcal{R}$ of N-dgas such that the induced isomorphism $\bar{\Phi}: A^{\bullet}_\mathcal{R}/\mathcal{R}_+ A^{\bullet}_\mathcal{R} \to B^{\bullet}_\mathcal{R}/\mathcal{R}_+ B^{\bullet}_\mathcal{R}$ satisfies $i_B\bar{\Phi} = i_A$, where i_A and i_B are the isomorphism $i_A : A^{\bullet}_{\mathcal{R}}/\mathcal{R}_+A^{\bullet}_{\mathcal{R}} \to A^{\bullet}$ and $i_B : B^{\bullet}_{\mathcal{R}} / \mathcal{R}_+ B^{\bullet}_{\mathcal{R}} \to A^{\bullet}.$

The core of Definition [14](#page-5-0) is to require that $d_{A_{\mathcal{R}}}$ reduces to d_A , and $m_{A_{\mathcal{R}}}$ reduces to m_A under the natural projection $\pi: A^{\bullet}_{\mathcal{R}} \to A^{\bullet}_{\mathcal{R}}/\mathcal{R}_+ A^{\bullet}_{\mathcal{R}} \cong A^{\bullet}$. Assume that $A^{\bullet}_{\mathcal{R}} = A^{\bullet} \otimes \mathcal{R}$ as graded algebras. We have the following decomposition

$$
A^{\bullet}_{\mathcal{R}} = A^{\bullet} \otimes \mathcal{R} = A^{\bullet} \otimes (\mathbf{k} \oplus \mathcal{R}_{+}) = (A^{\bullet} \otimes \mathbf{k}) \oplus (A^{\bullet} \otimes \mathcal{R}_{+}) = A^{\bullet} \oplus (A^{\bullet} \otimes \mathcal{R}_{+}).
$$

Thus, since $d_{A_{\mathcal{R}}}$ reduces to d_A under the projection π , we must have

$$
d_{A_{\mathcal{R}}}=d_{A}+e
$$

where $e \in Der(A^{\bullet} \otimes \mathcal{R}_{+})$ has degree 1. Moreover, the fact that $d_{A_{\mathcal{R}}}^{N}=0$ implies that e is required to satisfy an identity which we call the (M, N) -Maurer-Cartan equation. Next proposition is well known and considers the classical case, that is, the (2, 2)-Maurer-Cartan equation.

Proposition 15 Let A^{\bullet} be a 2-dga and $A^{\bullet}_{\mathcal{R}} = A^{\bullet} \otimes \mathcal{R}$ be a 2-deformation over \mathcal{R} , $d_{A_{\mathcal{R}}} = d_A + e$ where $e \in Der(A^{\bullet} \otimes \mathcal{R}_+)$, then e satisfies the $(2,2)$ -Maurer-Cartan equation given by

PROOF. We have
$$
d_{A_{\mathcal{R}}}(a) = (d_{A} + e)(d_{A} + e)(a)
$$

$$
= d_{A}^{2}(a) + d_{A}(e(a)) + e(d_{A}(a)) + e^{2}(a)
$$

$$
= d_{End}(e)(a) + e^{2}(a), \text{ for all } a \in A^{\bullet}.\blacklozenge
$$

Suppose that $N = 2k + n$, $n \in \{0, 1\}$ and $k \in \mathbb{N}$, then

$$
d_{A_{\mathcal{R}}}^{N} = d_{A_{\mathcal{R}}}^{2k+n} = (d_{A_{\mathcal{R}}}^{2})^{k} d_{A_{\mathcal{R}}}^{n} = (d_{End}(e) + e^{2})^{k} d_{A_{\mathcal{R}}}^{n}, \text{ thus}
$$

Theorem 16 Let A^{\bullet} be a 2-dga. $A^{\bullet}_{\mathcal{R}} = A^{\bullet} \otimes \mathcal{R}$ is an N-deformation over \mathcal{R} with $d_{A_{\mathcal{R}}} = d_A + e$ where $e \in Der(A^{\bullet} \otimes \mathcal{R}_+)$ of degree 1, iff e satisfies

$$
(d_{End}(e) + e^{2})^{\frac{N-1}{2}}(d_{A} + e) = 0 \quad \text{for } N \text{ odd},
$$

$$
(d_{End}(e) + e^{2})^{\frac{N}{2}} = 0 \quad \text{for } N \text{ even}.
$$

Theorem [16](#page-6-0) can be easily extended to study deformations of the differential of a 2-dgm M^{\bullet} over a 2-dga A^{\bullet} as follows

Theorem 17 Let M^{\bullet} be a 2-dgm over a 2-dga A^{\bullet} . Then $M^{\bullet}_{\mathcal{R}} = M^{\bullet} \otimes \mathcal{R}$ is an Ndeformation over R with $d_{A_{\mathcal{R}}} = d_A + e$ where $e \in End(M^{\bullet} \otimes R_+)$ has degree 1, iff e satisfies

$$
(d_{End}(e) + e^{2})^{\frac{N-1}{2}}(d_{M} + e) = 0 \quad \text{for } N \text{ odd},
$$

$$
(d_{End}(e) + e^{2})^{\frac{N}{2}} = 0 \quad \text{for } N \text{ even}.
$$

Let M be a 3-dimensional smooth manifold. The space $(\Omega^{\bullet}(M), d)$ of differential forms on M is a differential graded algebra with d the de Rham differential. Let $\pi : E \to M$ be a vector bundle, the space $(\Omega^{\bullet}(M, E), d_E)$ of E-valued forms is a differential graded module over $(\Omega^{\bullet}(M), d)$, where d_E is the differential induced by d. Let $A \in \Omega^1(M)$ and consider the endomorphism e_A induced by A, defined by $e_A(\omega) = A \wedge \omega$ for all $\omega \in \Omega^{\bullet}(M, E)$. The pair $(\Omega^{\bullet}(M, E), d_E + e_A)$ is a 4-dgm for any A. Moreover, according to Theorem [17](#page-6-1) $(\Omega^{\bullet}(M, E), d + e_A)$ is a 3-dgm if and only if for all ω

$$
d_{End}(e_A)(d+e_A)\omega = 0.
$$

Since $d_{End}(e_A)(d + e_A)$ is an operator of degree 3, the identity $d_{End}(e_A)(d + e_A)\omega = 0$ holds for any k-form $\omega, k \ge 1$. Thus $(\Omega^{\bullet}(M, E), d + e_A)$ is a 3-dgm if and only if for any 0-form ω

$$
d_{End}(e_A)(d + e_A)\omega = d(A) \wedge (d_E(\omega) + A \wedge \omega) = 0.
$$

Similarly, it is easy to deduce from Theorem [17](#page-6-1) that if M is an n-dimensional smooth manifold and $n < m$, then $(\Omega^{\bullet}(M, E), d + e_A)$ is a m-complex. Let now M be a 2ndimensional smooth manifold. Using local coordinates the 2-form $d_{End}(e_A)$ can be written as $F_{ij}dx^i \wedge dx^j$ where $F_{ij} = \partial_i A_j - \partial_j A_i$. Furthermore,

$$
(F_{ij}dx^i \wedge dx^j)^n = \bigg(\sum_{\alpha \in P(2n)} \prod_{i=1}^n sign(\alpha)F_{a_i,b_i}\bigg)dx^1 \wedge \ldots \wedge dx^{2n},
$$

where $P(2n)$ is the set of ordered pairings of $[2n] = \{1, ..., 2n\}$. Recall that a ordered pairing $\alpha \in P(2n)$ is a sequence $\{(a_i, b_i)\}_{i=1}^n$ such that $[2n] = \bigsqcup_{i=1}^n \{a_i, b_i\}$ and $a_i < b_i$. By Theorem [17,](#page-6-1) $(\Omega^{\bullet}(M, E), d + e_A)$ is a 2n-complex if and only if the 2-form $F_{ij}dx^{i} \wedge dx^{j}$ satisfies

$$
\sum_{\alpha \in P(2n)} sign(\alpha) \prod_{i=1}^{n} F_{a_i, b_i} = 0.
$$

Let M be a complex manifold and consider the differential graded algebra $(\Omega(M), \wedge, \bar{\partial}),$ where $\bar{\partial}$ is the Dolbault differential. Let $\pi : E \to M$ be a complex vector bundle, we consider $\Omega(M, E)$ the forms with values in E. Recall that a holomorphic structure on E is given by a left differential graded module structure $(\Omega(M, E), \wedge_E, \bar{\partial_E})$ over the 2-dga $(\Omega(M), \wedge, \bar{\partial})$. Suppose that on $(\Omega(M, E), \wedge_E, \bar{\partial}_E)$ there is a left N-differential graded module structure over the 2-dga $(\Omega(M), \wedge, \overline{\partial})$, then in this case we say that E carries an N-holomorphic structure.

3 Discrete quantum theory

Generally speaking the following data constitute the basic set up for a (non-relativistic) quantum mechanical system: A finite dimensional Riemannian manifold M which is thought as the configuration space of the quantum system; A Lagrangian function $L: TM \to \mathbb{R}$ which assigns weights to points in phase space.

Associated to this data is the Hilbert space $\mathcal H$ of quantum states which is usually taken to be $L^2(M)$, the space of square integrable functions on M. The dynamics of the quantum system is determined by operators $U_t : \mathcal{H} \to \mathcal{H}$, where $t \in \mathbb{R}$ represents time. The kernel ω_t of U_t is such that

$$
(U_t f)(y) = \int_M \omega_t(y, x) f(x) dx.
$$

The key insight of Feynman is that $\omega_t(y, x)$ admits an integral representation

$$
\omega_t(y,x) = \int e^{i \int_0^t L(\gamma,\dot{\gamma})dt} D(\gamma).
$$

The integral above runs over all paths $\gamma : [0, t] \to M$ such that $\gamma(0) = x$ and $\gamma(t) = y$. Making rigorous sense of this integral is the main obstacle in turning quantum mechanics a fully rigorous mathematical theory. Recall that a directed graph Γ is given by: *i*) A set V_{Γ} called the set of vertices, ii) A set E_{Γ} called the set of edges and iii) A map $(s, t) : E_{\Gamma} \to V_{\Gamma} \times V_{\Gamma}$. Following the pattern above, one may define a **discrete quantum** mechanical system as being given by the following data

- 1. A directed graph Γ (finite or infinite) which plays the role of configuration space.
- 2. A map $L: E_{\Gamma} \to \mathbb{R}$ called the Lagrangian map of the system.

The associated Hilbert space is $\mathcal{H} = \mathbb{C}^{V_{\Gamma}}$. The operators $U_n : \mathcal{H} \to \mathcal{H}$, where $n \in \mathbb{Z}$ represents discretized time are given by

$$
(U_n f)(y) = \sum_{x \in V_{\Gamma}} \omega_n(y, x) f(x),
$$

where the discretized kernel $\omega_n(y, x)$ admits the following representation

$$
\omega_n(y, x) = \sum_{\gamma \in P_n(\Gamma, x, y)} \prod_{e \in \gamma} e^{iL(e)}.
$$

Here $P_n(\Gamma, x, y)$ denotes the set of length n paths in Γ from x to y, i.e., sequences (e_1, \dots, e_n) of edges in Γ such that $s(e_1) = x$, $t(e_i) = s(e_{i+1}), i = 1, ..., n-1$ and $t(e_n) = y.$

In Section 4 we show that the generalized Maurer-Cartan equation controlling deformations of N-dgas is determined by the kernel of a discrete quantum mechanical system L which we proceed to introduce. Let us first explain our notation and conventions which generalize those introduced in [\[4\]](#page-14-4).

For $s = (s_1, ..., s_n) \in \mathbb{N}^n$ we set $l(s) = n$, the length of the vector s, and $|s| = \sum_i s_i$. For $1 \leq i \leq n$, $s_{>i}$ denotes the vector given by $s_{>i} = (s_{i+1},...,s_n)$, for $1 \leq i \leq n$, $s_{< i}$ stands for $s_{\leq i} = (s_1, ..., s_{i-1})$, we also set $s_{\geq n} = s_{\leq 1} = \emptyset$. $\mathbb{N}^{(\infty)}$ denotes the set $\bigsqcup_{n=0}^{(\infty)} \mathbb{N}^n$, where by convention $\mathbb{N}^{(0)} = \{\emptyset\}.$

We define maps δ_i , $\eta_i : \mathbb{N}^n \to \{0, 1\}$, for $1 \leq i \leq n$, as follows

$$
\delta_i(s) = \begin{cases} 1 & \text{if } s_i = 0, \\ 0 & \text{otherwise.} \end{cases} \qquad \qquad \eta_i(s) = \begin{cases} 1 & \text{if } s_i \ge 1, \\ 0 & \text{otherwise.} \end{cases}
$$

For an M-dga A^{\bullet} and $e \in End(A^{\bullet})$ and $s \in \mathbb{N}^n$ we define $e^{(s)} = e^{(s_1)} \cdots e^{(s_n)}$, where $e^{(l)} = d_{End}^l(e)$ if $l \geq 1$, $e^{(0)} = e$ and $e^{\emptyset} = 1$. In the case that $e_a \in End(A^{\bullet})$ is given by

$$
e_a(\phi) = a\phi
$$
, for $a \in A^1$ fixed and all $\phi \in A^{\bullet}$

,

then $e_a^{(l)} = d_{End}^l(e_a)$ reduces to $e_a^{(l)} = e_{d^l(a)}$, thus $e_a^{(s)} = e_a^{(s_1)} \cdots e_a^{(s_n)} = e_{d^{s_1}(a)} \cdots e_{d^{s_n}(a)}.$

Where [k] denotes the set $\{1, 2, ..., k\}$. For $N \in \mathbb{N}$ we define $E_N = \{s \in \mathbb{N}^{(\infty)}$: $|s| + l(s) \leq i$ and for $s \in E_N$ we define $N(s) \in \mathbb{Z}$ by $N(s) = N - |s| - l(s)$.

We introduce the discrete quantum mechanical system L by

- 1. $V_L = \mathbb{N}^{(\infty)}$.
- 2. There is a unique directed edge in L from vertex s to t if and only if $t \in \{(0, s), s, (s+$ (e_i) } where $e_i = (0, ..., 1)$ \sum_{i-th} $(a_1, \ldots, b_n) \in \mathbb{N}^{l(s)}$, in this case we set source $(e) = s$ and $target(e) = t$.
- 3. Edges in L are weighted according to the following table

The set $P_N(\emptyset, s)$ consists of all paths $\gamma = (e_1, ..., e_N)$, such that source $(e_1) = \emptyset$, $target(e_N) = s$ and $source(e_{l+1}) = target(e_l)$. For $\gamma \in P_N(\emptyset, s)$ we define the weight $\omega(\gamma)$ of γ as

$$
\omega(\gamma) = \prod_{i=1}^N \omega(e_i).
$$

4 The (M,N)-Maurer-Cartan equation

Lemma 18 Let A^{\bullet} be an M-dga and $\mathcal{R} \in Ob(Artin)$. We define $d_{A_{\mathcal{R}}} = d_A + e$ where $e \in Der(A^{\bullet} \otimes \mathcal{R}_{+})$ has degree 1, then

$$
(d_{A_{\mathcal{R}}})^N = \sum_{s \in E_N} c(s, N) e^{(s)} d_A^{N(s)},
$$

where the coefficient $c(s, N + 1)$ is equal to

$$
\delta_1(s)c(s_{>1}, N) + (-1)^{|s|+l(s)}c(s, N) + \sum_{i=1}^{l(s)} \eta_i(s)(-1)^{|s_{ (1)
$$

and $c(\emptyset, 1) = c(0, 1) = 1$.

Proof. We use an induction on N. For $N = 1$, since $E_1 = \{s = \emptyset, s = 0\}$

$$
d_{A_{\mathcal{R}}} = \sum_{s \in E_1} c(s, 1) e^{(s)} d_A^{1(s)} = c(\emptyset, 1) e^{(\emptyset)} d_A^{1 - |\emptyset| - l(\emptyset)} + c(0, 1) e^{(0)} d_A^{1 - |\emptyset| - l(0)}
$$

= $c(\emptyset, 1) d_A + c(0, 1) e$.

Suppose our formula holds for N and let us check it for $N+1$

$$
(d_{A_{\mathcal{R}}})^{N+1} = (d_{A} + e)(d_{A_{\mathcal{R}}})^{N}
$$

\n
$$
= (d_{A} + e) \left(\sum_{s \in E_{N}} c(s, N) e^{(s)} d_{A}^{N(s)} \right)
$$

\n
$$
= d_{A} \left(\sum_{s \in E_{N}} c(s, N) e^{(s)} d_{A}^{N(s)} \right) + e \left(\sum_{s \in E_{N}} c(s, N) e^{(s)} d_{A}^{N(s)} \right)
$$

\n
$$
= \sum_{s \in E_{N}} c(s, N) d_{A} (e^{(s)} d_{A}^{N(s)}) + \sum_{s \in E_{N}} c(s, N) e^{(s)} d_{A}^{N(s)}.
$$
 (2)

Consider the second term of the right hand side of [\(2\)](#page-10-0)

$$
\sum_{s \in E_N} c(s, N) e^{(s)} d_A^{N(s)} = \sum_{s \in E_N} c(s, N) e^{(0)} e^{(s)} d_A^{N(s)}
$$

=
$$
\sum_{\substack{t \in E_{N+1} \\ t_1 = 0}} c(t_{>1}, N) e^{(t)} d_A^{N - |t_{>1}| - l(t_{>1})}
$$
(3)

$$
= \sum_{s \in E_{N+1}} \delta_1(s) c(s_{>1}, N) e^{(s)} d_A^{N(s)+1}.
$$
 (4)

In [\(3\)](#page-10-1) we put $t = (0, s)$ thus $|t| = |s|$ and $l(t) = l(s) + 1$ and [\(4\)](#page-10-1) is obtained by rewriting and changing t by s .

Now consider the first term of the right hand side of [\(2\)](#page-10-0)

$$
\sum_{s \in E_N} c(s, N) d_A(e^{(s)} d_A^{N(s)}) = \sum_{\substack{s \in E_N \\ 1 \le i \le l(s)}} (-1)^{|s_{
$$
+ \sum_{s \in E_N} (-1)^{|s| + l(s)} c(s, N) e^{(s)} d_A^{N(s) + 1}
$$
(5)
$$
= \sum_{t \in E_{N+1}} \sum_{\substack{i=1 \\ i \le j}}^{l(t)} (-1)^{|t < i| + i - 1} c(t - e_i, N) e^{(t)} d_A^{N - |t - e_i| - l(t)}
$$
$$

$$
+\sum_{s\in E_N}(-1)^{|s|+l(s)}c(s,N)e^{(s)}d_A^{N(s)+1}
$$
\n(6)

$$
= \sum_{s \in E_{N+1}} \sum_{i=1}^{l(s)} \eta_i(s)(-1)^{|s \le i|+i-1} c(s - e_i, N) e^{(s)} d_A^{N(s)+1} + \sum_{s \in E_N} (-1)^{|s|+l(s)} c(s, N) e^{(s)} d_A^{N(s)+1}.
$$
 (7)

Putting $t = s + e_i$ in the first term of [\(5\)](#page-11-1) we obtain [\(6\)](#page-11-1) and rewriting and changing t by s we obtain (7) . Finally collecting similar terms in (4) and (7) , and using the recurrence formula we get

$$
(d_A + e)^{N+1} = \sum_{s \in E_{N+1}} c(s, N+1) e^{(s)} d_A^{N(s)+1},
$$

thus the proof is completed. \blacklozenge

The following result generalizes Theorem [16.](#page-6-0) It provides an explicit formulae for the coefficients of the generalized Maurer-Cartan equation introduced below.

Theorem 19 We have,

$$
(d_{A_{\mathcal{R}}})^N = \sum_{k=0}^{N-1} c_k d_A^k,
$$

where

$$
c_k = \sum_{\substack{s \in E_N \\ N(s) = k \\ s_i < M}} c(s, N) e^{(s)} \quad \text{and} \quad c(s, N) = \sum_{\gamma \in P_N(\emptyset, s)} \omega(\gamma).
$$

Proof. One checks that the coefficients $c(s, N) = \sum_{\gamma \in P_N(\emptyset, s)} \omega(\gamma)$ satisfy the recurrence formula of Lemma [18.](#page-9-0) For this one checks that $P_{N+1}(\emptyset, s)$ is naturally partitioned in three blocks. The first block contains paths that are the composition of a path $\gamma : \emptyset \to s$ in $P_N(\emptyset, s_{>1})$ with an edge $s_{>1} \to (0, s_{>1})$ and corresponds with the first term in [\(1\)](#page-9-1). The second block consists of paths that are the composition of a path $\gamma : \emptyset \to s$ in $P_N(\emptyset, s)$

with an edge $s \to s$ and corresponds with the second term in [\(1\)](#page-9-1), finally the last block consists of paths that are the composition of a path $\gamma : \emptyset \to s - e_i$ in $P_N(\emptyset, s - e_i)$ with an edge $s - e_i \rightarrow s$ and corresponds with the last term of [\(1\)](#page-9-1).

Let A^{\bullet} be an M-dga and $A^{\bullet}_{\mathcal{R}}$ an N-deformation over \mathcal{R} with $A^{\bullet}_{\mathcal{R}} = A^{\bullet} \otimes \mathcal{R}$. For $a \in A^1 \otimes \mathcal{R}_+$ we define $e_a: A^{\bullet}_{\mathcal{R}} \to A^{\bullet}_{\mathcal{R}}$ by

$$
e_a(b) = ab - (-1)^{\bar{b}}ba.
$$

We are assuming that the product is not graded commutative. It is easy to see that e_b is a derivation of degree 1 on $A^{\bullet} \otimes \mathcal{R}_+$. Then $d_{A_{\mathcal{R}}} = d_A + e_a$ is an N-deformation of d_A iff e_a satisfies the equation

$$
\sum_{\substack{s \in E_N \\ s_i < M}} c(s, N) e_a^{(s)} d_A^{N - |s| - l(s)} = 0. \tag{8}
$$

Equation [\(8\)](#page-12-0) will be called the (M, N) -Maurer-Cartan equation. We closed this section by formally introducing the (M, N) -Maurer-Cartan functor $MC_M^N(A)$ which controls deformations of the differential d_A of an N-dga A^{\bullet} .

Definition 20 For $N \geq M$, $a \in A^1 \otimes \mathcal{R}_+$ is said to be an (M, N) -Maurer-Cartan element of $A^{\bullet} \otimes \mathcal{R}$ if e_a satisfies the (M, N) -Maurer-Cartan equation (8). We say that a is homotopic to a', if e_a is homotopic to $e_{a'}$ as morphisms of N-dgas.

Definition 21 We define the (M, N) -Maurer-Cartan functor $MC_M^N(A)$: Artin \rightarrow Set for each M-dga A^{\bullet} over **k**. Functor $MC_M^N(A)$ is given by

- 1. Let R be an object of Artin. $MC_M^N(A)(R)$ is the set of homotopy classes of all (M, N) -Maurer-Cartan elements of $A^{\bullet} \otimes \mathcal{R}$.
- 2. If $\varphi : \mathcal{R} \to \mathcal{R}'$ be a morphism of the category **Artin** and a is an (M, N) -Maurer-Cartan element of $A^{\bullet} \otimes \mathcal{R}$, then $(1 \otimes \varphi)(a)$ is an (M, N) -Maurer-Cartan elements of $A^{\bullet} \otimes \mathcal{R}'$. Thus we obtain a map $\varphi_*: MC_M^N(A)(\mathcal{R}) \to MC_M^N(A)(\mathcal{R}')$.

Deformation theory of K-dgms over an M-dga can be defined similarly.

5 Chern-Simons actions

Let (A^{\bullet}, m_A, d_A) be a 2-dga over **k** and let (M^{\bullet}, m_M, d_M) be a 2-dgm over (A^{\bullet}, m_A, d_A) , consider its $2K$ -Maurer-Cartan equation, that is the equation that arises when we deform the 2-dgm (M^{\bullet}, m_M, d_M) into a 2K-dgm, $MC_{2K}(a) = (d_{End}(a) + a^2)^K = 0$, where $a \in$ $End(M^{\bullet})$ has degree 1. Let us assume that there exists a linear functional \int : $End(M^{\bullet}) \rightarrow$ **k** of degree $2K+1$, (i.e., $\int b = 0$ if $\bar{b} \neq 2K+1$) satisfying the following conditions:

- 1. \int is non degenerate, that is, $\int ab = 0$ for all a, then $b = 0$.
- 2. $\int d(a) = 0$ for all a, where $d = d_{End(M^{\bullet})}$.
- 3. \int is cyclic, this is $\int a_1 a_2 \cdots a_n = (-1)^{\bar{a_1}(\bar{a_2} \cdots \bar{a_n})} \int a_2 \cdots a_n a_1$.

We define the **Chern-Simons** functional $cs_{2,2K} : End(M^{\bullet}) \to \mathbf{k}$ by

$$
cs_{2,2K}(a) = 2K \int \pi(\#^{-1}(a(d_{End}(a) + a^2)^K)),
$$

where

- 1. $\mathbf{k} \leq a, d(a)$ denotes the free **k**-algebra generated by symbols a and $d(a)$.
- 2. $\# : \mathbf{k} < a, d(a) > \longrightarrow \mathbf{k} < a, d(a) >$ is the linear map defined by

$$
#(a^{i_1}d(a)^{j_1}...a^{i_k}d(a)^{j_k}) = (i_1 + ... + i_k + j_1 + ... + j_k)a^{i_1}d(a)^{j_1}...a^{i_k}d(a)^{j_k}.
$$

3. $\pi : \mathbf{k} < a, d(a) > \longrightarrow End(M^{\bullet})$ is the canonical projection.

For $K = 1$ we have that $cs_{2,2}(a)$ is equal to

$$
2\int \pi(\#^{-1}(a(d(a) + a^2))) = 2\int \pi(\#^{-1}(ad(a) + a^3))) = \int ad(a) + \frac{2}{3}a^3,
$$

which is the Chern-Simons functional. In general we have the following result

Theorem 22 Let $K \geq 1$ be an integer. The Chern-Simons functional $cs_{2,2K}$ is a Lagrangian for the 2K-Maurer-Cartan equation, i.e., $a \in End^1(M^{\bullet})$ is a critical point of $cs_{2,2K}$ if and only if $(d(a) + a^2)^K = 0$.

Proof. We check that $\frac{\partial}{\partial \varepsilon} c s_{2,2K+2}(a+b\varepsilon)|_{\varepsilon=0} = (2K+2) \int bMC_{2K+2}(a)$.

$$
\frac{\partial}{\partial \varepsilon} c s_{2,2K+2}(a+b\varepsilon) \Big|_{\varepsilon=0}
$$
\n
$$
= \frac{\partial}{\partial \varepsilon} (2K+2)\pi \int (\#^{-1}((a+b\varepsilon)MC_{2K+2}(a+b\varepsilon))|_{\varepsilon=0}
$$
\n
$$
= (2K+2) \int \pi (\#^{-1}(\frac{\partial}{\partial \varepsilon}(a+b\varepsilon)MC_{2K}(a+b\varepsilon)MC_{2}(a+b\varepsilon)))|_{\varepsilon=0}
$$
\n
$$
= (2K+2) \int \pi (\#^{-1}(\frac{\partial}{\partial \varepsilon}(a+b\varepsilon)MC_{2K}(a+b\varepsilon))|_{\varepsilon=0} MC_{2}(a)
$$
\n
$$
+ (2K+2) \int \pi (\#^{-1}(aMC_{2K}(a)\frac{\partial}{\partial \varepsilon}MC_{2}(a+b\varepsilon)))|_{\varepsilon=0}.
$$
\n(9)

By degree reasons the second term of [\(9\)](#page-13-0) vanishes, the inductive hypothesis yields

$$
\frac{\partial}{\partial \varepsilon} c s_{2,2K+2}(a+b\varepsilon)|_{\varepsilon=0} = (2K+2) \int bMC_{2K}(a)MC_2(a)
$$

$$
= (2K+2) \int bMC_{2K+2}(a).\blacklozenge
$$

For $K = 2$, 3 the Chern-Simons functional $cs_{2,2K}(a)$ is given by

$$
cs_{2,4}(a) = \int \frac{4}{3} a(d(a))^2 + 2a^3 d(a) + \frac{4}{5} a^5.
$$

$$
cs_{2,6}(a) = \int \frac{3}{2} a(d(a))^3 + \frac{12}{5} a^3 (d(a))^2 + \frac{6}{5} a d(a) a^2 d(a) + 3a^5 d(a) + \frac{6}{7} a^7.
$$

Acknowledgement

We thank Nicolás Andruskiewitsch, Edmundo Castillo, Eddy Pariguan, Sylvie Paycha and Jim Stasheff for helpful suggestions. Thanks also to an anonymous referee for precise corrections.

References

- [1] M. Angel, R. Díaz, N-flat connections.-To appear in Proceedings of the Conference on Geometric and Topological Methods for Quantum Field Theory, Edited by: Sylvie Paycha, Hernan Ocampo, and Bernardo Uribe (Villa de Leyva 2005), Contemp. Math. book series, A.M.S. Preprint [math.DG/0511242.](http://arxiv.org/abs/math/0511242)
- [2] M. Angel, R. Díaz, On the q-analogue of the Maurer-Cartan equation.-Advan. Studies in Contemporary Mathematics Vol 12 no. 2(2006), 315-322. Preprint [math.QA/0601698.](http://arxiv.org/abs/math/0601698)
- [3] M. Angel, R. Díaz, A_{∞}^{N} -algebras.-In preparation.
- [4] R. Díaz, E. Pariguan, *Symmetric Quantum Weyl algebras*.-Annales Mathematiques Blaise Pascal 11, 187-203, 2004.
- [5] M. Dubois-Violette, Lectures on differentials, generalized differentials and some examples related to theoretical physics.-Quantum symmetries in theoretical physics and mathematics (Bariloche 2000), pages 59-94, Contemp. Math. 294, A.M.S. 2002.
- [6] P. Freyd, Abelian Categories.-Harper International Edition, 1964.
- [7] K. Fukaya, Deformation theory, homological algebra, and mirror symmetry.-Preprint January 2002.
- [8] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*.-Wiley Interscience, 1978. Pure and Applied Mathematics.
- [9] M.M. Kapranov, On the q-analog of homological algebra.-Preprint [q-alg/9611005.](http://arxiv.org/abs/q-alg/9611005)
- [10] C. Kassel et M. Wambst, Algèbre homologique des N-complexes et homologie de Hochschild aux racines de l'unité.-Publ. Res. Inst. Math. Sci. Kyoto University 34 (1998), n 2, 91-114.
- [11] M. Kontsevich, Homological Algebra of mirror symmetry.-Proceedings of the International Congress of Mathematicians (Zürich, 1994), Birkhäser, 1995, pp. 120-139.
- [12] M. Kontsevich and Y. Soibelman, Homological mirror symmetry and torus fibrations.-Symplectic geometry and mirror symmetry, World Scientific, 2001, pp. 203-263.
- [13] A. Sitarz, On the tensor product construction for q-differential algebras.-Lett. Math. Phys. 44 1998.

Mauricio Angel. Universidad Central de Venezuela (UCV). mangel@euler.ciens.ucv.ve Rafael Díaz. Universidad Central de Venezuela (UCV). rdiaz@euler.ciens.ucv.ve