

q -Cartan matrices and combinatorial invariants of derived categories for skewed-gentle algebras

Christine Bessenrodt¹

Institut für Mathematik, Universität Hannover,
Welfengarten 1, D-30167 Hannover, Germany

Thorsten Holm²

Department of Pure Mathematics, University of Leeds,
Leeds LS2 9JT, England

April 26, 2005

Abstract

Cartan matrices are of fundamental importance in representation theory. For algebras defined by quivers (i.e. directed graphs) with relations the computation of the entries of the Cartan matrix amounts to counting nonzero paths in the quivers, leading naturally to a combinatorial setting. In this paper we study a refined version, so-called q -Cartan matrices, where each nonzero path is weighted by a power of an indeterminate q according to its length. Specializing $q = 1$ gives the classical Cartan matrix. Our main motivation are derived module categories and their invariants: the invariant factors, and hence the determinant, of the Cartan matrix are preserved by derived equivalences.

The paper deals with the important class of (skewed-) gentle algebras which occur naturally in representation theory, especially in the context of derived categories. These algebras are defined in purely combinatorial terms. We determine normal forms for the Cartan matrices of (skewed-) gentle algebras. In particular, we give explicit combinatorial formulae for the invariant factors and thus also for the determinant of the Cartan matrices of skewed-gentle algebras. This generalizes the results of [11].

As an application of our main results we show how one can use our formulae for the notoriously difficult problem of distinguishing derived equivalence classes.

¹e-mail: bessen@math.uni-hannover.de

²e-mail: tholm@maths.leeds.ac.uk

1 Introduction

This paper deals with combinatorial aspects in the representation theory of algebras. More precisely, for classes of algebras which are defined purely combinatorially by directed graphs we will characterize important representation-theoretic invariants in a nice combinatorial way. In particular, this will give new explicit invariants for the derived module categories of the algebras involved.

Derived categories are nowadays one of the most important objects in representation theory and in geometry, with numerous applications, e.g. in mathematical physics.

In representation theory of algebras, a notoriously difficult problem is to find invariants of the algebras which are preserved under derived equivalences. Only few such invariants are known so far. This makes it usually very difficult to distinguish algebras up to derived equivalence.

The invariants we consider in this paper are the invariant factors of the Cartan matrix. Cartan matrices play a fundamental rôle in all areas of representation theory, containing complete information on the multiplicities of simple modules as composition factors of projective indecomposable modules. The entries of the Cartan matrix are not known in general and some famous open problems are related to their computation.

The starting point for this article is that the unimodular equivalence class of the Cartan matrix of a finite dimensional algebra is invariant under derived equivalence. This seems to be folklore knowledge, however, there seems to be no precise reference in the literature, so we also provide a proof for the convenience of the reader.

In this way, being able to determine normal forms of Cartan matrices yields invariants of the derived category.

The class of algebras we study are the gentle algebras, and the related skewed-gentle algebras. Gentle algebras form a very important class of algebras which occur naturally in many places in the representation theory of finite dimensional algebras, especially in the context of derived categories. For instance, the algebras which are derived equivalent to hereditary algebras of type \mathbb{A} are precisely the gentle algebras whose underlying undirected graph is a tree [2]. The algebras which are derived equivalent to hereditary algebras of type $\tilde{\mathbb{A}}$ are certain gentle algebras whose underlying graph has exactly one cycle [3]. Remarkably, the class of gentle algebras is closed under derived equivalence [16].

A fundamental, classical distinction in the representation theory of algebras is given by the representation type, which can be either finite, tame or wild.

In the modern context of derived categories, also derived representation types have been defined. Again, gentle algebras occur in this context. D. Vossieck [18] showed that an algebra A has a discrete derived category if and only if either A is derived equivalent to a hereditary algebra of type \mathbb{A} , \mathbb{D} , \mathbb{E} or A is gentle with underlying graph having exactly one cycle plus some technical condition.

All gentle algebras are known to be of derived tame representation type. (For a definition of derived tameness, see [9].) The same holds for the more general skewed-gentle algebras which have been introduced in [10].

Gentle algebras are defined purely combinatorially in terms of a quiver (directed graph) and relations (for a precise definition see Section 2 below). The more general skewed-gentle algebras are defined from gentle algebras by specifying special vertices which then are split for the quiver of the skewed-gentle algebra. (For a precise definition see Section 4 below.) Note that in contrast to gentle algebras, the class of skewed-gentle algebras is no longer closed under derived equivalences.

It is a long-standing open problem to classify gentle algebras up to derived equivalence. A complete answer has only been obtained for the derived discrete case [7]. The main problem is to find good invariants of the derived categories.

In this paper we provide new, and easy-to-compute invariants of the derived categories of (skewed-) gentle algebras which are of a purely combinatorial nature. Our results are obtained from a detailed computation of the q -Cartan matrices of gentle and skewed-gentle algebras, respectively.

First we state our main result on gentle algebras.

Theorem 1. *Let $A = KQ/I$ be a gentle algebra, given by a gentle quiver Q with relations. Denote by c_k the number of oriented k -cycles in Q with full zero relations.*

Then the q -Cartan matrix $C_A(q)$ is unimodular equivalent (over $\mathbb{Z}[q]$) to a diagonal matrix with entries $(1 - (-q)^k)$, with multiplicity c_k , $k \geq 1$, and all further diagonal entries being 1.

This theorem has the following direct consequences.

Corollary 1. *Let $A = KQ/I$ be gentle, and denote by c_k the number of oriented k -cycles in Q with full zero relations. Then the q -Cartan matrix $C_A(q)$ has determinant*

$$\det C_A(q) = \prod_{k \geq 1} (1 - (-q)^k)^{c_k} .$$

The following special case of Corollary 1 was first proved in [11]. To state it we need the following two combinatorial invariants of the quiver of a gentle algebra A , given as a gentle quiver KQ/I with relations. We denote by $oc(A)$ the number of oriented cycles of odd length in Q having full zero relations, and by $ec(A)$ the number of analogous cycles of even length.

Corollary 2. *Let $A = KQ/I$ be a gentle algebra. Then for the determinant of its Cartan matrix C_A the following holds.*

$$\det C_A = \begin{cases} 0 & \text{if } ec(A) > 0 \\ 2^{oc(A)} & \text{else} \end{cases}$$

The most important application of Theorem 1 is the following corollary which gives for gentle algebras new, combinatorial and easy-to-check invariants of the derived category.

Corollary 3. *Let $A = KQ/I$ be a gentle algebra. If A is derived equivalent to a gentle algebra $A' = KQ'/I'$, then $ec(A) = ec(A')$ and $oc(A) = oc(A')$.*

As an illustration of how powerful these new invariants are, we give in Section 3 a complete derived equivalence classification of gentle algebras with two simple modules and of gentle algebras with three simple modules and Cartan determinant 0.

Our main result on skewed-gentle algebras determines the normal form of the q -Cartan matrix.

Theorem 2. *Let $\hat{A} = K\hat{Q}/\hat{I}$ be a skewed-gentle algebra, arising from choosing a suitable set of special vertices in the gentle pair (Q, I) . Denote by c_k the number of oriented k -cycles in Q with full zero relations.*

Then the q -Cartan matrix $C_{\hat{A}}(q)$ is unimodular equivalent to a diagonal matrix with entries $1 - (-q)^k$, with multiplicity c_k , $k \geq 1$, and all further diagonal entries being 1.

As an immediate consequence we obtain that the Cartan determinant of any skewed-gentle algebra is the same as the Cartan determinant for the underlying gentle algebra.

Corollary 4. *Let $\hat{A} = K\hat{Q}/\hat{I}$ be a skewed-gentle algebra, arising from choosing a suitable set of special vertices in the gentle algebra $A = KQ/I$. Then $\det C_{\hat{A}}(q) = \det C_A(q)$, and thus in particular, the determinants of the ordinary Cartan matrices coincide, i.e., $\det C_{\hat{A}} = \det C_A$.*

The paper is organized as follows. In Section 2 we collect all necessary background and definitions about quivers with relations and (q -)Cartan matrices. In Section 3 we prove all the main results about q -Cartan matrices for gentle algebras. Here we also give some extensive examples to illustrate our results. Section 4 contains the analogous main results for skewed-gentle algebras.

2 Quivers, q -Cartan matrices and derived invariants

Algebras can be defined naturally from a combinatorial setting by using directed graphs. A finite directed graph Q is called a *quiver*. The path algebra KQ , where K is any field, has as basis the set of all oriented paths in Q . The multiplication in the algebra KQ is defined by concatenation of paths, i.e., for a path p in Q let $s(p)$ denote its start vertex and $t(p)$ its end vertex. The product of two paths p and q is defined to be the concatenated path pq if $t(p) = s(q)$, and zero otherwise. Note that our convention is to write paths from left to right.

Such a path algebra KQ is finite dimensional precisely when Q does not contain an oriented cycle. Path algebras form an important, but rather restricted class of algebras. Far more general algebras are obtained by introducing relations on a path algebra.

An ideal $I \subseteq KQ$ is called admissible if $I \subseteq \text{rad}^2(KQ)$ where $\text{rad}(KQ)$ is the radical of the algebra KQ . Note that the radical of a path algebra is the ideal generated by the arrows.

A famous theorem of P. Gabriel states that if K is algebraically closed, any finite-dimensional K -algebra is Morita equivalent to a factor algebra KQ/I where I is an admissible ideal.

So for many representation-theoretic purposes it suffices to consider algebras of the form KQ/I , often referred to as *quivers with relations*.

Let $A = KQ/I$ be as above, where K is any field. By a slight abuse of notation we identify paths in the quiver Q with their cosets in A . Let Q_0 denote the set of vertices of Q . For any $i \in Q_0$ there is a path e_i of length zero. These are primitive orthogonal idempotents in A , the sum $\sum_{i \in Q_0} e_i$ is the unit element in A . In particular we get $A = 1 \cdot A = \bigoplus_{i \in Q_0} e_i A$, hence the (right) A -modules $P_i := e_i A$ are the indecomposable projective A -modules. The *Cartan matrix* $C = (c_{ij})$ of an algebra $A = KQ/I$ is the $|Q_0| \times |Q_0|$ -matrix defined by setting $c_{ij} := \dim_K \text{Hom}_A(P_j, P_i)$. Any homomorphism $\varphi : e_j A \rightarrow e_i A$ of right A -modules is uniquely determined by $\varphi(e_j) \in e_i A e_j$, the K -vector space generated by all paths in Q from vertex i to vertex j ,

which are nonzero in $A = KQ/I$. In particular, we have $c_{ij} = \dim_K e_i A e_j$. In this way, computing entries of the Cartan matrix for $A = KQ/I$ reduces to counting paths in the quiver Q which are nonzero in A . This is the key viewpoint in this paper, enabling us to obtain results on the representation-theoretic Cartan invariants by purely combinatorial methods.


In this paper we study a refined version of the Cartan matrix, which we call q -Cartan matrix. (In the literature also called filtered Cartan matrix, see for instance [8] and the references therein.)

Let $A = KQ/I$ be a finite-dimensional algebra. We assume that the ideal I is generated by homogeneous relations, i.e. by linear combinations of paths having the same length. (Actually, for the algebras considered in this paper, the ideal I will always be generated by monomials and commutativity (mesh) relations.)

Let q be an indeterminate. Instead of just counting paths in A , we now consider the q -weighted version, i.e. each path (non-zero in the algebra) of length n is weighted by q^n . We call the corresponding matrix the q -Cartan matrix $C_A(q)$ of A .

Note that setting $q = 1$ gives the usual Cartan matrix C_A . Even if we are mainly interested in the ordinary Cartan matrix at $q = 1$, the point of view of q -Cartan matrices provides some new insights as we take a closer look at the invariants of the Cartan matrix.

Example 2.1 We consider the following two quivers.



From these we define the following two algebras A and B by quivers with relations. Let $A = KQ_1/I_1$ be defined by the quiver Q_1 , where the ideal I_1 is generated by $\alpha\beta$, $\gamma\delta$ and $\delta\alpha - \beta\gamma$. The q -Cartan matrix of A can be obtained by counting nonzero paths in the quiver and weighting them according to their lengths. We get

$$C_A(q) = \begin{pmatrix} 1 + q^2 & q & 0 \\ q & 1 + q^2 & q \\ 0 & q & 1 + q^2 \end{pmatrix}.$$

The second algebra $B = KQ_2/I_2$ is defined by the quiver Q_2 , subject to the generating relations α^4 (i.e. all paths of length four are zero). The q -Cartan

matrix of B has the form

$$C_B(q) = \begin{pmatrix} 1+q^3 & q & q^2 \\ q^2 & 1+q^3 & q \\ q & q^2 & 1+q^3 \end{pmatrix}.$$

Our main motivation to study Cartan matrices in this paper comes from the context of derived module categories. For finite-dimensional algebras, a crucial problem for understanding derived module categories is to find invariants of the algebras which are preserved under derived equivalences. Cartan matrices provide such invariants, which is our main motivation to study normal forms, invariant factors and determinants of Cartan matrices in this paper. The following result is well-known.

Theorem 2.2 *Let A be a finite-dimensional algebra. The unimodular equivalence class of the Cartan matrix C_A is invariant under derived equivalence. In particular, the determinant of the Cartan matrix is invariant under derived equivalence.*

Remark 2.3 We emphasize that the above theorem only deals with ordinary Cartan matrices $C_A = C_A(1)$. Unfortunately, not even the determinant of the q -Cartan matrix is in general an invariant under derived equivalence. As an example, consider the algebras A and B from Example 2.1. The determinants of their Cartan matrices are different, namely $C_A(q) = 1+q^2+q^4+q^6$ and $C_B(q) = 1+q^3+q^6+q^9$. However, the algebras A and B are derived equivalent. In fact, both are Brauer tree algebras (for a definition see for instance [1]) for Brauer trees with the same number of edges and the same exceptional multiplicity (namely 1 in both cases). Hence, by a result of Rickard [15] they are derived equivalent. Note that when specializing $q = 1$ we indeed get the same determinants for the ordinary Cartan matrices, as predicted by Theorem 2.2

3 Gentle algebras

In this section, we extend and refine the main result on the determinant of the Cartan matrix of a gentle algebra from [11].

We first recall the definition of special biserial algebras and of gentle algebras, as these details will be crucial for what follows. We then prove a preparatory lemma about q -Cartan matrices which holds slightly more generally than only for gentle algebras. The main part of this section will follow, a proof

of Theorem 1 on the unimodular equivalence class of the q -Cartan matrix of an arbitrary gentle algebra.

Let K be a field and let Q be a quiver, with path algebra KQ (see Section 2 for background on path algebras and quivers with relations). A finite dimensional algebra A is called *special biserial* if A is Morita equivalent to an algebra KQ/I where I is an admissible ideal in KQ with the following properties.

- (i) Each vertex of Q is starting point of at most two arrows, and end point of at most two arrows.
- (ii) For each arrow α in Q there is at most one arrow β such that $\alpha\beta \notin I$, and at most one arrow γ such that $\gamma\alpha \notin I$.

Gentle algebras form a subclass of the class of special biserial algebras.

A finite dimensional algebra A is called *gentle* if it is special biserial and moreover the following holds.

- (iii) The ideal I is generated by paths of length 2.
- (iv) For each arrow α in Q there is at most one arrow β' with $t(\alpha) = s(\beta')$ such that $\alpha\beta' \in I$, and there is at most one arrow γ' with $t(\gamma') = s(\alpha)$ such that $\gamma'\alpha \in I$.

The following lemma will turn out to be very useful. It does not only hold for gentle algebras but for those where we have dropped the final condition (iv) in the definition of gentle quivers.

We will study matrices with entries in $\mathbb{Z}[q]$. We call two such matrices C , D unimodularly equivalent over $\mathbb{Z}[q]$ if there exist matrices P , Q over $\mathbb{Z}[q]$ of determinant 1 such that $D = PCQ$.

Lemma 3.1 *Let $A = KQ/I$ be a special biserial algebra, where I is generated by paths of length 2. Let α be an arrow in Q , not a loop, such that there is no arrow β with $s(\alpha) = t(\beta)$ and $\beta\alpha \in I$, or there is no arrow γ with $t(\alpha) = s(\gamma)$ and $\alpha\gamma \in I$. Let Q' be the quiver obtained from Q by removing the arrow α , I' the corresponding relation ideal and $A' = KQ'/I'$. Then the q -Cartan matrices $C_A(q)$ and $C_{A'}(q)$ are unimodularly equivalent (over $\mathbb{Z}[q]$).*

Proof. We consider the case where α is an arrow in Q such that there is no arrow β with $s(\alpha) = t(\beta)$ and $\beta\alpha \in I$; the second case is dual.

Let $\alpha = p_0 : v_0 \rightarrow v_1$. As A is special biserial, there is a unique maximal non-zero path starting with p_0 , say $p = p_0p_1 \dots p_t$, where $p_i : v_i \rightarrow v_{i+1}$, $i = 1, \dots, t$. As A is finite-dimensional, the condition on $\alpha = p_0$ guarantees that $v_i \neq v_0$ for all $i > 0$, but we may have $v_{t+1} = v_i$ for some $i > 0$. Now any

non-zero path of length j , say, ending at v_0 can uniquely be extended to a non-zero path of length $j + i$ ending at v_i , by concatenation with $p_0 \dots p_{i-1}$. Conversely, any non-zero path ending at v_i and involving p_0 arises in this way.

Now denote the column corresponding to a vertex v in the q -Cartan matrix $C_A(q)$ by s_v . We perform column transformations on $C_A(q)$ by replacing the columns s_{v_i} by $s_{v_i} - q^i s_{v_0}$, for $i = 1, \dots, t + 1$ (if $v_i = v_{t+1}$ for some $i < t + 1$, the column s_{v_i} will then be replaced by $s_{v_i} - (q^i + q^{t+1})s_{v_0}$). The resulting matrix $\tilde{C}(q)$ is then exactly the Cartan matrix $C_{A'}(q)$ to the algebra A' corresponding to the quiver Q' where $\alpha = p_0$ has been removed. \diamond

Theorem 3.2 *Let $A = KQ/I$ be gentle. Denote by c_k the number of oriented k -cycles in Q with full zero relations.*

Then the q -Cartan matrix $C_A(q)$ is unimodularly equivalent (over $\mathbb{Z}[q]$) to a diagonal matrix with entries $(1 - (-q)^k)$, with multiplicity c_k , $k \geq 1$, and all further diagonal entries being 1.

Proof. We want to prove the claim by double induction on the number of vertices and the number of arrows; clearly the result holds if Q has no arrows or if it consists of one vertex with a loop.

If Q has a vertex v of degree 1 or 3, or of degree 2 but with no zero relation at v , then we can use Lemma 3.1 to remove an arrow from Q ; note that by the conditions in Lemma 3.1 the removed arrow is not involved in any oriented cycle with full zero relations. Hence $C_A(q)$ is unimodular equivalent to $C_{A'}(q)$, where the corresponding quiver has one arrow less but the same number of oriented cycles with full zero relations, and hence the result holds by induction.

Hence we may now assume that all vertices are of degree 0, 2 or 4, and if a vertex is of degree 2, then there is a zero relation at the vertex. Also, if Q is not connected, we may use induction on the number of vertices to have the result for the components and thus for the whole quiver; hence we may assume that Q is connected. In particular, we now only have vertices v of degree 2 with a non-loop zero relation at v , and vertices of degree 4. As we do not have paths of arbitrary lengths, not all vertices can be of degree 4 (see also [11, Lemma 3]).

Now we take a vertex $v = v_1$ of degree 2, with incoming arrow $p_0 : v_0 \rightarrow v_1$ and outgoing arrow $p_1 : v_1 \rightarrow v_2$ with $p_0 p_1 = 0$ (here, $v_0 \neq v \neq v_2$).

As A is gentle, there is a unique maximal path p in Q with non-repeating arrows starting in v_0 with p_0 , such that the product of any two consecutive

arrows is zero in A ; in our present situation this path is an oriented cycle \mathcal{C} with full zero relations returning to v_0 . We denote the vertices on this path by $v_0, v_1 = v, v_2, \dots, v_s, v_{s+1} = v_0$, and the arrows by $p_i : v_i \rightarrow v_{i+1}$, $i = 0, \dots, s$ (also $p_s p_0 = 0$); note that the arrows on p are distinct, but the vertices are not necessarily distinct (but we point out that $v_i \neq v_0$ for all $i \in \{1, \dots, s\}$).

Denote by z_w the row of the q -Cartan matrix $C_A(q)$ corresponding to the vertex w . In $C_A(q)$, we now replace the row z_v by the linear combination

$$Z = \sum_{i=1}^{s+1} (-q)^{i-1} z_{v_i}$$

to obtain a new matrix $\tilde{C}(q)$ (note that this is a unimodular transformation over $\mathbb{Z}[q]$).

The careful choice of the coefficients is just made so that we can refine the argument in [11]. We recall some of the notation there. For any arrow α in Q let $\mathcal{P}(\alpha)$ be the set of paths starting with α which are non-zero in A . At each vertex v_i , $i \geq 1$, there is at most one outgoing arrow $r_i \neq p_i$; for this arrow we have $p_{i-1} r_i \neq 0$, as A is gentle.

Hence, cancelling p_i induces a natural bijection $\phi : \mathcal{P}(p_i) \rightarrow \{e_{v_{i+1}}\} \cup \mathcal{P}(r_{i+1})$, for $i = 1, \dots, s-1$, such that a path of q -weight q^j is mapped to a path of q -weight q^{j-1} .

As v is of degree 2, with a zero-relation at v , we also have the trivial bijection $\mathcal{P}(p_0) = \{p_0\} \rightarrow \{e_{v_1}\}$, again with a weight reduction by q . Now almost everything cancels in Z , apart from the one term $1 - (-q)^{s+1}$ that we obtain as the entry in the column corresponding to v .

In the next step, we use the dual (counter-clockwise) operation on the columns labelled by the vertices on the cycle \mathcal{C} , i.e., we set $v_{s+2} = v = v_1$ and replace the column s_v by the linear combination

$$S = \sum_{i=1}^{s+1} (-q)^{s+1-i} s_{v_{i+1}}.$$

Ordering vertices so that v corresponds to the first row and column of the Cartan matrix, we have thus unimodularly transformed $C_A(q)$ to a matrix of the form

$$\begin{pmatrix} 1 - (-q)^{s+1} & 0 & \dots & 0 \\ 0 & & & \\ \vdots & & C'(q) & \\ 0 & & & \end{pmatrix}$$

where $C'(q)$ is the q -Cartan matrix of the gentle algebra A' for the quiver Q' obtained from Q by removing v and the arrows incident with v . Note that in comparison with Q , the quiver Q' has one vertex less and one cycle with full relations of length $s + 1$ less; now by induction, the result holds for $C'(q) = C_{A'}(q)$, and hence the result for $C_A(q)$ follows immediately. \diamond

This result has several immediate nice consequences.

Corollary 3.3 *Let $A = KQ/I$ be gentle, and denote by c_k the number of oriented k -cycles in Q with full zero relations. Then the q -Cartan matrix $C_A(q)$ has determinant*

$$\det C_A(q) = \prod_{k \geq 1} (1 - (-q)^k)^{c_k}.$$

The following observation is a direct consequence of Theorem 3.2, but could also easily be proved directly by induction.

Remark 3.4 Let $A = KQ/I$ be gentle, with set of vertices Q_0 . Then there are at most $|Q_0|$ oriented cycles with full zero relations in the quiver.

Note that the property of being gentle is invariant under derived equivalence [16], and we now have some invariants to distinguish the derived equivalence classes. For an algebra $A = KQ/I$, recall that $ec(A)$ and $oc(A)$ denote the number of oriented cycles in Q with full zero relations of even and odd length, respectively.

As an immediate special case of Corollary 3.3 we obtain the following formula for the Cartan determinant which was the main result of the previous paper [11].

Corollary 3.5 ([11]) *Let $A = KQ/I$ be a gentle algebra. Then for the determinant of the Cartan matrix C_A the following holds.*

$$\det C_A = \begin{cases} 0 & \text{if } ec(A) > 0 \\ 2^{oc(A)} & \text{else} \end{cases}$$


Note that in combination with Remark 3.4 this implies that the Cartan determinant of a gentle algebra $A = KQ/I$ is at most $2^{l(A)}$, where $l(A) = |Q_0|$ is the number of simple modules of A .

The most important application of our Theorem 3.2 is the following corollary which gives for gentle algebras new, combinatorial and easy-to-check invariants of the derived category.

Corollary 3.6 *Let $A = KQ/I$ be a gentle algebra. If A is derived equivalent to a gentle algebra $A' = KQ'/I'$, then $ec(A) = ec(A')$ and $oc(A) = oc(A')$.*

We now illustrate our results and apply them to derived equivalence classifications of gentle algebras.

Example 3.7 Gentle algebras with two simple modules. There are nine connected gentle algebras with two simple modules, as given in the following list.



In [5] it was shown that these are precisely the basic connected algebras with two simple modules which are derived tame (in the sense of [9]). As a direct illustration of our results we show how to classify these algebras up to derived equivalence.

Recall that the property of being gentle is invariant under derived equivalence [16]. Moreover, the number of simple modules of an algebra is a derived invariant [13]. Thus we will be able to describe the complete derived equivalence classes.

We have shown above that the numbers $oc(A)$ and $ec(A)$ are derived invariants. In addition we look at two classical invariants, the center and the first Hochschild cohomology group HH^1 . Recall that the center of an algebra (and more generally the Hochschild cohomology ring) is invariant under derived equivalence [14]. If the quiver contains a loop, then the dimension of HH^1 depends on the characteristic being 2 or not. We indicate

the dimension in characteristic 2 in brackets in the table below. They can be computed using a method based on work of M. Bardzell [4] on minimal projective bimodule resolutions for monomial algebras. A very nice explicit combinatorial description is given in [17], Proposition 2.6.

Algebra	A_1	A_2	A_3	A_4	A_5	A_6	A_7	A_8	A_9
$oc(A)$	0	0	0	0	1	1	0	1	2
$ec(A)$	0	0	0	0	0	0	1	1	0
$\dim Z(A)$	1	1	1	2	1	1	1	2	1
$\dim \mathrm{HH}^1(A)$	0	3	2	1	1(2)	1(2)	1	2(3)	3(5)

The algebras A_1, A_2, A_3, A_4 are pairwise not derived equivalent. This can be deduced directly from the above table, since the dimensions of the first Hochschild cohomology groups are different.

Alternatively, one can argue as follows by using some classical results in tilting theory. By [2], the algebras derived equivalent to a hereditary algebra of type \mathbb{A} are the gentle algebras whose underlying undirected graph is a tree. So A_1 can not be derived equivalent to any of A_2, A_3, A_4 .

By [3], the algebras derived equivalent to a hereditary algebra of type $\tilde{\mathbb{A}}$ are the gentle algebras whose underlying graph has exactly one cycle and in this cycle the 'clock condition' is satisfied (i.e., the number of zero relations in clockwise direction and in anti-clockwise direction coincide). Hence, A_2 is not derived equivalent to A_3 or A_4 .

Finally, A_3 and A_4 are not derived equivalent since their centers are not isomorphic.

The algebras A_5 and A_6 are derived equivalent. (This can be shown by explicitly constructing a suitable tilting complex, similar to the detailed example given in the Appendix.)

The algebras A_7 and A_8 with Cartan determinant 0 are not derived equivalent, since their centers have different dimensions.

In summary, there are exactly eight derived equivalence classes of connected gentle algebras with two simple modules. They are indicated by the double vertical lines in the above table.


Example 3.8 Gentle algebras with three simple modules. As a second, more involved, illustration of our main result we consider (connected) gentle algebras with three simple modules. Using our results, we shall see how easily we can distinguish several gentle algebras up to derived equivalence which can not be distinguished by the more classical invariants. How-

ever, to complete the derived equivalence classification we also take some classical invariants into account.

Let A be a connected gentle algebra with three simple modules. By Corollary 3.4 we deduce that $\det C_A \in \{0, 1, 2, 4, 8\}$. Algebras with different Cartan determinant can not be derived equivalent, by Theorem 2.2.

We shall give a complete derived equivalence classification of those algebras with Cartan determinant 0. Recall that by Corollary 3.5, a gentle algebra has Cartan determinant 0 if and only if the quiver contains an even oriented cycle with full zero relations.

There are 18 connected gentle algebras with three simple modules having Cartan determinant 0, as listed in the following figure.



The main tool will be our Corollary 3.6 which states that the numbers $ec(A)$ and $oc(A)$ of oriented cycles with full zero relations of even (resp. odd) length are invariants of the derived category. This will already settle large parts of the classification, especially the notoriously difficult problem of distinguishing different derived equivalence classes.

In addition we will need to look at the centers and at the first Hochschild cohomology group. The following table collects all the necessary invariants. Again, in the cases where the quiver has loops, the dimension of HH^1 depends on the characteristic being 2 or not, and in these cases the dimension in characteristic 2 is given in brackets.

Algebra	$oc(\Lambda)$	$ec(\Lambda)$	$\dim Z(\Lambda)$	$\dim \mathrm{HH}^1(\Lambda)$
Λ_1	0	1	1	1
Λ_2	0	1	1	1
Λ_3	0	1	1	4
Λ_4	0	1	1	4
Λ_5	0	1	2	2
Λ_6	0	1	2	2
Λ_7	0	2	1	2
Λ_8	0	1	3	2
Λ_9	1	1	1	2(3)
Λ_{10}	1	1	1	2(3)
Λ_{11}	1	1	1	2(3)
Λ_{12}	1	1	1	2(3)
Λ_{13}	1	2	1	3(4)
Λ_{14}	2	1	1	4(6)
Λ_{15}	2	1	1	4(6)
Λ_{16}	0	1	1	4
Λ_{17}	0	1	1	3
Λ_{18}	0	1	1	3

For the derived equivalence classification, it only remains to consider those algebras having the same invariants. In the cases where the algebras are in fact derived equivalent, we leave out the details of the construction of a suitable tilting complex; in the appendix a detailed example is provided which serves to indicate the strategy which also works in all other cases.

The algebras Λ_1 and Λ_2 are derived equivalent. Moreover, the algebras Λ_3 and Λ_4 are derived equivalent.

Note that Λ_1 and Λ_3 represent different derived equivalence classes since their first Hochschild cohomology groups have different dimensions. (Alternatively, one can argue that Λ_1 has a discrete derived category (in the sense of [18], to which we also refer for the definition), whereas the derived category of Λ_3 is not discrete.)

The algebras Λ_5 and Λ_6 are derived equivalent. (The details for this case are provided in the appendix.)

Similarly, the algebras Λ_9 , Λ_{10} , Λ_{11} and Λ_{12} are derived equivalent, the algebras Λ_{14} and Λ_{15} are derived equivalent and moreover, the algebras Λ_{17} and Λ_{18} are derived equivalent.

The case of Λ_{16} is more subtle. This algebra has exactly the same invariants listed in the table as the algebras Λ_3 and Λ_4 . However, we claim that Λ_{16} is not derived equivalent to Λ_4 . In fact, the Lie algebra structures on HH^1 are not isomorphic. Note that with the Gerstenhaber bracket, the first Hochschild cohomology becomes a Lie algebra. By a result of B. Keller [12], this Lie algebra structure on HH^1 is invariant under derived equivalence. Based on work of M. Bardzell [4] on minimal projective bimodule resolutions for monomial algebras, there is an explicit way of computing HH^1 for a gentle algebra. A nice combinatorial version is given in [17], Proposition 2.6 (for the additive structure) and Theorem 2.7 (for the Lie algebra structure). With this method one can compute that the four-dimensional Lie algebras on $\mathrm{HH}^1(\Lambda_{16})$ and on $\mathrm{HH}^1(\Lambda_4)$ are not isomorphic. For instance, the Lie algebra center of $\mathrm{HH}^1(\Lambda_{16})$ is two-dimensional, whereas the Lie algebra center of $\mathrm{HH}^1(\Lambda_4)$ has dimension 1.

This completes the derived equivalence classification of connected gentle algebras with three simple modules and Cartan determinant 0. The ten derived equivalence classes are indicated in the above table by the horizontal double lines.

4 Skewed-gentle algebras

Skewed-gentle algebras were introduced in [10]; for the notation and definition we follow here mostly [6], but we try to explain how the construction works rather than repeating the technical definition from [6].

We start with a gentle pair (Q, I) . A set Sp of vertices of the quiver Q is an admissible set of *special* vertices if the quiver with relations obtained from Q by adding loops with square zero at these vertices is again gentle; we denote this gentle pair by $(Q^{\mathrm{sp}}, I^{\mathrm{sp}})$. The triple (Q, Sp, I) is then called *skewed-gentle*.

We want to point out that the admissibility of the set Sp of special vertices is both a local as well as a global condition. Let v be a vertex in the gentle quiver (Q, I) ; then we can only add a loop at v if v is of degree 1 or 0 or if it is of degree 2 with a zero relation, but not one coming from a loop. Hence only vertices of this type are potential special vertices. But for the choice of an admissible set of special vertices we also have to take care of the global condition that after adding all loops, the pair $(Q^{\text{sp}}, I^{\text{sp}})$ still does not have paths of arbitrary lengths.

Given a skewed-gentle triple (Q, Sp, I) , we now construct a new quiver with relations (\hat{Q}, \hat{I}) by doubling the special vertices, introducing arrows to and from these vertices corresponding to the previous such arrows and replacing a previous zero relation at the vertices by a mesh relation.

More precisely, we proceed as follows. The non-special (or: ordinary) vertices in Q are also vertices in the new quiver; any arrow between non-special vertices as well as corresponding relations are also kept. Any special vertex $v \in Sp$ is replaced by two vertices v^+ and v^- in the new quiver. An arrow a in Q from a non-special vertex w to v (or from v to w) will be doubled to arrows $a^\pm : w \rightarrow v^\pm$ (or $a^\pm : v^\pm \rightarrow w$, resp.) in the new quiver; an arrow between two special vertices v, w will correspondingly give four arrows between the pairs v^\pm and w^\pm . We say that these new arrows lie over the arrow a . Any relation $ab = 0$ where $t(a) = s(b)$ is non-special gives a corresponding zero relation for paths of length 2 with the same start and end points lying over ab . If v is a special vertex of degree 2 in Q , then the corresponding zero relation at v , say $ab = 0$ with $t(a) = v = s(b)$, is replaced by mesh commutation relations saying that any two paths of length 2 lying over ab , having the same start and end points but running over v^+ and v^- , respectively, coincide in the factor algebra to the new quiver with relations (\hat{Q}, \hat{I}) .

We will speak of (\hat{Q}, \hat{I}) as a skewed-gentle quiver *covering* the gentle pair (Q, I) .

A K -algebra is then called *skewed-gentle* if it is Morita equivalent to a factor algebra $K\hat{Q}/\hat{I}$, where (\hat{Q}, \hat{I}) comes from a skewed-gentle triple (Q, Sp, I) as above.

Remark. Let $A = KQ/I$ be gentle. In a gentle quiver, there is at most one non-zero cyclic path starting and ending at a given vertex; hence the diagonal entries in the q -Cartan matrix $C_A(q)$ are 1 or of the form $1 + q^j$, for some $j \in \mathbb{N}$.

If a vertex v in Q can be chosen as a special vertex for a covering skewed-gentle quiver \hat{Q} , then the corresponding diagonal entry in $C_A(q)$ is 1, as

otherwise we have paths of arbitrary lengths in Q^{sp} ; hence in the corresponding q -Cartan matrix for the skewed-gentle algebra \hat{A} we have $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ on the diagonal for the two split vertices v^\pm in \hat{Q} .

Theorem 4.1 *Let (Q, I) be a gentle quiver, and (\hat{Q}, \hat{I}) a covering skewed-gentle quiver. Let $\hat{A} = K\hat{Q}/\hat{I}$ be the corresponding skewed-gentle algebra. Denote by c_k the number of oriented k -cycles in (Q, I) with full zero relations.*

Then the q -Cartan matrix $C_{\hat{A}}(q)$ is unimodularly equivalent (over $\mathbb{Z}[q]$) to a diagonal matrix with entries $1 - (-q)^k$, with multiplicity c_k , $k \geq 1$, and all further diagonal entries being 1.

Proof. Again, we argue by induction on the number of vertices and arrows. We let $A = KQ/I$ be the gentle algebra and $C_A(q)$ the q -Cartan matrix as before.

If Q has no arrows, then \hat{Q} is just obtained by doubling the special vertices, and this still has no arrows, so the result clearly holds.

If Q has an arrow α as in Lemma 3.1, with a non-special $s(\alpha)$ in the first case, and a non-special $t(\alpha)$ in the second case, respectively, then we can argue as in the proof of Lemma 3.1 to remove α . Let us consider again the situation of the first case, so here $s(\alpha) = v_0$ is non-special.

Note that a maximal non-zero path p starting from v_0 with α or α^\pm (if v_1 is special) will end on a non-special vertex (and hence this maximal path is unique in \hat{A}); in general, this path will be longer than the one taken in A . In the column transformations, we only have to be careful at doubled vertices on the path p ; here we replace both corresponding columns $\hat{s}_{v_i^\pm}$ of $C_{\hat{A}}(q)$ by $\hat{s}_{v_i^\pm} - q^i \hat{s}_{v_0}$. This leads us to the Cartan matrix for the skewed-gentle algebra where α or α^\pm , respectively, has been removed from \hat{Q} , which is a skewed-gentle cover for the quiver obtained from Q by deleting α ; then the claim follows by induction.

Now assume Q has a source v which is special (w.l.o.g. the first vertex); the case of a sink is dual. Then the q -Cartan matrix for \hat{A} has the form

$$C_{\hat{A}}(q) = \begin{pmatrix} 1 & 0 & * & \cdots & * \\ 0 & 1 & * & \cdots & * \\ 0 & 0 & & & \\ \vdots & \vdots & & \hat{C}'(q) & \\ 0 & 0 & & & \end{pmatrix} \sim \tilde{C}(q) = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & & & \\ \vdots & \vdots & & \hat{C}'(q) & \\ 0 & 0 & & & \end{pmatrix}$$

where $\hat{C}'(q)$ is the q -Cartan matrix of the skewed-gentle algebra \hat{A}' for the quiver \hat{Q}' obtained from \hat{Q} by removing v^+, v^- and the arrows incident with v^\pm . Note that \hat{Q}' is the skewed-gentle cover for the quiver Q' which is obtained from Q by removing v and the arrow incident with v , and the choice $S_{p'} = Sp \setminus \{v\}$ as the set of special vertices; in short, we write this as $\hat{Q}' = \widehat{Q}'$. Again, using induction the claim follows immediately.

Thus again, we may now assume that Q has only vertices of degree 2 with a (non-loop) zero relation or vertices of degree 4; note that any special vertex in Q has to be of degree 2. As before, we may also assume that Q (and hence also \hat{Q}) are connected.

If there are no non-special vertices, or if all non-special vertices are of degree 4, then Q^{sp} is not gentle. Hence Q has a non-special vertex v of degree 2 with a zero relation at v . Let $p_0 : v_0 \rightarrow v$ be the (unique) incoming arrow. Again we consider the unique maximal path p in Q with non-repeating arrows starting in v_0 with p_0 , such that the product of any two consecutive arrows is zero in A ; as before, we note that in our current situation p has to be a cycle $\mathcal{C} = p_0 p_1 \dots p_s$, where $p_i : v_i \rightarrow v_{i+1}$, $i = 0, \dots, s$, and $v_{s+1} = v_0$. As in the previous situation, we note that the arrows are distinct, but vertices $v_i \neq v_0$ may be repeated.

For a vertex w in Q we denote by z_w the row of the q -Cartan matrix $C_A(q)$ corresponding to w .

If w is non-special, we denote by \hat{z}_w the corresponding row in the q -Cartan matrix $\hat{C}(q) = C_{\hat{A}}(q)$. If w is special, then for the two vertices w^\pm we have two corresponding rows \hat{z}_{w^\pm} in the Cartan matrix $\hat{C}(q)$, and we then set $\hat{z}_w = \hat{z}_{w^+} + \hat{z}_{w^-}$.

Before, we have transformed C by replacing z_v by $Z = \sum_{i=1}^{s+1} (-q)^{i-1} z_{v_i}$ and obtained a matrix $\tilde{C}(q)$. We now do a parallel transformation on $\hat{C}(q)$, that is, we replace \hat{z}_v by

$$\hat{Z} = \sum_{i=1}^{s+1} (-q)^{i-1} \hat{z}_{v_i},$$

and we obtain a matrix $\tilde{\hat{C}}(q)$. We have to compare the differences and check that everything stays under control for the induction argument.

If a vertex v_i , $1 \leq i \leq s$, is special, note that the doubled contribution in $\hat{z}_{v_i} = \hat{z}_{v_i^+} + \hat{z}_{v_i^-}$ is needed on the one hand for the cancellation with the previous row, and on the other hand to continue around the cycle \mathcal{C} . As v is non-special and of degree 2 with a zero-relation, we note that as before, in \hat{Z} we only have the contribution $1 - (-q)^{s+1}$ at v .

Following this by the parallel operation to the previous column operation we then replace the column \hat{s}_v by the linear combination

$$\hat{S} = \sum_{i=1}^{s+1} (-q)^{s+1-i} \hat{s}_{v_{i+1}} ,$$

where we use analogous conventions as before.

With v corresponding to the first row and column of the Cartan matrix, we have thus unimodularly transformed $\hat{C}(q)$ to a matrix of the form

$$\begin{pmatrix} 1 - (-q)^{s+1} & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & \hat{C}'(q) & \\ 0 & & & \end{pmatrix}$$

where $\hat{C}'(q)$ is the Cartan matrix of the skewed-gentle algebra \hat{A}' for the quiver \hat{Q}' obtained from \hat{Q} by removing v and the arrows incident with v . Note that in fact, $\hat{Q}' = \widehat{Q}'$ in the notation of our previous proof, i.e., as explained earlier, \widehat{Q}' is the skewed-gentle cover for the quiver Q' and the choice $Sp' = Sp \setminus \{v\}$ as the set of special vertices.

Thus the result follows by induction. \diamond

Remark 4.2 By comparing Theorem 3.2 and Theorem 4.1 we observe that the q -Cartan matrix $C_A(q)$ for the gentle algebra A to (Q, I) , and the q -Cartan matrix $C_{\hat{A}}(q)$ for a skewed-gentle cover \hat{A} are unimodularly equivalent to diagonal matrices which only differ by adding as many further 1's on the diagonal as there are special vertices chosen in Q . In particular, with notation as above,

$$\det C_{\hat{A}}(q) = \det C_A(q) = \prod_{k \geq 1} (1 - (-q)^k)^{c_k} .$$

This observation has the following immediate consequence when specializing to $q = 1$.

Corollary 4.3 *Let (Q, I) be a gentle quiver, and (\hat{Q}, \hat{I}) a covering skewed-gentle quiver. Then the determinant of the ordinary Cartan matrix of the skewed-gentle algebra $\hat{A} = K\hat{Q}/\hat{I}$ is the same as the one for the gentle algebra $A = KQ/I$, i.e., $\det C_{\hat{A}} = \det C_A$.*

Remark 4.4 A gentle algebra and a (proper) skewed-gentle algebra may have the same q -invariants but they cannot be derived equivalent by [16], Corollary 1.2.

5 Appendix: Tilting complexes and derived equivalences, a detailed example


This appendix is aimed at providing enough background on tilting complexes and explicit computations of their endomorphism rings so that the interested reader can fill in the details in the derived equivalence classifications of Examples 3.7 and 3.8. We explained there in detail how to distinguish derived equivalence classes (since this is the main topic of this paper), but have been fairly short on indicating why certain algebras in the lists are actually derived equivalent. In this section we will go through one example in detail; this will indicate the main strategy which also works in all other cases.

For an algebra A denote by $D^b(A)$ the bounded derived category and by $K^b(P_A)$ the homotopy category of bounded complexes of finitely generated projective A -modules.

Two algebras A and B are called derived equivalent if $D^b(A)$ and $D^b(B)$ are equivalent as triangulated categories. By J. Rickard's celebrated theorem [13], this happens if and only there exists a tilting complex T for A such that the endomorphism ring $\text{End}_{K^b(P_A)}(T)$ in the homotopy category is isomorphic to B . A bounded complex T of projective A -modules is called a tilting complex if the following conditions are satisfied.

- (i) $\text{Hom}_{K^b(P_A)}(T, T[i]) = 0$ for $i \neq 0$ (where $[\cdot]$ denotes the shift operator)
- (ii) $\text{add}(T)$, the full subcategory of $K^b(P_A)$ consisting of direct summands of direct sums of copies of T , generates $K^b(P_A)$ as a triangulated category.

In Example 3.8 we stated that the algebras Λ_5 and Λ_6 are derived equivalent. For the convenience of the reader we recall the definition of these algebras.



Recall from Section 2 our conventions to deal with right modules and to read paths from left to right. In particular, left multiplication by a nonzero path from vertex j to vertex i gives a homomorphism $P_i \rightarrow P_j$.

We define the following bounded complex $T := T_1 \oplus T_2 \oplus T_3$ of projective Λ_5 -modules. Let $T_1 . : 0 \rightarrow P_3 \rightarrow 0$ and $T_3 . : 0 \rightarrow P_1 \rightarrow 0$ be stalk complexes concentrated in degree 0. Moreover, let $T_2 . : 0 \rightarrow P_1 \oplus P_3 \xrightarrow{(\delta, \beta)} P_2 \rightarrow 0$ (in

degrees 0 and -1). We claim that T is a tilting complex. Property (i) above is obvious for all $|i| \geq 2$ since we are dealing with two-term complexes. Let $i = 1$, and consider possible maps $T_2 \rightarrow T_j[-1]$ where $j \in \{1, 2, 3\}$. This is given by a map of complexes as follows

$$\begin{array}{ccccccc} 0 & \longrightarrow & P_1 \oplus P_3 & \xrightarrow{(\delta, \beta)} & P_2 & \longrightarrow & 0 \\ & & & & \downarrow & & \\ & & 0 & \longrightarrow & Q & \longrightarrow & \dots \end{array}$$

where Q could be either of P_1 , P_3 , or $P_1 \oplus P_3$. But since we are dealing with gentle algebras, no nonzero map can be zero when composed with both δ and β . So the only homomorphism of complexes $T_2 \rightarrow T_j[-1]$ is the zero map, as desired. Directly from the definition we see that $\text{Hom}(T_1, T_j[-1]) = 0$ and $\text{Hom}(T_3, T_j[-1]) = 0$ (since they are stalk complexes). Thus we have shown that $\text{Hom}(T, T[-1]) = 0$.

Now let $i = 1$. We have to consider maps $T_j \rightarrow T_2[1]$; these are given as follows

$$\begin{array}{ccccccc} & & 0 & \longrightarrow & Q & \longrightarrow & \dots \\ & & & & \downarrow & & \\ 0 & \longrightarrow & P_1 \oplus P_3 & \xrightarrow{(\delta, \beta)} & P_2 & \longrightarrow & 0 \end{array}$$

where Q again can be either of P_1 , P_3 , or $P_1 \oplus P_3$. Now there certainly exist nonzero homomorphisms of complexes. But they are all homotopic to zero. In fact, every path in the quiver of Λ_5 from vertex 2 to vertex 1 or 3 either starts with δ or with β . Accordingly, every homomorphism $Q \rightarrow P_2$ can be factored through the map $(\delta, \beta) : P_1 \oplus P_3 \rightarrow P_2$.

It follows that $\text{Hom}_{K^b(P_A)}(T, T[1]) = 0$ (in the homotopy category).

It remains to show that the complex T also satisfies property (ii) of the definition of a tilting complex. It suffices to show that the projective indecomposable modules P_1 , P_2 and P_3 , viewed as stalk complexes, can be generated by $\text{add}(T)$. This is clear for P_1 and P_3 since they occur as summands of T . For P_2 , consider the map of complexes $\Psi : T_2 \rightarrow T_3 \oplus T_1$ given by the identity map on $P_1 \oplus P_3$ in degree 0. Then the stalk complex $P_2[0]$ with P_2 in degree 0 can be shown to be homotopy equivalent (i.e. isomorphic in $K^b(P_A)$) to the mapping cone of Ψ . Thus we have a distinguished triangle

$$\underbrace{T_2}_{\in \text{add}(T)} \rightarrow \underbrace{T_3 \oplus T_1}_{\in \text{add}(T)} \rightarrow P_2[0] \rightarrow \underbrace{T_2[1]}_{\in \text{add}(T)} .$$

By definition, $\text{add}(T)$ is triangulated, so it follows that also the stalk complex $P_2[0] \in \text{add}(T)$, which proves (ii).

Hence, T is indeed a tilting complex for Λ_5 .

By Rickard's theorem, the endomorphism ring of T in the homotopy category is derived equivalent to Λ_5 . We need to show that $E := \text{End}_{K^b(P_A)}(T)$ is isomorphic to Λ_6 . Note that the vertices of the quiver of E correspond to the summands of T .

For explicit calculations, the following formula is useful, which gives a general method for computing the Cartan matrix of an endomorphism ring of a tilting complex from the Cartan matrix of A .

Alternating sum formula. For a finite-dimensional algebra A , let $Q = (Q^r)_{r \in \mathbb{Z}}$ and $R = (R^s)_{s \in \mathbb{Z}}$ be bounded complexes of projective A -modules. Then

$$\sum_i (-1)^i \dim \text{Hom}_{K^b(P_A)}(Q, R[i]) = \sum_{r,s} (-1)^{r-s} \dim \text{Hom}_A(Q^r, R^s).$$

In particular, if Q and R are direct summands of a tilting complex then

$$\dim \text{Hom}_{K^b(P_A)}(Q, R) = \sum_{r,s} (-1)^{r-s} \dim \text{Hom}_A(Q^r, R^s).$$

Note that the Cartan matrix of Λ_5 has the form $\begin{pmatrix} 2 & 2 & 1 \\ 2 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}$. From that,

using the alternating sum formula, we can compute the Cartan matrix of E to be $\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 2 \end{pmatrix}$. Note that this is actually the Cartan matrix of Λ_6 .

Now we have to define maps of complexes between the summands of T , corresponding to the arrows of the quiver of Λ_6 . The final step then is to show that these maps satisfy the defining relations of Λ_6 , up to homotopy.

We define $\tilde{\alpha} : T_1 \rightarrow T_2$ by the map $(\alpha\beta, 0) : P_3 \rightarrow P_1 \oplus P_3$ in degree 0. Note that this is indeed a homomorphism of complexes since $\delta\alpha = 0$ in Λ_5 . Moreover, we define $\tilde{\beta} : T_2 \rightarrow T_3$ and $\tilde{\delta} : T_2 \rightarrow T_1$ by the projection onto the first and second summand in degree 0, respectively. Finally, we define $\tilde{\gamma} : T_3 \rightarrow T_1$ by $\gamma\delta : P_1 \rightarrow P_3$.

We now have to check the relations, up to homotopy. We write compositions from left to right (as in the relations of the quiver of E). Clearly, $\tilde{\alpha}\tilde{\delta} = 0$. The composition $\tilde{\beta}\tilde{\gamma} : T_2 \rightarrow T_1$ is given in degree 0 by $(\gamma\delta, 0) : P_1 \oplus P_3 \rightarrow P_3$. So it is not the zero map, but is homotopic to zero via the homotopy map $\gamma : P_2 \rightarrow P_3$ (use that $\gamma\beta = 0$ in Λ_5). Finally, consider $\tilde{\delta}\tilde{\alpha}$ on T_2 . It is given by $\begin{pmatrix} 0 & \alpha\beta \\ 0 & 0 \end{pmatrix}$ in degree 0 and the zero map in degree -1 . It is indeed

homotopic to zero via the homotopy map $(\alpha, 0) : P_2 \rightarrow P_1 \oplus P_3$. (Note that here we use that $\alpha\delta = 0$ and $\delta\alpha = 0$ in Λ_5 .)

Thus, we have defined maps between the summands of T , corresponding to the arrows of the quiver of Λ_6 . We have shown that they satisfy the defining relations of Λ_6 , and that the Cartan matrices of E and Λ_6 . From this we can conclude that $E \cong \Lambda_6$. Hence, Λ_5 and Λ_6 are derived equivalent, as desired.

All the other derived equivalences stated in Examples 3.7 and 3.8 can be verified exactly along these lines. In particular, they can also be realized by tilting complexes with non-zero entries in only two degrees.

References

- [1] J. Alperin, Local representation theory, Cambridge Studies in Advanced Mathematics, 11. Cambridge University Press, Cambridge, 1986
- [2] I. Assem, D. Happel, Generalized tilted algebras of type \mathbb{A}_n , Comm. Algebra 9 (1981), no.20, 2101-2125
- [3] I. Assem, A. Skowroński, Iterated tilted algebras of type $\tilde{\mathbb{A}}_n$, Math. Z. 195 (1987), 269-290
- [4] M. J. Bardzell, The alternating syzygy behavior of monomial algebras, J. Algebra 188 (1997), 69–89
- [5] V. Bekkert, Y. Drozd, Derived categories for some classes of algebras, Preprint. Available at <http://www.mat.ufmg.br/~bekkert/pub.html>
- [6] V. Bekkert, E.N. Marcos, H.A. Merklen, Indecomposables in derived categories of skewed-gentle algebras, Communications in Algebra (6) 31 (2003), 2615-2654
- [7] G. Bobiński, C. Geiss, A. Skowroński, Classification of discrete derived categories, Cent. Eur. J. Math. 2 (2004), no. 1, 19-49
- [8] K. Fuller, The Cartan determinant and global dimension of artinian rings, in: Azumaya algebras, actions, and modules, Proceedings 1990, Contemporary Mathematics 124 (AMS 1992) 51-72
- [9] C. Geiss, H. Krause, On the notion of derived tameness, J. Algebra Appl. 1 (2002), no. 2, 133-157

- [10] C. Geiss, J. A. de la Peña, Auslander-Reiten components for clans, *Boll. Soc. Mat. Mexicana* 5 (1999), 307-326
- [11] Th. Holm, Cartan determinants for gentle algebras, *Archiv Math.*, to appear
- [12] B. Keller, Derived invariance of higher structures on the Hochschild complex, Preprint (2003)
- [13] J. Rickard, Morita theory for derived categories, *J. London Math. Soc.* (2) 39 (1989) 436-456
- [14] J. Rickard, Derived equivalences as derived functors. *J. London Math. Soc.* (2) 43 (1991), 37-48
- [15] J. Rickard, Derived categories and stable equivalence. *J. Pure Appl. Algebra* 61 (1989), 303-317
- [16] J. Schröer, A. Zimmermann, Stable endomorphism algebras of modules over special biserial algebras, *Math. Z.* 244 (2003) 515-530
- [17] C. Strametz, The Lie algebra structure on the first Hochschild cohomology group of a monomial algebra, Preprint (2001), math.RT/0111060
- [18] D. Vossieck, The algebras with discrete derived category, *J. Algebra* 243 (2001), 168-176