Acta Arith. 127(2007), no. 2, 103–113.

MIXED SUMS OF SQUARES AND TRIANGULAR NUMBERS

ZHI-WEI SUN (NANJING)

ABSTRACT. By means of q-series, we prove that any natural number is a sum of an even square and two triangular numbers, and that each positive integer is a sum of a triangular number plus $x^2 + y^2$ for some $x, y \in \mathbb{Z}$ with $x \not\equiv y \pmod 2$ or x = y > 0. The paper also contains some other results and open conjectures on mixed sums of squares and triangular numbers.

1. Introduction

A classical result of Fermat asserts that any prime $p \equiv 1 \pmod{4}$ is a sum of two squares of integers. Fermat also conjectured that each $n \in \mathbb{N}$ can be written as a sum of three triangular numbers, where \mathbb{N} is the set $\{0,1,2,\ldots\}$ of natural numbers, and triangular numbers are those integers $t_x = x(x+1)/2$ with $x \in \mathbb{Z}$. An equivalent version of this conjecture states that 8n+3 is a sum of three squares (of odd integers). This follows from the following profound theorem (see, e.g., [G, pp. 38–49] or [N, pp. 17-23]).

Gauss-Legendre Theorem. $n \in \mathbb{N}$ can be written as a sum of three squares of integers if and only if n is not of the form $4^k(8l+7)$ with $k, l \in \mathbb{N}$.

Building on some work of Euler, in 1772 Lagrange showed that every natural number is a sum of four squares of integers.

For problems and results on representations of natural numbers by various quadratic forms with coefficients in \mathbb{N} , the reader may consult [Du] and [G].

Motivated by Ramanujan's work [Ra], L. Panaitopol [P] proved the following interesting result in 2005.

Theorem A. Let a, b, c be positive integers with $a \le b \le c$. Then every odd natural number can be written in the form $ax^2 + by^2 + cz^2$ with $x, y, z \in \mathbb{Z}$, if and only if the vector (a, b, c) is (1, 1, 2) or (1, 2, 3) or (1, 2, 4).

 $^{2000\} Mathematics\ Subject\ Classification:$ Primary 11E25; Secondary 05A30, 11D85, 11P99.

Supported by the National Science Fund for Distinguished Young Scholars (Grant No. 10425103) in China.

Z. W. SUN

According to L. E. Dickson [D2, p. 260], Euler already noted that any odd integer n > 0 is representable by $x^2 + y^2 + 2z^2$ with $x, y, z \in \mathbb{Z}$.

In 1862 J. Liouville (cf. [D2, p. 23]) proved the following result.

Theorem B. Let a, b, c be positive integers with $a \leq b \leq c$. Then every $n \in \mathbb{N}$ can be written as $at_x + bt_y + ct_z$ with $x, y, z \in \mathbb{Z}$, if and only if (a, b, c) is among the following vectors:

$$(1,1,1), (1,1,2), (1,1,4), (1,1,5), (1,2,2), (1,2,3), (1,2,4).$$

Now we turn to representations of natural numbers by mixed sums of squares (of integers) and triangular numbers.

Let $n \in \mathbb{N}$. By the Gauss-Legendre theorem, 8n+1 is a sum of three squares. It follows that $8n+1=(2x)^2+(2y)^2+(2z+1)^2$ for some $x,y,z\in\mathbb{Z}$ with $x\equiv y\pmod 2$; this yields the representation

$$n = \frac{x^2 + y^2}{2} + t_z = \left(\frac{x+y}{2}\right)^2 + \left(\frac{x-y}{2}\right)^2 + t_z$$

as observed by Euler. According to Dickson [D2, p. 24], E. Lionnet stated, and V. A. Lebesgue [L] and M. S. Réalis [Re] proved that n can also be written in the form $x^2 + t_y + t_z$ with $x, y, z \in \mathbb{Z}$. Quite recently, this was reproved by H. M. Farkas [F] via the theory of theta functions.

Using the theory of ternary quadratic forms, in 1939 B. W. Jones and G. Pall [JP, Theorem 6] proved that for any $n \in \mathbb{N}$ we have $8n + 1 = ax^2 + by^2 + cz^2$ for some $x, y, z \in \mathbb{Z}$ if the vector (a, b, c) belongs to the set

$$\{(1,1,16), (1,4,16), (1,16,16), (1,2,32), (1,8,32), (1,8,64)\}.$$

As $(2z+1)^2 = 8t_z + 1$, the result of Jones and Pall implies that each $n \in \mathbb{N}$ can be written in any of the following three forms with $x, y, z \in \mathbb{Z}$:

$$2x^2 + 2y^2 + t_z = (x+y)^2 + (x-y)^2 + t_z, \ x^2 + 4y^2 + t_z, \ x^2 + 8y^2 + t_z.$$

In this paper we establish the following result by means of q-series.

Theorem 1. (i) Any $n \in \mathbb{N}$ is a sum of an even square and two triangular numbers. Moreover, if n/2 is not a triangular number then

$$|\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \nmid x\}|$$

=|\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \mid x\}|. \tag{1}

(ii) If $n \in \mathbb{N}$ is not a triangular number, then

$$|\{(x,y,z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \not\equiv y \pmod{2}\}|$$

=|\{(x,y,z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \equiv y \pmod{2}\}| > 0. (2)

(iii) A positive integer n is a sum of an odd square, an even square and a triangular number, unless it is a triangular number t_m (m > 0) for which all prime divisors of 2m + 1 are congruent to $1 \mod 4$ and hence $t_m = x^2 + x^2 + t_z$ for some integers x > 0 and z with $x \equiv m/2 \pmod{2}$.

Remark. Note that $t_2 = 1^2 + 1^2 + t_1$ but we cannot write $t_2 = 3$ as a sum of an odd square, an even square and a triangular number.

Here are two more theorems of this paper.

Theorem 2. Let a, b, c be positive integers with $a \leq b$. Suppose that every $n \in \mathbb{N}$ can be written as $ax^2 + by^2 + ct_z$ with $x, y, z \in \mathbb{Z}$. Then (a, b, c) is among the following vectors:

$$(1,1,1), (1,1,2), (1,2,1), (1,2,2), (1,2,4), (1,3,1), (1,4,1), (1,4,2), (1,8,1), (2,2,1).$$

Theorem 3. Let a, b, c be positive integers with $b \ge c$. Suppose that every $n \in \mathbb{N}$ can be written as $ax^2 + bt_y + ct_z$ with $x, y, z \in \mathbb{Z}$. Then (a, b, c) is among the following vectors:

$$(1,1,1), (1,2,1), (1,2,2), (1,3,1), (1,4,1), (1,4,2), (1,5,2),$$

 $(1,6,1), (1,8,1), (2,1,1), (2,2,1), (2,4,1), (3,2,1), (4,1,1), (4,2,1).$

Theorem 1 and Theorems 2–3 will be proved in Sections 2 and 3 respectively. In Section 4, we will pose three conjectures and discuss the converses of Theorems 2 and 3.

2. Proof of Theorem 1

Given two integer-valued quadratic polynomials f(x, y, z) and g(x, y, z), by $f(x, y, z) \sim g(x, y, z)$ we mean

$$\{f(x,y,z): x,y,z \in \mathbb{Z}\} = \{g(x,y,z): x,y,z \in \mathbb{Z}\}.$$

Clearly \sim is an equivalence relation on the set of all integer-valued ternary quadratic polynomials.

The following lemma is a refinement of Euler's observation $t_y + t_z \sim y^2 + 2t_z$ (cf. [D2, p. 11]).

Lemma 1. For any $n \in \mathbb{N}$ we have

$$|\{(y,z) \in \mathbb{N}^2 : t_y + t_z = n\}| = |\{(y,z) \in \mathbb{Z} \times \mathbb{N} : y^2 + 2t_z = n\}|.$$
 (3)

Proof. Note that $t_{-y-1} = t_y$. Thus

$$\begin{aligned} &|\{(y,z)\in\mathbb{N}^2:\,t_y+t_z=n\}|=\frac{1}{4}|\{(y,z)\in\mathbb{Z}^2:\,t_y+t_z=n\}|\\ &=\frac{1}{4}|\{(y,z)\in\mathbb{Z}^2:\,4n+1=(y+z+1)^2+(y-z)^2\}|\\ &=\frac{1}{4}|\{(x_1,x_2)\in\mathbb{Z}^2:\,4n+1=x_1^2+x_2^2\text{ and }x_1\not\equiv x_2\pmod{2}\}|\\ &=\frac{2}{4}|\{(y,z)\in\mathbb{Z}^2:\,4n+1=(2y)^2+(2z+1)^2\}|\\ &=\frac{1}{2}|\{(y,z)\in\mathbb{Z}^2:\,n=y^2+2t_z\}|=|\{(y,z)\in\mathbb{Z}\times\mathbb{N}:\,n=y^2+2t_z\}|.\end{aligned}$$

This concludes the proof. \Box

Lemma 1 is actually equivalent to the following observation of Ramanujan (cf. Entry 25(iv) of [B, p. 40]): $\psi(q)^2 = \varphi(q)\psi(q^2)$ for |q| < 1, where

$$\varphi(q) = \sum_{n=-\infty}^{\infty} q^{n^2} \text{ and } \psi(q) = \sum_{n=0}^{\infty} q^{t_n}.$$
(4)

Let $n \in \mathbb{N}$ and define

$$r(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n\}|,$$
 (5)

$$r_0(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \mid x\}|,$$
 (6)

$$r_1(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \nmid x\}|.$$
 (7)

Clearly $r_0(n) + r_1(n) = r(n)$. In the following lemma we investigate the difference $r_0(n) - r_1(n)$.

Lemma 2. For m = 0, 1, 2, ... we have

$$r_0(2t_m) - r_1(2t_m) = (-1)^m (2m+1). (8)$$

Also, $r_0(n) = r_1(n)$ if $n \in \mathbb{N}$ is not a triangular number times 2.

Proof. Let |q| < 1. Recall the following three known identities implied by Jacobi's triple product identity (cf. [AAR, pp. 496–501]):

$$\varphi(-q) = \prod_{n=1}^{\infty} (1 - q^{2n-1})^2 (1 - q^{2n}) \quad \text{(Gauss)},$$

$$\psi(q) = \prod_{n=1}^{\infty} \frac{1 - q^{2n}}{1 - q^{2n-1}} \quad \text{(Gauss)},$$

$$\prod_{n=1}^{\infty} (1 - q^n)^3 = \sum_{n=0}^{\infty} (-1)^n (2n+1) q^{t_n} \quad \text{(Jacobi)}.$$

Observe that

$$\sum_{n=0}^{\infty} (r_0(n) - r_1(n))q^n$$

$$= \left(\sum_{x=-\infty}^{\infty} (-1)^x q^{x^2}\right) \left(\sum_{y=0}^{\infty} q^{t_y}\right) \left(\sum_{z=0}^{\infty} q^{t_z}\right) = \varphi(-q)\psi(q)^2$$

$$= \left(\prod_{n=1}^{\infty} (1 - q^{2n-1})^2 (1 - q^{2n})\right) \left(\prod_{n=1}^{\infty} \frac{1 - q^{2n}}{1 - q^{2n-1}}\right)^2$$

$$= \prod_{n=1}^{\infty} (1 - q^{2n})^3 = \sum_{m=0}^{\infty} (-1)^m (2m+1)(q^2)^{t_m}.$$

Comparing the coefficients of q^n on both sides, we obtain the desired result. \square

The following result was discovered by Hurwitz in 1907 (cf. [D2, p. 271]); an extension was established in [HS] via the theory of modular forms of half integer weight.

Lemma 3. Let n > 0 be an odd integer, and let p_1, \ldots, p_r be all the distinct prime divisors of n congruent to n mod n. Write $n = n_0 \prod_{0 < i \le r} p_i^{\alpha_i}$, where $n_0, \alpha_1, \ldots, \alpha_r$ are positive integers and n_0 has no prime divisors congruent to n mod n. Then

$$|\{(x,y,z) \in \mathbb{Z}^3: \ x^2 + y^2 + z^2 = n^2\}| = 6n_0 \prod_{0 < i \le r} \left(p_i^{\alpha_i} + 2 \frac{p_i^{\alpha_i} - 1}{p_i - 1} \right).$$
 (9)

Proof. We deduce (9) in a new way and use some standard notations in number theory.

By (4.8) and (4.10) of [G],

$$|\{(x, y, z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = n^2\}|$$

$$= \sum_{d|n} \frac{24}{\pi} d \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{-4d^2}{m}\right)$$

$$= \frac{24}{\pi} \sum_{d|n} d \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{2k-1} \sum_{c|\gcd(2k-1,d)} \mu(c)$$

$$= \frac{24}{\pi} \sum_{d|n} d \sum_{c|d} \mu(c) \sum_{k=1}^{\infty} \frac{(-1)^{((2k-1)c-1)/2}}{(2k-1)c}$$

$$= \frac{24}{\pi} \sum_{d|n} d \sum_{c|d} \frac{\mu(c)}{c} (-1)^{(c-1)/2} \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{2k-1}$$

$$= 6 \sum_{c|n} (-1)^{(c-1)/2} \frac{\mu(c)}{c} \sum_{q|\frac{n}{c}} cq = 6 \sum_{c|n} (-1)^{(c-1)/2} \mu(c) \sigma\left(\frac{n}{c}\right)$$

Z. W. SUN

and hence

$$\frac{1}{6} |\{(x,y,z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = n^2\}|
= \sum_{d_0|n_0} \sum_{d_1|p_1^{\alpha_1}, \dots, d_r|p_r^{\alpha_r}} \left(\frac{-1}{d_0 d_1 \cdots d_r}\right) \mu(d_0 d_1 \cdots d_r) \sigma\left(\frac{n_0}{d_0} \prod_{0 < i \leqslant r} \frac{p_i^{\alpha_i}}{d_i}\right)
= \sum_{d_0|n_0} \left(\frac{-1}{d_0}\right) \mu(d_0) \sigma\left(\frac{n_0}{d_0}\right) \times \prod_{0 < i \leqslant r} \sum_{d_i|p_i^{\alpha_i}} \left(\frac{-1}{d_i}\right) \mu(d_i) \sigma\left(\frac{p_i^{\alpha_i}}{d_i}\right)
= \sum_{d_0|n_0} \mu(d_0) \sigma\left(\frac{n_0}{d_0}\right) \times \prod_{0 < i \leqslant r} \left(\sigma(p_i^{\alpha_i}) + \left(\frac{-1}{p_i}\right) \mu(p_i) \sigma(p_i^{\alpha_i-1})\right)
= n_0 \prod_{0 < i \leqslant r} \left(p_i^{\alpha_i} + 2\sigma(p_i^{\alpha_i-1})\right) = n_0 \prod_{0 < i \leqslant r} \left(p_i^{\alpha_i} + 2\frac{p_i^{\alpha_i} - 1}{p_i - 1}\right).$$

This completes the proof. \Box

Proof of Theorem 1. (i) By the Gauss-Legendre theorem, 4n+1 is a sum of three squares and hence $4n+1=(2x)^2+(2y)^2+(2z+1)^2$ (i.e., $n=x^2+y^2+2t_z$) for some $x,y,z\in\mathbb{Z}$. Combining this with Lemma 1 we obtain a simple proof of the known result

$$r(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n\}| > 0.$$

Recall that $r_0(n) + r_1(n) = r(n)$. If n/2 is not a triangular number, then $r_0(n) = r_1(n) = r(n)/2 > 0$ by Lemma 2. If $n = 2t_m$ for some $m \in \mathbb{N}$, then we also have $r_0(n) > 0$ since $n = 0^2 + t_m + t_m$.

(ii) Note that

$$n = x^2 + y^2 + t_z \iff 2n = 2(x^2 + y^2) + 2t_z = (x+y)^2 + (x-y)^2 + 2t_z.$$

From this and Lemma 1, we get

$$\begin{aligned} &|\{(x,y,z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : \ x^2 + y^2 + t_z = n\}| \\ &= &|\{(x,y,z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : \ x^2 + y^2 + 2t_z = 2n\}| \\ &= &|\{(x,y,z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : \ x^2 + t_y + t_z = 2n\}| = r(2n) > 0; \end{aligned}$$

in the language of generating functions, it says that

$$\varphi(q)\psi(q)^{2} + \varphi(-q)\psi(-q)^{2} = 2\varphi(q^{2})^{2}\psi(q^{2}).$$

Similarly,

$$|\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \equiv y \pmod{2}\}|$$

$$=|\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + 2t_z = 2n \text{ and } 2 \mid x\}|$$

$$=|\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = 2n \text{ and } 2 \mid x\}| = r_0(2n) > 0$$

and

$$|\{(x,y,z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \not\equiv y \pmod{2}\}| = r_1(2n).$$
(10)

If n is not a triangular number, then $r_0(2n) = r_1(2n) = r(2n)/2 > 0$ by Lemma 2, and hence (2) follows from the above.

(iii) By Theorem 1(ii), if n is not a triangular number then $n=x^2+y^2+t_z$ for some $x,y,z\in\mathbb{Z}$ with $2\mid x$ and $2\nmid y$.

Now assume that $n = t_m$ (m > 0) is not a sum of an odd square, an even square and a triangular number. Then $r_1(2t_m) = 0$ by (10). In view of (ii) and (8),

$$|\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = t_m\}|$$

= $r_0(2t_m) + r_1(2t_m) = r_0(2t_m) - r_1(2t_m) = (-1)^m (2m+1).$

Therefore

$$(-1)^{m}(2m+1) = \frac{1}{2}|\{(x,y,z) \in \mathbb{Z}^{3}: x^{2} + y^{2} + t_{z} = t_{m}\}|$$

$$= \frac{1}{2}|\{(x,y,z) \in \mathbb{Z}^{3}: 2(2x)^{2} + 2(2y)^{2} + (2z+1)^{2} = 8t_{m} + 1\}|$$

$$= \frac{1}{2}|\{(x,y,z) \in \mathbb{Z}^{3}: (2x+2y)^{2} + (2x-2y)^{2} + (2z+1)^{2} = 8t_{m} + 1\}|$$

$$= \frac{1}{2}|\{(x_{1},y_{1},z) \in \mathbb{Z}^{3}: 4(x_{1}^{2} + y_{1}^{2}) + (2z+1)^{2} = (2m+1)^{2}\}|$$

$$= \frac{1}{6}|\{(x,y,z) \in \mathbb{Z}^{3}: x^{2} + y^{2} + z^{2} = (2m+1)^{2}\}|.$$

Since

$$(-1)^m 6(2m+1) = |\{(x,y,z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = (2m+1)^2\}| \geqslant 6(2m+1),$$

by Lemma 3 the odd number 2m+1 cannot have a prime divisor congruent to 3 mod 4. So all the prime divisors of 2m+1 are congruent to 1 mod 4, and hence

$$|\{(x,y) \in \mathbb{N}^2: x > 0 \text{ and } x^2 + y^2 = 2m + 1\}| = \sum_{d|2m+1} 1 > 1$$

by Proposition 17.6.1 of [IR, p. 279]. Thus 2m+1 is a sum of two squares of positive integers. Choose positive integers x and y such that $x^2 + y^2 = 2m+1$ with $2 \mid x$ and $2 \nmid y$. Then

$$8t_m + 1 = (2m+1)^2 = (x^2 - y^2)^2 + 4x^2y^2 = 8t_{(x^2 - y^2 - 1)/2} + 1 + 16\left(\frac{x}{2}\right)^2y^2$$

and hence

$$t_m = t_{(x^2 - y^2 - 1)/2} + \left(\frac{x}{2}y\right)^2 + \left(\frac{x}{2}y\right)^2.$$

As $x^2 = 2m + 1 - y^2 \equiv 2m \pmod{8}$, m is even and

$$\frac{m}{2} \equiv \left(\frac{x}{2}\right)^2 \equiv \frac{x}{2} \equiv \frac{x}{2}y \pmod{2}.$$

We are done. \square

3. Proofs of Theorems 2 and 3

Proof of Theorem 2. We distinguish four cases.

Case 1. a = c = 1. Write $8 = x_0^2 + by_0^2 + t_{z_0}$ with $x_0, y_0, z_0 \in \mathbb{Z}$, then $y_0 \neq 0$ and hence $8 \geqslant b$. Since $x^2 + 5y^2 + t_z \neq 13$, $x^2 + 6y^2 + t_z \neq 47$ and $x^2 + 7y^2 + t_z \neq 20$, we must have $b \in \{1, 2, 3, 4, 8\}$.

Case 2. a=1 and c=2. Write $5=x_0^2+by_0^2+2t_{z_0}$ with $x_0,y_0,z_0\in\mathbb{Z}$. Then $y_0\neq 0$ and hence $5\geqslant b$. Observe that $x^2+3y^2+2t_z\neq 8$ and $x^2+5y^2+2t_z\neq 19$. Therefore $b\in\{1,2,4\}$.

Case 3. a=1 and $c\geqslant 3$. Since $2=x^2+by^2+ct_z$ for some $x,y,z\in\mathbb{Z}$, we must have $b\leqslant 2$. If b=1, then there are $x_0,y_0,z_0\in\mathbb{Z}$ such that $3=x_0^2+y_0^2+ct_{z_0}\geqslant c$ and hence c=3. But $x^2+y^2+3t_z\neq 6$, therefore b=2. For some $x,y,z\in\mathbb{Z}$ we have $5=x^2+2y^2+ct_z\geqslant c$. Since $x^2+2y^2+3t_z\neq 23$ and $x^2+2y^2+5t_z\neq 10$, c must be 4.

Case 4. a > 1. As $b \ge a \ge 2$ and $ax^2 + by^2 + ct_z = 1$ for some $x, y, z \in \mathbb{Z}$, we must have c = 1. If a > 2, then $ax^2 + by^2 + t_z \ne 2$. Thus a = 2. For some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $4 = 2x_0^2 + by_0^2 + t_{z_0} \ge b$. Note that $2x^2 + 3y^2 + t_z \ne 7$ and $2x^2 + 4y^2 + t_z \ne 20$. Therefore b = 2.

In view of the above, Theorem 2 has been proven. \Box

Proof of Theorem 3. Let us first consider the case c>1. Since $1=ax^2+bt_y+ct_z$ for some $x,y,z\in\mathbb{Z}$, we must have a=1. Clearly $x^2+bt_y+ct_z\neq 2$ if $c\geqslant 3$. So c=2. For some $x_0,y_0,z_0\in\mathbb{Z}$ we have $5=x_0^2+bt_{y_0}+2t_{z_0}\geqslant b$. It is easy to check that $x^2+3t_y+2t_z\neq 8$. Therefore $b\in\{2,4,5\}$.

Below we assume that c=1. If a and b are both greater than 2, then $ax^2 + bt_y + t_z \neq 2$. So $a \leq 2$ or $b \leq 2$.

Case 1. a = 1. For some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $8 = x_0^2 + bt_{y_0} + t_{z_0} \ge b$. Note that $x^2 + 5t_y + t_z \ne 13$ and $x^2 + 7t_y + t_z \ne 20$. So $b \in \{1, 2, 3, 4, 6, 8\}$.

Case 2. a = 2. For some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $4 = 2x_0^2 + bt_{y_0} + t_{z_0} \ge b$. Thus $b \in \{1, 2, 4\}$ since $2x^2 + 3t_y + t_z \ne 7$.

Case 3. a > 2. In this case $b \le 2$. If b = 1, then for some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $5 = ax_0^2 + t_{y_0} + t_{z_0} \ge a$, and hence a = 4 since $3x^2 + t_y + t_z \ne 8$ and $5x^2 + t_y + t_z \ne 19$. If b = 2, then for some $x, y, z \in \mathbb{Z}$ we have $4 = ax^2 + 2t_y + t_z \ge a$ and so $a \in \{3, 4\}$.

The proof of Theorem 3 is now complete. \square

4. Some conjectures and related discussion

In this section we raise three related conjectures.

Conjecture 1. Any positive integer n is a sum of a square, an odd square and a triangular number. In other words, each natural number can be written in the form $x^2 + 8t_y + t_z$ with $x, y, z \in \mathbb{Z}$.

We have verified Conjecture 1 for $n \leq 15,000$. By Theorem 1(iii), Conjecture 1 is valid when $n \neq t_4, t_8, t_{12}, \ldots$

Conjecture 2. Each $n \in \mathbb{N}$ can be written in any of the following forms with $x, y, z \in \mathbb{Z}$:

$$x^{2} + 3y^{2} + t_{z}$$
, $x^{2} + 3t_{y} + t_{z}$, $x^{2} + 6t_{y} + t_{z}$, $3x^{2} + 2t_{y} + t_{z}$, $4x^{2} + 2t_{y} + t_{z}$.

Conjecture 3. Every $n \in \mathbb{N}$ can be written in the form $x^2 + 2y^2 + 3t_z$ (with $x, y, z \in \mathbb{Z}$) except n = 23, in the form $x^2 + 5y^2 + 2t_z$ (or the equivalent form $5x^2 + t_y + t_z$) except n = 19, in the form $x^2 + 6y^2 + t_z$ except n = 47, and in the form $2x^2 + 4y^2 + t_z$ except n = 20.

Both Conjectures 2 and 3 have been verified for $n \leq 10,000$.

The second statement in Conjecture 3 is related to an assertion of Ramanujan confirmed by Dickson [D1] which states that even natural numbers not of the form $4^k(16l+6)$ (with $k,l \in \mathbb{N}$) can be written as $x^2 + y^2 + 10z^2$ with $x, y, z \in \mathbb{Z}$. Observe that

$$n = x^2 + 5y^2 + 2t_z \text{ for some } x, y, z \in \mathbb{Z}$$

$$\iff 4n + 1 = x^2 + 5y^2 + z^2 \text{ for some } x, y, z \in \mathbb{Z} \text{ with } 2 \nmid z$$

$$\iff 8n + 2 = 2(x^2 + y^2) + 10z^2 = (x + y)^2 + (x - y)^2 + 10z^2$$
for some $x, y, z \in \mathbb{Z} \text{ with } 2 \nmid y$

$$\iff 8n + 2 = x^2 + y^2 + 10z^2 \text{ for some } x, y, z \in \mathbb{Z} \text{ with } x \not\equiv y \text{ (mod 4)}.$$

Below we reduce the converses of Theorems 2 and 3 to Conjectures 1 and 2. For convenience, we call a ternary quadratic polynomial f(x, y, z) essential if $\{f(x, y, z) : x, y, z \in \mathbb{Z}\} = \mathbb{N}$. (Actually, in 1748 Goldbach (cf. [D2, p. 11]) already stated that $x^2 + y^2 + 2t_z$, $x^2 + 2y^2 + t_z$, $x^2 + 2y^2 + 2t_z$ and $2x^2 + 2t_y + t_z$ are essential.)

Step I. We show that the 10 quadratic polynomials listed in Theorem 2 are essential except for the form $x^2 + 3y^2 + t_z$ appearing in Conjecture 2.

As $4x^2+y^2+2t_z\sim 4x^2+t_y+t_z$, the form $x^2+(2y)^2+2t_z$ is essential by Theorem 1(i). Both $x^2+(2y)^2+t_z$ and $2x^2+2y^2+t_z=(x+y)^2+(x-y)^2+t_z$ are essential by Theorem 1(ii) and the trivial fact $t_z=0^2+0^2+t_z$. We

10 Z. W. SUN

have pointed out in Section 1 that $x^2 + 2(2y)^2 + t_z$ is essential by [JP, Theorem 6], and we don't have an easy proof of this deep result.

Since

$$x^{2} + 2y^{2} + 4t_{z} \sim x^{2} + 2(t_{y} + t_{z}) \sim t_{x} + t_{y} + 2t_{z},$$

the form $x^2 + 2y^2 + 4t_z$ is essential by Theorem B (of Liouville). By the Gauss-Legendre theorem, for each $n \in \mathbb{N}$ we can write $8n + 2 = (4x)^2 + (2y+1)^2 + (2z+1)^2$ (i.e., $n = 2x^2 + t_y + t_z$) with $x, y, z \in \mathbb{Z}$. Thus the form $x^2 + 2y^2 + 2t_z$ is essential since $2x^2 + y^2 + 2t_z \sim 2x^2 + t_y + t_z$.

Step II. We analyze the 15 quadratic polynomials listed in Theorem 3.

By Theorem 1(i), $(2x)^2 + t_y + t_z$ and $x^2 + t_y + t_z$ are essential. Since

$$x^{2} + 2t_{y} + t_{z} \sim t_{x} + t_{y} + t_{z},$$

$$x^{2} + 2t_{y} + 2t_{z} \sim t_{x} + t_{y} + 2t_{z},$$

$$x^{2} + 4t_{y} + 2t_{z} \sim t_{x} + 4t_{y} + t_{z},$$

$$x^{2} + 5t_{y} + 2t_{z} \sim t_{x} + 5t_{y} + t_{z},$$

$$2x^{2} + 4t_{y} + t_{z} \sim 2t_{x} + 2t_{y} + t_{z},$$

the forms

$$x^{2} + 2t_{y} + t_{z}$$
, $x^{2} + 2t_{y} + 2t_{z}$, $x^{2} + 4t_{y} + 2t_{z}$, $x^{2} + 5t_{y} + 2t_{z}$, $2x^{2} + 4t_{y} + t_{z}$

are all essential by Liouville's theorem. For $n \in \mathbb{N}$ we can write $2n = x^2 + 4t_y + 2t_z$ with $x, y, z \in \mathbb{Z}$, and hence $n = 2x_0^2 + 2t_y + t_z$ with $x_0 = x/2 \in \mathbb{Z}$. So the form $2x^2 + 2t_y + t_z$ is also essential.

Recall that $2x^2 + t_y + t_z$ and $2x^2 + y^2 + 2t_z$ are essential by the last two sentences of Step I. For each $n \in \mathbb{N}$ we can choose $x, y, z \in \mathbb{Z}$ such that $2n + 1 = 2x^2 + (2y + 1)^2 + 2t_z$ and hence $n = x^2 + 4t_y + t_z$. So the form $x^2 + 4t_y + t_z$ is essential.

The remaining forms listed in Theorem 3 are $x^2 + 8t_y + t_z$ and four other forms, which appear in Conjectures 1 and 2 respectively. We are done.

Acknowledgment. The author is indebted to the referee for his/her helpful comments.

Added in proof. The second conjecture in Section 4 has been confirmed by Song Guo, Hao Pan and the author.

References

- [AAR] G. E. Andrews, R. Askey and R. Roy, *Special Functions*, Cambridge Univ. Press, Cambridge, 1999.
- [B] B. C. Berndt, Ramanujan's Notebooks, Part III, Springer, New York, 1991.
- [D1] L. E. Dickson, Integers represented by positive ternary quadratic forms, Bull. Amer. Math. Soc. **33** (1927), 63–70.
- [D2] L. E. Dickson, History of the Theory of Numbers, Vol. II, AMS Chelsea Publ., 1999.
- [Du] W. Duke, Some old problems and new results about quadratic forms, Notice Amer. Math. Soc. 44 (1997), 190–196.
- [F] H. M. Farkas, Sums of squares and triangular numbers, Online J. Anal. Combin.
 1 (2006), #1, 11 pp. (electronic).
- [G] E. Grosswald, Representation of Integers as Sums of Squares, Springer, New York, 1985.
- [HS] M. D. Hirschhorn and J. Sellers, On representations of a number as a sum of three squares, Discrete Math. 199 (1999), 85–101.
- [IR] K. Ireland and M. Rosen, A Classical Introduction to Modern Number Theory, 2nd ed., Grad. Texts in Math. 84, Springer, New York, 1990.
- [JP] B. W. Jones and G. Pall, Regular and semi-regular positive ternary quadratic forms, Acta Math. **70** (1939), 165–191.
- [L] V. A. Lebesque, Questions 1059,1060,1061 (Lionnet), Nouv. Ann. Math. 11 (1872), 516–519.
- [N] M. B. Nathanson, Additive Number Theory: The Classical Bases, Grad. Texts in Math. 164, Springer, New York, 1996.
- [P] L. Panaitopol, On the representation of natural numbers as sums of squares, Amer. Math. Monthly 112 (2005), 168–171.
- [Ra] S. Ramanujan, On the expression of a number in the form $ax^2 + by^2 + cz^2 + du^2$, in: Collected Papers of Srinivasa Ramanujan, Cambridge Univ. Press, 1927, 169–178.
- [Re] M. S. Réalis, Scolies pour un théoreme d'arithmétique, Nouv. Ann. Math. 12 (1873), 212–217.

Department of Mathematics
Nanjing University
Nanjing 210093
People's Republic of China
zwsun@nju.edu.cn

http://math.nju.edu.cn/~zwsun