ON THE NONEXISTENCE OF FAT PARTIALLY HYPERBOLIC HORSESHOES

JOSE F. ALVES AND VILTON PINHEIRO ´

ABSTRACT. We show that there are no partially hyperbolic horseshoes with positive Lebesgue measure for diffeomorphisms whose class of differentiability is higher than 1. This generalizes a classical result by Bowen for uniformly hyperbolic horseshoes.

CONTENTS

1. Introduction

Let M be a Riemannian manifold. We use Leb to denote a normalized volume form defined on the Borel sets of M that we call Lebesgue measure. Given a submanifold $\gamma \subset M$ we use Leb_{γ} to denote the measure on γ induced by the restriction of the Riemannian structure to γ .

Let $f : M \to M$ be a C^1 diffeomorphism, and let $\Lambda \subset M$ be a compact *invariant set*, i.e. $f(\Lambda) \subset \Lambda$. We say that Λ is a *hyperbolic set* if there is a Df-invariant splitting $T_\Lambda M = E^s \oplus E^u$ of the tangent bundle restricted to Λ and a constant $\lambda < 1$ such that (for some choice of a Riemannian metric on M) for every $x \in \Lambda$

$$
||Df||E_x^s|| < \lambda \quad \text{and} \quad ||Df^{-1}||E_x^u|| < \lambda.
$$

We say that an embedded disk $\gamma \subset M$ is an unstable manifold, or an unstable disk, if $dist(f^{-n}(x), f^{-n}(y)) \to 0$ exponentially fast as $n \to \infty$, for every $x, y \in \gamma$. Similarly, γ is called a *stable manifold*, or a *stable disk*, if $dist(f^n(x), f^n(y)) \to 0$ exponentially fast as $n \to \infty$, for every $x, y \in \gamma$. It is well-known that every point in a hyperbolic set possesses a local stable

Date: November 21, 2018.

Work carried out at the Federal University of Bahia. JFA is partially supported by CMUP and by a grant of FCT. VP is partially supported by PADCT/CNPq.

manifold $W^s_{loc}(x)$ and a local unstable manifold $W^u_{loc}(x)$ which are disks tangent to $\overline{E_x^s}$ and E_x^u at x respectively.

A hyperbolic set Λ is said to be a *horseshoe* if local stable and local unstable manifolds through points in Λ intersect Λ in a Cantor set. Horseshoes were introduced by Smale and appear naturally when one unfolds a homoclinic tangency associated to some hyperbolic periodic point of saddle type. It follows from [\[4,](#page-9-1) Theorem 4.11] that a $C^{1+\alpha}$ diffeomorphism cannot have a fat hyperbolic horseshoe, i.e. a hyperbolic horseshoe Λ with $Leb(\Lambda) > 0$; actually the result in [\[4\]](#page-9-1) is proved for basic sets. Nevertheless, here we obtain a generalization of that result to a much more general situation. Let us remark that fat hyperbolic horseshoes exist for $C¹$ diffeomorphisms, as shown in [\[3\]](#page-9-2).

We say that a compact invariant set Λ has a *dominated splitting* if there exists a continuous Df-invariant splitting $T_\Lambda M = E^{cs} \oplus E^{cu}$ of the tangent bundle restricted to Λ , and a constant $0 < \lambda < 1$ such that (for some choice of a Riemannian metric on M) for every $x \in \Lambda$

$$
||Df||E_x^{cs}|| \cdot ||Df^{-1}||E_{f(x)}^{cu}|| \le \lambda.
$$

We call E^{cs} the *centre-stable bundle* and E^{cu} the *centre-unstable bundle*. We say that f is non-uniformly expanding along the centre-unstable direction for $x \in \Lambda$ if

$$
\liminf_{n \to +\infty} \frac{1}{n} \sum_{j=1}^{n} \log \|Df^{-1} \| E_{f^j(x)}^{cu} \| < 0. \tag{NUE}
$$

Condition [NUE](#page-1-0) means that the derivative has expanding behavior in average over the orbit of x. This implies that x has $\dim(E^{cu})$ positive Lypaunov exponents in the E_x^{cu} direction. As shown in [\[2,](#page-9-3) Theorem C], if condition [NUE](#page-1-0) holds for every point in a compact invariant set Λ , then E^{cu} is necessarily uniformly expanding in Λ , i.e. there is $0 < \lambda < 1$ such that

$$
||Df^{-1} | E_{f(x)}^{cu} || \le \lambda, \text{ for every } x \in \Lambda.
$$

A class of diffeomorphisms with a dominated splitting $TM = E^{cs} \oplus E^{cu}$ for which [NUE](#page-1-0) holds Lebesgue almost everywhere in M and E^{cu} is not uniformly expanding can be found in [\[1,](#page-9-4) Appendix A].

Theorem A. Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism and let $\Lambda \subset M$ have a dominated splitting. If there is $H \subset \Lambda$ with $\text{Leb}(H) > 0$ such that [NUE](#page-1-0) holds for every $x \in H$, then Λ contains some local unstable disk.

We say that a compact invariant set Λ is partially hyperbolic if it has a dominated splitting $T_\Lambda M = E^{cs} \oplus E^{cu}$ for which E^{cs} is uniformly contracting or E^{cu} is uniformly expanding, meaning that there is $0 < \lambda < 1$ such that $0 < \lambda < 1$ such that $||Df||E_x^{cs}|| \leq \lambda$ for every $x \in \Lambda$, or $||Df^{-1}||E_{f(x)}^{cu}|| \leq \lambda$ for every $x \in \Lambda$.

The next result is a direct consequence of Theorem [A,](#page-1-1) whenever E^{cu} is uniformly expanding. If, on the other hand, E^{cs} is uniformly contracting, then we just have to apply Theorem [A](#page-1-1) to f^{-1} .

Corollary B. Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism and let $\Lambda \subset M$ be a partially hyperbolic set with $\text{Leb}(\Lambda) > 0$.

- (1) If E^{cs} is uniformly contracting, then Λ contains a local stable disk.
- (2) If E^{cu} is uniformly expanding, then Λ contains a local unstable disk.

In particular, $C^{1+\alpha}$ diffeomorphisms cannot have partially hyperbolic horseshoes with positive Lebesgue measure. The same holds for partially hyperbolic sets intersecting a local stable or a local unstable disk in a positive Lebesgue measure subset, as Corollary [D](#page-2-1) below shows.

Theorem C. Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism and let $\Lambda \subset M$ have a dominated splitting. Assume that there is a local unstable disk γ with Leb_{$\gamma(\gamma \cap \Lambda) > 0$ such that [NUE](#page-1-0) holds for every $x \in \gamma \cap \Lambda$. Then Λ contains} some local unstable disk.

The next result is a direct consequence of Theorem [C](#page-2-2) in the case that E^{cu} is uniformly expanding, and a consequence of the theorem applied to f^{-1} in the case that E^{cs} is uniformly contracting.

Corollary D. Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism and let $\Lambda \subset M$ be a partially hyperbolic set.

- (1) If E^{cs} is uniformly contracting and there is a local stable disk γ such that $Leb_{\gamma}(\gamma \cap \Lambda) > 0$, then Λ contains a local stable disk.
- (2) If E^{cu} is uniformly expanding and there is a local unstable disk γ such that $\text{Leb}_{\gamma}(\gamma \cap \Lambda) > 0$, then Λ contains a local unstable disk.

Theorems [A](#page-1-1) and [C](#page-2-2) are in fact corollaries of a slightly more general result that we present at the beginning of Section [4.](#page-6-0)

Acknowledgement. We are grateful to M. Viana for valuable references on this topic.

2. HÖLDER CONTROL OF TANGENT DIRECTION

This section is a survey of results in $[1, Section 2]$ concerning the Hölder control of the tangent direction of submanifolds. As observed in [\[1,](#page-9-4) Remark 2.3] those results are valid for diffeomorphisms of class $C^{1+\alpha}$. In this section we only use the existence of a dominated splitting $T_\Lambda M = E^{cs} \oplus E^{cu}$. We fix continuous extensions of the two bundles E^{cs} and E^{cu} to some neighborhood U of Λ , that we denote by \tilde{E}^{cs} and \tilde{E}^{cu} . We do not require these extensions to be invariant under Df . Given $0 < a < 1$, we define the *centre-unstable* cone field $C_a^{cu} = (C_a^{cu}(x))_{x \in U}$ of width a by

$$
C_a^{cu}(x) = \{v_1 + v_2 \in \tilde{E}_x^{cs} \oplus \tilde{E}_x^{cu} \text{ such that } ||v_1|| \le a||v_2||\}.
$$
 (1)

We define the *centre-stable cone field* $C_a^{cs} = (C_a^{cs}(x))_{x \in U}$ of width a in a similar way, just reversing the roles of the subbundles in [\(1\)](#page-2-3). We fix $a > 0$ and U small enough so that, up to slightly increasing $\lambda < 1$, the domination condition remains valid for any pair of vectors in the two cone fields:

$$
||Df(x)v^{cs}|| \cdot ||Df^{-1}(f(x))v^{cu}|| \leq \lambda ||v^{cs}|| \, ||v^{cu}||
$$

for every $v^{cs} \in C_a^{cs}(x)$, $v^{cu} \in C_a^{cu}(f(x))$, and any point $x \in U \cap f^{-1}(U)$. Note that the centre-unstable cone field is positively invariant:

 $Df(x)C_a^{cu}(x) \subset C_a^{cu}(f(x)), \quad \text{whenever } x, f(x) \in U.$

Indeed, the domination property together with the invariance of E^{cu} = $(E^{cu} | \Lambda)$ imply that

$$
Df(x)C_a^{cu}(x) \subset C_{\lambda a}^{cu}(f(x)) \subset C_a^{cu}(f(x)),
$$

for every $x \in K$. This extends to any $x \in U \cap f^{-1}(U)$ just by continuity.

We say that an embedded C^1 submanifold $N \subset U$ is tangent to the centreunstable cone field if the tangent subspace to N at each point $x \in N$ is contained in the corresponding cone $C_a^{cu}(x)$. Then $f(N)$ is also tangent to the centre-unstable cone field, if it is contained in U , by the domination property.

Our aim now is to express the notion of Hölder variation of the tangent bundle in local coordinates. We choose $\delta_0 > 0$ small enough so that the inverse of the exponential map \exp_x is defined on the δ_0 neighbourhood of every point x in U . From now on we identify this neighbourhood of x with the corresponding neighbourhood U_x of the origin in T_xN , through the local chart defined by \exp_x^{-1} . Reducing δ_0 , if necessary, we may suppose that \tilde{E}_x^{cs} is contained in the centre-stable cone $C_a^{cs}(y)$ of every $y \in U_x$. In particular, the intersection of $C_a^{cu}(y)$ with \tilde{E}_x^{cs} reduces to the zero vector. Then, the tangent space to N at y is parallel to the graph of a unique linear map $A_x(y) : T_xN \to \tilde{E}^{cs}_x$. Given constants $C > 0$ and $0 < \zeta \leq 1$, we say that the tangent bundle to N is (C, ζ) -Hölder if for every $y \in N \cap U_x$ and $x \in V_0$

$$
||A_x(y)|| \le C d_x(y)^{\zeta}, \tag{2}
$$

where $d_x(y)$ denotes the distance from x to y along $N \cap U_x$, defined as the length of the shortest curve connecting x to y inside $N \cap U_x$.

Recall that we have chosen the neighbourhood U and the cone width a sufficiently small so that the domination property remains valid for vectors in the cones $C_a^{cs}(z)$, $C_a^{cu}(z)$, and for any point z in U. Then, there exist $\lambda_1 \in (\lambda, 1)$ and $\zeta \in (0, 1]$ such that

$$
||Df(z)v^{cs}|| \cdot ||Df^{-1}(f(z))v^{cu}||^{1+\zeta} \le \lambda_1 < 1
$$
 (3)

for every norm 1 vectors $v^{cs} \in C_a^{cs}(z)$ and $v^{cu} \in C_a^{cu}(z)$, at any $z \in U$. Then, up to reducing $\delta_0 > 0$ and slightly increasing $\lambda_1 < 1$, condition [\(3\)](#page-3-0) remains true if we replace z by any $y \in U_x$, $x \in U$ (taking $\|\cdot\|$ to mean the Riemannian metric in the corresponding local chart).

We fix ζ and λ_1 as above. Given a C^1 submanifold $N \subset U$, we define

 $\kappa(N) = \inf\{C > 0 : \text{the tangent bundle of } N \text{ is } (C, \zeta) \text{-Hölder}\}.$ (4)

The next result appears in [\[1,](#page-9-4) Corollary 2.4].

Proposition 2.1. There exists $C_1 > 0$ such that, given any C^1 submanifold $N \subset U$ tangent to the centre-unstable cone field,

- (1) there exists $n_0 \geq 1$ such that $\kappa(f^n(N)) \leq C_1$ for every $n \geq n_0$ such that $f^k(N) \subset U$ for all $0 \leq k \leq n$;
- (2) if $\kappa(N) \leq C_1$, then the same is true for every iterate $f^n(N)$ such that $f^k(N) \subset U$ for all $0 \leq k \leq n$;
- (3) in particular, if N and n are as in (2) , then the functions

$$
J_k: f^k(N) \ni x \mapsto \log|\det\left(Df \mid T_x f^k(N)\right)|, \quad 0 \le k \le n,
$$

are (L, ζ) -Hölder continuous with $L > 0$ depending only on C_1 and f.

3. Hyperbolic times and bounded distortion

The following notion will allow us to derive *uniform behaviour* (expansion, distortion) from the non-uniform expansion.

Definition 3.1. Given $\sigma < 1$, we say that n is a σ -hyperbolic time for $x \in \Lambda$ if

$$
\prod_{j=n-k+1}^{n} \|Df^{-1} \mid E_{f^j(x)}^{cu} \| \le \sigma^k, \quad \text{for all } 1 \le k \le n.
$$

In particular, if *n* is a σ -hyperbolic time for x, then $Df^{-k} | E_{f^{n}(x)}^{cu}$ is a contraction for every $1 \leq k \leq n$:

$$
||Df^{-k} | E^{cu}_{f^n(x)}|| \leq \prod_{j=n-k+1}^n ||Df^{-1} | E^{cu}_{f^j(x)}|| \leq \sigma^k.
$$

Moreover, if $a > 0$ is taken sufficiently small in the definition of our cone fields, and we choose $\delta_1 > 0$ also small so that the δ_1 -neighborhood of Λ should be contained in U , then by continuity

$$
||Df^{-1}(f(y))v|| \le \frac{1}{\sqrt{\sigma}} ||Df^{-1}| E_{f(x)}^{cu} || ||v||,
$$
\n(5)

whenever $x \in \Lambda$, $dist(x, y) \leq \delta_1$, and $v \in C_a^{cu}(y)$.

Given any disk $\Delta \subset M$, we use dist_{Δ} (x, y) to denote the distance between $x, y \in \Delta$, measured along Δ . The distance from a point $x \in \Delta$ to the boundary of Δ is dist $\Delta(x, \partial \Delta) = \inf_{y \in \partial \Delta} \text{dist}_{\Delta}(x, y)$.

Lemma 3.2. Take any C^1 disk $\Delta \subset U$ of radius δ , with $0 < \delta < \delta_1$, tangent to the centre-unstable cone field. There is $n_0 \geq 1$ such that if $x \in \Delta$ with $dist_{\Delta}(x, \partial \Delta) \geq \delta/2$ and $n \geq n_0$ is a σ -hyperbolic time for x, then there is a neighborhood V_n of x in Δ such that:

- (1) f^n maps V_n diffeomorphically onto a disk of radius δ_1 around $f^n(x)$ tangent to the centre-unstable cone field;
- (2) for every $1 \leq k \leq n$ and $y, z \in V_n$,

$$
\mathrm{dist}_{f^{n-k}(V_n)}(f^{n-k}(y), f^{n-k}(z)) \le \sigma^{k/2} \mathrm{dist}_{f^n(V_n)}(f^n(y), f^n(z)).
$$

Proof. First we show that $f^{n}(\Delta)$ contains some disk of radius δ_1 around $f^{n}(x)$, as long as

$$
n > 2 \frac{\log(\delta/(2\delta_1))}{\log(\sigma)}.
$$
\n(6)

Assume that there is $y \in \partial \Delta$ with $\text{dist}_{f^n(\Delta)}(f^n(x), f^n(y)) < \delta_1$. Let η_0 be a curve of minimal length in $f^{n}(\Delta)$ connecting $f^{n}(x)$ to $f^{n}(y)$. For $0 \leq k \leq n$ we write $\eta_k = f^{n-k}(\eta_0)$. We prove by induction that length $(\eta_k) < \sigma^{k/2}\delta_1$, for $0 \leq k \leq n$. Let $1 \leq k \leq n$ and assume that

$$
length(\eta_j) < \sigma^{j/2} \delta_1, \quad \text{for } 0 \le j \le k - 1.
$$

Denote by $\dot{\eta}_0(w)$ the tangent vector to the curve η_0 at the point w. Then, by the choice of δ_1 in [\(5\)](#page-4-1) and the definition of σ -hyperbolic time,

$$
||Df^{-k}(w)\dot{\eta}_0(w)|| \leq \sigma^{-k/2} ||\dot{\eta}_0(w)|| \prod_{j=n-k+1}^n ||Df^{-1}| E_{f^j(x)}^{cu} || \leq \sigma^{k/2} ||\dot{\eta}_0(w)||.
$$

Hence,

length
$$
(\eta_k)
$$
 $\leq \sigma^{k/2}$ length $(\eta_0) < \sigma^{k/2} \delta_1$.

This completes our induction. In particular, we have $\text{length}(\eta_n) < \sigma^{n/2}\delta_1$. Note that η_n is a curve in Δ connecting x to $y \in \partial \Delta$, and so length $(\eta_n) \geq$ $\delta/2$. Thus we must have

$$
n < 2 \frac{\log(\delta/(2\delta_1))}{\log(\sigma)}.
$$

Hence $f^{n}(\Delta)$ contains some disk of radius δ_1 around $f^{n}(x)$ for n as in [\(6\)](#page-5-0).

Let now Δ_1 be the disk of radius δ_1 around $f^n(x)$ in $f^n(\Delta)$ and let $V_n = f^{-n}(\Delta_1)$, for n as in [\(6\)](#page-5-0). Take any $y, z \in V_n$ and let η_0 be a curve of minimal length in Δ_1 connecting $f^n(y)$ to $f^n(z)$. Defining $\eta_k = f^{n-k}(\eta_0)$, for $1 \leq k \leq n$, and arguing as before we inductively prove that for $1 \leq k \leq n$

length
$$
(\eta_k) \le \sigma^{k/2}
$$
 length $(\eta_0) = \sigma^{k/2}$ dist_{fⁿ(V_n)} $(fn(y), fn(z)),$

which implies that for $1 \leq k \leq n$

$$
dist_{f^{n-k}(V_n)}(f^{n-k}(y), f^{n-k}(z)) \le \sigma^{k/2} dist_{f^n(V_n)}(f^n(y), f^n(z)).
$$

This completes the proof of the lemma.

We shall sometimes refer to the sets V_n as *hyperbolic pre-balls* and to their images $f^n(V_n)$ as hyperbolic balls. Notice that the latter are indeed balls of radius δ_1 .

Corollary 3.3 (Bounded Distortion). There exists $C_2 > 1$ such that given Δ as in Lemma [3.2](#page-4-2) with $\kappa(\Delta) \leq C_1$, and given any hyperbolic pre-ball $V_n \subset \Delta$ with $n \geq n_0$, then for all $y, z \in V_n$

$$
\frac{1}{C_2} \le \frac{|\det Df^n \mid T_y \Delta|}{|\det Df^n \mid T_z \Delta|} \le C_2.
$$

$$
\qquad \qquad \Box
$$

Proof. For $0 \leq i < n$ and $y \in \Delta$, we denote $J_i(y) = |\det Df | T_{f^i(y)} f^i(\Delta)|$. Then,

$$
\log \frac{|\det Df^n | T_y \Delta|}{|\det Df^n | T_z \Delta|} = \sum_{i=0}^{n-1} \left(\log J_i(y) - \log J_i(z) \right).
$$

By Proposition [2.1,](#page-3-1) $\log J_i$ is (L, ζ) -Hölder continuous, for some uniform constant $L > 0$. Moreover, by Lemma [3.2,](#page-4-2) the sum of all dist $\Delta(f^j(y), f^j(z))^{\zeta}$ over $0 \leq j \leq n$ is bounded by $\delta_1/(1 - \sigma^{\zeta/2})$. Now it suffices to take $C_2 = \exp(L\delta_1/(1 - \sigma^{\zeta/2}))$ \Box)).

4. A LOCAL UNSTABLE DISK INSIDE Λ

Now we are able to prove Theorems [A](#page-1-1) and [C.](#page-2-2) These will be obtained as corollaries of the next result, as we shall see below.

Theorem 4.1. Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism and let $\Lambda \subset M$ have a dominated splitting. Assume that there is a disk Δ tangent to the centre-unstable cone field with $\text{Leb}_{\Delta}(\Delta \cap \Lambda) > 0$ such that [NUE](#page-1-0) holds for every $x \in \Delta \cap \Lambda$. Then Λ contains some local unstable disk.

Assume that there is $H \subset \Lambda$ with $\text{Leb}(H) > 0$ such that [NUE](#page-1-0) holds for every $x \in H$. Choosing a Leb density point of H, we laminate a neighborhood of that point into disks tangent to the centre-unstable cone field contained in U. Since the relative Lebesgue measure of the intersections of these disks with H cannot be all equal to zero, we obtain some disk Δ as in Theorem [4.1](#page-6-1) under the assumption of Theorem [A.](#page-1-1) For Theorem [C,](#page-2-2) observe that local stable manifolds are tangent to the centre-unstable spaces and these vary continuously with the points in Λ , thus being locally tangent to the centre-unstable cone field.

Let us now prove Theorem [4.1.](#page-6-1) Let Δ be a disk tangent to the centreunstable cone field intersecting Λ in a positive Leb_{Δ} subset such that [NUE](#page-1-0) holds for every $x \in \Delta \cap \Lambda$. Let $H = \Delta \cap \Lambda$. Taking a subset of H, if necessary, still with positive Leb Λ measure, we may assume that there is $c > 0$ such that for every $x \in H$

$$
\liminf_{n \to +\infty} \frac{1}{n} \sum_{j=1}^{n} \log \|Df^{-1} \| E_{f^j(x)}^{cu} \| \le -c. \tag{7}
$$

Since condition [\(7\)](#page-6-2) remains valid under iteration, by Proposition [2.1](#page-3-1) we may assume that $\kappa(\Delta) < C_1$. It is no restriction to assume that H intersects the sub-disk of Δ of radius $\delta/2$, for some $0 < \delta < \delta_1$, in a positive Leb_{Δ} subset, and we do so.

The following lemma is due to Pliss [\[6\]](#page-9-5), and a proof of it in the present form can be found in [\[1,](#page-9-4) Lemma 3.1].

Lemma 4.2 (Pliss). Given $A \ge c_2 > c_1 > 0$, let $\theta = (c_2 - c_1)/(A - c_1)$. Given any real numbers a_1, \ldots, a_N such that $a_j \leq A$ and

$$
\sum_{j=1}^{N} a_j \ge c_2 N, \quad \text{for every } 1 \le j \le N,
$$

then there are $l > \theta N$ and $1 < n_1 < \cdots < n_l \le N$ so that

$$
\sum_{j=n+1}^{n_i} a_j \ge c_1(n_i - n), \quad \text{for every } 0 \le n < n_i \text{ and } i = 1, \dots, l.
$$

Corollary 4.3. There is $\sigma > 0$ such that every $x \in H$ has infinitely many σ-hyperbolic times.

Proof. Given $x \in H$, by [\(7\)](#page-6-2) we have infinitely many values N for which

$$
\sum_{j=1}^{N} \log ||Df^{-1}| E_{f^j(x)}^{cu}|| \le -\frac{c}{2}N.
$$

Then it suffices to take $c_1 = c/2$, $c_2 = c$, $A = \sup | \log ||Df^{-1}| E^{cu} || |$, and $a_j = -\log ||Df^{-1}| E^{cu}_{f^j(x)}||$ in the previous lemma.

Note that under assumption [\(7\)](#page-6-2) we are unable to prove the existence of a positive frequency of hyperbolic times at infinity. This would be possible if we had lim sup instead of lim inf in (7) , as shown in $[1,$ Corollary 3.2. The existence of infinitely many hyperbolic times is enough for what comes next.

Lemma 4.4. There are an infinite sequence of integers $1 \leq k_1 < k_2 < \cdots$ and, for each $n \in \mathbb{N}$, a disk Δ_n of radius $\delta_1/4$ tangent to the centre-unstable cone field such that the relative Lebesgue measure of the set $f^{k_n}(H)$ in Δ_n converges to 1 as $n \to \infty$.

Proof. Let $\epsilon > 0$ be some small number. Let K be a compact subset of H and A be an open neighborhood of H in Δ such that

$$
Leb_{\Delta}(A \setminus K) < \epsilon Leb_{\Delta}(K).
$$

It follows from Corollary [4.3](#page-7-0) and Lemma [3.2](#page-4-2) that we can choose for each $x \in K$ a σ -hyperbolic time $n(x)$ and a hyperbolic pre-ball V_x such that $V_x \subset A$. Here V_x is the neighborhood of x associated to the hyperbolic time $n(x)$ constructed in Lemma [3.2,](#page-4-2) which is mapped diffeomorphically by $f^{n(x)}$ onto a ball $B_{\delta_1}(f^{n(x)}(x))$ of radius δ_1 around $f^{n(x)}(x)$ tangent to the centre-unstable cone field. Let $W_x \subset V_x$ be the pre-image of the ball $B_{\delta_1/4}(f^{n(x)}(x))$ of radius $\delta_1/4$ under this diffeomorphism.

By compactness we have $K \subset W_{x_1} \cup ... \cup W_{x_s}$ for some $x_1, ..., x_m \in K$. Writing

$$
\{n_1, ..., n_s\} = \{n(x_1), ..., n(x_m)\}, \text{ with } n_1 < n_2 < ... < n_s,\tag{8}
$$

let $\mathcal{U}_1 \subset \mathbb{N}$ be a maximal set of $\{1, ..., m\}$ such that if $u \in \mathcal{U}_1$ then $n(x_u) = n_1$ and $W_{x_u} \cap W_{x_a} = \emptyset$ for all $a \in \mathcal{U}_1$ with $a \neq u$. Inductively we define \mathcal{U}_j

for $1 < j \leq s$ as follows. Suppose that \mathcal{U}_{j-1} has already been defined. Let $U_j \subset \mathbb{N}$ be a maximal set of $\{1, \ldots, m\}$ such that if $u \in U_j$ then $n(x_u) = n_j$ and $W_{x_u} \cap W_{x_a} = \emptyset$ for all $a \in \mathcal{U}_j$ with $a \neq u$, and also $W_{x_u} \cap W_{x_a} = \emptyset$ for all $a \in \mathcal{U}_1 \cup ... \cup \mathcal{U}_{j-1}$.

Let $\mathcal{U} = \mathcal{U}_1 \cup \cdots \cup \mathcal{U}_s$. By maximality, each W_{x_i} , $1 \leq i \leq m$, intersects some W_{x_u} with $u \in \mathcal{U}$ and $n(x_i) \geq n(x_u)$. Thus, given any $1 \leq i \leq m$ and taking $u \in \mathcal{U}$ such that $W_{x_i} \cap W_{x_u} \neq \emptyset$ and $n(x_i) \geq n(x_u)$, we get

$$
f^{n(x_u)}(W_{x_i}) \cap B_{\delta_1/4}(f^{n(x_u)}(x_u)) \neq \emptyset.
$$

Lemma [3.2](#page-4-2) assures that

$$
\operatorname{diam}(f^{n(x_u)}(W_{x_i})) \le \frac{\delta_1}{2}\sigma^{(n(x_i)-n(x_u))/2} \le \frac{\delta_1}{2},
$$

and so

$$
f^{n(x_u)}(W_{x_i}) \subset B_{\delta_1}(f^{n(x_u)}(x_u)).
$$

This implies that $W_{x_i} \subset V_{x_u}$. Hence $\{V_{x_u}\}_{u \in \mathcal{U}}$ is a covering of K.

It follows from Corollary [3.3](#page-5-1) that there is a uniform constant $\gamma > 0$ such

$$
_{\rm that}
$$

$$
\frac{\text{Leb}_{\Delta}(W_{x_u})}{\text{Leb}_{\Delta}(V_{x_u})} \ge \gamma, \text{ for every } u \in \mathcal{U}.
$$

Hence

$$
Leb_{\Delta} (\cup_{u \in \mathcal{U}} W_{x_u}) = \sum_{u \in \mathcal{U}} Leb_{\Delta}(W_{x_u})
$$

\n
$$
\geq \sum_{u \in \mathcal{U}} \gamma Leb_{\Delta}(V_{x_u})
$$

\n
$$
\geq \gamma Leb_{\Delta} (\cup_{u \in \mathcal{U}} V_{x_u})
$$

\n
$$
\geq \gamma Leb_{\Delta}(K).
$$

Setting

$$
\rho = \min \left\{ \frac{\text{Leb}_{\Delta}(W_{x_u} \setminus K)}{\text{Leb}_{\Delta}(W_{x_u})} : u \in \mathcal{U} \right\},\,
$$

we have

$$
\varepsilon \operatorname{Leb}_{\Delta}(K) \geq \operatorname{Leb}_{\Delta}(A \setminus K)
$$

\n
$$
\geq \operatorname{Leb}_{\Delta}(\cup_{u \in \mathcal{U}} W_{x_u} \setminus K)
$$

\n
$$
\geq \sum_{u \in \mathcal{U}} \operatorname{Leb}_{\Delta}(W_{x_u} \setminus K)
$$

\n
$$
\geq \rho \operatorname{Leb}_{\Delta}(\cup_{u \in \mathcal{U}} W_{x_u})
$$

\n
$$
\geq \rho \gamma \operatorname{Leb}_{\Delta}(K).
$$

This implies that $\rho < \varepsilon/\gamma$. Since $\varepsilon > 0$ can be taken arbitrarily small, we may always choose W_{x_u} such that the relative Lebesgue measure of K in W_{x_u} is arbitrarily close to 1. Then, by bounded distortion, the relative Lebesgue measure of $f^{n(x_u)}(H) \supset f^{n(x_u)}(K)$ in $f^{n(x_u)}(W_{x_u})$, which is a disk of radius $\delta_1/4$ around $f^{n(x_u)}(x_u)$ tangent to centre-unstable cone field, is also arbitrarily close to 1. Observe that since points in H have infinitely many hyperbolic times, we may take the integer $n(x_u)$ arbitrarily large, as long as n_1 in [\(8\)](#page-7-1) is also taken large enough.

Proposition 4.5. There is a local unstable disk Δ_{∞} of radius $\delta_1/4$ inside Λ .

Proof. Let $(\Delta_n)_n$ be the sequence of disks given by Lemma [4.4,](#page-7-2) and consider $(x_n)_n$ the sequence of points at which these disks are centered. Up to taking subsequences, we may assume that the centers of the disks converge to some point x. Using Ascoli-Arzela, the disks converge to some disk Δ_{∞} centered at x. By construction, every point in Δ_{∞} is accumulated by some orbit of a point in $H \subset \Lambda$, and so $\Delta_{\infty} \subset \Lambda$.

Note that each Δ_n is contained in the k_n -iterate of Δ , which is a disk tangent to the centre-unstable cone field. The domination property implies that the angle between Δ_n and E^{cu} goes to zero as $n \to \infty$, uniformly on Λ. In particular, ∆[∞] is tangent to ^Ecu at every point in ∆[∞] [⊂] Λ. By Lemma [3.2,](#page-4-2) given any $k \geq 1$, then f^{-k} is a $\sigma^{k/2}$ -contraction on Δ_n for every large *n*. Passing to the limit, we get that f^{-k} is a $\sigma^{k/2}$ -contraction on Δ_{∞} for every $k \geq 1$.

In particular, we have shown that the subspace E_x^{cu} is uniformly expanding for Df. The fact that the Df-invariant splitting $T_\Lambda M = E^{cs} \oplus E^{cu}$ is a dominated splitting implies that any expansion Df may exhibit along the complementary direction E_x^{cs} is weaker than the expansion in the E_x^{cu} direction. Then, by [\[5\]](#page-9-6), there exists a unique unstable manifold $W^u_{loc}(x)$ tangent to E^{cu} and which is contracted by the negative iterates of f. Since Δ_{∞} is contracted by every f^{-k} , and all its negative iterates are tangent to centre-unstable cone field, then Δ_{∞} is contained in $W^u_{loc}(x)$.

REFERENCES

- [1] J. F. Alves, C. Bonatti, M. Viana, SRB measures for partially hyperbolic systems whose central direction is mostly expanding, Invent. Math. 140 (2000), 351-398.
- [2] J. F. Alves, V. Araújo, B. Saussol, On the uniform hyperbolicity of some nonuniformly hyperbolic systems, Proc. Amer. Math. Soc. 131, n.4 (2003), 1303-1309.
- [3] R. Bowen, A horseshoe with positive measure, Invent. Math. 29 (1975), 203-204.
- [4] R. Bowen, Equilibrium states and the ergodic theory of Axiom A diffeomorphisms, Lecture Notes in Mathematics 480 (1975), Springer.
- [5] Ya. Pesin, Families of invariant manifolds corresponding to non-zero characteristic exponents, Math. USSR Izv. 10 (1976) 1261-1302.
- [6] V. Pliss. On a conjecture due to Smale, Diff. Uravnenija 8 (1972) 262-268.

DEPARTAMENTO DE MATEMÁTICA PURA, FACULDADE DE CIÊNCIAS DO PORTO, RUA do Campo Alegre 687, 4169-007 Porto, Portugal

E-mail address: jfalves@fc.up.pt URL: http://www.fc.up.pt/cmup/jfalves

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA BAHIA, AV. ADEMAR de Barros s/n, 40170-110 Salvador, Brazil.

E-mail address: viltonj@ufba.br