The Julia set of Henon maps

John Erik Fornæss[∗] Mathematics Department, University of Michigan 530 Church Street, 2074 East Hall Ann Arbor, MI 48109-1043, USA

Abstract

In this paper we investigate the support of the unique measure of maximal entropy of complex Hénon maps, J^* . The main question is whether this set is the same as the analogue of the Julia set J.

1 Introduction

Let $H: \mathbb{C}^2 \to \mathbb{C}^2$ be an automorphism of the form $H(z, w) = (P(z) + aw, bz)$ where $P: \mathbb{C} \to \mathbb{C}$ is a polynomial of degree d at least 2 and a, b are nonzero complex constants. More generally, we define a (complex) Hénon map H : $\mathbb{C}^2 \to \mathbb{C}^2$ as $H = H_1 \circ \cdots \circ H_n$ where each $H_i = (P_i(z) + a_i w, b_i z)$ is of the above form and $n \geq 1$.

Associated to each Hénon map there is a natural invariant measure μ with compact support J^* . There is also a natural notion of Julia set, J . We recall the precise definitions in Section 2. The following is one of the basic questions in the theory of complex dynamics.

Problem 1 *Is* $J = J^*$?

It was proved in [2] that if H is uniformly hyperbolic when restricted to J then $J = J^*$. This leaves open the following interesting special case of the Problem:

Problem 2 If H is uniformly hyperbolic on J^* , is $J = J^*$?

For motivation, we recall that this question arose naturally in the author's investigation of sustainable complex H ϵ non maps, see [5] which also contains several references to the dynamics of complex Hénon maps and also includes an exposition of background material for this article. The author showed that

[∗]July 2, 2005. The author is supported by an NSF grant

complex Hénon maps which are uniformly hyperbolic on J are sustainable and that sustainable maps are uniformly hyperbolic on J^* . The last step in the characterization of sustainable complex H´enon maps was to show the equality of J and J^* , however, this used sustainability. We are left with the equivalent problem:

Problem 3 If H is uniformly hyperbolic on J^* , is H sustainable?

We also note in passing that it is an interesting open question whether there is a similar characterization of sustainable real Hénon maps.

Main Theorem 1 Let H be a complex Hénon map which is hyperbolic on J^* . If H is not volume preserving, then $J = J^*$.

In Section 2 we introduce notation and review background results. We prove the Main Theorem in Section 3. The author would like to thank Eric Bedford for valuable comments.

2 Notation and background results

We recall some standard notation for Henon maps which can be found in many sources, see for example [5]. Let H be a complex Hénon map. We denote by H^n the *n*-fold composition $H \circ \cdots \circ H$ for any positive integer $n \geq 1$. If $n < 0$ we write $H^n := (H^{-1})^{|n|}$ where H^{-1} denotes the inverse map. Also $H^0 =$ Id. We define the sets of bounded orbits and their boundaries.

 $K^+ := \{(z, w) \in \mathbb{C}^2; \{H^n(z, w)\}_{n \ge 1} \text{ is a bounded sequence}\}\$ $K^- := \{(z, w) \in \mathbb{C}^2; \{H^n(z, w)\}_{n \le -1} \text{ is a bounded sequence}\}\$ $K := K^+ \cap K^ J^+$:= ∂K^+ J^- := $\partial K^ J \; := \; J^+ \cap J^-$

We let d denote the degree of the highest order term in the polynomial mapping $H, d = \Pi_{i=1} \text{deg}(P_i)$. We define the escape functions $G^{\pm} : \mathbb{C}^2 \to \mathbb{R}$ by

$$
G^+(z, w) := \lim_{n \to \infty} \frac{\log^+ \|H^n(z, w)\|}{d^n}
$$

$$
G^-(z, w) := \lim_{n \to \infty} \frac{\log^+ \|H^n(z, w)\|}{d^n}
$$

We remark that the sets K^{\pm} are closed and the functions G^{\pm} are continuous and plurisubharmonic. Moreover G^{\pm} vanishes on K^{\pm} and is strictly positive and pluriharmonic on $\mathbb{C}^2 \setminus K^{\pm}$.

We define $\mu^{\pm} := dd^c G^{\pm}$ and $\mu := \mu^+ \wedge \mu^-$. The $(1,1)$ currents μ^{\pm} are supported on J^{\pm} and hence μ is a positive measure supported inside J. We set J^* equal to the support of μ . So we get immediately that $J^* \subset J$.

We recall the notion of uniform hyperbolicity. Let $F : M \to M$ be a holomorphic automorphism of a complex manifold M of dimension m with a metric d. Suppose that the compact set S, $S \subset M$ is completely invariant, i.e. $F(S) = S$.

We say that F is uniformly hyperbolic on S if there exists a continuous splitting $E_x^s \oplus E_x^u = T_x, x \in S$ of complex subspaces of the tangent spaces. Moreover $F'(E_x^s) = E_{F(x)}^s$ and $F'(E_x^u) = E_{F(x)}^u$. Also there exist constants $0 < c, \lambda < 1$ so that $||(F^n)'(v)|| \leq \frac{\lambda^n}{c}$ $\frac{c^n}{c} ||v||$ for every $v \in E_x^s, x \in S, n \ge 1$ and $||(F^{n})'(v)|| \geq \frac{c}{\lambda^{n}}||v||$ for every $v \in E_{x}^{u}, x \in S, n \geq 1$.

For $p \in S$, let $W_p^s := \{x \in M; d(F^n(x), F^n(p)) \to 0, n \to \infty\}$ and For $p \in S$, let $W_p : = \{x \in M, a(P^-(x), P^-(p)) \to 0, n \to \infty\}$, $n \to \infty$ and $W_p^u := \{x \in M; d(F^n(x), F^n(p)) \to 0, n \to -\infty\}$. Then W_p^s and W_p^u are immersed complex manifolds of the same dimension as E_p^s and E_p^u respectively. Similarly, we define $W_S^s := \{x \in M; d(F^n(x), S) \to 0, n \to \infty\}$ and $W_S^u := \{x \in M; d(F^n(x), S) \to 0, n \to -\infty\},\$ the stable and unstable sets of S.

Lemma 1 [3] Let \tilde{M} be a one dimensional complex manifold in \mathbb{C}^2 . Let $M \subset \subset \tilde{M}$ be smoothly bounded. If $\mu_{|\tilde{M}}^+(\partial M) = 0$, then

$$
\frac{1}{d^n} H_*^n([M]) \to c\mu^-, c = \mu_{|\tilde{M}}^+(M).
$$

Using this Lemma and the arguments of Lemma 9.1 and Theorem 9.6 in [1], it follows that for some large n, the manifold $Hⁿ(M)$ must intersect the stable manifold of any given periodic saddle point in J^* transversely in any given neighborhood of J^* . Hence the same is true for M. [There might also be some tangential intersections.] So we obtain:

Proposition 1 Suppose that H is a complex Hénon map. Let \tilde{M} be a one dimensional complex submanifold of \mathbb{C}^2 and let $M \subset\subset \tilde{M}$ such that $\mu_{|\tilde{M}}^+(M) > 0$. Let $U \supset J^*$ be some neighborhood. Then for any saddle point p there are arbitrarily large integers so that the manifold $H^n(M)$ intersects $W^s(p)$ transversally in U.

Proposition 2 Suppose that H is a complex Hénon map. Let $p \in J \setminus J^*$. Then there are arbitrarily small discs $M \subset \tilde{M}$ through p on which $\mu_{|\tilde{M}}^+(M)$ 0. For every transverse intersection x near p between M and a stable manifold of a saddle point, the function G^- vanishes identically in a neighborhood of the intersection in the stable manifold provided $G^-(x) = 0$, so in particular if G^- vanishes identically on M, which happens if M is an unstable manifold.

Proof: If not, then the local piece of stable manifold must contain a transverse intersection y with the unstable manifold of some other saddle point. But then the α and ω limit sets of y are periodic saddles. This implies that y is a limit of saddle points, so must be in J^* , a contradiction.

It is convenient to state the same results for the inverse map.

Proposition 3 Suppose that H is a complex Hénon map. Let \tilde{M} be a one dimensional complex submanifold of \mathbb{C}^2 and let $M \subset\subset \tilde{M}$ such that $\mu_{|\tilde{M}}^-(M) > 0$. Let $U \supset J^*$ be some neighborhood. Then for any saddle point p there are arbitrarily large integers n so that the manifold $H^{-n}(M)$ intersects $W^u(p)$ transversally in U.

Proposition 4 Suppose that H is a complex Hénon map. Let $p \in J \setminus J^*$. Then there are arbitrarily small discs $M \subset \tilde{M}$ through p on which $\mu_{|\tilde{M}}^-(M)$ 0. For every transverse intersection y near p between M and an unstable manifold of a saddle point, the function G^+ vanishes identically in a neighborhood of the intersection in the unstable manifold, provided that $G^+(y) = 0$, so in particular if G^+ vanishes identically on M, which happens if M is a stable manifold.

Theorem 1 Let H be any complex Hénon map. Then J^* is equal to the closure of the collection of periodic saddle orbits.

This is in [1].

3 Proof of the Main Theorem

We start by recalling a few standard lemmas which can for example be found in [5].

Lemma 2 If H is uniformly hyperbolic on J^* , then $W_{J^*}^s = \bigcup_{p \in J^*} W_p^s$.

This is [5], Lemma 4.15.

Lemma 3 Suppose H is uniformly hyperbolic on J^* and that $p \in W^s_{J^*} \cap W^u_{J^*}$. Then $p \in J^*$.

This is [5], Lemma 4.17.

Lemma 4 If H is uniformly hyperbolic on J^* , then J^* has local product structure.

This is [5], Lemma 4.14.

We investigate some consequences of assuming that $J \setminus J^*$ is nonempty. We give first some notation. Let $p \in J^*$ and let $\Phi_p^s : \mathbb{C} \to \mathbb{C}^2$ be a parametrization of the stable manifold W_p^s of p, $\Phi_p^s(0) = p$. This exists a.e. $d\mu$ on J^* (Oseledec regular points) and if H is uniformly hyperbolic on J^* , this exists for all $p \in J^*$. Also, let $J_{p,s}^* := (\Phi_p^s)^{-1}(J^*)$.

For the unstable manifold we use the analogous notation, $q, \Psi_q^u, J_{q,u}^*$.

We next define a notion describing how $J \setminus J^*$ might be attached to J ∗ . Namely we give a name to points whose stable manifolds or unstable manifolds belong (partly) to $J \setminus J^*$. We will say that a point in J^* is stably exposed (to $J \setminus J^*$) if its stable manifold enters into $J \setminus J^*$ and similarly for unstably exposed. More precisely:

Definition 1 We say that p is stably exposed if $0 \in \mathbb{C}$ is a boundary point of a simply connected open set $V_{s,p} \subset \mathbb{C}$ such that $\partial V_{s,p} \subset J^*_{p,s}$ and $\Phi^s_p(V_{s,p}) \subset$ $J \setminus J^*$. If we instead have $\Phi_p^s(V_{s,p}) \subset K \setminus J^*$, we say that p is weakly stably exposed.

We define unstably exposed similarly, using the notation $q, W_{u,q}$.

Remark 1 When the Jacobian of H has modulus 1, W_p^s , W_q^u cannot intersect the interiors of K^{\pm} so weakly (un)stable is equivalent to (un)stable. If the modulus is $\neq 1$, either K⁺ or K⁻ has empty interior.

Theorem 2 Suppose that H is a complex Hénon map which is uniformly hyperbolic on J^* . If $J \setminus J^*$ is nonempty, then there are points $p, q \in J^*$ so that p is weakly stably exposed and q is weakly unstably exposed.

Definition 2 Suppose that H is hyperbolic on J^* . Let $p \in J^*$ and let Φ_p^s : $\mathbb{C} \to W_p^s, \Psi_p^u : \mathbb{C} \to W_p^u$ parametrize the stable and unstable manifolds of $p, \ \Phi_p^s(0) = \Psi_p^u(0) = p$. Suppose that $0 \in \mathbb{C}$ is a boundary point of simply connected components $U \subset \mathbb{C} \setminus J^*_{p,s}$ and $V \subset \mathbb{C} \setminus J^*_{p,u}$. In this case, because of local product structure, J^* is locally contained on one side of each of the laminated local hypersurfaces which are the local unstable set of $\Phi_p^s(\partial U)$ and the local stable set of $\Psi_p^u(\partial V)$ respectively. We say in this case that p is a distinguished boundary point of J^* .

Remark 2 Heuristically one can think of the local stable set as $\{(z, w) \in$ \mathbb{C}^2 ; $|w|=1$ and the local unstable set as $\{(z,w)\in\mathbb{C}^2; |z|=1\}$ and that J^* contains the distinguished boundary of the bidisc and J^* is contained in the bidisc.

Theorem 3 Suppose that H is a complex Hénon map which is uniformly hyperbolic on J^* . If $J \setminus J^*$ is nonempty, then distinguished boundary points are dense in J^* .

Proposition 5 Suppose that H is uniformly hyperbolic on J^* . Let $p \in J^*$ and let $\Phi_p^s : \mathbb{C} \to \mathbb{C}^2$ parametrize the stable manifold of p, $\Phi_p^s(0) = p$. Fix a nonempty connected component V of $\mathbb{C}\setminus J^*_{p,s}$. The function $G^+\circ \Phi^s_p \equiv 0$ on \mathbb{C} . On the other hand $(G^- \circ \Phi_p^s)_{|V}$ is either identically zero or strictly positive.

Proof: The first part is obvious. Suppose there is a point $z \in V$ so that $G^{-} \circ \Phi_{p}^{s}(z) = 0$ but that $G^{-} \circ \Phi_{p}^{s}$ is not identically zero on V. Choosing another z if necessary we can assume that there are discs $z \in \Delta \subset\subset \Delta \subset\subset V$ so that $G^- \circ \Phi_p^s(z) = 0$ but that $G^- \circ \Phi_p^s$ is not vanishing identically on Δ . Then the nonnegative function $G^- \circ \Phi_p^s$ cannot be harmonic on Δ . Let $M := \Phi_p^s(\Delta)$. Then $\mu_{|M}^-$ has mass on M. Let U be a neighborhood of J^* and let $w \in J^*$ be a saddle point. By Proposition 3 there exists an integer $n \gg 1$ so that $H^{-n}(M)$ intersects the unstable manifold of w transversally in U. Let x be such an intersection point. Then $x \in W_p^s \cap W_w^u$. By Lemma 3 this implies that $x \in J^*$. Hence $H^n(x) \in J^*$. This contradicts that $(\Phi_p^s)^{-1}(H^n(x)) \in \Delta \subset \mathbb{C} \setminus (\Phi_p^s)^{-1}(J^*).$

Without the hypothesis of uniform hyperbolicity, the proof still works if we assume that p is a periodic saddle point. Hence:

Proposition 6 Let H be a complex Hénon map. Let p be a periodic saddle point. Suppose V is a connected component of $\mathbb{C} \setminus J^*_{p,s}$. Then $G^-_{|V}$ \bar{v} either vanishes identically or is strictly positive.

Proof: Suppose $x \in V$, $\Delta \subset V$, $x \in \Delta$ and $G^{-}(x) = 0$ while G^{-} does not vanish identically on Δ . Then there is a $y \in \Delta$ so that y is a transverse intersection with an unstable manifold of a periodic saddle point. But then $y \in J^*$, a contradiction.

The following result is obvious.

Lemma 5 Suppose that H is uniformly hyperbolic on J^* . Let $p, q \in J^*$. Suppose that $q \in W_p^u$. Then $W_q^u = W_p^u$. Likewise if $q \in W_p^s$, then $W_q^s = W_p^s$.

 \mathcal{L}

We can now prove Theorem 2.

Proof: Suppose that for every $p \in J^*, G^-$ is not vanishing on any open subset of $W^s(p)$. Then there are open subsets $J^* \subset U \subset\subset V$ so that $H(V) \supset U$, $J \setminus \overline{V} \neq \emptyset$ and $G^{-} > c > 0$ on $W_{loc}^{s}(J^{*}) \cap (V \setminus U)$. Pick $x \in J \setminus \overline{V}$. Then $x \in J^+$ and hence, if y is any point in J^* , then there is a sequence $\{y_n\} \subset W^s(y), y_n \to x$. In particular, $G^-(y_n) > c$ for all large n, so $G^{-}(x) \geq c$, a contradiction since $G^{-} \equiv 0$ on J.

Hence, by Proposition 6, there exists a $p \in J^*$ and a nonempty connected component $V \subset \mathbb{C}$ of $\mathbb{C} \setminus J^*_{p,s}$ on which $G^- \equiv 0$. Then V is simply connected by Lemma 5: In fact, if V contains a simple closed curve γ whose interior U contains $x \in J^*$, then $G^- \equiv 0$ on U by the maximum principle. Hence by the invariance of G^- it follows that $G^- \equiv 0$ on $H^{-n}(U)$ for all $n \geq 1$. This contradicts that $H^{-n}(U)$ has to be an unbounded sequence clustering all over J^+ . It follows that there is a weakly stably exposed point in J^* . Similarly we can find a point $q \in J^*$ which is weakly unstably exposed.

Г

 $\overline{}$

Next we prove the Theorem 3.

Proof: Pick a weakly stably exposed point $p \in J^*$. Then in a neighborhood of p , the unstable lamination contains a laminated hypersurface with J^* on one side. Let D denote a small unstable disc centered at p. The forward iterates of this disc become dense in J^* . Moreover, for every point $q \in H^{n}(D) \cap J^*$ there is a connected component $V_{s,q} \subset \mathbb{C} \setminus J^*_{q,s}$ obtained by following $V_{s,p}$ using the local product structure of J^* . Furthermore, by the local product structure each such $V_{s,q}$ is simply connected. Hence, arbitrarily close to any point in J^* there is a laminated hypersurface whis is contained in the local unstable set and with J^* on one side. One can do the same with a weakly unstably exposed point in J^* . Their intersection points give rise to a dense collection of distinguished boundary points.

Next we prove the Main Theorem.

Proof: We assume that $J \neq J^*$. The hypothesis that H is not volume preserving is only used at the end. As in the above proof, for every point $z \in H^{n}(D) \cap J^*$ there is a simply connected component of $W_z^s \setminus J^*$. Using the local product structure of J^* for a complex Hénon map which is hyperbolic on J^* , we get that in fact for every point $z \in J^*$, there is a simply connected component of $W^s_z \setminus J^*$. Also all connected components on which G^- has a zero are simply connected. By [4], if G^- is not identically zero on some simply connected component, then H is stably connected. In particular all connected components of all $W_x^s \setminus J^*$ for all $x \in J^*$ are simply connected and all $W_x^s \setminus J^*$ contain at least one component where $G^+ > 0$ and at least one has a component where $G^- = 0$. The same applies to the unstable set.

Lemma 6 Suppose that H is a complex Hénon map which is hyperbolic on J^* . Suppose that $J \neq J^*$. If H is not stably connected, then each stable manifold $W_z^s \setminus J^*$ considered as parametrized by C contains simply connected components in $\mathbb C$ on which $G^- \equiv 0$ and all such components are bounded in \mathbb{C} .

Proof: Each $W_z^s \setminus J^*$ contains connected components on which $G^- > 0$. However none of these are simply connected, hence they contain curves γ surrounding a nonempty part of J^* . Any simply connected component of some $W_x^s \setminus J^*$ can be followed to any other $W_z^s \setminus J^*$ using the local product structure and in particular can be followed into the interior of such a γ . However, unbounded components stay unbounded when they are followed in this way. This shows that all simply connected components are bounded.

Lemma 7 Let H be a complex Hénon map which is hyperbolic on J^* and suppose that $J \neq J^*$. Then H is stably connected and unstably connected.

Proof: We show that H is stably connected. Unstable connectedness is proved similarly. Suppose that H is not stably connected. We pick a point $z \in J^*$ and a bounded simply connected component V of $W_z^s \setminus J^*$. After changing z if necessary we can pick a curve $\lambda = \phi(t), t \in [0, 1]$ with $\lambda(t) \in V, t > 0$ and $\lambda(0) = z$. We may assume that $V \subset W_{\epsilon, loc}^{s}(J^*)$ for some small $\epsilon > 0$ after forward iteration if necessary. Next, consider the backward iterates, $V_n := H^{-n}(V), \gamma_n = H^{-n}(\gamma), z_n = H^{-n}(z)$. For $n \geq n_0$ there is a unique point $w_n = H^{-n}(\phi(t_n)) \in \gamma_n$ so that $w_n \in \partial W^s_{\epsilon,loc}(J^*)$ and the curve γ_n is contained in $W^s_{\epsilon,loc}(J^*)$ for $0 \leq t < t_n$.

Let w_0 be a cluster point of the sequence $\{w_n\}$. Then w_0 is in some $W^s_\eta, \eta \in J^*$ and belongs to some simply connected component U of $W^s_\eta \setminus J^*$ since $G^{-}(\eta) = 0$ by continuity. Since this component is bounded, we can after replacing with a forward iterate, assume that U is contained in the local stable manifold of η . Using the local product structure it follows that the diameter of V is arbitrarily small, a contradiction.

Using [4] Theorem 5.1 and Corollary 7.4 we get:

F

F

Theorem 4 Suppose that H is a complex H énon map which is hyperbolic on J^* . Moreover assume that $J \setminus J^* \neq \emptyset$. Then H is stably and unstably connected. Morever J is connected and H is volume preserving.

The Main Theorem is now an immediate consequence.

References

[1] E. Bedford, M. Lyubich, J. Smillie; Polynomial Diffeomorphisms of \mathbb{C}^2 . IV. The measure of maximal entropy and laminar currents, Inv. Math. 112 (1993), 77–125.

 \blacksquare

- [2] E. Bedford, J. Smillie; *Polynomial Diffeomorphisms of* \mathbb{C}^2 : Currents, equilibrium measure and hyperbolicity, Inv. Math. 103 (1991), 69–99.
- [3] E. Bedford, J. Smillie; *Polynomial Diffeomorphisms of* \mathbb{C}^2 . II. Stable manifolds and recurrence, J. Amer. Math. Soc. 4 (1991), 657–679.
- [4] E. Bedford, J. Smillie; *Polynomial Diffeomorphisms of* \mathbb{C}^2 : VI. Con*nectivity of J*, Ann. of Math. 148 (1998), 695–735.
- [5] J. E. Fornæss; Real Methods in Complex Dynamics, in Real Methods in Complex and CR Geometry, D. Zaitsev, G. Zampieri (ed.) Springer-Verlag, Lecture Notes in Mathematics 1848. C.I.M.E Summer Course in Martina Franca, Italy 2002.