

# Effective construction of irreducible curve singularities

A. Assi\*, M. Barile †

## Abstract <sup>1</sup>

By using the effective notion of the approximate roots of a polynomial, we describe the equisingularity classes of irreducible curve singularities with a given Milnor number.

## Introduction

Let  $\mathbf{K}$  be an algebraically closed field of characteristic zero. Let  $f$  be an irreducible monic polynomial of  $\mathbf{R} = \mathbf{K}[[x]][y]$ , say  $f = f(x, y) = y^n + a_1(x).y^{n-1} + \dots + a_n(x) \in \mathbf{R}$ . Up to a change of coordinates, we assume that  $a_1(x) = 0$ . For all  $g \in \mathbf{R}$  let  $\text{int}(f, g)$  denote the intersection multiplicity of  $f$  and  $g$ . Let  $\Gamma(f) = \{\text{int}(f, g); g \in \mathbf{R} - (f)\}$  be the semigroup of  $f$ . If  $f'$  is another irreducible polynomial of  $\mathbf{R}$ , then  $f$  and  $f'$  are said to be *equisingular* if  $\Gamma(f) = \Gamma(f')$  (for example,  $y^2 - x^3$  and  $y^3 - x^2$  are equisingular because they are both associated with the semigroup generated by 2, 3). In particular, two equisingular polynomials of  $\mathbf{R}$  need not have the same degree in  $y$ . It is well-known that in this case  $\mu(f) = \mu(f')$ , where  $\mu(f) = \text{int}(f_x, f_y)$  is called the *Milnor number* of  $f$ . The converse is false. The *equisingularity class* of the polynomial  $f$  is the set of irreducible polynomials of  $\mathbf{R}$  which are equisingular to  $f$ . It is of a certain interest to determine this equisingularity class, which gives a classification of the polynomials of  $\mathbf{R}$  in terms of subsemigroups of  $\mathbf{Z}$ . Another remarkable classification is obtained if one can characterize all polynomials whose Milnor number is equal to some fixed nonnegative integer  $m$ . The aim of this paper is to study the two questions from an effective point of view: we first give, for a fixed semigroup of an irreducible polynomial  $f$  of  $\mathbf{R}$ , all elements of the equisingularity class of  $f$ . Then, for a fixed  $m$  in  $\mathbf{N}$ , by similar methods we

\*Université d'Angers, Département de Mathématiques, 2 bd Lavoisier, 49045 Angers Cedex 01, France

†Università degli Studi di Bari, Dipartimento Interuniversitario di Matematica, Via E. Orabona 4, 70125 Bari, Italia.

The second author would like to thank the Departement of Mathematics of the University of Angers for hospitality and support during the preparation of the present paper.

<sup>1</sup>Mathematical Subject Classification: 14Q05, 14C17, 32S15

Keywords: Curve singularities, approximate roots, Milnor number.

construct the generic forms of all irreducible polynomials  $f$  of  $\mathbf{R}$  such that  $\mu(f) = m$ . The set of these polynomials is the union of a finite number of equisingularity classes. We think that this effective classification is useful in the study of problems and conjectures in the theory of irreducible curve singularities, particularly in the understand of their moduli spaces. Our approach uses the effective notion of approximate roots of a polynomial  $f$  of  $\mathbf{R}$  introduced by S.S. Abhyankar and T.T. Moh and the notion of generalized Newton polygon introduced by Abhyankar. The first one gives rise to an algorithm for the computation of the set of generators of the semigroup of  $f$  (and then the set of Newton-Puiseux pairs of  $f$ , see definition 1.3., and [1], [2]). The second one is used by Abhyankar to give an irreducibility criterion for the polynomial  $f$  (see [3]).

We would like to point out that our algorithms are intrinsic and that they have been implemented with *Mathematica* (see [8]), and *Maple*.

## 1 Characteristic sequences

In this Section we recall the notion of approximate roots of  $f$  as well as the characteristic sequences associated with an irreducible polynomial  $f = y^n + a_2(x).y^{n-2} + \dots + a_n(x)$  of  $\mathbf{R}$ .

**Definition 1.1** For any monic polynomial  $g \in \mathbf{R}$ , the **intersection multiplicity**  $\text{int}(f, g)$  of  $f$  with  $g$  is the  $x$ -order of the  $y$ -resultant of  $f$  and  $g$ .

The set  $\Gamma(f) = \{\text{int}(f, g); g \in \mathbf{R} - f\}$  is a subsemigroup of  $\mathbf{Z}$ , called the **semigroup of  $f$** .

**Definition 1.2** Let  $y(t) = \sum_j a_j t^j \in \mathbf{K}[[t]]$  be a root of  $f(t^n, y) = 0$ , according to Newton Theorem. Then set  $m_0 = d_1 = n, m_1 = \inf\{j; a_j \neq 0\}$ , and for all  $k \geq 1$ , let

$$m_{k+1} = \inf\{j; a_j \neq 0 \text{ and } d_k \text{ does not divide } j\}, \text{ and } d_{k+1} = \text{gcd}(m_{k+1}, d_k).$$

Since  $f$  is irreducible, there exists  $h$  such that  $d_{h+1} = 1$ . We set  $m_{h+1} = +\infty$ .

Finally set  $r_0 = m_0 = n, r_1 = O_x(a_n(x))$ -where  $O_x$  denotes the  $x$ -order-, and for all  $k = 1, \dots, h - 1$ :

$$r_{k+1} = r_k \left( \frac{d_k}{d_{k+1}} \right) + (m_{k+1} - m_k).$$

(Remark that, since  $a_1(x) = 0$ , then  $r_1 = m_1$ ).

We recall that with respect to these notations,  $r_0, \dots, r_h$  generates the semigroup  $\Gamma(f)$  of  $f$ . We denote  $\Gamma(f) = \langle r_0, \dots, r_h \rangle$ .

**Definition 1.3** For all  $k = 1, \dots, h$ , set  $e_k = \frac{d_k}{d_{k+1}}$ . The set  $\{(\frac{m_k}{d_{k+1}}, e_k), 1 \leq k \leq h\}$  is called the set of **Newton-Puiseux pairs** of  $f$ .

**Definition 1.4** Let  $d$  be a positive integer and assume that  $d$  divides  $n$ . Let  $g$  be a monic polynomial of  $\mathbf{R}$ , of degree  $\frac{n}{d}$  in  $y$ . We call  $g$  the  $d$ -th approximate root of  $f$  if one of the following holds:

i)  $\deg_y(f - g^d) < n - \frac{n}{d}$ .

ii) in the expansion  $f = g^d + \alpha_1 g^{d-1} + \dots + \alpha_d$  of  $f$  with respect to the powers of  $g$ ,  $\alpha_1 = 0$ .

Remark that i) and ii) are equivalent.

We denote the  $d$ -th approximate root of  $f$  by  $\text{App}_d(f)$ . It is clear that  $\text{App}_d(f)$  is unique, and also that it is effectively computable if the series  $a_k(x), k = 2, \dots, n$ , are polynomials.

**Remark 1.5** Given a divisor  $d$  of  $n$ , the  $d$ th approximate root  $\text{App}_d(f)$  of  $f$  can be effectively constructed from the equation of  $f$  in the following way:

Take  $G_0 = y^{n/d}$  and let  $f = G_0^d + \alpha_1^0 G_0^{d-1} + \dots + \alpha_d^0$  be the expansion of  $f$  with respect to the powers of  $G_0$ .

i) If  $\alpha_1^0 = 0$ , then  $G_0 = \text{App}_d(f)$ .

ii) If  $\alpha_1^0 \neq 0$ , then set  $G_1 = G_0 + \frac{\alpha_1^0}{d}$  and consider the expansion  $f = G_1^d + \alpha_1^1 G_1^{d-1} + \dots + \alpha_d^1$  of  $f$  with respect to the powers of  $G_1$ . If  $\alpha_1^1 \neq 0$ , then easy calculations show that  $\deg_y \alpha_1^0 > \deg_y \alpha_1^1$ . This process shall stop after a finite number of steps, constructing  $\text{App}_d(f)$ .

**Remark 1.6** If the characteristic of  $\mathbf{K}$  is not zero and if this characteristic does not divide  $n$ , then the construction above applies without any restriction. Otherwise, the theory of approximate roots does not work as it. Further information can be found in [9].

Let  $g_1, \dots, g_h, g_{h+1}$  be the  $d_k$ -th approximate roots of  $f$ , for  $k = 1, \dots, h+1$  (in particular  $g_1 = y$  and  $g_{h+1} = f$ ).

**Lemma 1.7** (see [1], (8.2) the Fundamental Theorem (part one)) For all  $k = 1, \dots, h$ , we have:

i)  $\text{int}(f, g_k) = r_k$ .

ii)  $g_k$  is irreducible in  $\mathbf{R}$  and  $\Gamma(g_k) = \langle \frac{r_0}{d_k}, \dots, \frac{r_{k-1}}{d_k} \rangle$ . Furthermore,  $g_1, \dots, g_{k-1}$  are the approximate roots of  $g_k$ .

**Lemma 1.8** (see [13]) The following formulas hold:

-  $\text{int}(f_x, f_y) = \sum_{i=1}^h (e_i - 1)r_i - n + 1$ . In particular  $\text{int}(f_x, f_y)$  is even.

- For all  $k = 2, \dots, h$ ,  $\text{int}(f_x, f_y) = d_k \cdot \text{int}(g_{k_x}, g_{k_y}) + \sum_{i=k}^h (e_i - 1)r_i - d_k + 1$ .

*Proof.* The proof of the first formula can be found in [13] (3.14., page 18). The second formula results from the first one by easy calculations.

**Remark 1.9** The intersection multiplicity  $\text{int}(f_x, f_y)$  is also called the Milnor number of  $f$ . It is an invariant of  $f$  and, by the formula above, it is common to the elements of the equisingularity class of  $f$ . It also coincides with the conductor of the semigroup  $\Gamma(f)$  -usually denoted by  $c$ - which has the following numerical characterization: for all  $p \geq c, p \in \Gamma(f)$ . Furthermore, given two integers  $a, b$ , if  $a + b = m - 1$  then exactly one of  $a, b \in \Gamma(f)$ . It follows that, since  $\Gamma(f)$  has no negative integers,  $\text{Card}(\mathbf{N} - \Gamma) = \frac{m}{2}$ . In fact,  $c$  is nothing but the order of the conductor of the quotient  $\frac{\mathbf{R}}{(f)}$  into its integral closure. Contrary to the Milnor number, the conductor can be defined without restriction on the characteristic of  $\mathbf{K}$ . An exhaustive exposition of this theory in positive characteristic can be found in [9].

## 2 Generalized Newton polygons and the irreducibility criterion of Abhyankar

Let  $f = y^n + a_2(x)y^{n-2} + \dots + a_n(x)$  be a monic polynomial, non necessarily irreducible in  $\mathbf{R}$ . In this section the notations introduced above will have a more general meaning:  $r = (r_0 = n, r_1, \dots, r_h)$  will denote any sequence of integers such that  $r_k < r_{k+1}$  for all  $k = 1, \dots, h - 1$ , and we shall set  $d_{k+1} = \text{gcd}(r_0, r_1, \dots, r_k)$  for all  $k = 0, \dots, h$ . For all  $k = 1, \dots, h$ , we set  $e_k = \frac{d_k}{d_{k+1}}$ ;  $g = (g_1, \dots, g_h, g_{h+1} = f)$  will be a sequence of monic polynomials of  $\mathbf{R}$  such that  $\deg_y g_k = \frac{n}{d_k}$  for all  $k = 1, \dots, h$ . We recall some important properties.

**Theorem 2.1** (see [1], (8.3) The fundamental Theorem (part two)) Let

$$B = \{b = (b_1, b_2, \dots, b_h, b_{h+1}) \in \mathbf{N}^{h+1}; b_1 < e_1, \dots, b_h < e_h, b_{h+1} < +\infty\}$$

For all  $b \in B$ , denote  $g^b = g_1^{b_1} \dots g_h^{b_h} \cdot f^{b_{h+1}}$ , then we have:

- i)  $\mathbf{R} = \sum_{b \in B} \mathbf{K}[[x]].g^b$ .
- ii) Let  $p$  be a polynomial of  $\mathbf{R}$  and write  $p = \sum_{k=1}^s a_k(x).g^{b^k}$ , where  $b^k \in B$  for all  $k = 1, \dots, s$ . Moreover let  $b_0^k = O_x a_k(x)$ , then associate with any "monomial"  $a_k(x).g^{b^k}$  the integer  $\langle (b_0^k, b_1^k, \dots, b_h^k), r \rangle = b_0^k.r_0 + \sum_{i=1}^h b_i^k.r_i$ . Finally let  $B' = \{b^k; b_{h+1}^k = 0\}$ . With these notations we have the following:

- 1) If  $B'$  contains at least two elements, then for all  $b^i, b^j \in B'$ ,

$$b^i \neq b^j \iff \langle (b_0^i, b_1^i, \dots, b_h^i), r \rangle \neq \langle (b_0^j, b_1^j, \dots, b_h^j), r \rangle$$

2)  $f$  doesn't divide  $p$  iff  $B' \neq \emptyset$ , and in this case there is a unique  $k_0$  such that  $\langle (b_0^{k_0}, b_1^{k_0}, \dots, b_h^{k_0}), r \rangle = \inf\{\langle (b_0^k, b_1^k, \dots, b_h^k), r \rangle; b^k \in B'\}$ .

**Definition 2.2** (see [3]) The integer  $\langle (b_0^{k_0}, b_1^{k_0}, \dots, b_h^{k_0}), r \rangle$  is called **formal intersection multiplicity** of  $p$  with respect to  $(r, g)$  and will be denoted by  $\text{fint}(p, r, g)$ .

Now we recall the notion of **generalized Newton polygon**. Let  $p$  be a monic polynomial of  $\mathbf{R}$  of degree  $n$  in  $y$  and consider a monic polynomial  $q$  of  $\mathbf{R}$  of degree  $\frac{n}{d}$  in  $y$ , where  $d$  is a divisor of  $n$ . Let

$$p = q^d + \alpha_1(x, y)q^{d-1} + \dots + \alpha_d(x, y)$$

be the expansion of  $p$  with respect to the powers of  $q$ , and consider the sequences  $r, g$  defined above. One associates with  $p$  the generalized Newton polygon which is defined as the union of all compact sides of the convex hull in  $\mathbf{R}^2$  of the set formed by the points  $(\text{fint}(\alpha_k, r, g), (d - k) \cdot \text{fint}(q, r, g))$  for all  $1 \leq k \leq d$ . It will be denoted by  $\text{GNP}(p, q, r, g)$  (see [3]).

With these notations one has the following:

**Irreducibility criterion** (see [3])

Write  $p = y^n + a_1(x)y^{n-1} + \dots + a_n(x) \in \mathbf{R}$  and assume, possibly after a change of variables, that  $a_1(x) = 0$ . Consider the sequences  $r_k, g_k, d_k$  defined in the following way:

$$r_0 = d_1 = n$$

$$g_1 = y, r_1 = \text{int}(p, g_1), \text{ and for all } k \geq 2:$$

$$d_k = \text{gcd}(r_0, r_1, \dots, r_{k-1}), g_k = \text{App}_{d_k}(p), r_k = \text{int}(p, g_k).$$

$p$  is irreducible if and only if the following conditions hold.

1) There is  $h \in \mathbf{N}$  such that  $d_{h+1} = 1$ .

2) For all  $k = 1, \dots, h - 1$ ,  $r_{k+1}d_{k+1} > r_k d_k$ .

3) Set  $p = g_{h+1}$  and let for all  $k = 1, \dots, h$ ,  $e_k = \frac{d_k}{d_{k+1}}$  and  $g^k = (g_1, \dots, g_k)$ . Let also for all  $k = 1, \dots, h$ ,  $\frac{r}{d_{k+1}} = (\frac{r_0}{d_{k+1}}, \dots, \frac{r_k}{d_{k+1}})$ . Then for all  $k = 1, \dots, h$ , the generalized Newton polygon  $\text{GNP}(g_{k+1}, g_k, \frac{r}{d_{k+1}}, g^k)$  is the line segment joining  $(0, \frac{r_k}{d_{k+1}} \cdot e_k)$  and  $(\frac{r_k}{d_{k+1}} \cdot e_k, 0)$ .

**Remarks 2.3:** i) (see [3]) Suppose that  $p$  is irreducible, and let  $r = (r_0, r_1, \dots, r_h)$  and  $g = (g_1 = y, g_2, \dots, g_h, g_{h+1} = p)$  be the sequences defined above. Let  $p'$  by a polynomial of  $\mathbf{R}$

and consider the expansion of  $p'$  with respect to the sequences  $r, g$  (see Theorem 2.1.). If the corresponding set  $B'$  is non empty, then  $\text{fint}(p', r, g) = \text{int}(p', p)$ .

ii) Part 3) of the criterion can be precised as follows: whenever  $p$  is irreducible, the generalized Newton polygon  $\text{GNP}(g_{k+1}, g_k, \frac{r}{d_{k+1}}, g^k)$  just contains the two extremal points for all  $k = 1, \dots, h-1$ . In fact let  $k \in \{1, \dots, h\}$  and let  $g_{k+1} = g_k^{e_k} + \alpha_2(x, y).g_k^{e_k-2} + \dots + \alpha_{e_k}(x, y)$  be the expansion of  $g_{k+1}$  w.r.t.  $g_k$ , then we have:

$$\text{a) } \text{int}(g_{k+1}, \alpha_{e_k}(x, y)) = \frac{r_k}{d_{k+1}}.e_k = \text{int}(g_{k+1}, g_k^{e_k}).$$

$$\text{b) } \text{For all } i = 2, \dots, e_k - 1, \text{int}(g_{k+1}, \alpha_i(x, y)) > \frac{r_k}{d_{k+1}}.i, \text{ In particular } \text{int}(g_{k+1}, \alpha_i(x, y)) + \text{int}(g_{k+1}, g_k^{e_k-i}) > \frac{r_k}{d_{k+1}}.e_k.$$

iii) (see [13]) As an immediate consequence of ii) a) we have that, for all  $k = 1, \dots, h$ ,

$$\frac{r_k}{d_{k+1}}.e_k \in e_k.\Gamma(g_k) = \langle \frac{r_0}{d_{k+1}}, \dots, \frac{r_{k-1}}{d_{k+1}} \rangle.$$

In particular

$$r_k.e_k \in \langle r_0, \dots, r_{k-1} \rangle.$$

### 3 Constructing the equisingularity class

In this Section we fix a semigroup of nonnegative integers  $\Gamma = \langle r_0, r_1, \dots, r_h \rangle$ , and we set  $d_1 = r_0$  and  $d_{k+1} = \text{gcd}(r_0, \dots, r_k)$  for all  $k = 1, \dots, h$  (we set by convention  $r_{h+1} = d_{h+2} = +\infty$ ). Moreover we assume that  $d_{k+1} = 1$  and that  $r_{k+1}.d_{k+1} > r_k.d_k$  for all  $k = 1, \dots, h$  (\*) (this condition appeared in the irreducibility criterion in Section 2). This implies that the sequence  $(r_1, \dots, r_h)$  is strictly increasing. This also holds if  $r_1$  is replaced by  $r_0$ .

Let  $\tilde{\mathbf{R}}$  denote the set of all irreducible monic polynomials  $f$  of  $\mathbf{R}$  of the form  $f = f(x, y) = y^n + a_2(x).y^{n-2} + \dots + a_n(x)$ . Condition (\*) implies that there exists a polynomial  $f \in \tilde{\mathbf{R}}$  such that  $\Gamma = \Gamma(f)$  (see [13]). Here we give the generic forms of all these polynomials, i.e., we describe the set of elements of  $\tilde{\mathbf{R}}$  having the semigroup  $\Gamma$ . The construction can be performed with respect to the arrangements  $(r_0, r_1, r_2, \dots, r_h)$  and  $(r_1, r_0, r_2, \dots, r_h)$ . We shall perform it with respect to the first arrangement. The polynomials that we would get with respect to the second arrangement are those obtained by exchanging  $x$  and  $y$ .

In this and in the following Sections we shall assume that  $r_0, \dots, r_h$  form a minimal system of generators for  $\Gamma$ . This condition can be reformulated equivalently as a numerical criterion. This is what we are going to do next. First we derive a useful identity: Set for all  $1 \leq i \leq h$ ,  $e_i = \frac{d_i}{d_{i+1}}$ .

For all  $2 \leq k \leq h$  we have:

$$\begin{aligned}
\sum_{i=1}^{k-1} (e_i - 1)r_i &= \sum_{i=1}^{k-1} (r_{i+1} - (m_{i+1} - m_i)) - \sum_{i=1}^{k-1} r_i \\
&= r_k - r_1 - \sum_{i=1}^{k-1} (m_{i+1} - m_i) = r_k - m_k \quad (**)
\end{aligned}$$

Now we are ready to prove the following

**Lemma 3.1** *Suppose that  $r_0, \dots, r_h$  satisfy condition (\*). These numbers form a minimal system of generators for the semigroup  $\Gamma$  if and only if  $d_1 > d_2 > \dots > d_h > d_{h+1}$ .*

*Proof.* Remark that in general  $d_k \geq d_{k+1}$  for all  $k = 1, \dots, h$ . Moreover, recall that the minimality of the system of generators is equivalent to the condition that  $r_k \notin \langle r_0, \dots, r_{k-1} \rangle$  for all  $k = 1, \dots, h$ . First suppose that this condition is not fulfilled for some index  $k$ . Then  $d_k = \gcd(r_0, \dots, r_{k-1})$  certainly divides  $r_k$ . Hence  $d_{k+1} = \gcd(d_k, r_k) = d_k$ . For the converse fix an index  $k$  and suppose that  $d_{k+1} = d_k$ . Then there exist some integers  $\alpha_0, \dots, \alpha_{k-1}$  such that

$$r_k = \sum_{i=0}^{k-1} \alpha_i r_i. \quad (1)$$

Now let  $\beta_{k-1}$  be the (nonnegative) remainder of the euclidean division of  $\alpha_{k-1}$  by  $e_{k-1} = \frac{d_{k-1}}{d_k}$ . Since the semigroup  $\Gamma$  verifies condition (\*),  $\Gamma$  is the semigroup of a polynomial of  $\tilde{\mathbf{R}}$ , in particular, by Remarks 2.3 iii),  $r_{k-1} \cdot (e_{k-1}) \in \langle r_0, \dots, r_{k-2} \rangle$ ; hence we can transform (1) in such a way that  $\alpha_{k-1} = \beta_{k-1}$ . If we successively perform the same procedure for the indices  $k-2, \dots, 1$ , we finally obtain that in (1)  $0 \leq \alpha_i < e_i$  for all  $i = 1, \dots, k-1$ . Now by (\*\*)

$$\alpha_0 r_0 = r_k - \sum_{i=1}^{k-1} \alpha_i r_i > r_k - \sum_{i=1}^{k-1} (e_i - 1)r_i = m_k > 0,$$

hence  $\alpha_0 > 0$ . This proves that  $r_k \in \langle r_0, \dots, r_{k-1} \rangle$  and completes the proof.

The construction of the generic form of all polynomials  $f \in \tilde{\mathbf{R}}$  having  $\Gamma$  as a semigroup is based on the notion of **generalized Newton polygons** introduced in Section 2. We shall recursively construct the sequence of approximate roots  $g_1, \dots, g_h, g_{h+1} = f$ .

Let  $g_1 = \text{App}_{d_1}(f)$  (and recall that, since  $a_1(x) = 0$ , then  $g_1 = y$ ). From Section 2 we know that  $g_2 = \text{App}_{d_2}(f)$  satisfies:

i)  $\Gamma(g_2) = \langle \frac{r_0}{d_2}, \frac{r_1}{d_2} \rangle$ .

ii)  $\text{GNP}(g_2, g_1, \frac{r}{d_2}, g^1)$  is the line segment joining the two points  $(0, \frac{r_1}{d_2} \cdot e_1)$  and  $(\frac{r_1}{d_2} \cdot e_1, 0)$ .

By virtue of part ii) of Remarks 2.3, this yields to the following generic form of  $g_2$ :

$$g_2 = y^{\frac{r_0}{d_2}} + a \cdot x^{\frac{r_1}{d_2}} + \sum_{i \cdot \frac{r_0}{d_2} + j \cdot \frac{r_1}{d_2} > \frac{r_0}{d_2}; 0 \leq j < \frac{d_1}{d_2} = \frac{r_0}{d_2}} a_{ij} x^i y^j,$$

where  $a \in \mathbf{K} - 0$  and for all  $(i, j)$ ,  $a_{ij} \in \mathbf{K}$ .

Suppose that we have the generic forms of  $g_1, \dots, g_k$  and consider the expansion of  $g_{k+1}$  with respect to  $g_k$ :

$$g_{k+1} = g_k^{e_k} + \alpha_2(x, y)g_k^{e_k-2} + \dots + \alpha_{e_k}(x, y).$$

From Section 2 we know that:

$$\text{i) } \Gamma(g_{k+1}) = \left\langle \frac{r_0}{d_{k+1}}, \dots, \frac{r_k}{d_{k+1}} \right\rangle.$$

$$\text{ii) } \text{GNP}(g_{k+1}, g_k, \frac{r}{d_{k+1}}, g^k) \text{ is the line segment joining the two points } (0, \frac{r_k}{d_{k+1}} \cdot e_k) \text{ and } (\frac{r_k}{d_{k+1}} \cdot e_k, 0).$$

It follows from Remarks 2.3 that

$$(1) \text{ int}(g_{k+1}, \alpha_{e_k}(x, y)) = \text{fint}(\alpha_{e_k}(x, y), \frac{r}{d_{k+1}}, g^k) = \frac{r_k}{d_{k+1}} \cdot e_k,$$

and that for all  $i = 2, \dots, e_k - 1$ :

$$(2) \text{ int}(g_{k+1}, \alpha_i(x, y)) = \text{fint}(\alpha_i(x, y), \frac{r}{d_{k+1}}, g^k) > \frac{r_k}{d_{k+1}} \cdot i,$$

Let  $B^k$  be the set of all  $\theta = (\theta_0, \dots, \theta_{k-1}) \in \mathbf{N}^k$  such that  $0 \leq \theta_j < e_j$  for all  $j = 1 \dots, k - 1$ , then associate with all  $\theta \in B^k$  the ‘‘monomial’’  $M_\theta = x^{\theta_0} \cdot g_1^{\theta_1} \cdot \dots \cdot g_{k-1}^{\theta_{k-1}}$ . For all  $i \in \mathbf{N}$  and for all  $\theta \in B^k$ , we say that  $M_\theta$  is of type  $(k, i, 1)$  (resp. of type  $(k, i, 2)$ ) if

$$\frac{r_k}{d_{k+1}} \cdot i = \theta_0 \cdot \frac{r_0}{d_{k+1}} + \theta_1 \cdot \frac{r_1}{d_{k+1}} + \dots + \theta_{k-1} \cdot \frac{r_{k-1}}{d_{k+1}}$$

(resp.

$$\frac{r_k}{d_{k+1}} \cdot i < \theta_0 \cdot \frac{r_0}{d_{k+1}} + \theta_1 \cdot \frac{r_1}{d_{k+1}} + \dots + \theta_{k-1} \cdot \frac{r_{k-1}}{d_{k+1}}).$$

Let  $E(k, i, 1)$  (resp.  $E(k, i, 2)$ ) be the set of monomials  $M_\theta, \theta \in B^k$ , of type  $(k, i, 1)$  (resp. of type  $(k, i, 2)$ ). Since  $\frac{r_k}{d_{k+1}} \cdot e_k \in \left\langle \frac{r_0}{d_{k+1}}, \dots, \frac{r_{k-1}}{d_{k+1}} \right\rangle$ , then  $E(k, e_k, 1)$  is reduced to one element.

If we write this element as  $M_{\theta^k} = x^{\theta_0} \cdot g_1^{\theta_1} \cdot \dots \cdot g_{k-1}^{\theta_{k-1}}$ , then  $(\theta_0, \theta_1, \dots, \theta_{k-1})$  can be calculated by euclidean division.

Using Remark 2.3., (1) and (2) this leads to the following generic forms of  $\alpha_2, \dots, \alpha_{e_k}$  :

$$\alpha_{e_k} = a \cdot M_{\theta^k} + \sum_{M_\theta \in E(k, e_k, 2)} a_\theta \cdot M_\theta,$$



(resp. for all  $i = 2, \dots, e_k - 1$ ,

$$\alpha_i = \sum_{M_\theta^i \in E(k,i,2)} a_\theta^i \cdot M_\theta^i),$$

where  $a \in \mathbf{K} - 0$ , and for all  $\theta, a_\theta \in \mathbf{K}$  (resp. for all  $\theta$  and for all  $i = 2, \dots, e_k - 1$ ,  $a_\theta^i \in \mathbf{K}$ ).

**Remark 3.2** We proved that, if  $\Gamma$  is the semigroup of a polynomial  $f \in \tilde{\mathbf{R}}$ , then  $f$  and its approximate roots  $g_1, \dots, g_h$  belong to the set of polynomials constructed above. Conversely, let  $(g_1, \dots, g_h, g_{h+1} = f)$  be as above, then part “only if” of the irreducibility criterion of Abhyankar shows that  $f$  is irreducible.

**Remark 3.3** Given  $1 \leq k \leq h$ , it follows from the above construction that a polynomial  $g_{k+1}$  may have an infinite number of monomials. In particular the above construction is not algorithmic. Remark however that  $g_{k+1}$  is obtained from the sum  $g_k^{e_k} + a \cdot M_{\theta^k}$ ,  $a \in \mathbf{K} - 0$  by adding monomials that verify some conditions. This suggests the introduction of the following set of polynomials: let  $G_1 = y$  and for all  $1 \leq k \leq h$ ,  $G_{k+1} = G_k^{e_k} - M_{\theta^k}$ . For all  $1 \leq k \leq h$ ,  $g_k$  is obtained from  $G_k$  in an obvious way. Set  $G = (G_1, \dots, G_h, G_{h+1})$  and call it the canonical element of the set of all  $(g_1, \dots, g_h, g_{h+1})$  constructed above. The above calculation leads to an algorithm that computes this canonical element. It is based on euclidean division in  $\mathbf{N}$ . The different steps can be summarized as follows:

- i) Consider a sequence of integers  $r_0 < \dots < r_h$ .
- ii) Compute the gcd sequence  $d = (d_1, \dots, d_h, d_{h+1})$  such that  $d_1 = r_0$  and for all  $2 \leq k \leq h + 1$ ,  $d_k = \gcd(r_{k-1}, d_{k-1})$ . Let  $e_k = \frac{d_k}{d_{k+1}}$  for all  $1 \leq k \leq h$ .
- iii) If either  $d_{h+1} > 1$ , or  $r_k \cdot e_k \geq r_{k+1}$  for at least one  $1 \leq k \leq h$ , then the sequence  $(r_0, \dots, r_h)$  is not the semigroup of an irreducible polynomial of  $\tilde{\mathbf{R}}$ .
- iv) Assume that  $d_{h+1} = 1$  and that  $r_k \cdot d_k < r_{k+1} \cdot d_{k+1}$  for all  $1 \leq k \leq h - 1$ .
  - a) If  $d_k = d_{k+1}$  for some  $1 \leq k \leq h$ , then eliminate  $r_k$  from the  $r$ -sequence of i).
  - b) Assume that  $d_1 > d_2 > \dots > d_h > d_{h+1} = 1$ . Then for all  $1 \leq k \leq h$ , compute (the unique)  $\theta^k = (\theta_0^k, \dots, \theta_{k-1}^k)$  such that  $0 \leq \theta_j^k < e_j$  for all  $j = 1, \dots, k - 1$  and  $\frac{r_k}{d_{k+1}} \cdot e_k = \theta_0^k \cdot \frac{r_0}{d_{k+1}} + \theta_1^k \cdot \frac{r_1}{d_{k+1}} + \dots + \theta_{k-1}^k \cdot \frac{r_{k-1}}{d_{k+1}}$ .
  - c) The canonical element is  $G = (G_1, \dots, G_h, G_{h+1})$  where  $G_1 = y$  and for all  $2 \leq k \leq h + 1$ ,  $G_k = G_{k-1}^{e_{k-1}} - x^{\theta_0^k} y^{\theta_1^k} \cdot \dots \cdot G_{k-1}^{\theta_{k-1}^k}$ .

This algorithm has been implemented with *Mathematica* (see [8]), and *Maple*: the input is an increasing sequence of positive integers. Then the output is “false” if this sequence does not

generate the semigroup of an irreducible polynomial of  $\tilde{\mathbf{R}}$ . Otherwise, we get the canonical element described above.

Note that our implementation is based on the following: given  $r_0, r_1, \dots, r_{k-1}$ , we need to compute the unique  $\theta^k = (\theta_0^k, \dots, \theta_{k-1}^k)$  such that  $0 \leq \theta_j^k < e_j$  for all  $j = 1 \dots, k-1$  and  $\frac{r_k}{d_{k+1}} \cdot e_k = \theta_0^k \cdot \frac{r_0}{d_{k+1}} + \theta_1^k \cdot \frac{r_1}{d_{k+1}} + \dots + \theta_{k-1}^k \cdot \frac{r_{k-1}}{d_{k+1}}$ . Instead of applying the Euclidean division, We have preferred to scan lists of values, namely the set of values  $(a_0, a_1, \dots, a_{k-1})$  where for all  $i \geq 1, 0 \leq a_i < e_i$  and  $0 \leq a_0 \leq \frac{r_k}{d_{k+1}} \cdot e_k \cdot \frac{d_{k+1}}{r_0} = \frac{r_k \cdot e_k}{r_0}$ . The cardinality of this set is:

$$\frac{r_k \cdot e_k}{r_0} \cdot \prod_{i=1}^{k-1} e_i = \frac{r_k \cdot e_k}{r_0} \cdot \frac{d_1}{d_k} = \frac{r_k}{d_{k+1}}$$

In conclusion the set of the values scanned in the algorithm is bounded by  $\sum_{k=1}^h \frac{r_k}{d_{k+1}}$ .

**Remark 3.4** An element  $f$  whose semigroup is  $\Gamma$  can also be calculated by using the theory of Gröbner bases: a reduced Gröbner basis with respect to any well-ordering on  $\mathbf{N}^3$  that eliminates  $t$  from the equations  $x - t^n, y - t^{m_1} - \dots - t^{m_r}$  contains a unique polynomial  $f(x, y)$ . If we consider  $f$  as an element of  $\mathbf{K}[[x, y]]$ , then obviously  $\Gamma = \langle r_0, \dots, r_h \rangle$  is the semigroup of  $f$ . It is well known that the complexity of a Gröbner basis is in general doubly exponential. Moreover, the algorithm computes more than we need. We think that our option is more natural in view of our situation, especially because of its complexity and that the output is expressed in terms of the polynomial  $f$ .

### Example:

Let  $\Gamma = \langle 8, 12, 50, 101 \rangle$ . Here  $h = 3$ , the  $r$ -sequence is  $r = (8, 12, 50, 101)$ , and the gcd-sequence is  $d = (8, 4, 2, 1)$ . Moreover,  $e_1 = e_2 = e_3 = 2$ . Let us construct the canonical element  $G = (G_1, G_2, G_3, G_4)$  following the algorithm above. Here we start directly from point iv), b):

$$k = 1 : \frac{r_1}{d_2} \cdot e_1 = 3 \cdot 2 = \theta_0^1 \cdot \frac{r_0}{d_2} = \theta_0^1 \cdot 2 \text{ implies that } \theta_0^1 = 3.$$

$$k = 2 : 50 = \frac{r_2}{d_3} \cdot e_2 = \theta_0^2 \cdot \frac{r_0}{d_3} + \theta_1^2 \cdot \frac{r_1}{d_3} = \theta_0^2 \cdot 4 + \theta_1^2 \cdot 6 \text{ with } 0 \leq \theta_1^2 < 2. \text{ This implies that } \theta_1^2 = 1, \text{ and } \theta_0^2 = 11.$$

$$k = 3 : 202 = \frac{r_3}{d_4} \cdot e_3 = \theta_0^3 \cdot \frac{r_0}{d_4} + \theta_1^3 \cdot \frac{r_1}{d_4} + \theta_2^3 \cdot \frac{r_2}{d_4} = \theta_0^3 \cdot 8 + \theta_1^3 \cdot 12 + \theta_2^3 \cdot 50 \text{ with } 0 \leq \theta_1^3, \theta_2^3 < 2.$$

This implies that  $\theta_2^3 = 1, \theta_1^3 = 0, \theta_0^3 = 19$ .

In particular,  $G_1 = y, G_2 = G_1^2 - x^3 = y^2 - x^3, G_3 = G_2^2 - x^{11} \cdot G_1 = (y^2 - x^3)^2 - x^{11} \cdot y, G_4 = G_3^2 - x^{19} \cdot G_2 = [(y^2 - x^3)^2 - x^{11} \cdot y]^2 - x^{19} \cdot (y^2 - x^3)$ .

With the same notations as above, the set of elements  $(g_1, g_2, g_3, g_4 = f)$  is then given by:

$$g_1 = y.$$

$$g_2 = y^2 + \alpha_2(x) = y^2 + ax^3 + \sum_{M_\theta \in E(1,2,2)} a_\theta \cdot M_\theta,$$

where  $a \in \mathbf{K} - 0$ , and for all  $\theta$ , one has  $a_\theta \in \mathbf{K}$  and  $M_\theta = x^{\theta_0}$ , with  $6 < 2\theta_0$ . Moreover

$$g_3 = g_2^2 + \alpha_2'(x, y) = g_2^2 + a'x^{11}y + \sum_{M'_\theta \in E(2,2,2)} a'_\theta \cdot M'_\theta,$$

where

- $a' \in \mathbf{K} - 0$ , and for all  $\theta$ ,  $a'_\theta \in \mathbf{K}$ ;
- for all  $\theta$ ,  $M'_\theta = x^{\theta'_0}y^{\theta'_1}$ , with  $50 < 4\theta'_0 + 6\theta'_1$ .

Finally,

$$f = g_3^2 + \alpha_2''(x, y) = g_3^2 + a''x^{19}g_2 + \sum_{M''_\theta \in E(3,2,2)} a''_\theta \cdot M''_\theta,$$

where

- $a'' \in \mathbf{K} - 0$ , and for all  $\theta$ ,  $a''_\theta \in \mathbf{K}$ ;
- for all  $\theta$ ,  $M''_\theta = x^{\theta''_0}y^{\theta''_1}g_2^{\theta''_2}$ , with  $202 < 8\theta''_0 + 12\theta''_1 + 50\theta''_2$ . Hence the generic form of all polynomials having  $\Gamma$  as a semigroup is the following:

$$f = [(y^2 + ax^3 + F)^2 + a'x^{11}y + F']^2 + a''x^{19}(y^2 + ax^3 + F) + F'',$$

where  $a, a', a'' \in \mathbf{K} - 0$  and  $F, F'$  and  $F''$  are arbitrary linear combinations of monomials from  $E(1, 2, 2)$ ,  $E(2, 2, 2)$  and  $E(3, 2, 2)$  respectively.

**Remark 3.5** i) The construction above does not depend on the choice of the coefficients in the field  $\mathbf{K}$  -provided that it is of characteristic zero-, in particular the algorithm described allows us to work over any subring  $A$  of  $\mathbf{K}$ . If  $A = k[t_1, \dots, t_m]$  is a polynomial ring over a field  $k$  of characteristic zero and  $\mathbf{K}$  is the algebraic closure of  $A$  into its fractions field, then we get the equisingularity class of the  $(t_1, \dots, t_m)$ -generic section.

ii) The restriction to the zero characteristic is made only because of the use of the approximate roots in the algorithm. If the characteristic of  $\mathbf{K}$  does not divide  $r_0$ , then everything above applies -see Remark 1.6.-. Note that a more general irreducibility criterion has been given by A. Granja (see [12]), but it does not seem to be in computational form.

## 4 Equisingularity classes with a given Milnor number

In this Section we generalize the results of Section 3 in the following way: let  $m \in \mathbf{N}$  be a fixed integer. If  $m \in 2\mathbf{N}$ , then there exists a polynomial  $f = y^n + a_2(x).y^{n-2} + \dots + a_n(x) \in \tilde{\mathbf{R}}$  such that  $\text{int}(f_x, f_y) = m$ . Here we shall give the generic forms of all these polynomials. Remark that if  $g$  is another polynomial of  $\tilde{\mathbf{R}}$ , then  $\Gamma(f) = \Gamma(g)$  implies that  $\text{int}(f_x, f_y) = \text{int}(g_x, g_y)$ . Thus the set of  $f = y^n + a_2(x).y^{n-2} + \dots + a_n(x) \in \tilde{\mathbf{R}}$  such that  $\text{int}(f_x, f_y) = m$  is the union of equisingularity classes. We shall first prove that this union is finite. This is an immediate application of the next Proposition. We recall that if a subsemigroup of  $\mathbf{Z}$  is minimally generated by  $h + 1$  elements, then  $h$  is called the *length* of the semigroup.

**Proposition 4.1** *Let  $h \in \mathbf{N}$  and consider a polynomial  $f \in \tilde{\mathbf{R}}$  such that  $h$  is the length of  $\Gamma(f)$ . Let  $\mu_{h+1} = \text{int}(f_x, f_y)$ , and let  $r_h$  be the last generator of  $\Gamma(f)$ . We have the following:*

- i)  $h = 1$  implies that  $r_h \geq 3$  and  $\mu_{h+1} \geq 2$ .
- ii)  $h = 2$  implies that  $r_h \geq 13$  and  $m_{h+1} \geq 16$ .
- iii) More generally we have:

$$1) r_h \geq 12.4^{h-2} + \sum_{i=0}^{h-2} 4^i = \frac{5}{3}.2^{2h-1} - \frac{1}{3}.$$

$$2) \mu_{h+1} \geq 2 + 2. \sum_{i=0}^{h-2} 4^i + 12. \sum_{i=h-2}^{2h-4} 2^i = \frac{5}{3}.2^{2h} - 3.2^h + \frac{4}{3}, \text{ assuming that the summation over negative exponents is 0.}$$

*Proof.* i) In this case, by Lemma 1.8.,  $\mu_1 = (r_0 - 1).(r_1 - 1)$ . Furthermore  $r_1 \geq 2$  and  $r_0 \geq 2$ ; otherwise  $\Gamma(f) = \langle 1 \rangle$ , and  $h = 0$ . On the other hand,  $\text{gcd}(r_0, r_1) = 1$ . This proves that  $\max(r_0, r_1) = r_1 \geq 3$  and  $\mu_2 \geq 2$ . Then our assertion follows. Remark that  $r_1 = 3$  and  $\mu_2 = 2$  holds for  $f = y^2 + ax^3$ , where  $a \in \mathbf{K} - 0$ .

ii) Let  $g_2$  be the second approximate root of  $f$ , then  $\mu_3 = d_2.\text{int}(g_{2x}, g_{2y}) + (d_2 - 1)(r_2 - 1)$ . It follows from i) that  $\frac{r_1}{d_2} \geq 3$  and that  $\text{int}(g_{2x}, g_{2y}) = (\frac{r_0}{d_2} - 1)(\frac{r_1}{d_2} - 1) \geq 2$ , and also that  $\frac{r_0}{d_2} + \frac{r_1}{d_2} \geq 5$ . In particular

$$(r_0 - d_2)(\frac{r_1}{d_2} - 1) \geq 2.d_2$$

and

$$r_0 + r_1 \geq 5.d_2.$$

Thus:

$$r_1.\frac{r_0}{d_2} \geq d_2 + r_1 + r_0 \geq 6.d_2 \geq 12.$$

But  $r_2 - 1 \geq r_1 \cdot \frac{d_1}{d_2} = r_1 \cdot \frac{r_0}{d_2}$ . Finally  $r_2 \geq r_1 \frac{r_0}{d_2} + 1 \geq 13$ , and  $\mu_{h+1} \geq 2 \cdot d_2 + (r_2 - 1) \geq 4 + 12 = 16$ . This implies our assertion. Note that the lower bounds 13 and 16 is sharp: they are satisfied for  $f = (y^2 + a \cdot x^3)^2 + b \cdot x^5 y$ , where  $a, b \in \mathbf{K} - 0$ , whose semigroup is  $\Gamma = \langle 4, 6, 13 \rangle$ .

iii) We prove the inequalities by induction on  $h$ . From i) and ii) both are satisfied for  $1 \leq h \leq 2$ . Assume that  $h \geq 3$  and that the formulas are true for  $h-1$ . We first prove the inequality 1): Remark first that  $r_h \geq (\frac{r_{h-1}}{d_h}) \cdot d_{h-1} + 1$ . The quotient  $\frac{r_{h-1}}{d_h}$  being the last generator of  $\Gamma(g_h)$  which is of length  $h-1$ , it follows by induction that  $\frac{r_{h-1}}{d_h} \geq 12 \cdot 4^{h-3} + \sum_{i=0}^{h-3} 4^i = \frac{5}{3} \cdot 2^{2h-3} - \frac{1}{3}$ . On the other hand,  $d_{h-1} \geq 4$ , thus  $r_h \geq 4 \cdot (\frac{5}{3} \cdot 4^{2h-3} - \frac{1}{3}) + 1 = \frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3}$ . This is the required inequality.

We now prove the inequality 2): Consider to this end the last approximate root  $g_h$  of  $f$ . We have:  $\mu_{h+1} = d_h \cdot \text{int}(g_{h_x}, g_{h_y}) + (d_h - 1)(r_h - 1)$ . But  $d_h \geq 2$  and  $r_h \geq \frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3}$ . On the other hand, the length of  $\Gamma(g_h)$  being  $h-1$ , it follows that

$$\text{int}(g_{h_x}, g_{h_y}) \geq 2 + 2 \cdot \sum_{i=0}^{h-3} 4^i + 12 \cdot \sum_{i=h-3}^{2h-6} 2^i = \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}$$

In particular

$$\mu_{h+1} \geq \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3} + \frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3} - 1 = \frac{5}{3} \cdot 2^{2h} - 3 \cdot 2^h + \frac{4}{3}$$

This is the required inequality.

**Remark 4.2** The bounds of the above Proposition are sharp. More precisely, for all  $h \geq 1$ , there is a polynomial  $f_h(x, y) \in \tilde{R}$  such that  $h$  is the length of  $\Gamma(f)$ , and that  $\text{int}(f_{h_x}, f_{h_y}) = \frac{5}{3} \cdot 2^{2h} - 3 \cdot 2^h + \frac{4}{3}$ , and if  $r_h$  denotes the last generator of  $\Gamma(f)$ , then  $r_h = \frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3}$ . Consider to this end the semigroup  $\Gamma_h$  generated by  $r_0 = 2^h$  and

$$r_k = 2^{h-k} \left( \frac{5}{3} \cdot 2^{2k-1} - \frac{1}{3} \right)$$

for all  $1 \leq k \leq h$  (equivalently  $r_1 = 2^{h-1} \cdot 3, r_2 = 2^h \cdot 3 + 2^{h-2}, \dots, r_{k+2} = 2^{h+k} \cdot 3 + \sum_{i=1}^{k+1} 2^{h+k-2i}$  for all  $1 \leq k \leq h-2$ ). Clearly  $r_h = \frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3}$ , and the  $d$ -sequence is given by  $d_k = 2^{h+1-k}, 1 \leq k \leq h+1$ . Furthermore,  $r_k d_k < r_{k+1} d_{k+1}$  for all  $1 \leq k \leq h$ . It follows that  $\Gamma_h$  is the semigroup of a polynomial of  $\tilde{R}$ . We shall prove by induction that the Milnor number of such a polynomial is  $\frac{5}{3} \cdot 2^{2h} - 3 \cdot 2^h + \frac{4}{3}$ . Denote this number by  $\mu_{h+1}$  and recall that  $\mu_{h+1} = \sum_{k=1}^h (\frac{d_k}{d_{k+1}} - 1) r_k - r_0 + 1$ .

Since  $\frac{d_k}{d_{k+1}} = 2$  for all  $1 \leq k \leq h$ , then:

$$\mu_{h+1} = \left( \sum_{k=1}^h r_k \right) - r_0 + 1 = \sum_{k=1}^h 2^{h-k} \left( \frac{5}{3} \cdot 2^{2k-1} - \frac{1}{3} \right) - 2^h + 1$$

Which is nothing but  $\frac{5}{3} \cdot 2^{2h} - 3 \cdot 2^h + \frac{4}{3}$ . This proves our assertion.

**Corollary 4.3** *Let  $m \in 2\mathbf{N}$ , then one can effectively compute the set of irreducible polynomials  $f \in \tilde{\mathbf{R}}$  such that  $m = \text{int}(f_x, f_y)$ .*

*Proof.* It follows from Proposition 4.1. that the length  $h$  of the semigroup of a polynomial  $f \in \tilde{\mathbf{R}}$  with  $m = \text{int}(f_x, f_y)$  takes a finite number of values. In fact, easy calculations show that  $h$  must verify the inequality:  $2^h \leq M = \frac{9 + \sqrt{1 + 60m}}{10}$ . In particular,  $h \leq \frac{\ln(M)}{\ln(2)}$ . Let  $H = \{h \in \mathbf{N}; h \leq \frac{\ln(M)}{\ln(2)}\}$ . Given  $h \in A$  we shall effectively construct the set  $\Sigma$  of all the sequences  $(r_0, r_1, \dots, r_h)$  which minimally generate a semigroup of a polynomial  $f \in \tilde{\mathbf{R}}$  of the required Milnor number. The steps of the algorithm can be summarized as follows:

Set  $m = \mu_{h+1}$ . We want to calculate the set of  $(\mu_h, r_h, d_h)$  with the following equality:

$$(E1) \quad \mu_{h+1} = \mu_h d_h + (r_h - 1)(d_h - 1)$$

Recall that we have the following restrictions:

i)  $d_h \geq 2$

ii)  $r_h \geq \frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3}$ .

iii)  $\text{gcd}(r_h, d_h) = 1$ .

iv)  $\mu_h \geq \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}$

v)  $\mu_1 = 0$ , and for all  $h \geq 2$ ,  $\mu_h = \frac{\mu_{h+1} - (d_h - 1)(r_h - 1)}{d_h}$  is an even integer.

Now equality (E1) gives  $(d_h - 1)(r_h - 1) = \mu_{h+1} - \mu_h d_h$ , and by iv)

$$-\mu_h d_h \leq -\left[ \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3} \right] d_h$$

in particular

$$(r_h - 1)(d_h - 1) \leq \mu_{h+1} - \left[ \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3} \right] \cdot d_h$$

This gives us the following upper bound for  $r_h$ :

$$r_h \leq \frac{\mu_{h+1}}{d_h - 1} - \left[ \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3} \right] \cdot \frac{d_h}{d_h - 1} + 1$$

**Corollary 4.4** *The above equality with (ii) give:*

$$(E2) \quad \frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3} \leq r_h \leq \frac{\mu_{h+1}}{d_h - 1} - \left[ \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3} \right] \cdot \frac{d_h}{d_h - 1} + 1$$

In the following we shall refine the lower bound of Corollary 4.4. We start with the following technical Lemma:

**Lemma 4.5**  $\sum_{i=1}^h (e_i - 1)r_i = r_h d_h - m_h$ . In particular,  $\mu_{h+1} = \sum_{i=1}^h (e_i - 1)r_i - r_0 + 1 = r_h d_h - m_h - r_0 + 1$ , where we recall that  $e_i = \frac{d_i}{d_{i+1}}$  for all  $1 \leq i \leq h$ .

*Proof.* Applying identity (\*\*\*) of Section 3 with  $k = h$  we get:

$$\sum_{i=1}^{h-1} (e_i - 1)r_i = r_h - m_h$$

Now adding  $(e_h - 1)r_h = (d_h - 1)r_h$  to the equality we get our assertion.

Lemma 4.5. with equality (E1) imply that  $r_h d_h = \mu_{h+1} + m_h + r_0 - 1$ . On the other hand, we have, with the notation  $m_0 = r_0$ , that for all  $1 \leq k \leq h$ ,  $m_k - m_{k-1} \geq d_{k+1}$ . Adding these inequalities we get

$$\mu_h \geq m_0 + d_2 + \dots + d_h + d_{h+1} = d_1 + d_2 + \dots + d_h + 1.$$

But for all  $1 \leq k \leq h$ ,  $d_k \geq 2^{h-k} \cdot d_h$ , so  $\mu_h \geq d_h \cdot (2^h - 1) + 1$ . Since  $r_0 = d_1 \geq 2^{h-1} \cdot d_h$ , we get the following:

$$r_h \geq \max\left(\frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3}, \frac{\mu_{h+1}}{d_h} + (3 \cdot 2^{h-1} - 1)\right)$$

Now equality (E1) implies that  $\frac{\mu_{h+1}}{d_h} = \mu_h + (r_h - 1)\left(1 - \frac{1}{d_h}\right)$ . But  $d_h \geq 2$ , thus, using inequalities of Proposition 4.1. we get:

$$\frac{\mu_{h+1}}{d_h} \geq \left(\frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}\right) + \left(\frac{5}{3} \cdot 2^{2h-1} - \frac{4}{3}\right) \cdot \frac{1}{2} = \frac{5}{3} \cdot 2^{2h-1} - 3 \cdot 2^{h-1} + \frac{2}{3}$$

In particular  $\max(\frac{5}{3} \cdot 2^{2h-1} - \frac{1}{3}, \frac{\mu_{h+1}}{d_h} + (3 \cdot 2^{h-1} - 1)) = \frac{\mu_{h+1}}{d_h} + (3 \cdot 2^{h-1} - 1)$ . This implies the following:

$$(E3) \quad \frac{\mu_{h+1}}{d_h} + (3 \cdot 2^{h-1} - 1) \leq r_h \leq \frac{\mu_{h+1}}{d_h - 1} - [\frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}] \cdot \frac{d_h}{d_h - 1} + 1$$

We shall now use inequality (E3) in order to give an upper bound for  $d_h$  (a lower bound being 2). Remark to this end that  $\frac{\mu_{h+1}}{d_h - 1} - [\frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}] \cdot \frac{d_h}{d_h - 1} + 1 - (\frac{\mu_{h+1}}{d_h} + (3 \cdot 2^{h-1} - 1)) \geq 0$ . If we set  $p = (\frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3})$  and  $q = 3 \cdot 2^{h-1} - 2$ , then an obvious analysis of the above inequality shows that it is equivalent to say that  $(p + q) \cdot d_h^2 - qd_h - \mu_{h+1} \leq 0$ , which is true if and only if the following holds:

$$(E4) \quad 2 \leq d_h \leq \frac{q + \sqrt{q^2 + 4\mu_{h+1} \cdot (p + q)}}{2 \cdot (p + q)} = \frac{3 \cdot 2^{h-1} - 2 + \sqrt{(3 \cdot 2^{h-1} - 2)^2 + 4\mu_{h+1} \cdot (\frac{5}{3} \cdot 2^{2h-2} - \frac{2}{3})}}{\frac{10}{3} \cdot 2^{2h-2} - \frac{4}{3}}$$

**The algorithm:** The two integers  $\mu_{h+1}$  and  $h$  being fixed, inequality (E4) determines the set  $D^h$  of possible values of  $d_h$ . Each value of  $d_h$  gives rise, using inequality (E3), to a set -denoted  $R_{d_h}^h$  - of possible values of  $r_h$  (Remark that  $\frac{\mu_{h+1} - (d_h - 1)(r_h - 1)}{d_h}$  should be an even integer).

We get this way the set -denoted  $P_{d_h}^h$  - of possible values of  $(\mu_h, r_h, d_h)$ . Now we restart with the set of  $\mu_h \dots$ . This process shall stop constructing a set of lists of length  $h$ . The set of semigroups corresponding to  $\mu_{h+1}$  is a subset of this list and can be easily calculated. Remark that if  $h = 1$ , then  $\mu_1 = 0$  and  $\mu_2 = (r_1 - 1)(d_1 - 1)$  by condition v). In this case, the values of  $(r_1, d_1 = r_0)$  can also be obtained from the set of divisors of  $\mu_2$ .

**Example 4.6** We perform an explicit computation for  $\mu_{h+1} = 28$ . In this case,

$$M = \frac{9 + \sqrt{1 + 60 \cdot 28}}{10} = 5, \text{ so } H = \{h; 1 \leq h \leq \frac{\ln(5)}{\ln(2)}\} = \{1, 2\}.$$

1)  $h = 1$ : In this case, since  $28 = 1 * 28 = 2 * 14 = 4 * 7$ , then  $(r_1, d_1) \in \{(2, 29), (3, 15), (5, 8)\}$  and condition iii) eliminates  $(3, 15)$ . We get this way the semigroups  $\langle 2, 29 \rangle$  and  $\langle 5, 8 \rangle$ . The canonical representative of the equisingularity class of the first one (resp. the second one) is  $y^2 - x^{29}$  (resp.  $y^5 - x^8$ ).

2)  $h = 2$ : Inequality (E4) implies in this case that  $2 \leq d_2 \leq \frac{4 + \sqrt{688}}{12} < 3$ . In particular  $D^2 = \{2\}$ .

Now inequality (E3) implies that  $\frac{28}{2} + 5 = 19 \leq r_2 \leq 28 - 4 + 1 = 25$ , and with conditions iii), v), we get  $R_2^2 = \{21, 25\}$ . If  $r_2 = 25$  (resp.  $r_2 = 21$ ), then  $\mu_2 = 2$  (resp.  $\mu_2 = 4$ ). Thus  $P_2^2 = \{(2, 25, 2), (4, 21, 2)\}$ .



i)  $(\mu_2, r_2, d_2) = (2, 25, 2)$ . In this case, applying to construction above to  $\mu_2 = 2$ , we get  $\frac{d_1}{d_2} = 2, \frac{r_1}{d_2} = 3$ . This leads to the semigroup  $\langle 4, 6, 25 \rangle$ . The canonical representative of the equisingularity class of this semigroupe is  $(y^2 - x^3)^2 - x^{11}y$ .

ii)  $(\mu_2, r_2, d_2) = (4, 21, 2)$ . In this case, applying to construction above to  $\mu_2 = 4$ , we get  $\frac{d_1}{d_2} = 2, \frac{r_1}{d_2} = 5$ . This leads to the semigroup  $\langle 4, 10, 21 \rangle$ . The canonical representative of the equisingularity class of this semigroupe is  $(y^2 - x^5)^2 - x^8y$ .

Let  $m$  be an even integer, and let  $H$  is the set of positive integers not exceeding  $\frac{\ln(M)}{\ln(2)}$ , where

$M = \frac{9 + \sqrt{1 + 60m}}{10}$ . Assume that  $H$  is not reduced to 0 and let  $h$  be a nonzero element of  $H$ . Set  $m = \mu_{h+1}$  and let

$$a_h = \frac{q + \sqrt{q^2 + 4\mu_{h+1} \cdot (p + q)}}{2 \cdot (p + q)} = \frac{3 \cdot 2^{h-1} - 2 + \sqrt{(3 \cdot 2^{h-1} - 2)^2 + 4\mu_{h+1} \cdot (\frac{5}{3} \cdot 2^{2h-2} - \frac{2}{3})}}{\frac{10}{3} \cdot 2^{2h-2} - \frac{4}{3}}.$$

Let  $D^h$  be the set positive integers between 2 and  $a_h$  (we easily verify that the condition  $a_h \geq 2$  is equivalent to the numerical condition  $\mu_{h+1} \geq \frac{5}{3} \cdot 2^{2h} - 3 \cdot 2^h + \frac{1}{3}$  proved in Proposition 4.1., in particular  $D^h$  is not the emptyset). Set  $P^h = \cup_{d \in D^h} P_d^h$  and denote by  $C_{h+1}$  the cardinality of  $P^h$ . In the following we shall give an upper bound for  $C_{h+1}$ . Set  $b_d^h = \frac{\mu_{h+1}}{d} + (3 \cdot 2^{h-1} - 1)$ , and  $c_d^h = \frac{\mu_{h+1}}{d-1} - [\frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{4}{3}] \cdot \frac{d}{d-1} + 1$ . The set  $R_d^h$  of possible values of  $r_h$  is a subset of the set of integers between  $b_d^h$  and  $c_d^h$ . Its cardinality is then bounded by  $c_d^h - b_d^h + 1$ . Furthermore, we easily verify that if  $r \in R_d^h$ , then

$$\frac{\mu_{h+1} - (r-1)(d-1)}{d} \geq \frac{5}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} + \frac{1}{3}.$$

In particular, if  $h \geq 2$ , then  $(\mu_h = \frac{\mu_{h+1} - (r-1)(d-1)}{d}, r, d)$  is an element of  $P_d^h$ .

Now

$$\begin{aligned} c_d^h - b_d^h + 1 &= \frac{\mu_{h+1}}{d(d-1)} - (p_h - q_h) \cdot \frac{d}{d-1} - q_h + 1 \\ &= \frac{\mu_{h+1}}{d-1} - \frac{\mu_{h+1}}{d} - (p_h - q_h) \cdot (1 + \frac{1}{d-1}) - q_h + 1 \end{aligned}$$

Consequently, if  $a = [a_h]$ , then the cardinality  $C_{h+1}$  of  $P_h$  is bounded by:

$$\sum_{d=2}^a (c_d^h - b_d^h + 1) = (\mu_{h+1})\left(1 - \frac{1}{a}\right) - p_h(a-1) - (p_h - q_h) \cdot \sum_{d=1}^{a-1} \frac{1}{d} + (a-1)$$

But  $1 - \frac{1}{a} < 1$ , and substituting 2 to  $a$  in the other members of the above formula we get:

$$C_{h+1} \leq \mu_{h+1} - 2p_h + q_h + 1 = \mu_{h+1} - \left(\frac{10}{3} \cdot 2^{2h-2} - 3 \cdot 2^{h-1} - \frac{1}{3}\right) = \mu_{h+1} - \left(\frac{B_{h+1}}{2} - 1\right)$$

Where  $B_{h+1} = \frac{5}{3} \cdot 2^{2h} - 3 \cdot 2^h + \frac{4}{3}$  is the lower bound of  $\mu_{h+1}$  in Proposition 4.1.

**Remark 4.7** Note that the bound above is not the optimal one, indeed, given  $d \in D^h$ , the cardinality of the set of  $r \in R_d^h$  such that  $\gcd(r, d) = 1$  can be bounded by  $\frac{c_d^h - b_d^h}{d} + 1$ , but in view of our algorithm, all values of  $R_d^h$  are used, in particular the value above bounds also the nombre of operations used in the first step of the algorithm.

Let  $(\mu_h, r, d)$  be an element of  $P_h$  and recall that  $\mu_h = \frac{\mu_{h+1}}{d} - (r-1)\frac{d-1}{d}$ . Since  $b_d^h \leq r \leq c_d^h$ , then

$$\begin{aligned} \mu_h &\leq \frac{\mu_{h+1}}{d} - (b_d^h - 1)\frac{d-1}{d} \leq \frac{\mu_{h+1}}{d} - (b_d^h - 1)\frac{d-1}{d} = \frac{b_d^h - 1}{d} - (3 \cdot 2^{h-1} - 2) \\ &= \frac{\mu_{h+1}}{d^2} + \left(\frac{1}{d} - 1\right)(3 \cdot 2^{h-1} - 2) \leq \frac{\mu_{h+1}}{4} - \frac{1}{2}(3 \cdot 2^{h-1} - 2) \leq \frac{\mu_{h+1}}{4} - (3 \cdot 2^{h-2} - 1) \end{aligned}$$

Let  $A_{h+1} = 3 \cdot 2^{h-2} - 1$ . It follows by induction that for all  $0 \leq k \leq h-1$ , if  $\mu_{h-k}$  is a possible value of the Milnor number at the step  $k+1$ , then we have:

$$\mu_{h-k} \leq \frac{\mu_{h+1}}{4^{k+1}} - \sum_{i=0}^k \frac{A_{h+1-i}}{4^{k-i}} = 3 \cdot 2^{h-2k-2}(2^{k+1} - 1) - \frac{4}{3} - \frac{1}{3 \cdot 4^k}$$

(Remark that the above inequality is valid if  $k = h-1$  because  $\mu_1 = 0$ ). Thus, we can obtain a bound of the set of values calculated at the step  $k+1$ ,  $0 \leq k \leq h-2$  in the following way: let  $\mu_{h-k}$  be a possible value of the Milnor number obtained by reiterating the algorithm above  $k+1$  times, and denote by  $C_{h-k}(\mu_{h-k})$  the cardinality of the set -denoted  $P_{h-k-1}(\mu_{h-k})$ - of the 3-uplets  $(\mu, r, d)$  obtained by applying the algorithm above to  $\mu_{h-k}$  instead of  $\mu_{h+1}$ . It follows from the discussion above that

$$C_{h-k} \leq \mu_{h-k} - \frac{B_{h-k}}{2} + 1 \leq \frac{\mu_{h+1}}{4^{k+1}} - \sum_{i=0}^k \frac{A_{h+1-i}}{4^{k-i}} - \frac{B_{h-k}}{2} + 1$$

$$= \frac{\mu_{h+1}}{4^{k+1}} - 3 \cdot 2^{h-k-2} + 3 \cdot 2^{h-2k-2} - \frac{5}{3} \cdot 2^{2h-2k-1} - \frac{1}{3 \cdot 4^k} + \frac{5}{3}$$

In particular, the cardinality of the set of semigroups corresponding to the given Milnor number  $m = \mu_{h+1}$  is bounded by  $\prod_{i=2}^h C_{h+1-i}$  which is a polynomial in  $m$  bounded by its leading coefficient  $\frac{m^h}{2^{h(h-1)}}$ . Note that in view of Remark 4.8., the number of operations used in the algorithm is then bounded by  $\sum_{i=0}^{h-1} \prod_{k=0}^i C_{h+1-k}$ .

The above algorithm has been implemented with *MAPLE*. The input is an integer  $m$ , and the output is the list of semigroups whose conductor is  $m$ . In the implementation work we followed the ideas explained above, with the following simplification: at the last step, the set of values we are interested in is calculated by using the factorization of the given Milnor number. The algorithm is an iterating of the following:

*Input:*  $m \in 2 \cdot \mathbf{N}$

*Output:* The set  $P^h$ .

*Step I:* Compute the set  $H$ .

*Step II:* Take  $h \in H$ .

*Step III:* Compute the set  $D^h$ .

*Step IV:* Take  $d \in D^h$ .

*Step V:* Compute  $R_d^h$

(\*) if  $r \in R_d^h$  and  $\gcd(r, d) = 1$  and  $\frac{m - (d-1)(r-1)}{d} \in 2 \cdot \mathbf{N}$  (resp.  $(d-1)(r-1) = m$  if  $h=1$ ) then add  $(\frac{m - (d-1)(r-1)}{d}, r, d)$  to  $P_d^h$ .

*Step VI:*  $P^h = \bigcup_{d \in D^h} P_d^h$

The main operation of the algorithm is the one described in the line (\*). We experimented it on various values of  $m$ : the computation takes around 0.2sec for  $m = 160$ , 0.7sec for  $m = 300$ , 1.5sec for  $m = 500$ , and 3sec for  $m = 1000$ .

## References

- [1] S.S. Abhyankar.- Lectures on expansion techniques in Algebraic Geometry, Tata Institute of Fundamental research, Bombay, 1977.
- [2] S.S. Abhyankar.- On the semigroup of a meromorphic curve, Part 1, in Proceedings of International Symposium on Algebraic Geometry, Kyoto, pp. 240-414, 1977.
- [3] S.S. Abhyankar.- Irreducibility criterion for germs of analytic functions of two complex variables, Advances in Mathematics 74, pp. 190-257, 1989.

- [4] S.S. Abhyankar.- Some remarks on the Jacobian problem, Proc.Indian Acad.Sci., vol 104, n<sup>o</sup> 3, pp. 515-542, 1994.
- [5] S.S. Abhyankar and T.T. Moh.- Newton Puiseux expansion and generalized Tschirnhausen transformation, J.Reine Angew.Math, 260, pp. 47-83 and 261, pp. 29-54, 1973.
- [6] A. Assi.-Deux remarques sur les racines approchées d'Abhyankar-Moh, C.R.A.S., t.319, Serie 1, 1994, 1191-1196.
- [7] A. Assi.-Meromorphic plane curves, Math.Z., vol 230, 1999, 165-183.
- [8] A. Assi, M. Barile.- Computing irreducible curve singularities with Mathematica, in: L. GonzalzVega, T. Recio (eds.) Actas del 9 Encuentro de Algebra Computacional y Aplicaciones, Universidad de Cantabria, Santander, 1-3 julio 2004. pp. 17-21.
- [9] A. Campillo.- Algebroid curves in positive characteristic, Lect. Not. Math., 813. Springer-Verlag, 1980.
- [10] V. Cossart, G. Moreno-Socías.- Racines approchées et suffisance des jets, To appear in Annales de Toulouse.
- [11] V. Cossart, G. Moreno-Socías.- Irreducibility criterion: a geometric point of view, in Valuation theory and its applications, Fields Institute Communications Series, AMS, 2002.
- [12] A. Granja.- Irreducible polynomials with coefficients in a complete discrete valuation field, Advances in Mathematics 109, n 1, pp. 75-87, 1994.
- [13] O. Zariski.-Le problème des modules pour les branches planes, Hermann, 1986.