

q – Catalan numbers and q – Narayana polynomials

Johann Cigler

Fakultät für Mathematik
Universität Wien
A-1090 Wien, Nordbergstraße 15
e-mail: johann.cigler@univie.ac.at

The q – Catalan numbers

$1, 1, 1 + q, 1 + 2q + q^2 + q^3, 1 + 3q + 3q^2 + 3q^3 + 2q^4 + q^5 + q^6, \dots$,

which were invented by Carlitz and Riordan ([2], cf. also [7] and [10]), are defined by

$$C_n(q) = \sum_{k=0}^{n-1} q^k C_k(q) C_{n-k-1}(q) \text{ with initial value } C_0(q) = 1.$$

Let $f(z, q) = \sum_{k \geq 0} C_k(q) z^k$ be their generating function, which is uniquely determined by

the functional equation $f(z, q) = 1 + zf(z, q)f(qz, q)$. This implies the well known fact that it can be represented in the form

$$f(z, q) = \frac{E_2(-qz)}{E_2(-z)}, \quad (1)$$

where $E_r(z)$ denotes the generalized q – exponential function

$$E_r(z) = \sum_{k \geq 0} q^{\binom{k}{2}} \frac{z^k}{(1-q)(1-q^2) \cdots (1-q^k)}. \quad (2)$$

In this note we want to sketch some extensions of this interesting result.

a) First observe that

$$E_r(z) - E_r(qz) = zE_r(q^r z). \quad (3)$$

For $n \in \mathbb{N}$ define

$$G_r(z, n) = \sum_{k \geq 0} G(k, n, r) z^k := \frac{E_r(-q^n z)}{E_r(-z)}. \quad (4)$$

Then we have

$$G_r(z, n+1) = G_r(z, n) + q^n z G_r(z, n+r). \quad (5)$$

Comparing coefficients we get

$$\frac{G(k, n+1, r) - G(k, n, r)}{q^n} = G(k-1, n+r, r) \quad (6)$$

with $G(k, 0, r) = [k = 0]$ and $G(0, n, r) = 1$.

This implies

$$G_r(z, 1) = 1 + zG_r(z, r). \quad (7)$$

These are the characteristic properties of the q – Gould polynomials (cf. [4]).

For $q = 1$ they have the explicit formula $G(k, n, r) = \frac{n}{n + rk} \binom{n + rk}{k}$ (cf. e.g. [9]).

For general q special values are $G(k, n, 0) = q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}$ and $G(k, n, 1) = \begin{bmatrix} n + k - 1 \\ k \end{bmatrix}$, where $\begin{bmatrix} n \\ k \end{bmatrix}$

denotes a q – binomial coefficient. For $r > 1$ no explicit formulas are known.

Note that $G_2(z, 1) = 1 + zG_2(z, 2) = 1 + zG_2(z, 1)G_2(qz, 1) = f(z, q)$ is the generating function of the q – Catalan numbers.

$$\text{From } \frac{E_r(-q^n z)}{E_r(-z)} = \frac{E_r(-qz)}{E_r(-z)} \frac{E_r(-q^2 z)}{E_r(-qz)} \dots \frac{E_r(-q^n z)}{E_r(-q^{n-1} z)}$$

we get

$$G_r(z, n) = G_r(z, 1)G_r(qz, 1) \dots G_r(q^{n-1}z, 1) \quad (8)$$

and

$$G_r(z, m + n) = G_r(z, m)G_r(q^m z, n). \quad (9)$$

b) Let now $(a \dot{+} b)^k := (a + b)(a + qb) \dots (a + q^{k-1}b)$

and consider the modified exponential function

$$h(z, a, b, q) = \sum_{k \geq 0} q^{\binom{k}{2}} \frac{(a \dot{+} b)^k}{(1 \dot{-} q)^k} (-z)^k, \quad (10)$$

which for $(a, b) = (0, 1)$ reduces to $E_2(-z)$.

We want to study the series

$$f(z, a, b, q) = \frac{h(qz, a, b, q)}{h(z, a, b, q)}. \quad (11)$$

From

$$\begin{aligned} h(z, a, b, q) - h(qz, a, b, q) &= \sum_{k \geq 0} q^{\binom{k}{2}} \frac{(a \dot{+} b)^k}{(1 \dot{-} q)^k} (-z)^k (1 - q^k) = \\ &= -z \sum_{k \geq 0} q^{\binom{k-1}{2}} \frac{(a \dot{+} b)^{k-1} (a + q^{k-1}b)}{(1 \dot{-} q)^{k-1}} (-qz)^{k-1} = -azh(qz, a, b, q) - bzh(q^2 z, a, b, q) \end{aligned}$$

we deduce that

$$f(z, a, b, q) = 1 + azf(z, a, b, q) + bzf(z, a, b, q)f(qz, a, b, q). \quad (12)$$

If we write

$$f(z, a, b, q) = \sum_{n \geq 0} C_n(a, b, q) z^n, \quad (13)$$

then $C_n(a, b, q)$ is a q -analogue of $C_n(a, b) = \frac{1}{n} \sum_{k=1}^n \binom{n}{k} \binom{n}{k-1} b^{n-k} (a+b)^k$.

For it is easy to verify that the uniquely determined formal power series $f(z, a, b)$, which satisfies the functional equation $f(z, a, b) = 1 + azf(z, a, b) + bzf(z, a, b)^2$ has the series expansion

$$f(z, a, b) = 1 + \sum_{n \geq 1} \sum_{k=1}^n \frac{1}{n} \binom{n}{k} \binom{n}{k-1} b^{n-k} (a+b)^k z^n, \quad (14)$$

Of course $C_n(0, 1) = \frac{1}{n+1} \binom{2n}{n} = C_n$ are the well known Catalan numbers.

The numbers $N(n, k) = \frac{1}{n} \binom{n}{k} \binom{n}{k-1}$ are called Narayana numbers. (cf. e.g. [10]).

Therefore we call $C_n(a, b, q)$ a q -Narayana polynomial. It satisfies

$$C_n(a, b, q) = aC_{n-1}(a, b, q) + b \sum_{k=0}^{n-1} q^k C_k(a, b, q) C_{n-k-1}(a, b, q) \quad (15)$$

with initial value $C_0(a, b, q) = 1$.

The first values are $1, a+b, a^2 + (2+q)ab + (1+q)b^2, \dots$.

From the definition is clear that $C_n(a, b, q)$ is a polynomial in a, b which is homogeneous of degree n . Therefore $C_n(a, b, q)$ has a unique representation in the form

$$C_n(a, b, q) = \sum_{k=0}^n N(n, k, q) (a+b)^k b^{n-k}. \quad (16)$$

There does not seem to be a simple formula for $N(n, k, q)$. All we can say is that they are

polynomials in q with integer coefficients and that $N(n, k, 1) = N(n, k) = \frac{1}{n} \binom{n}{k} \binom{n}{k-1}$.

The first values of these q -Narayana numbers are given in the following table:

1						
0	1					
0	1	1				
0	1	$2+q$	1			
0	1	$3+3q$	$3+2q+q^2$	1		
0	1	$4+6q+q^2-q^3$	$6+8q+5q^2+q^3$	$4+3q+2q^2+q^3$	1	

E.g. we get

$$C_0(a, b, q) = 1,$$

$$C_1(a, b, q) = a + b,$$

$$C_2(a, b, q) = (a+b)b + (a+b)^2,$$

$$C_3(a, b, q) = (a+b)b^2 + (2+q)(a+b)^2b + (a+b)^3.$$

It is clear that $C_n(a, b, 0) = (a + b)^n$. Therefore we have $N(n, k, 0) = \binom{n-1}{k-1}$.

We shall need also another relation:

$$\begin{aligned} h(z, a, b, q) - h(qz, a, b, q) &= \sum_{k \geq 0} q^{\binom{k}{2}} \frac{(a + b)^k}{(1 - q)^k} (-z)^k (1 - q^k) = \\ &= -(a + b)z \sum_{k \geq 0} q^{\binom{k-1}{2}} \frac{(a + qb)^{k-1}}{(1 - q)^{k-1}} (-qz)^{k-1} = -(a + b)zh(qz, a, qb, q). \end{aligned}$$

This implies

$$\frac{f(qz, a, b, q) - 1}{f(z, a, b, q) - 1} = \frac{h(q^2z, a, b, q) - h(qz, a, b, q)}{h(qz, a, b, q) - h(z, a, b, q)} = q \frac{f(qz, a, qb, q)}{f(z, a, b, q)}. \quad (17)$$

It is well known (cf. e.g. [10]) that the Catalan numbers $C_n = \frac{1}{n+1} \binom{2n}{n}$ can be

characterized by the values of their Hankel determinants. They satisfy $\det(C_{i+j})_{i,j=0}^n = 1$ and $\det(C_{i+j+1})_{i,j=0}^n = 1$.

We want to show that the q -Narayana polynomials can also be characterized by the values of their Hankel determinants. This shows that these polynomials are in some sense a natural generalization of the q -Catalan numbers.

Theorem

The polynomials $C_n(a, b, q)$ are characterized by their Hankel determinants

$$\det(C_{i+j}(a, b, q))_{i,j=0}^n = q^{\frac{n^2(n+1)}{2}} b^{\binom{n+1}{2}} (a + b)^n (a + qb)^{n-1} (a + q^2b)^{n-2} \cdots (a + q^{n-1}b) \quad (18)$$

and

$$\det(C_{i+j+1}(a, b, q))_{i,j=0}^n = q^{\frac{n(n+1)^2}{2}} b^{\binom{n+1}{2}} (a + b)^{n+1} (a + qb)^n (a + q^2b)^{n-1} \cdots (a + q^n b). \quad (19)$$

Remark

For the q -Catalan numbers this reduces to

$$\det(C_{i+j}(q))_{i,j=0}^n = q^{\frac{n(n+1)(4n-1)}{6}} \quad \text{and} \quad \det(C_{i+j+1}(q))_{i,j=0}^n = q^{\frac{n(n+1)(4n+5)}{6}}.$$

These special cases have been proved by another method in [5].

In order to prove the theorem we need the following well-known (cf. e.g. [1] or [8])

Lemma

Let

$$\sum_{k \geq 0} \mu_k z^k = \frac{1}{1 - s_0 z - \frac{t_0 z^2}{1 - s_1 z - \frac{t_1 z^2}{1 - \dots}}} = \frac{1}{1 - s_0 z} \frac{t_0 z^2}{1 - s_1 z} \frac{t_1 z^2}{1 - s_2 z} \dots. \quad (20)$$

Then the Hankel determinants have the following values

$$\det(\mu_{i+j})_{i,j=0}^n = t_0^n t_1^{n-1} \dots t_{n-1}$$

and

$$\det(\mu_{i+j+1})_{i,j=0}^n = d_{n+1} t_0^n t_1^{n-1} \dots t_{n-1},$$

where $d_n = s_{n-1} d_{n-1} - t_{n-2} d_{n-2}$, $d_0 = 1$, $d_1 = s_0$.

Let F be the linear functional defined by $F(z^n) = \mu_n$.

Then the polynomials $p_n(z)$, defined by

$$p_0(z) = 1, p_1(z) = z - s_0,$$

and $p_k(z) = (z - s_{k-1})p_{k-1}(z) - t_{k-2}p_{k-2}(z)$

satisfy $F(p_n p_m) = t_0 \dots t_{n-1} [n = m]$,

i.e. are orthogonal with respect to the linear functional F .

Proof of the Theorem

We set

$$f(z, a, b, q) = 1 + (a + b)z g\left(\frac{z}{q}, a, b, q\right). \quad (21)$$

Then we can write (12)

in the form

$$f(z, a, b, q) = 1 + (a + b)z f(z, a, b, q) + q(a + b)bz^2 f(z, a, b, q)g(z, a, b, q), \quad (22)$$

where the series $g(z, a, b, q)$ satisfies

$$g(z, a, b, q) = 1 + q(a + b)z g(z, a, b, q) + q^2 bz g(qz, a, b, q) + q^3 b(a + b)z^2 g(z, a, b, q)g(qz, a, b, q).$$

We want to show that this is equivalent with

$$g(z, a, b, q) = 1 + q(a + b + qb)z g(z, a, b, q) + q^4 b(a + qb)z^2 g(z, a, b, q)g(qz, a, qb, q). \quad (23)$$

This means that

$$\begin{aligned} & q^2 bz g(qz, a, b, q) + q^3 b(a + b)z^2 g(z, a, b, q)g(qz, a, b, q) = \\ & = qb z g(z, a, b, q) + q^4 b(a + qb)z^2 g(z, a, b, q)g(qz, a, qb, q) \end{aligned}$$

or

$$g(qz, a, b, q)(1 + q(a + b)z g(z, a, b, q)) = g(z, a, b, q)(1 + (a + qb)q^2 z g(qz, a, qb, q)).$$

By the definition of $g(z, a, b, q)$ this reduces to

$$g(qz, a, b, q)f(qz, a, b, q) = g(z, a, b, q)f(q^2z, a, qb, q),$$

which is equivalent with

$$\frac{g(qz, a, b, q)}{g(z, a, b, q)} = \frac{1}{q} \frac{(a+b)q^2zg(qz, a, b, q)}{(a+b)qzg(z, a, b, q)} = \frac{f(q^2z, a, qb, q)}{f(qz, a, b, q)}.$$

But this is an immediate consequence of (17).

Now (23) is equivalent with

$$g(z, a, b, q) = \frac{1}{1 - s_1z - t_1z^2g(qz, a, qb, q)}$$

with $s_1 = q(a + b + qb)$ and $t_1 = q^4b(a + qb)$.

This gives us a representation of $g(z, a, b, q)$ as a continued fraction of the form (20) with $s_n = q^n(a + q^{n-1}b + q^n b)$ and $t_n = q^{3n+1}b(q^n b + a)$.

From (22) we conclude that $f(z, a, b, q)$ has also a representation as a continued fraction of the form (20) with the same s_n, t_n together with $s_0 = a + b$ and $t_0 = q(a + b)b$.

From this the theorem immediately follows.

It is clear that the sequence $(C_n(a, b, q))$ is uniquely determined by these determinants.

Remark: *It can be shown that the corresponding orthogonal polynomials are*

$$p_n(z, a, b, q) = \sum_{k=0}^n (-1)^{n-k} q^{\binom{n-k}{2}} z^k \sum_{j=k}^n q^{\binom{n+1}{2} - \binom{n+k+1-j}{2}} \begin{bmatrix} n+k-j \\ k \end{bmatrix} \begin{bmatrix} j-1 \\ k-1 \end{bmatrix} b^{j-k} (a+b)^{n-j}$$

and

$$p_n(z, 0, 1, q) = \sum_{k=0}^n (-1)^{n-k} q^{\binom{n-k}{2}} \begin{bmatrix} n+k \\ 2k \end{bmatrix} z^k.$$

But we don't need this result.

c) Let now

$$h^*(z, a, b, q) := \sum_{k \geq 0} q^{\binom{k}{2}} \frac{r_k(a, b)}{(1-q)^k} (-z)^k, \quad (24)$$

where $r_n(a, b, q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} a^k b^{n-k}$ is a Rogers-Szegö polynomial, which satisfies

$$r_n(a, b, q) = (a+b)r_{n-1}(a, b, q) + ab(q^{n-1} - 1)r_{n-2}(a, b, q) \quad (\text{cf. e.g. [3]}).$$

Therefore we get

$$h^*(qz, a, b, q) - h^*(z, a, b, q) = -z(a+b)h^*(qz, a, b, q) - qabz^2h^*(q^2z, a, b, q).$$

If we define

$$f^*(z, a, b, q) = \frac{h^*(qz, a, b, q)}{h^*(z, a, b, q)}, \quad (25)$$

we see that $f^*(z, a, b, q)$ satisfies the functional equation

$$f^*(z, a, b, q) = 1 + (a + b)zf^*(z, a, b, q) + qabz^2f^*(z, a, b, q)f^*(qz, a, b, q). \quad (26)$$

It is easy to see (compare (14) and (23)) that the series $f^*(z, a, b) = g(z, a - b, b, 1)$ satisfying $f^*(z, a, b) = 1 + (a + b)zf^*(z, a, b) + abz^2f^*(z, a, b)^2$ has the expansion

$$azf^*(z, a, b) = \sum_{n \geq 1} \left(\sum_{k=1}^n N(n, k) a^k b^{n-k} \right) z^n.$$

Therefore we write $af^*(z, a, b, q) = \sum_{n \geq 0} C_{n+1}^*(a, b, q) z^n$

with

$$C_n^*(a, b, q) = \sum_{k=1}^n N^*(n, k, q) a^k b^{n-k}. \quad (27)$$

Again we have $N^*(n, k, 0) = \binom{n-1}{k-1}$ and $D_n^*(1, 1, 0) = 2^{n-1}$.

The first values of $(N^*(n, k, q))_{k=1}^n, n \geq 1$, are

$$\begin{array}{ccccccc} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 2 + q & 1 & & & & \\ 1 & 3 + 2q + q^2 & 3 + 2q + q^2 & 1 & & & \\ 1 & 4 + 3q + 2q^2 + q^3 & 6 + 6q + 5q^2 + 2q^3 + q^4 & 4 + 3q + 2q^2 + q^3 & 1 & & \end{array}$$

Since $f^*(z, a, b) = f^*(z, b, a)$ we see that $N^*(n, k, q) = N^*(n, n - k + 1, q)$.

From this we get

$$C_1^*(a, b, q) = a$$

$$C_2^*(a, b, q) = ab + a^2,$$

$$C_3^*(a, b, q) = ab^2 + (2 + q)a^2b + a^3,$$

$$C_4^*(a, b, q) = ab^3 + (3 + 2q + q^2)a^2b^2 + (3 + 2q + q^2)a^3b + a^4, \dots$$

Let now $F(z, a, b, q) = 1 + azf^*(z, a, b, q)$. Then by (26)

$$F(z, a, b, q) = 1 + azF(z, a, b, q) - bzF(qz, a, b, q) + bzF(z, a, b, q)F(qz, a, b, q) \quad (28)$$

If we set $C_0^*(a, b, q) = 1$, then (28) implies the recurrence

$$C_n^*(a, b, q) = aC_{n-1}^*(a, b, q) + b \sum_{k=0}^{n-2} q^k C_k^*(a, b, q) C_{n-1-k}^*(a, b, q). \quad (29)$$

Comparing with [7] (5.5) we see that $C_n^*(1, s, q)$ are the Pólya-Gessel q – Catalan numbers, the first values of which are

$$1, 1, 1+s, 1+2s+qs+s^2, 1+3s+2qs+q^2s+3s^2+2qs^2+q^2s^2+s^3, \dots$$

By (26) the corresponding s_k, t_k are given by

$s_k = q^k(a+b)$ and $t_k = q^{2k+1}ab$. Therefore their Hankel determinants are

$$\det(C_{i+j+1}^*(a, b, q))_{i,j=0}^n = (ab)^{\binom{n+1}{2}} q^{\sum_{i=0}^n i^2} = (ab)^{\binom{n+1}{2}} q^{\frac{n(n+1)(2n+1)}{6}} \quad (30)$$

and

$$\det(C_{i+j+2}^*(a, b, q))_{i,j=0}^n = (abq)^{\binom{n+1}{2}} q^{\frac{n(n+1)(2n+1)}{6}} \frac{a^{n+2} - b^{n+2}}{a-b}. \quad (31)$$

It is easy to verify that the corresponding orthogonal polynomials are

$$p_n(z, a, b) = \sum_{k=0}^n (-1)^{n-k} q^{\binom{n-k}{2}} z^k \sum_{j=k}^n \begin{bmatrix} n+k-j \\ k \end{bmatrix} \begin{bmatrix} j \\ k \end{bmatrix} a^{j-k} b^{n-j}.$$

In order to compute

$$\det(C_{i+j}^*(a, b, q))_{i,j=0}^n$$

we observe that from (28) we get

$$\begin{aligned} F(z, a, b, q) &= \frac{1 - bzF(qz, a, b, q)}{1 - bzF(qz, a, b, q) - az} = \frac{1}{1 - \frac{az}{1 - bzF(qz, a, b, q)}} = \frac{1}{1 - 1 - 1 - 1 - 1} \dots = \\ &= \frac{1}{1 - azF(z, b, qa, q)} \end{aligned}$$

Therefore we have

$$F(z, a, b, q) = 1 + azF(z, a, b, q)F(z, b, qa, q)$$

or

$$F(z, a, b, q) = 1 + azF(z, a, b, q) + abz^2F(z, a, b, q)f^*(z, b, qa, q)$$

This gives us $t_k = q^{2k}ab$ for all $k \geq 0$ and therefore we have

$$\det(C_{i+j}^*(a, b, q))_{i,j=0}^n = (ab)^{\binom{n+1}{2}} q^{\frac{n(n+1)(n-1)}{3}}.$$

In this case the orthogonal polynomials are

$$p_n(z, a, b) = \sum_{k=0}^n (-1)^{n-k} q^{\binom{n-k}{2}} z^k \sum_{j=k}^n \begin{bmatrix} n+k-j \\ k \end{bmatrix} \begin{bmatrix} j-1 \\ j-k \end{bmatrix} b^{j-k} a^{n-j}.$$

For the special case $(a, b) = (1, s)$ these results have been proved by other methods in [6].

The generating function $f(z, q)$ of the q – Catalan numbers $C_n(q)$ has the continued fraction expansion

$$f(z, q) = \frac{1}{1 - zf(qz, q)} = \frac{1}{1 - \frac{z}{1 - qzf(q^2z, q)}} = \frac{1 - qzf(q^2z, q)}{1 - qzf(q^2z, q) - z}.$$

Therefore we get

$$f(z, q) = 1 + zf(z, q) - qzf(q^2z, q) + qzf(z, q)f(q^2z, q).$$

This implies $f(z, q) = F(z, 1, q, q^2)$ and thus the well known (cf. [7]) result that

$$C_n(q) = C_n^*(1, q, q^2).$$

This can also directly be seen:

$$h^*(z, 1, q, q^2) = \sum_{k=0}^n q^{2\binom{k}{2}} \left(\sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_{q^2} q^k \right) \frac{(-z)^k}{(1 - q^2)^k} = E_2(-z),$$

since

$$\sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_{q^2} q^k = (1 + q)^k \text{ (cf. [3]).}$$

From Gauss's formula $r_{2n+1}(1, -1) = 0, r_{2n}(1, -1) = (1 - q)(1 - q^3) \cdots (1 - q^{2n-1})$ (cf e.g.[3])

we conclude in the same way that

$$C_{2n+1}^*(1, -1, q) = (-1)^n q^n C_n(q^2)$$

and $C_{2n+2}^*(1, -1, q) = 0$.

d) Another interesting special case is given by the q - Motzkin numbers $M_n(q)$ which have been considered in [5]. Their generating function $M(z) = \sum_{n \geq 0} M_n(q)z^n$ satisfies

$$M(z) = 1 + zM(z) + qz^2M(z)M(qz). \text{ Therefore } M_n(q) = C_n^*\left(\frac{1 + \sqrt{-3}}{2}, \frac{1 - \sqrt{-3}}{2}, q\right).$$

The first values are

$$1, 1, 1 + q, 1 + 2q + q^2, 1 + 3q + 3q^2 + q^3 + q^4, \dots$$

The Hankel determinants are easily seen to be

$$\det(M_{i+j}(q))_{i,j=0}^n = q^{\frac{n(n+1)(2n+1)}{6}}$$

and

$$\det(M_{i+j+1}(q))_{i,j=0}^n = q^{\frac{n(n+1)(2n+1)}{6} + \binom{n+1}{2}} d_{n+1} = q^{2\binom{n+2}{3}} d_{n+1}.$$

Here $(d_n)_{n \geq 0} = (1, 1, 0, -1, -1, 0, 1, 1, 0, \dots)$ is periodic with period 6.

References

- [1] G.E. Andrews, R. Askey and R. Roy, Special Functions, Cambridge Univ. Press, 1999
- [2] L. Carlitz and J. Riordan, Two element lattice permutation numbers and their q -generalization, Duke J. Math. **31** (1964), 371-388
- [3] J. Cigler, Elementare q -Identitäten, Sem. Lotharingien Comb. B05a (1981)
- [4] J. Cigler, Operatormethoden für q -Identitäten IV: Eine Klasse von q -Gould-Polynomen, Sitzungsber. ÖAW **205** (1996), 169-174.
http://hw.oeaw.ac.at/sitzungsberichte_und_anzeiger
- [5] J. Cigler, q -Catalan- und q -Motzkinzahlen, Sitzungsber. ÖAW **208** (1999), 3-20
http://hw.oeaw.ac.at/sitzungsberichte_und_anzeiger
- [6] J. Cigler, Einige q -Analoge der Catalan-Zahlen, Sitzungsber. ÖAW **209** (2000), 19-46
http://hw.oeaw.ac.at/sitzungsberichte_und_anzeiger
- [7] J. Fürlinger and J. Hofbauer, q -Catalan numbers, J. Comb. Th. A **40** (1985), 248-264
- [8] C. Krattenthaler, Advanced Determinant Calculus: A Complement, ArXiv math.CO/0503507
- [9] G.-C. Rota, Finite Operator Calculus, Academic Press 1975
- [10] R. P. Stanley, Enumerative Combinatorics, Vol.2, Cambridge University Press 1999