CRAMER'S THEOREM FOR NONNEGATIVE MULTIVARIATE POINT PROCESSES WITH INDEPENDENT INCREMENTS

KLEBANER F. AND R. LIPTSER

ABSTRACT. We consider a continuous time version of Cramer's theorem with nonnegative summands $S_t = \frac{1}{t} \sum_{i:\tau_i \leq t} \xi_i, t \to \infty$, where $(\tau_i, \xi_i)_{i\geq 1}$ is a sequence of random variables such that tS_t is a random process with independent increments.

1. Introduction and main result

The following version of the Cramer theorem [1] can be extracted from Dembo and Zeitouni, [2].

Theorem 1. Let $(\xi_i)_{i\geq 1}$ be a sequence of nonnegative identically distributed and independent random variables with ξ_1 admitting the Laplace transform:

$$\mathscr{L}(\lambda) = \mathsf{E}e^{\lambda\xi_1}, \ \lambda \in (-\infty, \Lambda), \quad \exists \ \Lambda = \inf\{\lambda > 0 : \mathscr{L}(\lambda) = \infty\}.$$

Then, the family

$$S_n = \frac{1}{n} \sum_{i=1}^n \xi_i, \quad n \to \infty$$

obeys the Large Deviation principle (LDP) in the metric space (\mathbb{R}_+, ϱ) (for the Euclidean metric ϱ) with the rate $\frac{1}{n}$ and the rate function

$$I(u) = \begin{cases} \sup_{\lambda \in (-\infty,\Lambda)} [\lambda u - g(\lambda)], & u > 0\\ -\log \mathsf{P}(\xi_1 = 0), & u = 0, \end{cases}$$

where $g(\lambda)$ is the log moment generation function,

$$g(\lambda) = \log \mathsf{E} e^{\lambda \xi_1}, \ \lambda < \Lambda.$$

In this paper, we study a "continuous time version" of this theorem. For $t \ge 0$, set

$$S_t = \frac{1}{t} \sum_{i:\tau_i \le t} \xi_i,$$

Date: October 17, 2006.

¹⁹⁹¹ Mathematics Subject Classification. 60F10, 60J27.

Key words and phrases. Nonnegative summands, boundary effect.

where $(\xi_i, \tau_i)_{i \ge 1}$ is a sequence of random pairs, where ξ_i 's and τ_i 's are random variables:

$$\xi_i \ge 0$$
 and $\tau_0 = 0 < \tau_1 < \tau_2 < \dots \tau_i < \dots$

defined on a probability space $(\Omega, \mathcal{F}, \mathsf{P})$. Let $(\mathscr{G}_n)_{n\geq 0}$ be the filtration with $\mathscr{G}_0 = (\emptyset, \Omega)$ and $\mathscr{G}_n := \sigma\{(\tau_i, \xi_i)_{i\leq n}\}$. Random variables ξ_i are assumed to be identically distributed and independent of \mathscr{G}_{i-1} with the distribution function

$$G(x) = \mathsf{P}(\xi_1 \le x), \ x \ge 0.$$

The conditional distribution of τ_i given \mathscr{G}_{i-1} is exponential:

$$\mathsf{P}\big(\tau_i \le t | \mathscr{G}_{i-1}\big) = \big(1 - e^{-r(t - \tau_{i-1})}\big), \ t \ge \tau_{i-1},$$

where r is a positive number. Moreover, we assume that

$$\mathsf{P}(\xi_i \le x, \tau_i \le t | \mathscr{G}_{i-1}) = G(x) (1 - e^{-r(t - \tau_{i-1})}), \ i \ge 1.$$

The following theorem is an analogue of Theorem 1.

Theorem 2. The family

$$S_t = \frac{1}{t} \sum_{i:\tau_i \le t} \xi_i, \quad t \to \infty$$

obeys the LDP in the metric space (\mathbb{R}_+, ϱ) with the rate $\frac{1}{t}$ and the rate function

$$I(u) = \begin{cases} \sup_{\lambda \in (-\infty,\Lambda)} [\lambda u - r \int_0^\infty (e^{\lambda z} - 1) dG(z)], & u > 0\\ r[1 - G(0+)], & u = 0. \end{cases}$$

We give two examples illustrating compatibility with Theorems 1 and 2. For both discrete and continuous time cases, let

$$\mathsf{P}(\xi_1 \le x) = 1 - e^{-x}, \ x \ge 0,$$

so that, ξ_1 has the Laplace transform with $\Lambda = 1$. The log moment generating function is

$$g(\lambda) = -\log(1-\lambda), \ \lambda < 1.$$

For $G(x) = 1 - e^{-x}, x \ge 0$.

$$\int_0^\infty r(e^{\lambda z} - 1) dG(z) = \frac{r\lambda}{1 - \lambda}, \ \lambda < 1.$$

In both cases, rate functions are explicitly computable (see also figures (1) and (2)),

$$I^{d}(u) = \begin{cases} u - 1 - \log(u), & u > 0\\ \infty, & u = 0 \end{cases} \text{ (discrete time case)}$$
$$I^{c}(u) = \begin{cases} \left(\sqrt{r} - \sqrt{u}\right)^{2}, & u > 0\\ r, & u = 0. \end{cases} \text{ (continuous time case)}$$



FIGURE 1. The rate function $I^d(u)$



FIGURE 2. The rate function $I^{c}(u)$ for r = 1

Remark 1. Related topics to Theorem 2 can be found e.g. in Georgii and Zessin, [3], serving a class of marked point random fields. Probably, the proof of Theorem 2 can be adapted with arguments from proofs in [3] provided that many details not related to our setting have to be omitted and other ones concerning to the boundary effect have to be added.

We prefer to give a complete and direct proof of Theorem 2.

2. Counting random measure, its compensator, Laplace transform

We consider $(\xi_i, \tau_i)_{i \ge 1}$ as a multivariate (marked) point process (see, e.g. [4], [5]) with the counting measure

$$\mu(dt, dy) = \sum_{i \ge 1} \mathbf{I}_{\{\tau_i < \infty\}} \delta_{\{\tau_i, \xi_i\}}(t, y) dt dy,$$

where $\delta_{\{\tau_i,\xi_i\}}$ is the Dirac delta-function on $\mathbb{R}_+ \times \mathbb{R}_+$. Parallel to $(\mathscr{G}_n)_{n\geq 0}$, we introduce one more filtration $(\mathscr{G}_t)_{t\geq 0}$ related to $(\xi_i, \tau_i)_{i\geq 1}$:

$$\mathscr{G}_t := \sigma(\mu([0, t'] \times \Gamma) : t' \le t, \Gamma \in \mathscr{B}(\mathbb{R}_+)),$$

where $\mathscr{B}(\mathbb{R}_+)$ is the Borel σ -algebra on \mathbb{R}_+ , and assume that \mathscr{G}_0 is augmented by P-zero sets from \mathcal{F} (notice that, then, $(\mathscr{G}_t)_{t\geq 0}$ satisfies the general conditions). With the help of the counting measure μ , one can present tS_t in a form of a stochastic integral with respect to μ :

$$tS_t = \int_0^t \int_{x>0} x\mu(ds, dx).$$
 (2.1)

Then, the Levy measure $\nu(ds, dx)$, related to μ , is explicitly computed (see, e.g. Theorem III.1.33, [5])

$$\nu(ds, dx) = \sum_{i \ge 1} I_{[]\tau_{i-1}, \tau_i]}(t) \frac{dG(x)de^{-r(s-\tau_{i-1})}}{e^{-r(s-\tau_{i-1})}} = rdsdG(x)$$
(2.2)

and is not random. It is well known (see, e.g. Corollary to Theorem 1 in §4, Ch.4, [6] that under the deterministic Levy process tS_t is the random process with independent increments. We recall a useful property: for any nonnegative and (\mathscr{G}_t) -predictable function $f(\omega, x, t)$,

$$\mathsf{E}\int_0^t \int_{x>0} f(\omega, x, s) \mu(ds, dx) = \mathsf{E}\int_0^t \int_{x>0} f(\omega, x, s) \nu(ds, dx)$$

"\overline = \overline ".

with " $\infty = \infty$ ".

Lemma 1. [Laplace transform] For any $\lambda < \Lambda$ and t > 0,

$$\mathsf{E}e^{\lambda t S_t} = e^{rt \int_{x>0} [e^{\lambda z} - 1] dG(z)}$$

Proof. Though the direct computation of Laplace's transform is permissible, we prefer to apply the stochastic calculus. The process $U_t = e^{\lambda t S_t}$ has right-continuous piece-wise constant paths with jumps

$$\Delta U_s = (U_s - U_{s-}) = U_{s-} \int_{x>0} [e^{\lambda x} - 1] \mu(\{s\}, dx),$$

so that, for any t > 0,

$$U_t = 1 + \int_0^t \int_{x>0} U_{s-}[e^{\lambda x} - 1]\mu(ds, dx).$$

Since the function $f(\omega, x, s) := U_{s-}[e^{\lambda x} + 1]$ is nonnegative and predictable, the following equality with $\lambda < \Lambda$ holds true:

$$E \int_0^t \int_{x>0} U_{s-}[e^{\lambda x} + 1]\mu(ds, dx) = E \int_0^t \int_{x>0} U_{s-}[e^{\lambda x} + 1]\nu(ds, dx) (<\infty).$$

Then, we also have

$$E \int_0^t \int_{x>0} U_{s-}[e^{\lambda x} - 1]\mu(ds, dx) = E \int_0^t \int_{x>0} U_{s-}[e^{\lambda x} - 1]\nu(ds, dx) \ (\in \mathbb{R}).$$

Since $\nu(ds, dx) = rdsdG(x)$, the later provides

$$(\mathsf{E}U_t) = 1 + \int_0^t \int_{x>0} (EU_s) [e^{\lambda x} - 1] dG(x) r ds.$$

This can be written in an equivalent form of differential equation

$$\frac{d(\mathsf{E}U_t)}{dt} = (\mathsf{E}U_t) \int_{x>0} [e^{\lambda x} - 1] dG(x) r$$

subject to $(\mathsf{E}U_0) = 1$.

Thus, the desired result holds.

3. The proof of Theorem 2

We verify the necessary and sufficient conditions for the LDP to hold (for more details, see Puhalskii, [7]):

1) exponential tightness,

$$\lim_{j \to \infty} \overline{\lim_{t \to \infty}} \, \frac{1}{t} \log \mathsf{P} \big(S_t \in \mathbb{R}_+ \setminus \mathcal{K}_j \big) = -\infty,$$

where \mathcal{K}_j 's are compacts increasing to \mathbb{R}_+ ;

2) local LDP, defining the rate function $I(u), u \in \mathbb{R}_+$

$$\lim_{\delta \to 0} \lim_{t \to \infty} \frac{1}{t} \log \mathsf{P}(|S_t - u| \le \delta). = -I(u).$$

3.1. The exponential tightness. By choosing

 $K_j = \{x \in \mathbb{R}_+ : x \in [0, j]\}$

and applying Chernoff's inequality with parameter 0.5Λ , we find that

 $\mathsf{P}(S_t > j) \le e^{-0.5\Lambda j + \log \mathsf{E}e^{0.5\Lambda t S_t}}.$

By Lemma 1,

$$\mathsf{E}e^{0.5\Lambda tS_t} = e^{rt\int_{x>0} [e^{0.5\Lambda x} - 1]dG(z)}$$

and, therefore,

$$\frac{1}{t}\log \mathsf{P}(S_t > j) \le -0.5\Lambda j + r \int_{x>0} [e^{0.5\Lambda x} - 1] dG(x) \xrightarrow[j \to \infty]{} -\infty$$

and 1) is done.

3.2. The local LDP. We begin with computation of I(0) and prove

$$\underbrace{\lim_{\delta \to 0} \lim_{t \to \infty} \frac{1}{t} \log \mathsf{P}(S_t \le \delta) \ge -r[1 - G(0+)]}_{\lim_{\delta \to 0} \lim_{t \to \infty} \frac{1}{t} \log \mathsf{P}(S_t \le \delta) \le -r[1 - G(0+)]}$$
(3.1)

By (2.2), $\{tS_t = 0\} = \{\mu((0, t] \times \{x > 0\}) = 0$. Consequently for any t > 0,

$$\mathsf{P}(S_t \le \delta) \ge \mathsf{P}(S_t = 0) = \mathsf{P}(tS_t = 0) = \mathsf{P}(\mu((0, t], \{x > 0\}) = 0).$$

The counting process $\pi_t := \mu((0, t], \{x > 0\})$ has independent increments and the rate

$$\mathsf{E}\mu\big((0,t],\{x>0\}\big)=\nu\big((0,t],\{x>0\}\big)=r[1-G(0+)]t.$$

It is a counting process with the compensator $\nu((0,t], \{x > 0\}) = r[1 - G(0+)]t$. Therefore, by the Watanabe theorem, [8], π_t is a Poisson process with parameter r[1 - G(0+)]. Hence, due to well known property of the Poisson process

$$\mathsf{P}(\pi_t = 0) = e^{-tr[1 - G(0+)]}.$$

We find that

$$\frac{1}{t}\log \mathsf{P}(S_t \le \delta) \ge \frac{1}{t}\log \mathsf{P}(\pi_t = 0) = -r[1 - G(+)]$$

implying the lower bond from (3.1).

The upper bound from (3.1) is derived with the help of Laplace's transform with $0 < \lambda < \Lambda$. To this end, we use identity

$$1 = \mathsf{E} \exp\left(\lambda t S_t - tr \int_{x>0} [e^{\lambda x} - 1] dG(x)\right)$$

implying the inequality

$$1 \ge \mathsf{E}I_{\{S_t \le \delta\}} \exp\left(t \left[\lambda \delta - r \int_{x>0} [e^{\lambda x} - 1] dG(x)\right]\right)$$

being equivalent to

$$\frac{1}{t}\log \mathsf{P}(S_t \le \delta) \le -\lambda\delta + r \int_{x>0} [e^{-\lambda x} - 1] dG(x).$$

Now, passing $t \to \infty$, we obtain the following upper bound depending on δ and λ :

$$\overline{\lim_{t \to \infty} \frac{1}{t}} \log \mathsf{P}(S_t \le \delta) \le -\lambda \delta + r \int_{x>0} [e^{\lambda x} - 1] dG(x).$$

Now, passing $\delta \to 0$ and λ to $-\infty$ we find that

$$\overline{\lim_{\delta \to 0} \lim_{t \to \infty} \frac{1}{t} \log \mathsf{P}(S_t \le \delta)} \le -r \int_{x>0} dG(x) = -r[1 - G(0+)].$$

We continue the proof by checking the formula for I(u) when u > 0, i.e.

$$\overline{\lim_{\delta \to 0} \lim_{t \to \infty} \frac{1}{t}} \log \mathsf{P}(|S_t - u| \le \delta) \le -I(u)$$
$$\underline{\lim_{\delta \to 0} \lim_{t \to \infty} \frac{1}{t}} \log \mathsf{P}(|S_t - u| \le \delta) \ge -I(u),$$

with

$$I(u) = \sup_{\lambda \in (-\infty,\Lambda)} \left[\lambda u - r \int_0^\infty (e^{\lambda x} - 1) dG(x) \right].$$

The Laplace transform

$$1 = \mathsf{E} \exp\left(\lambda t S_t - tr \int_{x>0} [e^{\lambda x} - 1] dG(x)\right), \ \lambda < \Lambda,$$

implies the inequality

$$1 \ge \mathsf{E}I_{\{|S_t-u|\le\delta\}} \exp\left(-t\delta u + \lambda tu - tr \int_{x>0} [e^{\lambda x} - 1] dG(x)\right)$$

prviding the following upper bound depending on λ :

$$\overline{\lim_{\delta \to 0} \lim_{t \to \infty} \frac{1}{t} \log \mathsf{P}(|S_t - u| \le \delta) \le -\Big(\lambda u - r \int_{x>0} [e^{\lambda x} - 1] dG(x)\Big).$$

A further minimization of the right hand side of the above inequality in λ over $(-\infty, \Lambda)$ gives the desired result.

The lower bound proof uses a standard approach of changing "probability measure". Denote by $\lambda^* = \operatorname{argmax}_{\lambda < \Lambda} \left(\lambda u - r \int_{x>0} [e^{\lambda x} - 1] dG(x) \right)$. Since λ^* solves the equation (with u > 0)

$$u - r \int_{x>0} x e^{\lambda x} dG(x) = 0, \qquad (3.2)$$

 λ^* is a proper number strictly less than Λ . Set

$$\mathfrak{L}_t(\lambda^*) = \exp\left(\lambda^* t S_t - \int_0^t \int_{x>0} r[e^{\lambda^* x} - 1] dG(x) ds\right).$$
(3.3)

First of all we notice that the Laplace transform for tS_t with λ^* guarantees $\mathsf{E}\mathfrak{L}_t(\lambda^*) = 1$. Moreover, taking into account (2.1) and applying the Itô formula to $\mathfrak{L}_t(\lambda^*)$ one can see that $(\mathfrak{L}_t(\lambda^*), \mathscr{G}_t)_{t\geq 0}$ is a positive local martingale with paths from the Skorokhod space $\mathbb{D}_{[0,\infty)}$. Then a measure $\widetilde{\mathsf{P}}_t$, defined by $d\widetilde{\mathsf{P}}_t = \mathfrak{L}_t(\lambda^*)d\mathsf{P}_t$, where P_t is a restriction of P on \mathscr{G}_t , is the probability measure. We introduce the probability space $(\Omega, \mathcal{F}, \widetilde{\mathsf{P}}_t)$. Since $\mathfrak{L}_t(\lambda^*) > 0$, P -a.s., not only $\widetilde{\mathsf{P}}_t \ll \mathsf{P}_t$ but also $\widetilde{\mathsf{P}}_t \ll \mathsf{P}_t$ with $d\mathsf{P}_t = \mathfrak{L}_t^{-1}(\lambda^*)\widetilde{\mathsf{P}}_t$. This property and (3.3) provide a lower bound

$$\mathsf{P}(|S_t - u| \le \delta) = \mathsf{P}_t(|S_t - u| \le \delta) = \int_{\{|S_t - u| \le \delta\}} \mathfrak{L}_t^{-1}(\lambda^*) d\widetilde{\mathsf{P}}_t$$
$$\ge e^{-\lambda^* t \delta - tI(u)} \widetilde{\mathsf{P}}_t(|S_t - u| \le \delta).$$

Therefore, we find that

$$\lim_{t \to \infty} \frac{1}{t} \log \mathsf{P}(|S_t - u| \le \delta) \ge -I(u) - \lambda^* \delta + \lim_{t \to \infty} \frac{1}{t} \log \widetilde{\mathsf{P}}_t(|S_t - u| \le \delta).$$

Obviously, the desired lower bound to obtain it is left to prove that

$$\lim_{t \to \infty} \widetilde{\mathsf{P}}_t (|S_t - u| \le \delta) = 1$$

or, equivalently,

$$\lim_{t \to \infty} \widetilde{\mathsf{P}}_t (|S_t - u| > \delta) = 0.$$
(3.4)

Thus the last step of the proof deal with (3.4). To this end, we show that $(\widetilde{\mathsf{E}}_t \text{ denotes the expectation relative to } \widetilde{\mathsf{P}}_t)$

$$\lim_{t \to \infty} \widetilde{\mathsf{E}}_t |S_t - u|^2 = 0.$$

Since $\mathsf{E}\mathfrak{L}_t(\lambda) = 1$, it holds

$$0 = \frac{\partial^2 \mathsf{E}\mathfrak{L}_t(\lambda)}{\partial \lambda^2}|_{\lambda = \lambda^*}$$

= $t^2 \mathsf{E} \Big(S_t - r \int_{\{x>0\}} x e^{\lambda^* x} dG(x) \Big)^2 \mathfrak{L}_t(\lambda^*) - t \underbrace{r \int_{\{x>0\}} x^2 e^{\lambda^* x} dG(x)}_{=u \text{ (see (3.2))}}$
= $t^2 \widetilde{\mathsf{E}} \Big(S_t - r \int_{\{x>0\}} x e^{\lambda^* x} dG(x) \Big)^2 - tu.$

Hence,

$$\widetilde{\mathsf{E}}(S_t - u)^2 = \frac{1}{t} r \mathsf{E} \int_{\{x > 0\}} x^2 e^{\lambda^* x} dG(x) \xrightarrow[t \to \infty]{} 0.$$

References

- Cramer, H. (1938) Sur un nouveau theoreme-limite de la theoriedes probabilites. Actualites Santifiques et Industrielles. Colloque cosacrea la throrie des probabilitrs. 3, 736:2–23. Hermann, Paris.
- [2] Dembo, A. and Zeitouni, O. (1993) Large Deviations Techniques and Applications. Jones and Bartlet, 1993.
- [3] Georgii, Hans-Otto and Zessin, Hans. (1993) Large deviations and the maximum entropy principle of marked point random fields. *Probability Theory and Related Fields.* 96, 177–204.
- [4] Jacod, J. Multivariate point processes: predictable projection, Radon–Nikodym derivatives, representation martingales. Z. Wahrsch. Verw. Gebiete, 31 (1975), 235–53.
- [5] Jacod, J. and Shiryaev, A. N. Limit Theorems for Stochastic Processes. Springer-Verlag, New York, Heidelberg, Berlin, (1987).
- [6] Liptser, R.S. and Shiryaev, A.N. (1989): Theory of Martingales. Kluwer, Dordrecht (Russian edition 1986)
- [7] Puhalskii, A. (2001) Large Deviations and Idempotent Probability, Chapman & Hall/CRC Press.
- [8] Watanabe S. (1964) On discontinuous additive functionals and Lévy measures of Markov process. Japan J. Math., 34, 53–70.

School of Mathematical Sciences, Building 28M, Monash University, Clayton Campus, Victoria 3800, Australia.

 $E\text{-}mail \ address: \texttt{fima.klebaner@sci.monash.edu.au}$

DEPARTMENT OF ELECTRICAL ENGINEERING SYSTEMS, TEL AVIV UNIVER-SITY, 69978 TEL AVIV, ISRAEL

E-mail address: liptser@eng.tau.ac.il