# ON REALIZATION OF THE KREIN-LANGER CLASS  $N_{\kappa}$  OF MATRIX-VALUED FUNCTIONS IN HILBERT SPACES WITH INDEFINITE METRIC

YURI ARLINSKII, SERGEY BELYI, VLADIMIR DERKACH, AND EDUARD TSEKANOVSKII

*Dedicated to Henk de Snoo on the occasion of his 60th birthday*

Abstract. In this paper the realization problems for the Krein-Langer class  $N_{\kappa}$  of matrix-valued functions are being considered. We found the criterion when a given matrix-valued function from the class  $N_{\kappa}$  can be realized as linear-fractional transformation of the transfer function of canonical conservative system of the M. Livsic type (Brodskii-Livsic rigged operator colligation) with the main operator acting on a rigged Pontryagin space  $\Pi_{\kappa}$  with indefinite metric. We specify three subclasses of the class  $N_{\kappa}(R)$  of all realizable matrixvalued functions that correspond to different properties of a realizing system, in particular, when the domains of the main operator of a system and its conjugate coincide, when the domain of the hermitian part of a main operator is dense in Πκ. Alternatively we show that the class  $N_{\kappa}(R)$  can be realized as transfer matrix-functions of some canonical impedance systems with selfadjoint main operators in rigged spaces  $\Pi_{\kappa}$ . The case of scalar functions of the class  $N_{\kappa}(R)$  is considered in details and some examples are presented.

#### 1. INTRODUCTION

Realizations and corresponding operator models of different classes of holomorphic matrix-valued functions in the open right half-plane, unit circle and upper half-plane play important role in spectral analysis of different classes of linear operators in Hilbert spaces, interpolation problems and system theory, and we refer in this matter to [\[1\]](#page-18-0), [\[2\]](#page-18-1), [\[5\]](#page-18-2)-[\[8\]](#page-19-0), [\[10\]](#page-19-1), [\[11\]](#page-19-2)-[\[13\]](#page-19-3), [\[16\]](#page-19-4), [\[21\]](#page-19-5), [\[22\]](#page-19-6), [\[24\]](#page-19-7), [\[26\]](#page-19-8), [\[33\]](#page-20-0), [\[37\]](#page-20-1)- [\[40\]](#page-20-2). In this paper we continue the investigation of various problems that arise in the study of linear stationary conservative dynamic systems (operator colligations). Relying on the results and technique developed in [\[11\]](#page-19-2), [\[12\]](#page-19-9) we keep dealing with linear stationary conservative dynamic systems (l.s.c.d.s)  $\theta$  of the form

$$
\begin{cases}\n(\mathbb{A} \quad -zI) = K\mathcal{J}\varphi_{-} & (\text{Im }\mathbb{A} = K\mathcal{J}K^*) \\
\varphi_{+} = \varphi_{-} - 2iK^*x & \qquad \qquad \mathcal{J} \\
\theta = \begin{pmatrix} \mathbb{A} & K & \mathcal{J} \\
\mathfrak{H}^{+} \subset \Pi_{\kappa} \subset \mathfrak{H}_{-} & E\n\end{pmatrix}.\n\end{cases}
$$

In the system  $\theta$  above A is a bounded linear operator acting from  $\mathfrak{H}^+$  into  $\mathfrak{H}_-,$ where  $\mathfrak{H}^+ \subset \Pi_{\kappa} \subset \mathfrak{H}_-$  is a rigged Pontryagin space, K is a linear bounded operator from a Hilbert space E into  $\mathfrak{H}_-, \mathcal{J} = \mathcal{J}^* = \mathcal{J}^{-1}$  is acting in E,  $\varphi_\pm \in E, \varphi_-$  is an

or

*Date*: November 5, 2018.

<sup>1991</sup> *Mathematics Subject Classification.* Primary 47A10; Secondary 47N50, 81Q10.

input vector,  $\varphi_+$  is an output vector, and  $x \in \mathfrak{H}^+$  is a vector of the inner state of the system  $\theta$ . The operator-valued function

$$
W_{\theta}(z) = I - 2iK^*(\mathbb{A} - zI)^{-1}K\mathcal{J} \qquad (\varphi_+ = W_{\theta}(z)\varphi_-),
$$

is the transfer operator-valued function of the system  $\theta$ . It was shown in [\[12\]](#page-19-9) that a Herglotz-Nevanlinna matrix-valued function  $V(z)$  acting on a Hilbert space E can be represented and realized in the form

$$
V(z) = i[W_{\Theta}(z) + I]^{-1}[W_{\Theta}(z) - I]\mathcal{J} = K^*(A_R - zI)^{-1}K,
$$

where  $W_{\Theta}(z)$  is a transfer function of some canonical system  $\Theta$ ,

<span id="page-1-1"></span>
$$
\Theta = \begin{pmatrix} A & K & \mathcal{J} \\ \mathfrak{H}_+ \subset \mathfrak{H} \subset \mathfrak{H}_- & K & \mathcal{J} \end{pmatrix},
$$

if certain conditions on integral representation of  $V(z)$  are met. Alternatively, one can realize a Herglotz-Nevanlinna matrix-valued  $V(z)$  as a transfer mapping of an impedance system  $\Delta$  of the form

$$
\begin{cases} (\mathbb{D} - zI)x = K\varphi_-, \\ \varphi_+ = K^*x, \end{cases} \tag{1}
$$

where D is a self-adjoint operators acting from  $\mathfrak{H}_+$  into  $\mathfrak{H}_-$  (see [\[12\]](#page-19-9), [\[13\]](#page-19-3)). In this case associated transfer function is given by

$$
V_{\Delta}(z) = K^*(\mathbb{D} - zI)^{-1}K.
$$

In this paper we study similar realization problems but utilize a new type of realizing systems whose main operator is acting on a rigged Pontryagin space  $\Pi_{\kappa}$ . The set of realizable functions appears to be a subclass of the well known Krein-Langer's class  $N_{\kappa}$  also known as generalized Nevanlinna functions. In Section [6](#page-11-0) we specify three subclasses of the class  $N_{\kappa}(R)$  of all realizable matrix-valued functions that yield different properties of operators in the realizing systems. It is worth mentioning that in the case when  $\kappa = 0$  all the subclasses coincide with the similar subclasses of realizable Herglotz-Nevanlinna functions described in [\[12\]](#page-19-9), [\[13\]](#page-19-3). Section [7](#page-13-0) uses a factorization formula from [\[25\]](#page-19-10) to provide applications of  $N_{\kappa}(R)$  realizations to the scalar case when  $E = \mathbb{C}$  while establishing a connection with the class  $N(R)$ of realizable Herglotz-Nevanlinna functions. The paper is concluded with several examples.

#### 2. OPERATORS IN PONTRYAGIN SPACES  $\Pi_{\kappa}$

<span id="page-1-0"></span>We start with the basic construction following some results from the theory of operators in  $\Pi_{\kappa}$  spaces [\[9\]](#page-19-11), [\[29\]](#page-19-12), [\[32\]](#page-19-13), [\[33\]](#page-20-0). Let  $\Pi_{\kappa}$  be a Pontryagin space [29], i.e., a Hilbert space  $\mathcal H$  where along with the usual scalar product  $(x, y)$  there is an indefinite scalar product

$$
[x, y] = (Jx, y),\tag{2}
$$

where  $J = P_+ - P_-$  is a bounded linear operator such that  $J = J^*$ ,  $J^2 = I$ , and  $P_+$ and P<sub>-</sub> are complementary orthoprojections,  $P_+ + P_- = I$ . Putting  $\Pi_{\pm} = P_{\pm} \Pi_{\kappa}$ we have

$$
\Pi_{\kappa} = \Pi_{+} \boxplus \Pi_{-}, \quad \dim \Pi_{-} = \kappa.
$$

Here and below the direct orthogonal sum with respect to an indefinite scalar prod-uct [\(2\)](#page-1-0) is denoted by  $\boxplus$  and called  $\pi\text{-}orthogonal sum$ . Similarly, the  $\pi\text{-}orthogonal$ 

*complement* of a lineal L will be denoted by  $L^{[\perp]}$ . The positive definite  $(x, y)$  and indefinite  $[x, y]$  scalar products are related by

$$
(x, y) = [x_{+}, y_{+}] - [x_{-}, y_{-}],
$$
  

$$
[x, y] = (x_{+}, y_{+}) - (x_{-}, y_{-}),
$$

where  $x = x_+ + x_-, y = y_+ + y_-, x_+, y_+ \in \Pi_+$ , and  $x_-, y_- \in \Pi_-$ .

The set of vectors  $f \in L$  that are  $\pi$ -orthogonal to L, i.e.  $f[\perp]L$  is called [\[29\]](#page-19-12) the *isotropic part* of the linear manifold  $L$ . If the isotropic part of  $L$  has non-zero elements we say that the scalar product [ $\cdot$ , $\cdot$ ] is *degenerate* [\[32\]](#page-19-13) on L. L<sub>+</sub> (respectively,  $L_-, L_0$ ) will denote the set of all  $x \in \Pi_{\kappa}$  for which  $[x, x] > 0$  (respectively,  $[x, x] < 0$ ,  $[x, x] = 0$  and is called *positive (negative, neutral)* part of L. Every subspace  $\mathcal{L} \in \Pi_{\kappa}$  can be decomposed into a direct sum of  $\pi$ -orthogonal subspaces

$$
\mathcal{L}=\mathcal{L}_+\boxplus\mathcal{L}_0\boxplus\mathcal{L}_-,
$$

where  $\mathcal{L}_+$ ,  $\mathcal{L}_0$ , and  $\mathcal{L}_-$  are, respectively, positive, neutral, and negative subspaces, some of which may degenerate into null subspaces. For a subspace  $\mathcal L$  above we write  $sign \mathcal{L} = (l_+, l_0, l_-)$  where  $l_{\pm} = \dim \mathcal{L}_{\pm}$  and  $l_0 = \dim \mathcal{L}_0$  [\[32\]](#page-19-13).

Recall [\[1\]](#page-18-0) that a *linear relation* in  $\Pi_{\kappa}$  is a subspace  $\mathcal{A}$  in  $\Pi_{\kappa} \times \Pi_{\kappa}$ . The *domain* of a linear relation  $\mathcal A$  is

$$
\mathcal{D}(\mathcal{A}) = \{ f \in \Pi_{\kappa} : \langle f, f' \rangle \in \mathcal{A} \quad \text{for some} \quad f' \in \Pi_{\kappa} \},
$$

and the range of  $\mathcal A$  is

$$
\mathcal{R}(\mathcal{A}) = \{f' \in \Pi_{\kappa} : \langle f, f' \rangle \in \mathcal{A} \text{ for some } f \in \Pi_{\kappa} \}.
$$

The subspace

$$
\operatorname{mul} \mathcal{A} = \{ g \in \Pi_{\kappa} : \langle 0, g \rangle \in \mathcal{A} \}
$$

is called the *multivalued part* of a linear relation  $A$ . A linear relation  $A$  is the graph of a linear operator in  $\Pi_{\kappa}$  if and only if  $mul \mathcal{A} = \{0\}.$ 

Let us associate with a linear operator A in  $\Pi_{\kappa}$  the linear relation  $\mathcal{A} := Gr(A)$ , the graph of the operator A.

For a linear relation  $\mathcal A$  in  $\Pi_{\kappa}$  its  $\pi$ -adjoint  $\mathcal A^+$  is defined by

$$
\mathcal{A}^+ = \{ \langle h, h' \rangle \in \Pi_{\kappa} \times \Pi_{\kappa} : [f', h] = [f, h'] \text{ for all } \langle f, f' \rangle \in \mathcal{A} \}.
$$

A linear relation A (operator A) is called  $\pi$ -symmetric if  $\mathcal{A} \subset \mathcal{A}^+$  and  $\pi$ -selfadjoint if  $\mathcal{A} = \mathcal{A}^+$ .

We recall [\[29\]](#page-19-12) that a  $\pi$ -symmetric operator A in  $\Pi_{\kappa}$  can not have more than  $\kappa$ eigenvalues, counting multiplicities, in the upper (lower) half-plane. If the operator  $\Lambda$  is  $\pi$ -self-adjoint, then these non-real eigenvalues are located symmetrically with respect to the real axis. For an arbitrary complex number  $z$  and a  $\pi$ -symmetric operator A in  $\Pi_{\kappa}$  we set [\[29\]](#page-19-12)

$$
\mathcal{M}_z = (A - z)\mathcal{D}(A), \qquad \mathfrak{N}_{\bar{z}} = \mathcal{M}_z^{[\perp]}.
$$

If  $\lambda$  (Im $\lambda \neq 0$ ) is not an eigenvalue of A, then  $\mathcal{M}_{\lambda}$  is a subspace of  $\Pi_{\kappa}$  and  $\mathfrak{N}_{\lambda}$ is called [\[29\]](#page-19-12) a *deficiency subspace* corresponding to  $\lambda$  with dim  $\mathfrak{N}_{\lambda}$  maintaining a constant value as a *deficiency index* of A in  $\Pi_{\kappa}$ . Let  $\Delta_A$  be the set of all non-real  $\lambda$  for which the scalar product [·, ·] is degenerate on  $\mathfrak{N}_{\lambda}$ . According to [\[32\]](#page-19-13) the set  $\Delta_A$  of a  $\pi$ -symmetric operator A contains no interior points, its complement  $(\mathbb{C}_{+} \cup \mathbb{C}_{-}) \setminus \Delta_A$  is an open set, and on every component of this open set  $sign \mathfrak{N}_{\lambda}$ is constant.

It was shown in [\[19\]](#page-19-14) that every  $\pi$ -symmetric operator A in the space  $\Pi_{\kappa}$  admits  $\pi$ -self-adjoint extensions in  $\Pi_{\kappa}$  if and only if its deficiency indices coincide. An operator A is called prime if it has no non-real eigenvalues and

<span id="page-3-0"></span>
$$
\mathbf{c}.\mathbf{1}.\mathbf{s}.\{\mathfrak{N}_z,\,z\neq\bar{z}\}=\Pi_{\kappa}.\tag{3}
$$

In what follows we denote  $\text{Re}(T) = (T + T^+)/2$ ,  $\text{Im}(T) = (T - T^+)/2i$  for linear operators T in  $\Pi_{\kappa}$  with  $\mathcal{D}(T) = \mathcal{D}(T^+)$ . Similarly, for a linear operator Q with  $\mathcal{D}(Q) = \mathcal{D}(Q^*)$  in a Hilbert space we use the same notation to denote  $\text{Re}(Q) = (Q + Q^*)/2$  and  $\text{Im}(Q) = (Q - Q^*)/2i$ .

### 3. Bi-extensions in Rigged Pontryagin Space

<span id="page-3-1"></span>Let consider A as an operator from the Hilbert space  $\overline{\mathcal{D}(A)}$  into the Hilbert space H. Then its adjoint  $A^*$  is defined on a set  $\mathcal{D}(A^*)$  that is dense in H and has the range in  $\overline{\mathcal{D}(A)}$ . This allows us to introduce the Hilbert spaces  $\mathfrak{H}_+ = \mathcal{D}(A^*)$  and  $\mathfrak{H}^+ = J\mathcal{D}(A^*)$  with corresponding inner products

$$
(f,g)_{+} = (f,g) + (A^*f, A^*g), \quad f, g \in \mathfrak{H}_+, (f,g)^{+} = (f,g) + (A^*Jf, A^*Jg), \quad f, g \in \mathfrak{H}^+.
$$

Next we construct two rigged Hilbert spaces [\[15\]](#page-19-15), [\[18\]](#page-19-16), [\[5\]](#page-18-2), [\[12\]](#page-19-9)

$$
\mathfrak{H}_{+}\subset\mathcal{H}\subset\mathfrak{H}_{-}\quad\text{and}\quad\mathfrak{H}^{+}\subset\mathcal{H}\subset\mathfrak{H}^{-}.
$$

Let  $\mathcal{R}_1 \in [\mathfrak{H}_+, \mathfrak{H}_+]$  and  $\mathcal{R}_2 \in [\mathfrak{H}^+, \mathfrak{H}^-]$  be the isometric Riesz-Berezanskiĭ operators [\[15\]](#page-19-15) corresponding to the above triplets. We introduce

$$
J_+ = J \Big|_{\mathfrak{H}_+},
$$

the linear operator mapping  $\mathfrak{H}_+$  isometrically onto  $\mathfrak{H}^+$  and let  $J^{\times}_+ \in [\mathfrak{H}^-, \mathfrak{H}_-]$  be its dual, which isometrically maps  $\mathfrak{H}^-$  onto  $\mathfrak{H}_-$ . It is easy to see that

$$
(J_+^{\times})^{-1} = \mathcal{R}_2 J_+ \mathcal{R}_1^{-1}.
$$

Since  $\mathfrak{H}_-$  is isomorphic to  $\mathfrak{H}^-$  it can be considered as the space of anti-linear functionals on  $\mathfrak{H}^+$  defined by

$$
\alpha(f) = (\alpha, Jf) = [\alpha, f], \quad \alpha \in \mathfrak{H}_-, f \in \mathfrak{H}^+.
$$

Thus, we can form a rigged  $\Pi_{\kappa}$  space

$$
\mathfrak{H}^+\subset\Pi_\kappa\subset\mathfrak{H}_-.
$$

Consequently, if  $\mathbb{A} \in [\mathfrak{H}^+, \mathfrak{H}_-]$  then  $\mathbb{A}^\times \in [\mathfrak{H}^+, \mathfrak{H}_-]$  and  $[\mathbb{A}f, g] = [f, \mathbb{A}^\times g]$  for all  $f, g \in \mathfrak{H}^+$ .

**Definition 1.** An operator  $A \in [\mathfrak{H}^+, \mathfrak{H}_-]$  is called bi-extension of a closed  $\pi$ symmetric operator A if  $A \supseteq A$  and  $A^{\times} \supseteq A$ . A bi-extension A is called  $\pi$ -selfadjoint if  $\mathbb{A} = \mathbb{A}^{\times}$ .

Let A be a bi-extension of a  $\pi$ -symmetric operator A. The operator  $\hat{A} = A$  with  $\mathcal{D}(\hat{A}) = \{f \in \mathfrak{H}^+ \mid \mathbb{A}f \in \Pi_{\kappa}\}\$ is called *quasi-kernel* [\[5\]](#page-18-2) of the operator A. If  $\mathbb{A} = \mathbb{A}^{\times}$ and  $\hat{A}$  is a quasi-kernel of A such that  $A \neq \hat{A}$ ,  $\hat{A}^{+} = \hat{A}$  then A is said to be a strong  $\pi$ -self-adjoint bi-extension of A.

In what follows we will assume that the closed and non-densely defined  $\pi$ -symmetric operator A is J-regular [\[5\]](#page-18-2), i.e., the operator  $PA$  is closed, where P is the orthogonal projection onto  $\overline{J\mathcal{D}(A)}$  in H. The analog of von Neumann's formula for the operator  $JA$  (see [\[40\]](#page-20-2))

$$
\mathfrak{H}^+ = \mathcal{D}(A) \oplus \mathfrak{N}'_i \oplus \mathfrak{N}'_{-i} \oplus \mathfrak{N}
$$

holds in the space  $\mathfrak{H}^+$ , where  $\mathfrak{N}'_{\pm i}$  are the *semi-deficiency subspaces* of the operator JA [\[31\]](#page-19-17), i.e.

$$
\mathfrak{N}'_{\pm i} = \overline{JD(A)} \ominus (PJA \mp iI)\mathcal{D}(A)),
$$

 $\mathcal{R} = J_+^{\times} \mathcal{R}_2$ ,  $\mathfrak{N} = \mathcal{R}^{-1} \mathfrak{L}$ , and  $\mathfrak{L} = \Pi_{\kappa} \boxminus \overline{\mathcal{D}(A)}$ . The condition of A being J-regular is equivalent to the subspace  $\mathfrak L$  being closed in  $\mathfrak H_-\,$  and a sufficient condition of J-regularity is dim  $\mathfrak{L} < \infty$  (see [\[40\]](#page-20-2), [\[5\]](#page-18-2)). Let  $P_{\mathcal{D}(A)}^+$ ,  $P_{\mathfrak{N}'_i}^+$ ,  $P_{\mathfrak{N}'_{-i}}^+$ ,  $P_{\mathfrak{N}}^+$ , and  $P_{\mathfrak{M}}^+$  be the orthogonal projections in  $\mathfrak{H}^+$  onto  $\mathcal{D}(A)$ ,  $\mathfrak{N}'_i$ ,  $\mathfrak{N}'_{-i}$ ,  $\mathfrak{N}$ , and  $\mathfrak{M} = \mathfrak{H}^+ \ominus \mathcal{D}(A)$ , respectively. Then the set of all bi-extensions of a J-regular operator A is described by the formula [\[40\]](#page-20-2)

$$
\mathbb{A} = AP_{\mathcal{D}(A)}^+ + \left(A^+ + \mathcal{R}S_{\mathbb{A}}\right)P_{\mathfrak{M}}^+,
$$

where  $S_{\mathbb{A}} \in [\mathfrak{M}, \mathfrak{M}]$ . Moreover,  $\mathbb{A}$  is  $\pi$ -self-adjoint if and only if

$$
S_{\mathbb{A}} - S_{\mathbb{A}}^* = -i P_{\mathfrak{N}'_i}^+ + i P_{\mathfrak{N}'_{-i}}^+.
$$

The Hilbert space version of the class  $\Omega_A$  in the definition below is found in [\[3\]](#page-18-3), [\[4\]](#page-18-4), [\[40\]](#page-20-2), [\[11\]](#page-19-2).

**Definition 2.** We say that a closed densely defined linear operator  $T$  acting in a Pontryagin space  $\Pi_\kappa$  belongs to the  $class \ \Omega_A$  if:

- (1)  $T \supseteq A$ ,  $T^+ \supseteq A$  where A is a closed Hermitian operator;
- $(2)$  T has a regular point in the lower half-plane;
- (3)  $PT$  and  $PT$ <sup>+</sup> are closed operators.

Note that a closed and non-densely defined  $J$ -regular  $\pi$ -symmetric operator  $A$ admits  $\pi$ -selfadjoint extensions of the class  $\Omega_A$  if and only if its semi-deficiency indices coincide [\[7\]](#page-19-18), [\[40\]](#page-20-2).

An operator A in  $[5^+,\mathfrak{H}_-]$  is called a  $(*)$ -extension [\[11\]](#page-19-2) of an operator T of the class  $\Omega_A$  if both  $A \supset T$  and  $A^{\times} \supset T^+$ . This (\*)-extension is called *correct* [\[11\]](#page-19-2) (regular [\[4\]](#page-18-4)), if an operator  $\text{Re } A = \frac{1}{2}(A + A^{\times})$  is a strong  $\pi$ -self-adjoint bi-extension of an operator A. It is easy to show that if  $\mathbb A$  is a  $(*)$ -extension of T, the T and T<sup>+</sup> are quasi-kernels of  $\mathbb A$  and  $\mathbb A^+$ , respectively.

**Definition 3.** We say the operator T of the class  $\Omega_A$  belongs to the class  $\Lambda_A$  if

- (1)  $T$  admits a correct  $(*)$ -extension;
- (2)  $Gr(A) = Gr(T) \cap Gr(T^+).$

It can be shown (see [\[5\]](#page-18-2)) that if  $T \in \Lambda_A$  then the equation

 $(\mathbb{A} - \lambda I)x = q,$ 

is solvable for all  $\lambda \in \rho(T)$  and all  $g \in \text{Im} \mathbb{A} = \frac{1}{2} (\mathbb{A} - \mathbb{A}^{\times}).$ 

Remark 4. A survey of theory of bi-extensions of symmetric operator in a Hilbert space and its application to characteristic functions of operators of the class  $\Lambda_A$ is presented in [\[40\]](#page-20-2). Bi-extensions of  $\pi$ -symmetric operators in Pontryagin spaces were studied in [\[18\]](#page-19-16).

#### 4. OPERATOR COLLIGATIONS IN  $\Pi_{\kappa}$

In this section we consider linear stationary conservative dynamic systems (l. s. c. d. s.)  $\theta$  of the form

$$
\begin{cases}\n(\mathbb{A} - zI) = K \mathcal{J} \varphi_{-} \\
\varphi_{+} = \varphi_{-} - 2iK^{+}x\n\end{cases}
$$
\n(Im  $\mathbb{A} = K \mathcal{J} K^{*}$ ).

In the system  $\theta$  above A is a bounded linear operator acting from  $\mathfrak{H}^+$  into  $\mathfrak{H}_-,$ where  $\mathfrak{H}^+ \subset \Pi_{\kappa} \subset \mathfrak{H}_-$  is a rigged Pontryagin space, K is a linear bounded operator from a Hilbert space E into  $\mathfrak{H}_-, \mathcal{J} = \mathcal{J}^* = \mathcal{J}^{-1}$  is acting in E,  $\varphi_{\pm} \in E, \varphi_{-}$  is an input vector,  $\varphi_+$  is an output vector, and  $x \in \mathfrak{H}^+$  is a vector of the inner state of the system  $\theta$ .

<span id="page-5-0"></span>For our purposes we need the following more precise definition:

#### <span id="page-5-3"></span>Definition 5. The array

$$
\theta = \begin{pmatrix} A & K & \mathcal{J} \\ \mathfrak{H}^+ \subset \Pi_{\kappa} \subset \mathfrak{H}_- & E \end{pmatrix}
$$
 (4)

is called a linear stationary conservative dynamic system (l.s.c.d.s.) or Brodskii-Livšic rigged operator colligation if

- (1) A is a correct (\*)-extension of an operator T of the class  $\Lambda_A$  for some J-regular operator A with finite and equal deficiency indices;
- (2)  $\mathcal{J} = \mathcal{J}^* = \mathcal{J}^{-1} \in [E, E], \quad \dim E < \infty;$
- (3)  $\mathbb{A} \mathbb{A}^{\times} = 2iK \mathcal{J} K^{+}$ , where  $K \in [E, \mathfrak{H}_{-}]$  ker  $K = \{0\}$   $(K^{+} \in [\mathfrak{H}^{+}, E])$ .

In this case, the operator K is called a *channel operator* and  $\mathcal J$  is called a *direction* operator [\[11\]](#page-19-2). We associate with the system  $\theta$  an operator-valued function

$$
W_{\theta}(z) = I - 2iK^{+}(\mathbb{A} - zI)^{-1}K\mathcal{J}
$$

which is called a transfer operator-valued function of the system  $\theta$  or a characteristic operator-valued function of Brodski-Livšic rigged operator colligations [\[11\]](#page-19-2).

Following [\[16\]](#page-19-4), [\[12\]](#page-19-9) we call an l.s.c.d. system  $\theta$  minimal if the  $\pi$ -symmetric operator  $A$  is such that there are no nontrivial invariant subspaces on which  $A$ induces  $\pi$ -self-adjoint operators. Clearly, the l.s.c.d. system  $\theta$  is minimal if the operator A satisfies the condition [\(3\)](#page-3-0).

Let  $\theta$  be a l.s.c.d.s. of the form [\(4\)](#page-5-0). We consider an operator-valued function

<span id="page-5-2"></span><span id="page-5-1"></span>
$$
V_{\theta}(z) = K^{+} (\mathbb{A}_R - zI)^{-1} K.
$$
 (5)

The transfer operator-function  $W_{\theta}(z)$  of the system  $\theta$  and an operator-function  $V_{\theta}(z)$  of the form [\(5\)](#page-5-1) are connected by the relation

$$
V_{\theta}(z) = i[W_{\theta}(z) + I]^{-1}[W_{\theta}(z) - I]\mathcal{J}
$$
\n(6)

# 5. CLASS  $N_{\kappa}$ . REALIZATION THEOREMS.

Let E be a Hilbert space with an inner product  $(\cdot, \cdot)$  and an operator-valued function  $Q(z)$  belong to  $[E, E]$ .

**Definition 6.** [\[33\]](#page-20-0) An operator-valued function  $V(z) \in [E, E]$  belongs to the class  $N_{\kappa}$  if it is meromorphic in  $\mathbb{C}\setminus\mathbb{R}$  and such that  $V(z) = V(z)^{*}$ ,  $z \in Z_V$ , and the kernel  $\mathbf{r}$  ≠  $\mathbf{r}$  +  $\mathbf{r}$ 

$$
N_V(z,\zeta) = \frac{V(\zeta) - V(z)^*}{\zeta - \bar{z}}, \quad z, \zeta \in Z_V, \quad \zeta \neq \bar{z},
$$
  
\n
$$
N_V(z,\bar{z}) = V'(z), \quad z \in Z_V,
$$
\n(7)

has  $\kappa$  negative squares, i.e. for all  $z_i$  in the domain of holomorphy  $Z_V$  of the meromorphic (in  $\mathbb{C} \setminus \mathbb{R}$ ) function  $V(z)$  and  $h_j \in E$   $(j = 0, 1, ..., n)$ , the form

$$
\sum_{j,k=0}^{n} \left( N_V(z_j, z_k) h_j, h_k \right) \xi_j \bar{\xi}_k \tag{8}
$$

contains at most  $\kappa$  negative squares and for one such a set exactly  $\kappa$  negative squares.

Mention, that the kernel  $N_V(z,\zeta)$  for a function  $V \in N_\kappa$  restricted to the upper half-plane has the same number  $\kappa$  of negative squares, see [\[32\]](#page-19-13).

Class  $N_{\kappa}$  was introduced in [\[33\]](#page-20-0) and studied further in [\[34\]](#page-20-3), [\[17\]](#page-19-19). Different operator models corresponding to  $N_{\kappa}$ -functions are constructed in [\[33\]](#page-20-0), [\[21\]](#page-19-5), [\[22\]](#page-19-6), [\[23\]](#page-19-20), [\[28\]](#page-19-21).

<span id="page-6-0"></span>**Definition 7.** [\[11\]](#page-19-2) An operator-valued function  $V(z)$  in a finite-dimensional Hilbert space E is called realizable if, in some domain  $\mathcal{D} \subset \mathbb{C}_-$ ,  $V(z)$  can be represented in the form

$$
V_{\theta}(z) = i[W_{\theta}(z) + I]^{-1}[W_{\theta}(z) - I],
$$
\n(9)

where  $W_{\theta}(z)$  is a transfer operator-function of some l.s.c.d.s.  $\theta$  with the direction operator  $\mathcal{J} = I$ .

<span id="page-6-4"></span>**Definition 8.** An operator-function  $V(z) \in [E, E]$  (dim  $E < \infty$ ) belongs to the class  $N_{\kappa}(R)$  if the following conditions are met:

<span id="page-6-1"></span>
$$
(1) V \in N_{\kappa};
$$

(2) for all  $f \in E$ 

$$
\lim_{y \uparrow \infty} \frac{\left(V(iy)f, f\right)}{y} = 0; \tag{10}
$$

<span id="page-6-2"></span>(3) For all  $z \in Z_V$ 

<span id="page-6-3"></span>
$$
\bigcap_{\zeta \in Z_V} \ker N_V(\zeta, z) = \{0\};\tag{11}
$$

(4) for all 
$$
f \in \mathcal{B} = \{ f \in E \mid \lim_{y \uparrow \infty} y(\operatorname{Im} V(iy)f, f) < \infty \}
$$
  

$$
\lim_{y \uparrow \infty} V(iy)f = 0.
$$
 (12)

<span id="page-6-5"></span>**Theorem 9.** Let  $\theta$  be a minimal l.s.c.d.s. of the form [\(4\)](#page-5-0) with  $\mathcal{J} = I$  in E,  $\dim E < \infty$ . Then the operator-function  $V_{\theta}(z)$  of the form [\(9\)](#page-6-0) admits a holomorphic continuation to a function  $V(z)$  which belongs to the class  $N_{\kappa}(R)$ .

*Proof.* For a l.s.c.d.s  $\theta$  of the form [\(4\)](#page-5-0) consider

$$
V(z) = K^+(\mathbb{A}_R - zI)^{-1}K, \quad z \in \rho(\hat{A}_R),
$$

where  $A_R$  is the quasi-kernel of the operator  $A_R$ . As follows from [\(6\)](#page-5-2)  $V(z)$  is a holomorphic continuation of the function  $V_{\theta}(z)$ . We set

$$
\Gamma_z = (\mathbb{A}_R - zI)^{-1}K, \quad z \in \rho(\hat{A}_R).
$$

It can be seen that the operator  $\Gamma_z$  is invertible and the following relation is valid

$$
\Gamma_z = (\mathbb{A}_R - \zeta I)(\mathbb{A}_R - zI)^{-1} \Gamma_{\zeta}, \quad z, \quad z \in \rho(\hat{A}_R).
$$

Let  $z_0 \in \rho(\hat{A}_R)$  and let  $\tilde{A}$  be a  $\pi$  - symmetric operator defined as follows  $\mathcal{D}(\tilde{A}) = \left\{ f \in \mathcal{D}(\hat{A}_R) \mid \left[ (\hat{A}_R - \bar{z}_0 I) f, \Gamma_{z_0} g \right] = 0, \forall g \in E \right\}, \ \tilde{A}f = \hat{A}_R f, \ f \in \mathcal{D}(\hat{A}_R),$ 

<span id="page-7-0"></span>Since

$$
\frac{V(z) - V^*(\zeta)}{z - \bar{\zeta}} = \frac{K^+(\mathbb{A}_R - zI)^{-1}K - K^+(\mathbb{A}_R - \bar{\zeta}I)^{-1}K}{z - \bar{\zeta}}
$$
\n
$$
= K^+(\mathbb{A}_R - \bar{\zeta}I)^{-1}(\mathbb{A}_R - zI)^{-1}K
$$
\n
$$
= \Gamma_{\zeta}^+\Gamma_z, \quad z, \quad z \in \rho(\hat{A}_R), \tag{13}
$$

the operator-function  $V(z)$  is a Krein-Langer Q-function [\[32\]](#page-19-13), [\[33\]](#page-20-0) for the  $\pi$  - symmetric operator  $\tilde{A}$  and its  $\pi$ -self-adjoint extension  $\hat{A}_R$ . Let  $f \in \mathcal{D}(\tilde{A})$ . Then for any  $g \in \mathfrak{H}^+$ 

$$
[\text{Im } \mathbb{A}f, g] = [f, \text{Im } \mathbb{A}g] = [f, (\mathbb{A}_R - \bar{z}_0 I)\Gamma_{z_0} K^+ g] = [(\hat{A}_R - \bar{z}_0 I)f, \Gamma_{z_0} K^+ g] = 0.
$$

Thus, Im  $\mathbb{A}f = 0$  for  $f \in \mathcal{D}(\tilde{A})$  and  $\mathbb{A} \supset T \supset \tilde{A}$ ,  $\mathbb{A}^{\times} \supset T^{+} \supset \tilde{A}$ . But this is possible only when  $\tilde{A} = A$  since A is the maximal  $\pi$ -symmetric part of the operator T. Consequently,  $\Gamma_z E = \mathfrak{N}_z$  and  $V_\theta(z)$  is a Krein-Langer Q-function for a prime  $\pi$  - symmetric operator A. It was shown in [\[33\]](#page-20-0) that a Q-function of a prime  $\pi$  symmetric operator (in the upper half-plane) belongs to the class  $N_{\kappa}$  and condition [\(10\)](#page-6-1) holds.

In order to prove  $(11)$  we note that since the operator A is prime then for all  $f \in \bigcap_{\zeta \in Z_V} \ker N_V(\zeta, z)$  and all  $g \in E$  we have

$$
[\Gamma_z f, \Gamma_\zeta g] = \left(\frac{V_\theta(z) - V_\theta^*(\zeta)}{z - \overline{\zeta}} f, g\right) = 0,
$$

and hence  $f = 0$ .

<span id="page-7-4"></span><span id="page-7-3"></span>It was shown in [\[33\]](#page-20-0) that

$$
\mathfrak{N}_z \cap \mathcal{D}(\hat{A}_R) = \Gamma_z \mathcal{B}, \quad z \in \rho(\hat{A}_R). \tag{14}
$$

It is easy to see that

<span id="page-7-2"></span>
$$
(\mathbb{A}_R - zI)(\mathfrak{N}_z \cap \mathcal{D}(\hat{A}_R)) = \mathfrak{L},\tag{15}
$$

where  $\mathfrak{L} = \mathcal{D}(A)^{[\perp]}$ . Clearly, dim  $\mathcal{B} < \infty$ , since  $\mathcal{B} \subset E$ .

In order to prove [\(12\)](#page-6-3) we need to use the spectral decomposition of the  $\pi$ -selfadjoint operator  $\mathbb{A}_R$ . It was shown in [\[5\]](#page-18-2) that for all  $g \in \mathcal{D}(\mathbb{A}_R)$ 

$$
\mathbb{A}_R g = \lim_{z = iy \uparrow \infty} (-z^2) \left[ (\mathbb{A}_R - zI)^{-1} + \frac{1}{z} \right] g.
$$
 (16)

It follows from [\(13\)](#page-7-0) that

<span id="page-7-1"></span>
$$
(-z^2)\Gamma_{z_0}^+\left[ (\mathbb{A}_R - zI)^{-1} + \frac{1}{z} \right] \Gamma_{z_0} = \frac{-z^2[V_\theta(z) - V_\theta^*(z_0)]}{(z - z_0)(z - \bar{z}_0)} + \frac{zz_0}{z - z_0} \Gamma_{z_0}^+ \Gamma_{z_0} \tag{17}
$$

Taking the limit in [\(17\)](#page-7-1) and using [\(16\)](#page-7-2), we obtain for all  $f \in \mathcal{B}$ 

$$
\Gamma_{z_0}^+ A_R \Gamma_{z_0} f = - \lim_{z = iy \uparrow \infty} V_\theta(z) f + V_\theta^*(z_0) f + z_0 \Gamma_{z_0}^+ \Gamma_{z_0} f
$$
\n
$$
= - \lim_{z = iy \uparrow \infty} V_\theta(z) f + \text{Re } V_\theta(z_0) f + \text{Re } z_0 \Gamma_{z_0}^+ \Gamma_{z_0} f.
$$
\n(18)

Taking into account that  $\text{Re } V_{\theta}(z_0) = \Gamma_{z_0}^+ \mathbb{A}_R \Gamma_{z_0} f - \text{Re } z_0 \Gamma_{z_0}^+ \Gamma_{z_0}$  we get [\(12\)](#page-6-3).  $\Box$ 

<span id="page-8-7"></span>**Theorem 10.** Let an operator-valued function  $V(z)$  belong to the class  $N_{\kappa}(R)$ . Then  $V(z)$  admits a minimal realization by a system  $\theta$  of the form [\(4\)](#page-5-0) with  $\mathcal{J} = I$ .

*Proof.* It was shown in [\[33\]](#page-20-0) that if an operator-function  $V(z)$  in [E, E] satisfies the conditions 1) and 2) of Definition [8](#page-6-4) then it is possible to construct a Pontryagin space  $\Pi_{\kappa}(V)$  and a symmetric operator A with a  $\pi$ -self-adjoint extension  $A_V$  in  $\Pi_{\kappa}(V)$ . The strictness condition 3) guarantees that the function  $V(z)$  is a Krein-Langer  $Q$ -function of a pair  $A$  and  $A_V$ . We will use a reproducing kernel space model for the operators A,  $A_V$  elaborated in [\[2\]](#page-18-1), [\[20\]](#page-19-22) and [\[22\]](#page-19-6).

Step 1. Let us consider the reproducing kernel Pontryagin space  $\Pi_{\kappa}(V)$  corresponding to the kernel  $N(z,\zeta) := N_V(z,\zeta)$ . The latter means see [\[1\]](#page-18-0) that  $\Pi_{\kappa}(V)$ consists of functions holomorphic on  $Z_V$  and for each  $z \in Z_V$  and  $h \in E$  the followings hold:

- (a)  $N(z,\zeta)h$  belongs to  $\Pi_{\kappa}(V)$  as a function on  $\zeta$ ;
- (b)  $[f(\cdot), N(z,\zeta)h] = (f(z),h)_E$  for every  $f(\cdot)$  in  $\Pi_{\kappa}(V)$ .

In particular, it follows from (b) that the evaluation operator  $f \mapsto (f(z), h)_E$  is continuous in  $\Pi_{\kappa}(V)$ . The multiplication operator

<span id="page-8-5"></span>
$$
A: f(\zeta) \mapsto \zeta f(\zeta), \tag{19}
$$

with the domain dom  $A = \{f \in \Pi_{\kappa}(V) : \zeta f(\zeta) \in \Pi_{\kappa}(V)\}\$ is a prime closed symmetric operator in  $\Pi_{\kappa}(V)$ . Let  $\mathcal{A} := Gr(A)$  As was shown in [\[22\]](#page-19-6) the adjoint linear relation  $\mathcal{A}^+$  takes the form

<span id="page-8-2"></span>
$$
\mathcal{A}^+ = \{ \{f, \tilde{f}\} \in \Pi_{\kappa}(V)^2 : \tilde{f}(\zeta) - \zeta f(\zeta) = h_1 - V(\zeta)h_0; \ h_0, h_1 \in E \}
$$

and the linear relation

$$
\mathcal{A}_V = \{ \{f, \tilde{f}\} \in \Pi_{\kappa}(V)^2 : \tilde{f}(\zeta) - \zeta f(\zeta) = h_1 \in E \}
$$
(20)

is a self-adjoint extension of A with  $\rho(A_V) = Z_V$ .

The deficiency subspace  $\mathfrak{N}_z$  of  $A$  ( $z \in Z_V$ ) consists of vector-functions  $\gamma(z)h :=$  $N(\bar{z},\cdot)h, h \in E$ . The mapping  $\gamma(z): E \to \mathfrak{N}_z$  is injective since the assumption  $\gamma(z)h = 0$  and the equality

<span id="page-8-4"></span>
$$
[N(\bar{z}, \cdot)h, N(\bar{w}, \cdot)g] = (N(\bar{z}, \bar{w})h, g)_E = \left(\frac{V(z) - V(w)^*}{z - \bar{w}}h, g\right)_E
$$
(21)

imply that  $h \in \bigcap$  $\bigcap_{w \in Z_V} \ker(V(z) - V(w)^*)$ . It follows from the hypothesis 3) that  $h = 0$ . The linear span of deficiency subspaces  $\mathfrak{N}_z$  ( $z \in Z_V$ ) is dense in  $\Pi_{\kappa}(V)$  and, hence, the operator V is a prime symmetric operator in  $\Pi_{\kappa}(V)$ .

Let us show that  $\gamma(z)$  satisfies the identity

<span id="page-8-3"></span>
$$
\gamma(z) = (\mathcal{A}_V - z_0)(\mathcal{A}_V - z)^{-1} \gamma(z_0), \quad z, z_0 \in Z_V.
$$
 (22)

<span id="page-8-0"></span>This is straightforward from the identities

<span id="page-8-6"></span><span id="page-8-1"></span>
$$
zN(\bar{z},\zeta)h - \zeta N(\bar{z},\zeta)h = V(z)h - V(\zeta)h,\tag{23}
$$

$$
z_0 N(\bar{z}_0, \zeta)h - \zeta N(\bar{z}_0, \zeta)h = V(z_0)h - V(\zeta)h.
$$
 (24)

Subtracting [\(24\)](#page-8-0) from [\(23\)](#page-8-1) one obtains due to [\(20\)](#page-8-2)

$$
\{N(\bar{z},\zeta)h - N(\bar{z}_0,\zeta), zN(\bar{z},\zeta)h - z_0N(\bar{z}_0,\zeta)h\} \in \mathcal{A}_V.
$$
 (25)

The latter equality is equivalent to [\(22\)](#page-8-3).

The identities [\(22\)](#page-8-3) and [\(21\)](#page-8-4) show that  $V(z)$  is the Kreĭn-Langer Q-function of the pair A,  $A_V$ . The assumption [\(10\)](#page-6-1) implies that mul  $A_V = \{0\}$ , see [\[32\]](#page-19-13), and, therefore,  $\mathcal{A}_V$  is the graph of an operator  $A_V$ .

<span id="page-9-0"></span>Step 2. The operator A need not be densely defined. Let us calculate vectorfunctions from the subspace  $\mathcal{L} = \mathcal{D}(A)^{[\perp]}$  explicitly. As follows from [\(14\)](#page-7-3) and[\(15\)](#page-7-4)

<span id="page-9-1"></span>
$$
\gamma(z)\mathcal{B} = \mathfrak{N}_z \cap \mathcal{D}(A_V) = (A_V - z)^{-1}\mathcal{L}.
$$
 (26)

This implies that for every  $h \in \mathcal{B}$  there exists a strong limit of the vector-function

<span id="page-9-2"></span>
$$
h_{\infty}(\cdot) := \lim_{y \uparrow \infty} (iy)\gamma(-iy)h = \lim_{y \uparrow \infty} (iy)N(iy, \cdot)h \tag{27}
$$

as  $y \to \infty$ . Since the evaluation operator is continuous in  $\Pi_{\kappa}(V)$  one obtains from the hypothesis 4) for every  $z \in Z_V$ 

$$
h_{\infty}(z) = \lim_{y \uparrow \infty} (iy) N(iy, z)h
$$
  
= 
$$
\lim_{y \uparrow \infty} (iy) \frac{V(z) - V(-iy)}{z + iy} h = V(z)h.
$$
 (28)

Therefore, the subspace  $\mathcal L$  takes the form

<span id="page-9-5"></span>
$$
\mathcal{L} = \{V(\cdot)h : h \in \mathcal{B}\}.
$$

It follows from [\(23\)](#page-8-1) and [\(20\)](#page-8-2) that for every  $h \in \mathcal{B}$ 

<span id="page-9-3"></span>
$$
(Av - z)^{-1}V(\cdot)h = N(\bar{z}, \cdot)h \quad (z \in Z_V). \tag{29}
$$

One can derive the same equality from [\(26\)](#page-9-0), [\(27\)](#page-9-1), [\(28\)](#page-9-2) and [\(22\)](#page-8-3).

<span id="page-9-4"></span>Step 3. Let us show that for every  $g \in E$  the function  $V(\cdot)g$  generates a functional on  $\mathcal{D}(A^+) = \mathcal{D}(A_V) + \mathfrak{N}_i$  by the formulas

$$
[(A_V - z_0)^{-1} f(\cdot), V(\cdot)g] = (f(z_0), g)_E, \quad f \in \Pi_{\kappa}(V),
$$
 (30)

$$
[N(\bar{z}_0, \cdot)h, V(\cdot)g] = (V(z_0)h, g)_E, \quad h \in E,
$$
\n(31)

which is continuous in the norm of  $\mathfrak{H}^+ = \mathcal{D}(A^+).$ 

Mention first that the formulas [\(30\)](#page-9-3) and [\(31\)](#page-9-4) are consistent since for  $f(\cdot) = V(\cdot)h$  $(h \in \mathcal{B})$  one obtains from [\(29\)](#page-9-5) and [\(30\)](#page-9-3) the formula

$$
[(A_V - z_0)^{-1}V(\cdot)h, V(\cdot)g] = (V(z_0)h, g)_E,
$$

which agrees with [\(31\)](#page-9-4).

Next, it follows from  $(30)$ ,  $(31)$ ,  $(22)$  and the identity

$$
(A_V - z)^{-1} = (A_V - z_0)^{-1} (A_V - z_0)(A_V - z)^{-1}
$$

<span id="page-9-6"></span>that

$$
[N(\bar{z}, \cdot)h, V(\cdot)g] = \left( \left[ V(z_0)v + (z - z_0) \frac{V(z) - V(z_0)}{z - z_0} h \right], g \right)_E = (V(z)h, g)_E.
$$
\n(32)

Let now a sequence  $\varphi_n \in \mathcal{D}(A_V)$  converges to  $\varphi \in \mathcal{D}(A_V)$  in  $\mathfrak{H}^+$ -norm. Then

$$
f_n(\cdot) = (A_V - z_0)\varphi_n(\cdot) \to f(\cdot) = (A_V - z_0)\varphi(\cdot)
$$
 strongly in  $\Pi_{\kappa}(V)$ 

and by continuity of the evaluation operator one has

$$
f_n(z) \to f(z) \quad \forall z \in Z_V.
$$

Then it follows from [\(30\)](#page-9-3) that

$$
[\varphi_n(\cdot),V(\cdot)g]\to [\varphi(\cdot),V(\cdot)g] \quad (n\to\infty)
$$

and, therefore, the functional generated by  $V(\cdot)g$  via [\(30\)](#page-9-3) and [\(31\)](#page-9-4) is continious, since dim  $\mathfrak{H}^+ \text{(mod } \mathcal{D}(A_V)) < \infty$ .

Step 4. Using the operator A defined in  $(19)$  we construct a rigged Pontryagin space  $\mathfrak{H}^+ \subset \Pi_{\kappa}(V) \subset \mathfrak{H}_-$  the way it was described in the section [3.](#page-3-1) The functional  $V(\cdot)h, h \in E$  considered above can be viewed as an element from  $\mathfrak{H}_-$ . Let us define a linear operator  $K : E \to \mathfrak{H}_-$  by the equality

<span id="page-10-2"></span><span id="page-10-1"></span>
$$
Kh := V(\cdot)h, \quad h \in E. \tag{33}
$$

<span id="page-10-0"></span>Clearly the operator K is invertible, otherwise  $V(z) \equiv 0$  and definition [8](#page-6-4) is violated. Let us extend the operator  $A_V$  to the linear operator  $A_R : \mathfrak{H}^+ = \mathcal{D}(A_V) + \mathfrak{N}_{z_0} \to$  $\mathfrak{H}_-$  by the equality

$$
\mathbb{A}_R N(\bar{z}_0, \cdot) g = z_0 N(\bar{z}_0, \cdot) g + V(\cdot) g, \quad g \in E. \tag{34}
$$

This definition agrees with [\(29\)](#page-9-5) for  $g \in \mathcal{B} \subset E$ . The operator  $\mathbb{A}_R$  is a  $\pi$ -self-adjoint bi-extension of A with the quasi-kernel  $A_V$ . It follows from [\(25\)](#page-8-6) and [\(34\)](#page-10-0) that

$$
(\mathbb{A}_R - z)N(\bar{z}, \cdot)g = V(\cdot)g, \quad g \in E. \tag{35}
$$

Making use of [\(33\)](#page-10-1) and [\(35\)](#page-10-2) one obtains

$$
K^+(\mathbb{A}_R - z)^{-1} Kg = K^+ N(\bar{z}, \cdot)g, \quad z \in Z_V.
$$

An application of [\(32\)](#page-9-6) and [\(33\)](#page-10-1) implies

$$
(K^+N(\bar{z},\cdot)g,h)=[N(\bar{z},\cdot)g,V(\cdot)h]=(V(z)g,h)_E,
$$

and, therefore,

$$
K^+(\mathbb{A}_R - z)^{-1}K = V(z), \quad z \in Z_V.
$$
  
Step 5. We set  $\mathbb{A}_I = KK^+$  and define

$$
\mathbb{A} = \mathbb{A}_R + i\mathbb{A}_I.
$$

It is obvious that the l.s.c.d.s.

$$
\theta = \begin{pmatrix} A & I \\ \mathfrak{H}^+ \subset \Pi_{\kappa}(V) \subset \mathfrak{H}_- & E \end{pmatrix}
$$
 (36)

satisfies conditions (2) and (3) of Definition [5.](#page-5-3) What remains to show then is that the quasi-kernel T of the operator A belongs to the class  $\Lambda_A$ . To do this it suffices to show that  $\rho(T) \cap \mathbb{C}_-$  is nonempty and check that A is the maximal symmetric part of T and  $T^+$ . Since  $V \in N_{\kappa}$  it follows from [\[34,](#page-20-3) Theorem 2.2] that  $V(z) - iI$  has at most  $\kappa$  zeros in  $\mathbb{C}_-$ . Let us assume without loss of generality that the operator  $(V(z_0) - iI)$  is invertible. Consequently, the operator

$$
H = I + iK^{+}(\mathbb{A}_{R} - z_{0}I)^{-1}K = I + iV(z_{0}) \in [E, E],
$$

is invertible as well. It follows from [\(28\)](#page-9-2), [\(34\)](#page-10-0), [\(35\)](#page-10-2) that

$$
\mathbb{A}N(\bar{z}_0,\cdot)g=z_0N(\bar{z}_0,\cdot)g+V(\cdot)Hg,\quad g\in E.
$$

Since H is invertible this implies  $\Re(A - z_0 I) \supset \Re(K)$ .

Let  $f \in \mathfrak{N}_z$ . Then  $(A - z_0) f = g + (z - z_0) f$ , where  $g = (A_R - zI) f + iA_I f$ . Since  $\Re(\mathbb{A}_I) \subset \Re(K)$  and  $(\mathbb{A}_R - zI)\mathfrak{N}_z = \Re(K)$ , we have  $g \in \Re(K)$  and therefore there is an x such that  $(A - z_0I)x = g$ . Thus

$$
(\mathbb{A} - z_0 I)(f - x) = (z - z_0)f,
$$

i.e.,  $\Re(\mathbb{A}-z_0I)\supset \mathfrak{N}_z$ . Since the operator A is prime, c.l.s.  $\{\mathfrak{N}_z, z\neq \bar{z}\} = \Pi_{\kappa}(V)$ , we have  $\overline{\mathfrak{R}(T - z_0 I)} = \Pi_{\kappa}(V)$ . On the other hand, the relations

$$
\Re(T-z_0I)\supset \mathfrak{M}_{z_0}, \quad \dim \mathfrak{R}(T-z_0I)(\mod(\mathfrak{M}_{z_0}))<\infty,
$$

imply that  $\Re(T - z_0 I)$  is closed and hence coincides with  $\Pi_{\kappa}(V)$ . Similarly, one proves that  $\Re(T^+ - z_0 I) = \Pi_{\kappa}(V), T^+$  is a quasi-kernel of  $\mathbb{A}^{\times}$ , and concludes that A is a correct  $(*)$ -extension of T.

To prove that A is the maximal symmetric part of T and  $T^+$ , we assume the contrary. Then there exists such a  $\pi$ -symmetric operator  $A_0$  that

$$
T \supset A_0 \supset A, \quad T^+ \supset A_0 \supset A.
$$

Consequently, for all  $f \in \mathcal{D}(A_0)$ ,  $\mathbb{A}f = \mathbb{A}^{\times}f = A_0f$ ,

$$
\mathbb{A}_I f = \frac{(\mathbb{A} - \mathbb{A}^{\times})}{2i} f = 0, \quad \mathbb{A}_R f = (\mathbb{A} - \mathbb{A}_I) f = A_0 f \in \Pi_{\kappa}(V).
$$

Thus,  $f \in \mathcal{D}(A_V)$  and  $A_V f = A_0 f$ . For any  $g \in \mathfrak{H}^+$  we have

$$
[(A_V - \bar{z}_0 I)f, \gamma(z_0)K^+g] = [f, A_I g] = [A_I f, g] = 0.
$$

But since  $\Re(\gamma(z_0)K^+) = \mathfrak{N}_{z_0}$ , then  $f \in \mathcal{D}(A)$  and  $\mathcal{D}(A_0) = \mathcal{D}(A)$ . Thus A is the maximal symmetric part of  $T$  and  $T^+$ .

 $\Box$ 

Remark 11. We should mention that the realization results obtained in Theorem [10](#page-8-7) can be interpreted as realization with impedance systems  $\Delta$  of the form [\(1\)](#page-1-1) with  $\mathbb{D} = \mathbb{A}_R$  and

$$
V_{\Delta}(z) = K^+(\mathbb{A}_R - zI)^{-1}K.
$$

Remark 12. In the recent paper [\[35\]](#page-20-4) the authors derive an alternative integral representation for matrix-functions of the class  $N_{\kappa}$ . It can be easily shown that all functions of the class  $N_{\kappa}(R)$  fall into the special class described in Theorem 4.1 of [\[35\]](#page-20-4) and permit a reduced Krein-Langer integral representation developed in that theorem.

# 6. SUBCLASSES OF THE CLASS  $N_{\kappa}(R)$

<span id="page-11-0"></span>In this section we follow [\[11\]](#page-19-2) and introduce three distinct subclasses of the class of realizable operator-valued functions  $N_{\kappa}(R)$ .

**Definition 13.** An operator-function  $V(z)$  of the class  $N_{\kappa}(R)$  belongs to the subclass  $N_{\kappa}^{0}(R)$  if in the definition [8](#page-6-4) the subspace  $\mathcal{B}$  is trivial, i.e.,

$$
\lim_{y \uparrow \infty} y(\operatorname{Im} V(iy)f, f) = \infty, \quad \forall f \in E, f \neq 0.
$$
\n(37)

**Definition 14.** An operator-function  $V(z)$  of the class  $N_{\kappa}(R)$  belongs to the subclass  $N_{\kappa}^1(R)$  if in the definition [8](#page-6-4) the subspace  $\mathcal{B} = E$ , i.e.,

$$
\lim_{y \uparrow \infty} y(\operatorname{Im} V(iy)f, f) < \infty, \quad \forall f \in E, f \neq 0. \tag{38}
$$

**Definition 15.** An operator-function  $V(z)$  of the class  $N_{\kappa}(R)$  belongs to the subclass  $N_{\kappa}^{01}(R)$  if in the definition [8](#page-6-4) the subspace  $\mathcal{B}$  is neither trivial nor equals E, i.e.

<span id="page-11-1"></span>
$$
\{0\} \varsubsetneq \mathcal{B} \varsubsetneq E.
$$

One may notice that  $N(R)$  is a union of three distinct subclasses  $N_0(R)$ ,  $N_1(R)$ and  $N_{01}(R)$ . Now we prove the direct and inverse realization theorems in each of the subclasses.

<span id="page-12-1"></span>**Theorem 16.** Let  $\theta$  be a l.s.c.d.s. of the form [\(4\)](#page-5-0) such that  $\mathcal{J} = I$  and A is an operator with dense domain. Then operator-function  $V_{\theta}(z)$  of the form [\(5\)](#page-5-1), [\(6\)](#page-5-2) has a holomorphic continuation  $V(z)$  which belongs to the class  $N_{\kappa}^{0}(R)$ .

Conversely, let an operator-valued function  $V(z)$  belong to the class  $N_{\kappa}^{0}(R)$ . Then  $V(z)$  admits a minimal realization by a system  $\theta$  of the form [\(4\)](#page-5-0) with a densely defined operator A.

*Proof.* Since the operator A is densely defined  $\mathcal{D}(\tilde{A}) \cap \mathfrak{N}_z = \{0\}$  for every selfadjoint extension  $\widetilde{A}$  of A. In particular, one obtains  $\mathcal{D}(\widehat{A}_R) \cap \mathfrak{N}_z = \{0\}$ . Due to [\(14\)](#page-7-3) this implies  $\mathcal{B} = \{0\}.$ 

Conversely, if  $\mathcal{B} = \{0\}$  then it follows from [\(14\)](#page-7-3) and [\(15\)](#page-7-4) that  $\mathcal{L} = \{0\}$ , and, nce, the operator A is densely defined. hence, the operator A is densely defined.

In order to proceed with the similar results in the class  $N_{\kappa}^1(R)$  we need to recall the definition of O-operator [\[12\]](#page-19-9) and give its analogue for the spaces with indefinite metric. A J-regular  $\pi$ -symmetric operator A is called an O-operator if its semi-deficiency indices are equal to zero. As it was shown in [\[5\]](#page-18-2) for an operator  $T \in \Lambda_A$ there exist linear operators

$$
\begin{aligned} M_T: \mathfrak{N}_i' \boxplus \mathfrak{N} &\rightarrow \mathfrak{N}_{-i}' \boxplus \mathfrak{N}, \\ M_{T^+}: \mathfrak{N}_{-i}' \boxplus \mathfrak{N} &\rightarrow \mathfrak{N}_i' \boxplus \mathfrak{N}, \end{aligned}
$$

<span id="page-12-0"></span>such that  $\mathcal{M}_T = Gr(M_T)$  and  $\mathcal{M}_{T^+} = Gr(M_{T^+})$  have trivial intersections with the manifold  $\{x, x\} : x \in \mathfrak{N}\}\$ and

$$
\mathcal{D}(T) = \mathcal{D}(A) \boxplus \{x - y \mid \langle x, y \rangle \in \mathcal{M}_T\},\tag{39}
$$

$$
\mathcal{D}(T^+) = \mathcal{D}(A) \boxplus \{x - y \mid \langle x, y \rangle \in \mathcal{M}_{T^+}\}.
$$
 (40)

If A is an O-operator then  $M_T$  and  $M_{T^+}$  are operators in  $\mathfrak{N}$  (dim  $\mathfrak{N} < \infty$ ) with  $1 \in$  $\rho(M_T) \cap \rho(M_{T^+})$  and the relations [\(39\)](#page-12-0)-[\(40\)](#page-12-0) imply that  $\mathcal{D}(T) = \mathcal{D}(T^+) = \mathfrak{N}, \text{Im}(T)$ is a bounded selfadjoint operator in  $\Pi_{\kappa}$ , and  $\text{Re}(T)$  is a  $\pi$ -selfadjoint extension of A in  $\Pi_{\kappa}$ .

<span id="page-12-2"></span>**Theorem 17.** Let  $\theta$  be a l.s.c.d.s. of the form [\(4\)](#page-5-0) such that  $\mathcal{J} = I$ . A is an Ooperator, and  $\mathcal{D}(T) = \mathcal{D}(T^+)$ . Then operator-function  $V_{\theta}(z)$  of the form [\(5\)](#page-5-1), [\(6\)](#page-5-2) has a holomorphic continuation  $V(z)$  which belongs to the class  $N_{\kappa}^1(R)$ .

Conversely, let an operator-valued function  $V(z)$  belongs to the class  $N_{\kappa}^1(R)$ . Then  $V(z)$  admits a minimal realization by a system  $\theta$  of the form [\(4\)](#page-5-0) with a non-densely defined O-operator A.

*Proof.* Once again using Theorem [9](#page-6-5) we have that  $V(z) \in N_{\kappa}(R)$  and need to show [\(38\)](#page-11-1). Since A is an O-operator,  $\mathfrak{N}'_z = 0$ , and hence dim  $\mathfrak{N}_z = \dim \mathfrak{N}$ . Using [\(14\)](#page-7-3) we have

$$
\dim(\Gamma_z \mathcal{B}) = \dim(\mathcal{D}(A_R) \cap \mathfrak{N}_z) = \dim \mathfrak{N}.
$$

On the other hand,  $\mathfrak{N}_z = \Gamma_z E$  and thus dim  $\Gamma_z E = \dim \mathfrak{N}$ . Consequently, since  $\mathcal{B}$ is a subspace of E and dim  $E = \dim \mathcal{B}$  we have  $\mathcal{B} = E$ .

Conversely, if  $\mathcal{B} = E$ , then using [\(14\)](#page-7-3) we get  $\Gamma_z \mathcal{B} = \Gamma_z E = \mathcal{D}(\widehat{A}_R) \cap \mathfrak{N}_z = \mathfrak{N}_z$ . Applying [\(15\)](#page-7-4) yields

$$
\dim ((\mathbb{A}_R - zI)\mathfrak{N}_z) = \dim \mathfrak{L} = \dim \mathfrak{N}, \quad z \in \rho(\hat{A}_R).
$$

Since  $(A_R - zI)$  is an invertible operator for  $z \in \rho(A_R)$  we conclude that dim  $\mathfrak{N}_z =$ dim  $\mathfrak{N}$ . It can be shown (see [\[5\]](#page-18-2), [\[40\]](#page-20-2)) that the operator  $P_{\mathfrak{M}}^{+}$  described in the section [3](#page-3-1) is a bijective mapping from  $\mathfrak{N}_{\pm i}$  onto  $\mathfrak{N}_{\pm i}^{\prime} \boxplus \mathfrak{N}$ . Considering the above we have then  $\dim \mathfrak{N}'_{\pm i} \boxplus \mathfrak{N} = \dim \mathfrak{N}$ . This proves that  $\mathfrak{N}'_{\pm i} = 0$  and thus A is an O-operator.  $\Box$ 

Remark 18. If  $V \in N_{\kappa}^{0}(R)$  then the operator A in the realization [\(4\)](#page-5-0) is densely defined and this implies that  $\mathcal{D}(T) \neq \mathcal{D}(T^+)$  for the operator T mutually disjoint with  $T^+$ . When the operator A is nondensely defined even mutually disjoint operators T and  $T^+$  may have the same domain. In fact the equality  $\mathcal{D}(T) = \mathcal{D}(T^+)$ holds if  $V \in N^1_{\kappa}(R)$ . In this case we may not consider the bi-extensions of T in the rigged Pontryagin space and the corresponding l.s.c.d.s. can be written as follows

$$
\theta = \begin{pmatrix} T & K & \mathcal{J} \\ \Pi_{\kappa} & & E \end{pmatrix}.
$$

**Theorem 19.** Let  $\theta$  be a l.s.c.d.s. of the form [\(4\)](#page-5-0) such that  $\mathcal{J} = I$ ,  $\overline{\mathcal{D}(A)} \neq \Pi_{\kappa}$ , and  $\mathcal{D}(T) \neq \mathcal{D}(T^+)$ . Then operator-function  $V_{\theta}(z)$  of the form [\(5\)](#page-5-1), [\(6\)](#page-5-2) has a holomorphic continuation  $V(z)$  which belongs to the class  $N_{\kappa}^{01}(R)$ .

Conversely, let an operator-valued function  $V(z)$  belongs to the class  $N_{\kappa}^{01}(R)$ . Then  $V(z)$  admits a minimal realization by a system  $\theta$  of the form [\(4\)](#page-5-0) with a non-densely defined operator A and  $\mathcal{D}(T) \neq \mathcal{D}(T^+)$ .

The proof is immediate from Theorem [16](#page-12-1) and Theorem [17.](#page-12-2)

## 7. Applications to the scalar case

<span id="page-13-0"></span>In this section we consider scalar functions of the class  $N_\kappa.$  We will establish the link between scalar  $(E = \mathbb{C})$  realizable functions of the class  $N_{\kappa}(R)$  and a class of realizable Nevanlinna functions [\[12\]](#page-19-9).

The realization problems of the present type for Nevanlinna operator-valued functions were studied in details in [\[12\]](#page-19-9) and [\[13\]](#page-19-3) where similar subclass structure was developed. In particular, it was shown that any realizable Nevanlinna operatorfunction  $V(z)$  admits an integral representation

<span id="page-13-2"></span>
$$
V(z) = Q + F \cdot z + \int_{-\infty}^{+\infty} \left( \frac{1}{t - z} - \frac{t}{1 + t^2} \right) dG(t), \tag{41}
$$

in the Hilbert space E. In this representation  $Q = Q^*, F = 0, G(t)$  is non-decreasing operator-function on  $(-\infty, +\infty)$  for which

$$
\int_{-\infty}^{+\infty} \frac{dG(t)}{1+t^2} \in [E, E],
$$

and

$$
Qe = \int_{-\infty}^{+\infty} \frac{t}{1+t^2} \, dG(t)e
$$

<span id="page-13-1"></span>for all  $e \in E_{\infty}$  where

$$
E_{\infty} = \left\{ e \in E : \int_{-\infty}^{+\infty} \left( dG(t)e, e \right)_E < \infty \right\}.
$$
 (42)

The class of all realizable Nevanlinna operator-functions is called  $N(R)$  (see [\[12\]](#page-19-9)). The three subclasses of the class  $N(R)$  were introduced in [\[13\]](#page-19-3) and are called  $N^0(R)$ ,  $N^1(R)$ , and  $N^{01}(R)$ . Each subclass is described in terms of the subspace  $E_{\infty}$  in [\(42\)](#page-13-1) determined by the variation of measure  $G(t)$  in the representation [\(41\)](#page-13-2). In particular,  $E_{\infty} = \{0\}$  for the class  $N^{0}(R)$ ,  $E_{\infty} = E$  for  $N^{1}(R)$ , and  $\{0\} \subsetneq E_{\infty} \subsetneq E$ for  $N^{01}(R)$ .

Now let us recall the following factorization result from [\[25\]](#page-19-10) (see also [\[22\]](#page-19-6)). Every scalar function  $V(z) \in N_{\kappa}$  admits a unique factorization

<span id="page-14-0"></span>
$$
V(z) = \frac{p(z)p^{\sharp}(z)}{q(z)q^{\sharp}(z)}V_0(z),
$$
\n(43)

where  $V_0$  belongs to the class  $N_0$ ,  $p(z)$  and  $q(z)$  are relatively prime monic polynomials such that max(deg p, deg q) =  $\kappa$ ,  $p^{\sharp}(z) = \overline{p(\bar{z})}$ . Recall also that the point  $\infty$ is called a generalized pole of nonpositive type of  $Q$  if

$$
-\infty \le \lim_{z \to \infty} \frac{Q(z)}{z} < 0,\tag{44}
$$

and the point  $\infty$  is called a generalized zero of nonpositive type of Q if

$$
0 \le \lim_{z \to \infty} zQ(z) < \infty. \tag{45}
$$

In terms of the factorization [\(43\)](#page-14-0) one can consider the following three possibilities:

- (1)  $\infty$  is a generalized pole of non-positive type of the function  $V(z)$  if and only if deg  $p > \deg q$ ;
- (2)  $\infty$  is a generalized zero of non-positive type of the function  $V(z)$  if and only if deg  $p <$  deg q;
- (3)  $\infty$  is neither the generalized pole of non-positive type nor generalized zero of non-positive type of the function  $V(z)$  if and only if deg  $p = \deg q$ .

In the first case V does not belong to the class  $N_{\kappa}(R)$ , in the second case V is definitely in the class  $N_{\kappa}(R)$ , and in the third case the inclusion  $V \in N_{\kappa}(R)$  can be characterized in terms of the function  $V_0$ .

**Theorem 20.** Let  $V(z) \in N_{\kappa}$  with  $E = \mathbb{C}$  and let  $\infty$  be neither the generalized pole nor generalized zero of non-positive type of the function  $V(z)$ . Then  $V(z)$  belongs to the class  $N_{\kappa}(R)$  if and only if the function  $V_0$  in the factorization [\(43\)](#page-14-0) belongs to the class  $N(R)$ .

<span id="page-14-1"></span>*Proof.* Suppose  $V_0(z) \in N(R)$  and let V admits the factorization [\(43\)](#page-14-0) with p and q such that deg  $p = \text{deg } q = \kappa$ . Then V belongs to the class  $N_{\kappa}$  (see [\[25\]](#page-19-10)). Also, since  $V_0(z) \in N(R)$  then  $F = 0$  in [\(41\)](#page-13-2). It follows from the representation (41) (see  $[30]$ ,  $[27]$  that in this case

$$
\lim_{y \uparrow \infty} \frac{V_0(iy)}{y} = F = 0. \tag{46}
$$

Combining [\(43\)](#page-14-0) and [\(46\)](#page-14-1) we get [\(10\)](#page-6-1). In order to prove the third item in the definition of the class  $N_{\kappa}(R)$  we first notice that it is equivalent to the function  $V(z)$  not being an identical constant. Let us assume the contrary, i.e.,

$$
V(z) = \frac{p(z)p^{\sharp}(z)}{q(z)q^{\sharp}(z)}V_0(z) \equiv k, \quad k \in \mathbb{C}.
$$

This immediately contradicts that  $V_0(z)$  is holomorphic in the upper half-plane. Therefore, the condition (3) of the definition of the class  $N_{\kappa}(R)$  is satisfied.

It was shown in [\[30\]](#page-19-23) that if  $V_0(z) \in N(R)$  then

<span id="page-15-0"></span>
$$
\lim_{y \uparrow \infty} y V_0(iy) = \int_{-\infty}^{\infty} dG(t),\tag{47}
$$

where  $G(t)$  is the function from the representation [\(41\)](#page-13-2) of  $V_0(z)$ . Then we see that the subspace B defined by [\(12\)](#page-6-3) for the class  $N_{\kappa}(R)$  coincides with the definition of the subspace  $E_{\infty}$  in [\(42\)](#page-13-1) written for the function  $V_0(z)$ . We notice that since  $E = \mathbb{C}$  then either  $E_{\infty} = \{0\}$  or  $E_{\infty} = E = \mathbb{C}$ . Finally, if  $E_{\infty} = \mathbb{C}$ 

$$
\lim_{y \uparrow \infty} V(iy) = \lim_{y \uparrow \infty} V_0(iy) = 0,
$$

(see [\[30\]](#page-19-23)) and hence  $V(z) \in N_{\kappa}(R)$ .

Conversely, let  $V(z) \in N_{\kappa}(R)$ . Then [\(46\)](#page-14-1) will provide us with  $F = 0$  in the integral representation [\(41\)](#page-13-2) of  $V_0(z)$ . Furthermore, since  $V(z) \in N_{\kappa}(R)$  then

<span id="page-15-1"></span>
$$
\lim_{y \uparrow \infty} yV(iy) = \lim_{y \uparrow \infty} yV_0(iy),\tag{48}
$$

and is either finite or infinite. If the limit is infinite then  $E_{\infty} = \{0\}$  for  $V_0(z)$  and  $V_0(z) \in N(R)$ . If the limit is finite then  $E_{\infty} = E = \mathbb{C}$ , [\(47\)](#page-15-0) holds for  $V_0(z)$ , and (see [\[30\]](#page-19-23))

$$
Q = \int_{-\infty}^{+\infty} \frac{t}{1+t^2} dG(t).
$$

Therefore,  $V_0(z) \in N(R)$ .

**Corollary 21.** A function  $V(z)$  belongs to the class  $N_{\kappa}^{0}(R)$  with  $E = \mathbb{C}$  if and only if the function  $V_0$  in the equation [\(43\)](#page-14-0) belongs to the class  $N^0(R)$ .

**Corollary 22.** A function  $V(z)$  belongs to the class  $N_{\kappa}^1(R)$  with  $E = \mathbb{C}$  if and only if the function  $V_0$  in the equation [\(43\)](#page-14-0) belongs to the class  $N^1(R)$ .

*Proof.* The proofs of both corollaries immediately follow from [\(47\)](#page-15-0) and [\(48\)](#page-15-1).  $\Box$ 

# 8. Examples

We conclude the paper with simple illustrations.

**Example 23.** Let us define  $\Pi_1$  as a set of all  $L^2([0, 2\pi], dx)$  functions with the scalar product

$$
[f,g] = \int_{0}^{2\pi} f(x)\overline{g(x)} dx - \frac{1}{\pi} \int_{0}^{2\pi} f(x) dx \int_{0}^{2\pi} \overline{g(x)} dx
$$

Let also A be a  $\pi$ -symmetric operator defined by

$$
A f = \frac{1}{i} \frac{df}{dx}
$$

with

$$
\mathcal{D}(A) = \left\{ f \in \Pi_1 \mid f, f' \in AC_{loc}([0, 2\pi]), \ f(0) = f(2\pi) = 0 \right\}.
$$

Let T be an operator in  $\Pi_1$  defined by

$$
T f = \frac{1}{i} \frac{df}{dx} - \frac{1}{\pi i} f(2\pi),
$$

$$
\Box
$$

where

$$
\mathcal{D}(T) = \left\{ f \in \Pi_1 \mid f, f' \in AC_{loc}([0, 2\pi]), f(0) = 0 \right\}.
$$

One can check that the operator  $T \supset A$ ,  $T^+ \supset A$ , and A is a maximal  $\pi$ -symmetric part of T and  $T^+$ , i.e.,  $T \in \Omega_A$ . The following two formulas define a (\*)-extension of T.

$$
\mathbb{A}f = \frac{1}{i}\frac{df}{dx} - \frac{1}{\pi i}\left(f(2\pi) - f(0)\right) - if(0)\left[\delta(x - 2\pi) + \delta(x) + \frac{2}{\pi}\right],
$$
  

$$
\mathbb{A}^{\times}f = \frac{1}{i}\frac{df}{dx} - \frac{1}{\pi i}\left(f(2\pi) - f(0)\right) + if(2\pi)\left[\delta(x - 2\pi) + \delta(x) - \frac{2}{\pi}\right].
$$

By straightforward calculations we get

$$
\frac{\mathbb{A} - \mathbb{A}^{\times}}{i} f = (f(0) + f(2\pi)) \left[ \frac{2}{\pi} - \delta(x - 2\pi) - \delta(x) \right].
$$

Now we can include A into a l.s.c.d.s.

$$
\theta = \left( \begin{array}{cc} \mathbb{A} & K & 1 \\ \mathfrak{H}^+ \subset \Pi_1 \subset \mathfrak{H}_- & \mathbb{C} \end{array} \right),
$$

where

$$
Ke = \frac{1}{\sqrt{2}} \left[ \frac{2}{\pi} - \delta(x - 2\pi) - \delta(x) \right] e, \quad e \in \mathbb{C}.
$$

Then we can derive

$$
W_{\theta}(z) = 1 + 2i[(\mathbb{A} - zI)^{-1}Ke, Ke] = \frac{(z\pi i - 1)e^{2\pi zi} + 1}{e^{2\pi zi} - z\pi i - 1}.
$$

Consequently, the function

$$
V_{\theta}(z) = i[W_{\theta}(z) + I]^{-1}[W_{\theta}(z) - I]\mathcal{J} = \frac{2 + \pi i z - (2 - \pi i z)e^{2\pi i z}}{\pi z(e^{2\pi z i} - 1)},
$$

belongs to the class  $N_1^0$ .

Example 24. Now we consider a construction which leads to examples of functions of the class  $N_{\kappa}^1(R)$ . Let H and  $\mathfrak{N}$  be two Hilbert spaces. Suppose that  $A_0$  is possibly unbounded operator in H with nonempty resolvent set  $\rho(A_0)$ . Let the operator T in the Hilbert space  $\mathcal{H} = \mathcal{H} \oplus \mathfrak{N}$  is given by the block-operator matrix

$$
T=\begin{pmatrix} A_0 & C \\ B & D \end{pmatrix}
$$

with bounded entries  $B, C$  and  $D$ . Recall that the operator valued function

$$
S(z) = D - B(A_0 - zI)^{-1}C, \ z \in \rho(A_0)
$$

is called the transfer function of the system determined by the matrix  $T$ . It is well known that the number  $z \in \rho(A_0)$  belongs to  $\rho(T)$  if and only if

<span id="page-16-0"></span>
$$
X(z) := S(z) - zI = D - B(A_0 - zI)^{-1}C - zI
$$
\n(49)

has bounded inverse defined on  $\mathfrak{N}$ . If this is a case then the resolvent  $(T - zI)^{-1}$  is given by the Schur-Frobenius formula

$$
\begin{pmatrix}\n(A_0 - zI)^{-1} \left( I + CX^{-1}(z)B(A_0 - zI)^{-1} \right) & -(A_0 - zI)^{-1}CX^{-1}(z) \\
-X^{-1}(z)B(A_0 - zI)^{-1} & X^{-1}(z)\n\end{pmatrix}
$$
\n(50)

Note that  $X(z)$  is called the Schur complement of the block-matrix  $T - zI$ . Let  $\dim \mathfrak{N} = \kappa < \infty$ . Suppose that  $A_0$  is a selfadjoint operator in H. In this case if

 $|\text{Im}z|$  is sufficiently large then the norm of the operator  $z^{-1}(D - B(A_0 - zI)^{-1}C)$ is less then one, therefore  $X^{-1}(z)$  exists and  $X^{-1}(z)$  is a meromorphic function in  $\mathbb{C}_+ \cup \mathbb{C}_-$ . Equip the Hilbert space H by the indefinite inner product

$$
[h,g] := (P_{\mathcal{H}_0}h, P_{\mathcal{H}_0}g) - (P_{\mathfrak{N}}h P_{\mathfrak{N}}g), h, g \in \mathcal{H},
$$

where  $P_{\mathcal{H}_0}$  and  $P_{\mathfrak{N}}$  are the orthogonal projections in H onto  $\mathcal{H}_0$  and  $\mathfrak{N}$ , respectively. Then H becomes a Pontryagin space  $\Pi_{\kappa}$ . Let A be a linear operator in the Hilbert space  $H$  defined as follows

$$
\mathcal{D}(A) = \mathcal{D}(A_0), \quad Ah := A_0h + Bh, \quad h \in \mathcal{D}(A).
$$

The operator A is a non-densely defined and closed  $\pi$ -Hermitian operator in  $\Pi_{\kappa}$ . Since the operator  $P_{\mathcal{H}_0}A = A_0$  is selfadjoint, the operator A is J regular O-operator, where  $J = P_{\mathcal{H}_0} - P_{\mathfrak{N}}$ . Evidently, the operator

$$
T = \begin{pmatrix} A_0 & -B^* \\ B & D \end{pmatrix} \tag{51}
$$

meets the conditions

$$
\mathcal{D}(T) = \mathcal{D}(T^+) = \mathcal{D}(A_0) \oplus \mathfrak{N}, \ T \supset A, \ T^+ \supset A
$$

In particular,  $T$  is  $\pi$ -selfadjoint if and only if D is a selfadjoint in the Hilbert space N.

Let  $D = \text{Re}(D) + i\text{Im}(D)$ . Then

$$
\text{Re}(T) = \frac{1}{2}(T + T^+) = \begin{pmatrix} A_0 & -B^* \\ B & \text{Re}(D) \end{pmatrix}, \text{ Im}(T) = \frac{1}{2i}(T - T^+) = \begin{pmatrix} 0 & 0 \\ 0 & \text{Im}(D) \end{pmatrix}
$$

Assume also that  $D_I = K \mathcal{J} K^+$ , where K acts from the Hilbert space  $\mathfrak{N}$  into the negative subspace  $\mathfrak N$  of the Pontryagin space  $\Pi_{\kappa}$  and K is invertible. Note that  $K^+ = -K^*$  where  $K^*$  is the Hilbert space adjoint to  $K : \mathfrak{N} \to \mathfrak{N}$ . Then

$$
\theta = \begin{pmatrix} T & K & \mathcal{J} \\ \Pi_{\kappa} & & \mathfrak{N} \end{pmatrix}
$$

is the l.s.c.d.s. The transfer function  $W(z)$  of  $\theta$  is given by

$$
W(z) = I - 2iK^{+}(T - zI)^{-1}KJ = I - 2iK^{+}P_{\mathfrak{N}}(T - zI)^{-1}KJ, \ z \in \rho(T)
$$

and its fractional-linear transformation is

 $V(z) = i[W(z) + I]^{-1}[W(z) - I] \mathcal{J} = K^+ P_{\mathfrak{N}}(\text{Re}(T) - zI)^{-1}K, \ z \in \rho(T) \cap \rho(\text{Re}(T)).$ Let

$$
X_T(z) = D + B(A_0 - zI)^{-1}B^* - zI, \ X_{T_R}(z) = \text{Re}(D) + B(A_0 - zI)^{-1}B^* - zI.
$$

Then from [\(50\)](#page-16-0) it follows that

$$
W(z) = I - 2iK^{+}X_{T}^{-1}(z)K\mathcal{J}, V(z) = K^{+}X_{T_{R}}^{-1}(z)K.
$$

Let us take Im(D) =  $-I$ ,  $K = I$  then  $K^+ = -I$  and  $-X_{T_R}^{-1}(z)$  belongs to the class  $N_{\kappa}^1(R)$  in the Hilbert space  $\mathfrak{N}$ . Thus we obtain the following theorem.

<span id="page-17-0"></span>**Theorem 25.** Let  $\mathcal{H}_0$  and  $\mathfrak{N}$  be Hilbert spaces, dim  $\mathfrak{N} = \kappa < \infty$ . Let  $A_0$  be a selfadjoint operator in  $\mathcal{H}_0$ , B is a bounded operator from  $\mathcal{H}_0$  into  $\mathfrak{N}$ , and let D be a selfadjoint operator in N. Then the operator valued function

$$
V(z) = -\left(D + B(A_0 - zI)^{-1}B^* - zI\right)^{-1}, \ z \in \mathbb{C}_+ \cup \mathbb{C}_-
$$

belongs to the class  $N_{\kappa}^1(R)$  in the Hilbert space  $\mathfrak{N}$ .

Using Theorem [25](#page-17-0) let us give some concrete example of scalar function from the class  $N_1^1(R)$ .

Let H be a weighted Hilbert space  $\mathcal{L}_2([-1,1], \rho(t))$  with the weight

$$
\rho(t) = \frac{2}{\pi} \sqrt{1 - t^2}.
$$

Let the operator  $A_0$  in  $\mathcal{L}_2([-1,1], \rho(t))$  be defined as follows:

$$
(A_0 f)(t) = tf(t), \ f(t) \in \mathcal{L}_2([-1,1], \rho(t)).
$$

Then  $A_0$  is a selfadjoint contraction. Let  $e_0(t) = 1, t \in [-1,1]$  The function  $e_0(t)$  belongs to  $\mathcal{L}_2([-1,1], \rho(t))$  and  $||e_0|| = 1$ . Let  $\mathfrak{N} = \mathbb{C}$ . Define the operator  $B: \mathcal{L}_2([-1,1], \rho(t)) \to \mathbb{C}$  as follows

$$
Bf(t) = \frac{2\gamma}{\pi} \int_{-1}^{1} f(t) \sqrt{1 - t^2} dt, \ f(t) \in \mathcal{L}_2 \left( [-1, 1], \rho(t) \right),
$$

where  $\gamma \neq 0$ . Then

$$
B^*c = \overline{\gamma} c e_0(t), \ c \in \mathbb{C}.
$$

Let D be the operator of multiplication on a real number d in the space  $\mathbb{C}$ . It is known [\[15\]](#page-19-15) that

$$
\frac{2}{\pi} \int_{-1}^{1} \frac{\sqrt{1-t^2}}{t-z} dt = 2(\sqrt{z^2-1} - z), \ z \notin [-1,1],
$$

where the branch of the function  $\sqrt{z^2-1}$  is taken such that Im  $\sqrt{z^2-1} > 0$  for Im  $z > 0$ . It follows that the function  $V(z) = -(D + B(A_0 - zI)^{-1}B^* - zI)^{-1}$ takes the form

$$
V(z) = \frac{1}{z - 2|\gamma|^2(\sqrt{z^2 - 1} - z) - d} = \frac{1}{(1 + 2|\gamma|^2)z - 2|\gamma|^2\sqrt{z^2 - 1} - d}
$$

According to Theorem [25](#page-17-0) the function  $V(z)$ ,  $z \in \mathbb{C}_+$  belongs to the class  $N_1^1(R)$ . If  $|\gamma|^2 > \max\{0, (d^2 - 1)/4\}$  the function  $V(z)$  has a simple pole

$$
z_0 = \frac{d(1+2|\gamma|^2) + 2|\gamma|^2 i \sqrt{4|\gamma|^2 + 1 - d^2}}{1 + 2|\gamma|^2}
$$

in  $\mathbb{C}_+$ .

### **REFERENCES**

- <span id="page-18-0"></span>[1] D. Alpay, A. Dijksma, J. Rovnyak, and H.S.V. de Snoo, *Schur functions, operator colligations, and reproducing kernel Pontryagin spaces*, Oper. Theory: Adv. Appl., 96, Birkhäuser Verlag, Basel, 1997.
- <span id="page-18-1"></span>[2] Alpay D., Bruinsma P., Dijksma A., and de Snoo H.S.V., A Hilbert space associated with a Nevanlinna function, in: *Proc. MTNS Meeting, Amsterdam* (1989), pp. 115-122.
- <span id="page-18-3"></span>[3] Yu. M. Arlinskiı̆. The inverse problem of the theory of characteristic functions of unbounded operator colligations. *Dokl. Akad. Nauk Ukrain. SSR, Ser. A* (1976), no. 2, 105–109.
- <span id="page-18-4"></span>[4] Yu. M. Arlinskiĭ. Regular (\*)-extensions of quasi-Hermitian operators in rigged Hilbert spaces. *Izv. Akad. Nauk Armyan. SSR, Ser. Mat.*, 14 (1979), no. 4, 297–312.
- <span id="page-18-2"></span>[5] Yu. M. Arlinskii and V. A. Derkach. Inverse problem in the theory of characteristic operator functions of unbounded operator bundles in a space  $\Pi_{\kappa}$ , *Ukrain. Math. J.*, **31**, no. 2, (1979), 89–94.
- [6] Yu. M. Arlinskii and V. A. Derkach. Reconstruction of an unbounded operator colligation from characterstic function in the space  $\Pi_{\kappa}$ , *Preprint VINITI*, **555-79**, (1978), 2–72.
- <span id="page-19-18"></span>[7] Yu.M. Arlinskiĭ and E.R. Tsekanovskiĭ. The method of rigged spaces in the theory of extensions of Hermitian operators with a nondense domain of definition. *Sib. Math. J.* 15, (1974), 169-182.
- <span id="page-19-0"></span>[8] D. Arov, "Passive linear systems and scattering theory", in Dynamical Systems, Control Coding, Computer Vision, vol.25 of Progress in Systems and Control Theory, (1999), Birhäuser Verlag, 27–44.
- <span id="page-19-11"></span>[9] T. Ya. Azizov and I.S. Iokhvidov, "Linear operators in spaces with an indefinite metric", Wiley, New York, 1989.
- <span id="page-19-1"></span>[10] J.A. Ball and O.J. Staffans, "Conservative state-space realizations of dissipative system behaviors", Report No. 37, Institute Mittag-Leffler, (2002/2003), 55 pp.; IEOT (to appear)
- <span id="page-19-2"></span>[11] S.V. Belyi and E.R. Tsekanovskii. Realization and factorization problems for J-contractive operator-valued functions in half-plane and systems with unbounded operators, *Systems and Networks: Mathematical Theory and Applications*, Akademie Verlag, 2, (1994), 621–624.
- <span id="page-19-9"></span>[12] S. V. Belyi and E. Tsekanovskii. Realization theorems for operator-valued R-functions, Op*erator theory: Advances and Applications*, 98, Birkhäuser Verlag Basel, (1997), 55–91.
- <span id="page-19-3"></span>[13] S.V. Belyi and E.R. Tsekanovskii. On classes of realizable operator-valued R-functions, Op*erator theory: Advances and Applications*, 115, Birkhäuser Verlag Basel, (2000), 85–112.
- [14] S.V. Belyi and E.R. Tsekanovski˘i. On Krein's formula in indefinite metric spaces, *Linear Algebra and Applications*, 389 C, (2004), 305-322.
- <span id="page-19-15"></span>[15] Yu.M. Berezanskii, Expansions in Eighenfunctions of Self-Adjoint Operators, Transl. Math. Mono. 17, Amer. Math. Soc., Providence, RI, 1968.
- <span id="page-19-4"></span>[16] M.S. Brodskiĭ, *Triangular and Jordan representations of linear operators*, Moscow, Nauka, 1969 (Russian).
- <span id="page-19-19"></span><span id="page-19-16"></span>[17] K. Daho and H. Langer. Matrix functions of the class Nκ. *Math. Nachr., 120 (1985), 275–294*
- [18] V.A. Derkach. π-selfadjoint biextensions of π-Hermitian operators, *Dopovidi Akad. Nauk Ukraine. SSR, Ser. A.*, (1975), no. 4, 304–306.
- <span id="page-19-14"></span>[19] V.A. Derkach. On signature of defect subspaces of  $\pi$ -Hermitian operators with nondense domains in spaces Πκ, *Dopovidi Akad. Nauk Ukraine. SSR, Ser. A.*, (1977), no. 7, 579–581.
- <span id="page-19-22"></span>[20] V. Derkach and M. Malamud, "The extension theory of hermitian operators and the moment problem", J. Math. Sci., 73 (1995), 141–242.
- <span id="page-19-5"></span>[21] V.A. Derkach and S. Hassi, A reproducing kernel space model for Nκ-functions. *Proc. Amer. Math. Soc.*, 131 (2003), no. 12, 3795–3806.
- <span id="page-19-6"></span>[22] V. Derkach, S. Hassi, and H.S.V. de Snoo, "Operator models associated with Kac subclasses of generalized Nevanlinna functions", Methods of Functional Analysis and Topology, 5 (1999), 65–87.
- <span id="page-19-20"></span>[23] V.A. Derkach, S. Hassi, and H. de Snoo, Operator models associated with singular perturbations, *Methods Funct. Anal. Topology*, 7, no. 3, (2001), 1–21.
- <span id="page-19-7"></span>[24] V.A. Derkach and E.R Tsekanovskiı̆. A class of operator-functions that are realizable by accretive operator colligations as transfer mappings of linear systems. (Russian)*Theory of mappings, its generalizations and applications*, (1982) 74–83, "Naukova Dumka", Kiev.
- <span id="page-19-10"></span>[25] A. Dijksma, H. Langer, A. Luger, and Yu. Shondin. A factorization result for generalizaed Nevanlinna functions of the class  $N_{\kappa}$ , *Integr. Eq. Oper. Th.*, **36**, (2000), 121–125.
- <span id="page-19-8"></span>[26] F. Gesztesy, N. J. Kalton, K. A. Makarov, E. Tsekanovskii, Some Applications of Operator-Valued Herglotz Functions, *Operator Theory: Advances and Applications*, 123, Birkhäuser, Basel, (2001), 271–321.
- <span id="page-19-24"></span>[27] F. Gesztesy and E. Tsekanovskii, On matrix-valued Herglotz functions, *Math. Nach.* 218, (2000), 61–138.
- <span id="page-19-21"></span>[28] F. Gesztesy and M. Zinchenko, On Spectral Theory for Schrödinger Operators with Strongly Singular Potentials, [arXiv:math.SP/0505120,](http://arxiv.org/abs/math/0505120) vol 1, (2005).
- <span id="page-19-12"></span>[29] I. S. Iokhvidov and M. G. Krein, Spectral theory of operators in spaces with an indefinite metric, *Amer. Math. Soc. Translations*, 2, 13, (1960), 105–175.
- <span id="page-19-23"></span>[30] I. S. Kac and M. G. Krein, R*-functions–analytic functions mapping the upper halfplane into itself*, Amer. Math. Soc. Transl. (2) 103, 1-18 (1974).
- <span id="page-19-17"></span>[31] M.A. Krasnosel'ski˘ı, On selfadjoint extensions of Hermitian operators, *Ukrain. Math. J.*,1 (1949), 21-38 (Russian).
- <span id="page-19-13"></span>[32] M. G. Krein and H. K. Langer, Defect subspaces and generalized resolvents of a Hermitian operator in the space  $\Pi_{\kappa}$ , *Func. Anal. Appl.*, **5**, 2, (1971), 136–146.
- <span id="page-20-0"></span>[33] M. G. Krein and H. K. Langer, Über die  $Q$ -Funktion eines  $\pi$ -hermiteschen Operators im Raume Πκ, *Acta Sci. Math*, 34, (1973), 191–230.
- <span id="page-20-3"></span>[34] M. G. Krein and H. K. Langer, Some propositions on analytic matrix functions related to the theory of operators in the space  $\Pi_{\kappa}$ , *Acta Sci. Math*, **43**, (1981), 181–205.
- <span id="page-20-4"></span>[35] J. Rovnyak and L.A. Sakhnovich, On the Krein-Langer integral representation of generalizaed Nevanilinna functions, *Electronic Journal of Linear Algebra*, 11, (2004), pp. 1–15.
- <span id="page-20-1"></span>[36] Ju. L. Šmuljan, Extension theory for operators and spaces with indefinite metric, *Math. USSR Izvestija*, 8, no. 4, (1974), 895–907.
- [37] O.J. Staffans, *Well-Posed Linear Systems:Part I* Book manuscript, available at [http://www.abo.fi/](http://www.abo.fi/~staffans/)∼staffans/, 2002
- [38] O.J. Staffans, "Passive and conservative continuous time impedance and scattering systems, Part I:Well posed systems", Math. Control Signals Systems, 15, (2002), 291–315.
- [39] O.J. Staffans and G. Weiss, "Transfer functions of regular linear systems, Part II: the system operator and the Lax-Phillips semigroup", Trans. Amer. Math. Soc., 354, (2002), 3229–3262.
- <span id="page-20-2"></span>[40] E. R. Tsekanovskii and Yu. L. Shmul'jan. The theory of bi-extensions of operators on rigged Hilbert spaces. Unbounded operator colligations and characteristic functions, *Russ. Math. Surv.*, 32, no. 5, (1977), 73–131.

Department of Mathematics, East Ukrainian National University, Kvartal Molodyozhny, 20-A, 91034 Lugansk, Ukraine

*E-mail address*: yma@snu.edu.ua

Department of Mathematics, Troy State University, Troy, AL 36082, USA *E-mail address*: sbelyi@trojan.troyst.edu, *URL:*[http://spectrum.troyst.edu/](http://spectrum.troyst.edu/~belyi)∼belyi

Department of Mathematics, Donetsk National University, Universitetskaya str, 24, 83055 Donetsk, Ukraine

*E-mail address*: derkach@univ.donetsk.ua

Department of Mathematics, Niagara University, NY 14109, USA *E-mail address*: tsekanov@niagara.edu, *URL:*<http://faculty.niagara.edu/tsekanov/>