# Modified scattering for a wave equation with weak dissipation

Jens Wirth\*<sup>†</sup>

October 29, 2018

#### Abstract

We consider the Cauchy problem for the weakly dissipative wave equation

$$\Box u + \frac{\mu}{1+t}u_t = 0$$

with parameter  $\mu \geq 2$ .

Based on the explicit representations of solutions provided in [Math. Meth. Appl. Sci. 2004; **27**:101-124] sharp decay estimates for data from a dense subspace of the energy space are derived. Furthermore, sharpness is discussed in terms of a modified scattering theory.

AMS subject classification: 35L50, 35L15, 35B45

## 1 Introduction

We are interested in a precise description of the behaviour of solutions to the Cauchy problem

$$\Box u + \frac{\mu}{1+t}u_t = 0, \qquad u(0, \cdot) = u_1, \quad u_t(0, \cdot) = u_2 \qquad (1.1)$$

with  $\Box = \partial_t^2 - \Delta$  and for data  $u_1 \in H^1(\mathbb{R}^n)$ ,  $u_2 \in L^2(\mathbb{R}^n)$ . It is well known that this problem is well-posed in the sense that it has a unique solution in the space

 $C^{1}([0,\infty), L^{2}(\mathbb{R}^{n})) \cap C([0,\infty), H^{1}(\mathbb{R}_{+})).$ 

It is known from the fundamental works of A.Matsumura, [1], and H.Uesaka, [2], that the energy of the solutions to this equation,

$$E(u;t) = \frac{1}{2} \int (u_t^2 + |\nabla u|^2) \mathrm{d}x, \qquad (1.2)$$

satisfies an estimate of the form

$$E(u;t) = \mathcal{O}(t^{-\alpha}), \qquad \alpha = \min\{2,\mu\}.$$
(1.3)

\*Department of Applied Analysis, TU Bergakademie Freiberg, 09596<br/> Freiberg, Germany

<sup>&</sup>lt;sup>†</sup>email: wirth@math.tu-freiberg.de

Following F.Hirosawa and H.Nakazawa, [3], it is possible to improve the last estimate in the case  $\mu > 2$  to

$$E(u;t) = o(t^{-2}), (1.4)$$

which means that the above given estimate is not sharp (with respect to the strong topology).

In [4] the author provided explicit representations of solutions to the Cauchy problem (1.1) in terms of special functions. Furthermore, from that treatment it follows that the estimate  $E(u;t) = \mathcal{O}(t^{-2})$  is sharp in the sense of a norm estimate for the energy operator

$$\mathbb{E}(t) : (\langle \mathbf{D} \rangle u_1, u_2)^T \mapsto (|\mathbf{D}| u(t, \cdot), \partial_t u(t, \cdot))^T$$
(1.5)

associated to the solution representation. As usual we denote  $\langle \xi \rangle = \sqrt{1 + |\xi|^2}$ .

In this paper we will use this explicit representation to describe a dense subspace of the energy space such that for all data from that space we obtain a sharp two-sided estimate of the energy in the form

$$E(u;t) \sim t^{-\mu}.\tag{1.6}$$

The result can be formulated in terms of a modified scattering theory. Especially it describes the discrepancy between the estimates of Hirosawa/Nakazawa and the norm estimate from [4].

# 2 Preliminaries

Let us recall the representations of solutions to equation (1.1). With the notation

$$\Psi_{k,s,\rho,\delta}(t,\xi) = |\xi|^k \langle \xi \rangle^{s+1-k} \begin{vmatrix} \mathcal{H}_{\rho}^{-}(|\xi|) & \mathcal{H}_{\rho+\delta}^{-}((1+t)|\xi|) \\ \mathcal{H}_{\rho}^{+}(|\xi|) & \mathcal{H}_{\rho+\delta}^{+}((1+t)|\xi) \end{vmatrix}$$
(2.1a)  
$$= 2i \csc(\rho\pi) |\xi|^k \langle \xi \rangle^{s+1-k} \begin{vmatrix} \mathcal{J}_{-\rho}(|\xi|) & \mathcal{J}_{-\rho-\delta}((1+t)|\xi) \\ (-)^{\delta} \mathcal{J}_{\rho}(|\xi|) & \mathcal{J}_{\rho+\delta}((1+t)|\xi|) \end{vmatrix},$$
(2.1b)

the last line analytically continued to  $\rho \in \mathbb{Z}$ , the following theorem is valid, [4, Theorem 2.1].

**Theorem 2.1.** The spatial Fourier transform of the solution to problem (1.1) can be represented in the form

$$\hat{u}(t,\xi) = \sum_{j=1,2} \Phi_j(t,\xi) \hat{u}_j(\xi), \qquad (2.2)$$

where

$$\Phi_1(t,\xi) = \frac{i\pi}{4} (1+t)^{\rho} \Psi_{1,0,\rho-1,1}(t,\xi), \qquad (2.3a)$$

$$\partial_t \Phi_1(t,\xi) = \frac{i\pi}{4} \Psi_{2,1,\rho-1,0}(t,\xi), \qquad (2.3b)$$

$$\Phi_2(t,\xi) = -\frac{i\pi}{4}(1+t)^{\rho}\Psi_{0,-1,\rho,0}(t,\xi), \qquad (2.3c)$$

$$\partial_t \Phi_2(t,\xi) = -\frac{i\pi}{4} (1+t)^{\rho} \Psi_{1,0,\rho-1,0}(t,\xi)$$
(2.3d)

with  $\rho = (1 - \mu)/2$ .

We want to obtain estimates in  $L^2$ -scale, which correspond by Plancherel's theorem to  $L^{\infty}$  estimates for the Fourier multiplier. Therefore we recall the fundamental estimate from [4].

**Lemma 2.2.** It holds  $\Psi_{k,s,\rho,\delta}(t,\cdot) \in L^{\infty}(\mathbb{R}^n)$  for all t if and only if  $s \leq 0$  and  $k \geq |\delta|$ . Furthermore, the estimate

$$||\Psi_{k,s,\rho,\delta}(t,\cdot)||_{\infty} \sim \begin{cases} (1+t)^{-\frac{1}{2}}, & \rho \neq 0, |\rho| - k \leq -\frac{1}{2}, \\ (1+t)^{|\rho|-k}, & \rho \neq 0, |\rho| - k \geq -\frac{1}{2}, \\ (1+t)^{-k} \log(e+t), & \rho = 0, k \leq \frac{1}{2} \end{cases}$$
(2.4)

is valid.

For later use we will introduce a notation. Let

$$[\xi] = \frac{|\xi|}{\langle \xi \rangle}.$$
(2.5)

By the aid of this symbol we can control the vanishing order in the frequency  $\xi = 0$ . We denote

$$[D]^{\kappa}L^{2}(\mathbb{R}^{n}) = \{ [D]^{\kappa}f \mid f \in L^{2} \}, \qquad ||g||_{[D]^{\kappa}L^{2}} = ||[D]^{-\kappa}g||_{2}.$$
(2.6)

It turns out that the usage of date from these spaces allows an improvement of the decay order of the energy.

# 3 Improvements of the energy decay

**Theorem 3.1.** Let  $\langle D \rangle u_1, u_2 \in [D]^{\kappa} L^2(\mathbb{R}^n)$ . Then the solution u = u(t, x) to (1.1) with  $\mu > 2$  satisfies

$$|||\mathbf{D}|u(t,\cdot)||_{2} + ||u_{t}(t,\cdot)||_{2} \lesssim ||\langle \mathbf{D}\rangle u_{1}, u_{2}||_{[\mathbf{D}]^{\kappa}L^{2}} \begin{cases} (1+t)^{-1}, & \kappa = 0, \\ (1+t)^{-1-\kappa}, & 0 \le \kappa \le \frac{\mu-2}{2}, \\ (1+t)^{-\frac{\mu}{2}}, & \kappa \ge \frac{\mu-2}{2}. \end{cases}$$
(3.1)

*Proof.* The proof is a direct application of Lemma 2.2. We use the canonical isomorphism between  $[D]^{\kappa}L^2$  and  $L^2$  to simplify the estimates. The mapping  $(\langle D \rangle u_1, u_2)^T \mapsto (|D|u(t, \cdot), u_t(t, \cdot)^T \text{ can be represented as matrix Fourier multiplier in terms of <math>\Psi_{k,s,\rho,\delta}$ , because it includes the identification of spaces we include the number  $\kappa$  in the notation. It holds

$$\mathbb{E}^{(\kappa)}(t,\xi) = \frac{i\pi}{4} (1+t)^{\rho} \begin{pmatrix} \Psi_{2+\kappa,0,\rho-1,1}(t,\xi) & -\Psi_{1+\kappa,0,\rho,0}(t,\xi) \\ \Psi_{2+\kappa,0,\rho-1,0}(t,\xi) & -\Psi_{1+\kappa,0,\rho,-1}(t,\xi) \end{pmatrix}$$

It remains to check the conditions on the indices. For  $\mu > 2$  we have  $\rho < -1/2$  and therefore we have to compare  $1 + \kappa - \rho$  with -1/2.

Some remarks for the interpretation of the result are necessary. The decay rate  $t^{-\mu/2}$  corresponds to the rate for high frequencies. The cut-off by  $[\xi]^{\kappa}$  dampes out the small frequencies around the exceptional frequency  $\xi = 0$ .

For  $\mu > 2$  the decay rate obtained in [4] was determined by the small frequencies around  $\xi = 0$ . This is the reason for the improvement of this theorem. An improvement over the decay rate for high frequencies is not possible.

### 4 Sharpness

The aim of this section is to discuss the sharpness of the estimate in the limit case  $\kappa = (\mu - 2)/2$ . For this we compare the operator family  $\mathbb{E}^{(\kappa)}(t)$  with the unitary evolution  $\mathcal{E}_0(t)$  of free waves in energy space,

$$\mathcal{E}_0(t,\xi) = \begin{pmatrix} \cos t|\xi| & \sin t|\xi| \\ -\sin t|\xi| & \cos t|\xi| \end{pmatrix},\tag{4.1}$$

modified by the decay rate.

**Theorem 4.1.** Let  $\kappa = \frac{\mu-2}{2}$  and  $\mu \ge 2$ . Then the strong limit

$$Z_{+} = \underset{t \to \infty}{\operatorname{s-lim}} (1+t)^{\mu/2} \mathcal{E}_{0}(-t) \mathbb{E}^{(\kappa)}(t)$$

$$(4.2)$$

exists in the operator space  $[D]^{\kappa}L^2 \to L^2$ .

Let us first explain the main strategy of the proof. Theorem 3.1 yields a uniform bound for the operator family. In order to obtain the strong convergence we employ Banach-Steinhaus theorem for the dense subspace

$$M = \{ f \in L^2 \mid 0 \not\in \operatorname{supp} \hat{f} \}.$$

For this it is sufficient to prove (uniform) convergence of the Fourier multiplier for  $|\xi| \ge c$  with c > 0. This can be done by the given explicit representations using known asymptotic expansions for Bessel functions, [5]. Because of its simplicity we will use the representation by real valued functions valid for  $\rho \notin \mathbb{Z}$ . The multiplier  $Z_+$  will be analytic in  $\rho$  and therefore the statement will follow in all cases.

The result is based on the following asymptotic formula, [5].

Proposition 4.2. For Bessel functions of first kind it holds

$$\mathcal{J}_{\rho}(z) = \sqrt{\frac{2}{\pi z}} \cos\left(z - \frac{\rho\pi}{2} - \frac{\pi}{4}\right) + \mathcal{O}(z^{-3/2}), \qquad z \to \infty.$$
(4.3)

Proof of Theorem 4.1. Let  $|\xi| \ge c > 0$ . Then it holds

$$(1+t)^{\mu/2}\mathbb{E}^{(\kappa)}(t,\xi) = \frac{i\pi}{4}(1+t)^{1/2} \begin{pmatrix} \Psi_{2+\kappa,0,\rho-1,1}(t,\xi) & -\Psi_{1+\kappa,0,\rho,0}(t,\xi) \\ \Psi_{2+\kappa,0,\rho-1,0}(t,\xi) & -\Psi_{1+\kappa,0,\rho,-1}(t,\xi) \end{pmatrix}$$

such that

$$(1+t)^{\mu/2} \mathcal{E}_{0}(-t,\xi) \mathbb{E}^{(\kappa)}(t,\xi) = \frac{i\pi}{4} (1+t)^{1/2} \begin{pmatrix} \cos t|\xi| & -\sin t|\xi| \\ \sin t|\xi| & \cos t|\xi| \end{pmatrix} \\ \begin{pmatrix} \Psi_{2+\kappa,0,\rho-1,1}(t,\xi) & -\Psi_{1+\kappa,0,\rho,0}(t,\xi) \\ \Psi_{2+\kappa,0,\rho-1,0}(t,\xi) & -\Psi_{1+\kappa,0,\rho,-1}(t,\xi) \end{pmatrix} \\ = \begin{pmatrix} m_{1,1}(t,\xi) & m_{1,2}(t,\xi) \\ m_{2,1}(t,\xi) & m_{2,2}(t,\xi) \end{pmatrix}.$$

We proceed with calculating  $m_{1,1}(t,\xi)$ . It holds

$$\begin{split} m_{1,1}(t,\xi) &= \frac{\pi}{2} \csc(\rho\pi) |\xi|^{2+\kappa} \langle\xi\rangle^{-\kappa-1} (1+t)^{1/2} \\ & \left( \cos t |\xi| \left( \mathcal{J}_{1-\rho}(|\xi|) \mathcal{J}_{\rho}((1+t)|\xi|) + \mathcal{J}_{\rho-1}(|\xi|) \mathcal{J}_{-\rho}((1+t)|\xi|) \right) \right) \\ & - \sin t |\xi| \left( \mathcal{J}_{1-\rho}(|\xi|) \mathcal{J}_{\rho-1}((1+t)|\xi|) - \mathcal{J}_{\rho-1}(|\xi|) \mathcal{J}_{1-\rho}((1+t)|\xi|) \right) \right) \\ & = \sqrt{\frac{\pi}{2}} \csc(\rho\pi) |\xi|^{2+\kappa-\frac{1}{2}} \langle\xi\rangle^{-\kappa-1}(1+t)^{1/2} \\ & \left( \cos t |\xi| \left( \mathcal{J}_{1-\rho}(|\xi|) \cos((1+t)|\xi| - \frac{\rho\pi}{2} - \frac{\pi}{4} \right) \right) \\ & + \mathcal{J}_{\rho-1}(|\xi|) \cos((1+t)|\xi| + \frac{\rho\pi}{2} - \frac{\pi}{4} \right) \right) \\ & - \sin t |\xi| \left( \mathcal{J}_{1-\rho}(|\xi|) \cos((1+t)|\xi| - \frac{(1-\rho)\pi}{2} - \frac{\pi}{4} \right) \\ & - \mathcal{J}_{\rho-1}(|\xi|) \cos((1+t)|\xi| - \frac{(1-\rho)\pi}{2} - \frac{\pi}{4} ) \right) \right) \\ & + \mathcal{O}((1+t)^{-1}|\xi|^{-1}) \\ & = \sqrt{\frac{\pi}{2}} \csc(\rho\pi) [\xi]^{\kappa+1} |\xi|^{1/2} \\ & \left( \cos(|\xi| - \frac{\rho\pi}{2} - \frac{\pi}{4} \right) \mathcal{J}_{1-\rho}(|\xi|) - \cos(|\xi| + \frac{\rho\pi}{2} - \frac{\pi}{4} ) \mathcal{J}_{\rho-1}(|\xi|) \right) \\ & + \mathcal{O}((1+t)^{-1}|\xi|^{-1}) \end{split}$$

as  $t \to \infty$ ,  $|\xi| \ge c$ . Hence, as  $t \to \infty$  the function  $m_{1,1}(t,\xi)$  tends uniformly (on  $|\xi| \ge c$ ) to the limit  $m_{1,1}(\infty,\xi)$  which extends continuously (and analytically) up to  $\xi = 0$ . The last statement follows from the generalized power series expansion of  $J_{\rho}(z) = z^{\rho} \Lambda_{\rho}(z)$ ,  $\Lambda_{\rho}(0) \ne 0$  together with  $\kappa + 3/2 = (\mu + 1)/2 = 1 - \rho$ .

A similar calculation yields for the other entries of the matrix limit expressions of the form

$$\begin{split} m_{1,2}(\infty,\xi) &= \sqrt{\frac{\pi}{2}} \csc(\rho\pi) [\xi]^{\kappa} |\xi|^{1/2} \\ &\left( \cos(|\xi| - \frac{\rho\pi}{2} - \frac{\pi}{4}) \mathcal{J}_{-\rho}(|\xi|) + \cos(|\xi| + \frac{\rho\pi}{2} - \frac{\pi}{4}) \mathcal{J}_{\rho}(|\xi|) \right) \\ m_{2,1}(\infty,\xi) &= \sqrt{\frac{\pi}{2}} \csc(\rho\pi) [\xi]^{\kappa+1} |\xi|^{1/2} \\ &\left( \sin(|\xi| - \frac{\rho\pi}{2} - \frac{\pi}{4}) \mathcal{J}_{1-\rho}(|\xi|) + \sin(|\xi| + \frac{\rho\pi}{2} - \frac{\pi}{4}) \mathcal{J}_{\rho-1}(|\xi|) \right) \\ m_{2,2}(\infty,\xi) &= \sqrt{\frac{\pi}{2}} \csc(\rho\pi) [\xi]^{\kappa} |\xi|^{1/2} \\ &\left( \sin(|\xi| - \frac{\rho\pi}{2} - \frac{\pi}{4}) \mathcal{J}_{-\rho}(|\xi|) - \sin(|\xi| + \frac{\rho\pi}{2} - \frac{\pi}{4}) \mathcal{J}_{\rho}(|\xi|) \right). \end{split}$$

By analytic continuation these formulas are valid for all  $\xi \in \mathbb{R}^n$  and  $m_{i,j}(\infty, \cdot) \in L^{\infty}(\mathbb{R}^n)$ . Furthermore, all entries are non-zero for  $\xi = 0$  (due to the exact cancellation).

Furthermore the representations are analytic in  $\rho$  using the definition of the Weber functions  $\mathcal{Y}_{\rho}(z) = \csc(\rho \pi) (\mathcal{J}_{\rho}(z) \cos(\rho \pi) - \mathcal{J}_{-\rho}(z)).$ 

An application of Liouville theorem to the differential equation for  $(|\mathbf{D}|u, u_t)^T$  yields

$$\begin{vmatrix} \Psi_{1,0,\rho-1,1} & -\Psi_{1,0,\rho,0} \\ \Psi_{1,0,\rho-1,0} & -\Psi_{1,0,\rho,-1} \end{vmatrix} = (1+t)^{-\mu}$$
(4.4)

and therefore  $\det(1+t)^{\mu/2}\mathbb{E}^{(\kappa)}(t,\xi) = [\xi]^{1+2\kappa}$ .

**Corollary 4.3.** The operator  $Z_+ : [D]^{\kappa} L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$  is injective and its symbol satisfies det  $Z_+(\xi) = [\xi]^{1+2\kappa}$ .

Remark. What have we obtained so far? The existence of nontrivial Cauchy data  $(\langle D \rangle u_1, u_2)$  in the null space of  $Z_+$  is equivalent to the fact that the energy of the corresponding solution  $||\mathbb{E}(t)(\langle D \rangle u_1, u_2)^T||_2^2$  decays faster than  $t^{-\mu}$ . But we have shown, that such Cauchy data do not exist. This means, we have proven a two-sided energy estimate.

**Corollary 4.4.** Let  $(\langle D \rangle u_1, u_2) \in [D]^{\kappa} L^2(\mathbb{R}^n)$ . Then

$$E(u;t) \sim (1+t)^{-\mu}.$$
 (4.5)

Furthermore, the solutions behave in energy space as solutions to the free wave equation multiplied by the decay rate  $(1 + t)^{-\mu/2}$ . In the case of high frequencies we can even say more. The operator  $Z_+$  almost preserves high frequencies. Using again the asymptotic representation of Bessel functions for large arguments, Proposition 4.2, we conclude

**Corollary 4.5.** It holds  $\lim_{|\xi|\to\infty} Z_+(\xi) = I$ .

For  $\mu = 2$  we have even more. In this case the Bessel functions are trigonometric ones and  $Z_{+}(\xi) = I$ .

In order to conclude this article we will give one further interpretation to the assumptions on the Cauchy data we made. In case  $0 \le \kappa < \frac{n}{2}$  the Sobolev-Hardy inequality can be used to obtain an embedding of the space of weighted  $L^2$ -functions into  $[D]^{\kappa}L^2$ .

**Lemma 4.6.** Let 
$$0 \leq \kappa < \frac{n}{2}$$
. Then  $\langle x \rangle^{-\kappa} L^2(\mathbb{R}^n) \subseteq [D]^{\kappa} L^2(\mathbb{R}^n)$ .

*Proof.* Let  $\chi \in C_0^{\infty}(\mathbb{R}^n)$  satisfy  $\chi(\xi) = 1$  near  $\xi = 0$ . Then

$$\begin{aligned} ||f||_{[\mathbb{D}]^{\kappa}L^{2}} &= ||[\xi]^{-\kappa}\hat{f}||_{2} \sim ||\,|\xi|^{-\kappa}\chi(\xi)\hat{f}||_{2} + ||(1-\chi(\xi))f||_{2} \\ &\lesssim ||\hat{f}||_{H^{\kappa}} + ||\hat{f}||_{2} \sim ||\langle x \rangle^{\kappa}f||_{2} \end{aligned}$$

by Sobolev-Hardy inequality and Plancherel's theorem.

#### References

- A. Matsumura. Energy decay of solutions of dissipative wave equations. Proceedings of the Japan Academy 1977; 53:232-236
- H. Uesaka. The total energy decay of solutions for the wave equation with a dissipative term. Journal of Mathematics, Kyoto University 1979; 20(1): 57-65
- [3] F. Hirosawa and H. Nakazawa. Rapid decay of the total energy for dissipative wave equations. Tsukuba Journal of Mathematics 2003; 27(2):217-232
- [4] J. Wirth. Solution representations for a wave equation with weak dissipation. Mathematical Methods in the Applied Sciences 2004; 27:101-124
- [5] G.N. Watson. A treatise on the theory of Bessel functions. Cambridge University Press 1922