Second Order Backward Stochastic Differential Equations and Fully Non-Linear Parabolic PDEs

Patrick Cheridito * H. Mete Soner † Nizar Touzi ‡ Nicolas Victoir §

September 12, 2005

Abstract

For a d-dimensional diffusion of the form $dX_t = \mu(X_t)dt + \sigma(X_t)dW_t$, and continuous functions f and g, we study the existence and the uniqueness of adapted processes Y, Z, Γ and A solving the second order backward stochastic differential equation (2BSDE)

$$dY_{t} = f(t, X_{t}, Y_{t}, Z_{t}, \Gamma_{t}) dt + Z'_{t} \circ dX_{t}, \quad t \in [0, T),$$

$$dZ_{t} = A_{t} dt + \Gamma_{t} dX_{t}, \quad t \in [0, T),$$

$$Y_{T} = g(X_{T}).$$

If the associated PDE

$$-v_t(t,x) + f(t,x,v(t,x),Dv(t,x),D^2v(t,x)) = 0$$
, $(t,x) \in [0,T) \times \mathbb{R}^d$, $v(T,x) = g(x)$,

admits a C^3 -solution, it follows directly from Itô's lemma that the processes

$$v(t, X_t), Dv(t, X_t), D^2v(t, X_t), \mathcal{L}Dv(t, X_t), \quad t \in [0, T],$$

solve the 2BSDE, where \mathcal{L} is the Dynkin operator of X without the drift term.

The main result of the paper shows that if the PDE has comparison as in the theory of viscosity solutions and if f is Lipschitz in Y and decreasing in Γ , the existence of a solution (Y, Z, Γ, A) to the 2BSDE implies that the associated PDE has a unique continuous viscosity solution v, and $Y_t = v(t, X_t)$, $t \in [0, T]$. In particular, the 2BSDE has at most one solution. This provides a stochastic representation for solutions of fully non-linear parabolic PDEs. As a consequence, the numerical treatment of such PDE's can now be approached by Monte Carlo methods.

Key words: Second order backward stochastic differential equations, Fully non-linear parabolic partial differential equations, Viscosity solutions, Scaling limits.

AMS 2000 subject classifications: 60H10, 35K55, 60H30, 60H35.

^{*}Princeton University, Princeton USA, dito@princeton.edu.

[†]Koç University, Istanbul, Turkey, msoner@ku.edu.tr. Member of the Turkish Academy of Sciences and this work was partly supported by the Turkish Academy of Sciences.

[‡]CREST, Paris, France, touzi@ensae.fr, and Tanaka Business School, Imperial College London, England, n.touzi@imperial.ac.uk.

[§]Oxford University, Oxford, England, victoir@gmail.com.

1 Introduction

Since their introduction, backward stochastic differential equations (BSDEs) have received considerable attention in the probability literature. Interesting connections to partial differential equations (PDEs) have been obtained and the theory has found wide applications in areas like stochastic control, theoretical economics and mathematical finance.

BSDEs were introduced by Bismut (1973) for the linear case and by Pardoux and Peng (1990) for the general case. According to these authors, a solution to a BSDE consists of a pair of adapted processes (Y, Z) taking values in \mathbb{R}^n and $\mathbb{R}^{d \times n}$, respectively, such that

$$dY_{t} = f(t, Y_{t}, Z_{t})dt + Z'_{t}dW_{t}, \quad t \in [0, T),$$

$$Y_{T} = \xi,$$
(1.1)

where T is a finite time horizon, $(W_t)_{t\in[0,T]}$ a d-dimensional Brownian motion on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\in[0,T]}, P)$, f a progressively measurable function from $\Omega \times [0,T] \times \mathbb{R}^n \times \mathbb{R}^{d\times n}$ to \mathbb{R}^n and ξ an \mathbb{R}^n -valued, \mathcal{F}_T -measurable random variable.

The key feature of BSDEs is the random terminal condition ξ that the solution is required to satisfy. Due to the adaptedness requirement on the processes Y and Z, this condition poses certain difficulties in the stochastic setting. But these difficulties have been overcome, and now an impressive theory is available; see for instance, Bismut (1973, 1978), Arkin and Saksonov (1979), Pardoux and Peng (1990, 1992, 1994), Peng (1990, 1991, 1992a, 1992b, 1992c, 1993), Antonelli (1993), Ma, et al. (1994, 1999, 2002), Douglas et al. (1996), Cvitanić and Ma (1996), Cvitanić and Karatzas (1996), Chevance (1997), Cvitanić et al. (1999), Pardoux and Tang (1999), Delarue (2002), Bouchard and Touzi (2004), or the overview paper El Karoui et al. (1997).

If the randomness in the parameters f and ξ in (1.1) is coming from the state of a forward SDE, then the BSDE is referred to as a forward-backward stochastic differential equation (FBSDE) and its solution can be written as a deterministic function of time and the state process. Under suitable regularity assumptions, this function can be shown to be the solution of a parabolic PDE. FBSDEs are called uncoupled if the solution to the BSDE does not enter the dynamics of the forward SDE and coupled if it does. The corresponding parabolic PDE is semi-linear in case the FBSDE is uncoupled and quasilinear if the FBSDE is coupled; see Peng (1991, 1992b), Pardoux and Peng (1992), Antonelli (1993), Ma et al. (1994), Pardoux and Tang (1999), Ma and Yong (1999). These connections between FBSDEs and PDEs have led to interesting stochastic representation results for solutions of semi-linear and quasi-linear parabolic PDEs, generalizing the Feynman–Kac representation of linear parabolic PDEs and opening the way to Monte Carlo methods for the numerical treatment of such PDEs, see for instance, Zhang (2001), Bally and Pagès (2002), Ma et al (1994, 1999, 2002), Bouchard and Touzi (2004), Delarue and Menozzi (2004). However, PDEs corresponding to standard FBSDEs cannot be non-linear in the second order derivatives because the second order terms only arise linearly through Itô's formula from the quadratic variation of the underlying state process.

In this paper we introduce FBSDEs with second order dependence in the generator f. We call them second order backward stochastic differential equations (2BSDEs) and show how

they are related to fully non-linear parabolic PDEs. This extends the range of connections between stochastic equations and PDEs. In particular, it opens the way for the development of Monte Carlo methods for the numerical solution of fully non-linear parabolic PDEs. Our approach is motivated by results in Cheridito et al. (2005a, 2005b) which show how second order trading constraints lead to non-linear parabolic Hamilton–Jacobi–Bellman equations for the super-replication cost of European contingent claims.

The structure of the paper is as follows: In Section 2, we explain the notation and introduce 2BSDEs together with their associated PDEs. In Section 3, we show that the existence of a C^3 solution to the associated PDE implies the existence of a solution to the 2BSDE. Our main result in Section 4 shows the converse: If the PDE satisfies comparison as in the theory of viscosity solutions and suitable Lipschitz and monotonicity (parabolicity) conditions, then the existence of a solution to the 2BSDE implies that the PDE has a unique continuous viscosity solution v. Moreover, the solution of the 2BSDE can then be written in terms of v and the underlying state process. This implies that the solution of the 2BSDE is unique, and it provides a stochastic representation result for fully non-linear parabolic PDEs. In Section 5 we discuss Monte Carlo schemes for the numerical solution of such PDEs. In Section 6 we shortly discuss how the results of the paper can be adjusted to the case of PDEs with boundary conditions.

Acknowledgements. Parts of this paper were completed during a visit of Soner and Touzi to the Department of Operations Research and Financial Engineering at Princeton University. They would like to thank Princeton University, Erhan Çinlar and Patrick Cheridito for the hospitality. We also thank Shige Peng and Arturo Kohatsu-Higa for stimulating and helpful discussions.

2 Notation and definitions

Let $d \geq 1$ be a natural number. We denote by \mathcal{M}^d the set of all $d \times d$ matrices with real components. B' is the transpose of a matrix $B \in \mathcal{M}^d$ and Tr[B] its trace. By $\mathcal{M}^d_{\text{inv}}$ we denote the set of all invertible matrices in \mathcal{M}^d , by \mathcal{S}^d all symmetric matrices in \mathcal{M}^d , and by \mathcal{S}^d_+ all positive semi-definite matrices in \mathcal{M}^d . For $B, C \in \mathcal{M}^d$, we write $B \geq C$ if $B - C \in \mathcal{S}^d_+$. For $x \in \mathbb{R}^d$, we set

$$|x| := \sqrt{x_1^2 + \dots x_d^2}$$

and for $B \in \mathcal{M}^d$,

$$|B| := \sup_{x \in \mathbb{R}^d, |x| \le 1} Bx$$

Equalities and inequalities between random variables are always understood in the almost sure sense. We fix a finite time horizon $T \in (0, \infty)$ and let $(W_t)_{t \in [0,T]}$ be a d-dimensional Brownian motion on a complete probability space (Ω, \mathcal{F}, P) . For $t \in [0,T]$, we set $W_s^t := W_s - W_t$, $s \in [t,T]$ and denote by $\mathbb{F}^{t,T} = (\mathcal{F}_s^t)_{s \in [t,T]}$ the smallest filtration satisfying the usual conditions and containing the filtration generated by $(W_s^t)_{s \in [t,T]}$.

The coefficients $\mu : \mathbb{R}^d \to \mathbb{R}^d$ and $\sigma : \mathbb{R}^d \to \mathcal{M}_{inv}^d$ are continuous functions, such that for all $N \geq 0$, there exists a constant L_N with

$$|\mu(x) - \mu(y)| + |\sigma(x) - \sigma(y)| \le L_N|x - y|$$
 for all $x, y \in \mathbb{R}^d$ with $|x|, |y| \le N$, (2.1)

and constants $L \geq 0$ and $p_1 \in [0, 1]$ such that

$$|\mu(x)| + |\sigma(x)| \le L(1+|x|^{p_1})$$
 for all $x \in \mathbb{R}^d$. (2.2)

Then, for every initial condition $(t, x) \in [0, T] \times \mathbb{R}^d$, the forward SDE

$$dX_s = \mu(X_s)ds + \sigma(X_s)dW_s, \quad s \in [t, T],$$

$$X_t = x,$$
(2.3)

has a unique strong solution $(X_s^{t,x})_{s\in[t,T]}$. This follows for instance, from Theorems 2.3, 2.4 and 3.1 in Chapter IV of Ikeda and Watanabe (1989). Note that for existence and uniqueness of a strong solution to the SDE (2.3), $p_1 = 1$ in condition (2.2) is enough. But for $p_1 \in [0,1)$, we will get a better growth exponent p in Proposition 4.5 below. In any case, the process $(X_s^{t,x})_{s\in[t,T]}$ is adapted to the filtration $\mathbb{F}^{t,T}$, and by Itô's formula we have for all $s \in [t,T]$ and $\varphi \in C^{1,2}([t,T] \times \mathbb{R}^d)$,

$$\varphi\left(s, X_s^{t,x}\right) = \varphi(t, x) + \int_t^s \mathcal{L}\varphi\left(r, X_r^{t,x}\right) dr + \int_t^s D\varphi\left(r, X_r^{t,x}\right)' dX_r^{t,x}, \qquad (2.4)$$

where

$$\mathcal{L}\varphi(t,x) = \varphi_t(t,x) + \frac{1}{2} \text{Tr}[D^2 \varphi(t,x) \sigma(x) \sigma(x)'],$$

and $D\varphi$, $D^2\varphi$ are the gradient and the matrix of second derivatives of φ with respect to the x variables.

In the whole paper, $f:[0,T)\times\mathbb{R}^d\times\mathbb{R}\times\mathbb{R}^d\times\mathcal{S}^d\to\mathbb{R}$ and $g:\mathbb{R}^d\to\mathbb{R}$ are continuous functions.

Definition 2.1 Let $(t,x) \in [0,T) \times \mathbb{R}^d$ and $(Y_s,Z_s,\Gamma_s,A_s)_{s\in[t,T]}$ a quadruple of $\mathbb{F}^{t,T}$ progressively measurable processes taking values in \mathbb{R} , \mathbb{R}^d , \mathcal{S}^d and \mathbb{R}^d , respectively. Then
we call (Y,Z,Γ,A) a solution to the second order backward stochastic differential equation
(2BSDE) corresponding to $(X^{t,x},f,g)$ if

$$dY_s = f(s, X_s^{t,x}, Y_s, Z_s, \Gamma_s) ds + Z_s' \circ dX_s^{t,x}, \quad s \in [t, T),$$
(2.5)

$$dZ_s = A_s ds + \Gamma_s dX_s^{t,x}, \quad s \in [t, T),$$

$$(2.6)$$

$$Y_T = g\left(X_T^{t,x}\right), (2.7)$$

where $Z_s' \circ dX_s^{t,x}$ denotes Fisk-Stratonovich integration, which is related to Itô integration by

$$Z'_{s} \circ dX_{s}^{t,x} = Z'_{s} dX_{s}^{t,x} + \frac{1}{2} d \langle Z, X^{t,x} \rangle_{s} = Z'_{s} dX_{s}^{t,x} + \frac{1}{2} \operatorname{Tr} [\Gamma_{s} \sigma(X_{s}^{t,x}) \sigma(X_{s}^{t,x})'] ds.$$

The equations (2.5)–(2.7) can be viewed as a whole family of 2BSDEs indexed by $(t, x) \in [0, T) \in \mathbb{R}^d$. In the following sections, we will show relations between this family of 2BSDEs and the associated PDE

$$-v_t(t,x) + f(t,x,v(t,x),Dv(t,x),D^2v(t,x)) = 0 \text{ on } [0,T) \times \mathbb{R}^d,$$
 (2.8)

with terminal condition

$$v(T,x) = g(x), \quad x \in \mathbb{R}. \tag{2.9}$$

Since Z is a semi-martingale, the use of the Fisk–Stratonovich integral in (2.5) means no loss of generality, but it simplifies the notation in the PDE (2.8). Alternatively, (2.5) could be written in terms of the Itô integral as

$$dY_s = \tilde{f}(s, X_s^{t,x}, Y_s, Z_s, \Gamma_s) ds + Z_s' dX_s^{t,x}$$
(2.10)

for

$$\tilde{f}(t, x, y, z, \gamma) = f(t, x, y, z, \gamma) + \frac{1}{2} \operatorname{Tr} \left[\gamma \sigma(x) \sigma(x)' \right].$$

In terms of \tilde{f} , the PDE (2.8) reads as follows:

$$-v_t(t,x) + \tilde{f}(t,x,v(t,x),Dv(t,x),D^2v(t,x)) - \frac{1}{2}\text{Tr}[D^2v(t,x)\sigma(x)\sigma(x)'] = 0.$$

Note that the form of the PDE (2.8) does not depend on the functions μ and σ determining the dynamics in (2.3). So, we could restrict our attention to the case where $\mu \equiv 0$ and $\sigma \equiv I_d$, the $d \times d$ identity matrix. But the freedom to choose μ and σ from a more general class provides additional flexibility in the design of the Monte Carlo schemes discussed in Section 5 below.

3 From a solution of the PDE to a solution of the 2BSDE

Assume $v:[0,T]\times\mathbb{R}^d\to\mathbb{R}$ is a continuous function such that

$$v_t, Dv, D^2v, \mathcal{L}Dv$$
 exist and are continuous on $[0, T) \times \mathbb{R}^d$,

and v solves the PDE (2.8) with terminal condition (2.9). Then it follows directly from Itô's formula (2.4) that for each pair $(t, x) \in [0, T) \times \mathbb{R}^d$, the processes

$$\begin{aligned} Y_s &= v\left(s, X_s^{t,x}\right), \quad s \in [t, T], \\ Z_s &= Dv\left(s, X_s^{t,x}\right), \quad s \in [t, T], \\ \Gamma_s &= D^2v\left(s, X_s^{t,x}\right), \quad s \in [t, T], \\ A_s &= \mathcal{L}Dv\left(s, X_s^{t,x}\right), \quad s \in [t, T], \end{aligned}$$

solve the 2BSDE corresponding to $(X^{t,x}, f, g)$.

4 From a solution of the 2BSDE to a solution of the PDE

In all of Section 4 we assume that

$$f:[0,T)\times\mathbb{R}^d\times\mathbb{R}\times\mathbb{R}^d\times\mathcal{S}^d\to\mathbb{R}$$
 and $g:\mathbb{R}^d\to\mathbb{R}$

are continuous functions that satisfy the following Lipschitz and growth assumptions:

(A1) For every $N \ge 1$ there exists a constant F_N such that

$$|f(t, x, y, z, \gamma) - f(t, x, \tilde{y}, z, \gamma)| \le F_N |y - \tilde{y}|$$

for all $(t, x, y, \tilde{y}, z, \gamma) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^2 \times \mathbb{R}^d \times \mathcal{S}^d$ with max $\{|x|, |y|, |\tilde{y}|, |z|, |\gamma|\} \leq N$. (A2) There exist constants F and $p_2 \geq 0$ such that

$$|f(t, x, y, z, \gamma)| \le F(1 + |x|^{p_2} + |y| + |z|^{p_2} + |\gamma|^{p_2})$$

for all $(t, x, y, z, \gamma) \in [0, T] \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{S}^d$.

(A3) There exist constants G and $p_3 \ge 0$ such that

$$|g(x)| \le G(1+|x|^{p_3})$$
 for all $x \in \mathbb{R}^d$.

4.1 Admissible strategies

We fix constants $p_4, p_5 \ge 0$ and denote for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $m \ge 0$ by $\mathcal{A}_m^{t, x}$ the class of all processes of the form

$$Z_s = z + \int_t^s A_r dr + \int_t^s \Gamma_r dX_r^{t,x}, \quad s \in [t, T],$$

where $z \in \mathbb{R}^d$, $(A_s)_{s \in [t,T]}$ is an \mathbb{R}^d -valued, $\mathbb{F}^{t,T}$ -progressively measurable process, $(\Gamma_s)_{s \in [t,T]}$ is an \mathcal{S}^d -valued, $\mathbb{F}^{t,T}$ -progressively measurable process such that

$$\max\{|Z_s|, |A_s|, |\Gamma_s|\} \le m(1 + |X_s^{t,x}|^{p_4}) \quad \text{for all } s \in [t, T], \tag{4.1}$$

and

$$|\Gamma_r - \Gamma_s| \le m(1 + |X_r^{t,x}|^{p_5} + |X_s^{t,x}|^{p_5})(|r - s| + |X_r^{t,x} - X_s^{t,x}|)$$
 for all $r, s \in [t, T]$. (4.2)

Set $\mathcal{A}^{t,x} := \bigcup_{m \geq 0} \mathcal{A}^{t,x}_m$. It follows from the assumptions (A1) and (A2) on f and condition (4.1) on Z that for all $y \in \mathbb{R}$ and $Z \in \mathcal{A}^{t,x}$, the forward SDE

$$dY_s = f(s, X_s^{t,x}, Y_s, Z_s, \Gamma_s) ds + Z_s' \circ dX_s^{t,x}, \quad s \in [t, T],$$
(4.3)

$$Y_t = y, (4.4)$$

has a unique strong solution $\left(Y_s^{t,x,y,Z}\right)_{s\in[t,T]}$ (this can, for instance, be shown with the arguments in the proofs of Theorems 2.3, 2.4 and 3.1 in Chapter IV of Ikeda and Watanabe, 1989).

4.2 Auxiliary stochastic target problems

For every $m \geq 0$, we define the functions $V^m, U_m : [0, T] \times \mathbb{R}^d \to \mathbb{R}$ as follows:

$$V^m(t,x) := \inf\{y \in \mathbb{R} : Y_T^{t,x,y,Z} \ge g(X_T^{t,x}) \text{ for some } Z \in \mathcal{A}_m^{t,x}\},$$

and

$$U_m(t,x) := \sup\{y \in \mathbb{R} : Y_T^{t,x,y,Z} \le g(X_T^{t,x}) \text{ for some } Z \in \mathcal{A}_m^{t,x}\}.$$

Notice that these problems do not fit into the class of stochastic target problems studied by Soner and Touzi (2002) and are more in the spirit of Cheridito et al. (2005a, b).

Lemma 4.1 (Partial Dynamic Programming Principle)

Let
$$(t, x, m) \in [0, T) \times \mathbb{R}^d \times \mathbb{R}_+$$
 and $(y, Z) \in \mathbb{R} \times \mathcal{A}_m^{t, x}$ such that $Y_T^{t, x, y, Z} \geq g(X_T^{t, x})$. Then

$$Y_s^{t,x,y,Z} \geq V^m(s,X_s^{t,x}) \text{ for all } s \in (t,T).$$

Proof. Fix $s \in (t,T)$ and denote by $C^d[t,s]$ the set of all continuous functions from [t,s] to \mathbb{R}^d . Since $X_s^{t,x}$ and $Y_s^{t,x,y,Z}$ are \mathcal{F}_s^t -measurable, there exist measurable functions

$$\xi: C^d[t,s] \to \mathbb{R}^d$$
 and $\psi: C^d[t,s] \to \mathbb{R}$

such that

$$X_s^{t,x} = \xi(W^{t,s})$$
 and $Y_s^{t,x,y,Z} = \psi(W^{t,s})$,

where we denote $W^{t,s} := (W_r^t)_{r \in [t,s]}$. The process Z is of the form

$$Z_r = z + \int_t^r A_u du + \int_t^r \Gamma_u dX_u^{t,x}, \quad r \in [t, T],$$

for $z \in \mathbb{R}$, $(A_r)_{r \in [t,T]}$ an \mathbb{R}^d -valued, $\mathbb{F}^{t,T}$ -progressively measurable process, and $(\Gamma_r)_{r \in [t,T]}$ an \mathcal{S}^d -valued, $\mathbb{F}^{t,T}$ -progressively measurable process. Therefore, there exist progressively measurable functions (see Definition 3.5.15 in Karatzas and Shreve, 1991)

$$\zeta, \phi : [t, T] \times C^d[t, T] \to \mathbb{R}^d$$

 $\chi : [t, T] \times C^d[t, T] \to \mathcal{S}^d$

such that

$$Z_r = \zeta(r, W^{t,T}) \,, \quad A_r = \phi(r, W^{t,T}) \quad \text{and} \quad \Gamma_r = \chi(r, W^{t,T}) \quad \text{for } r \in [t,T] \,,$$

where $W^{t,T} := (W_r^t)_{r \in [t,T]}$. With obvious notation, we define for every $w \in C^d[t,s]$ the \mathbb{R}^d -valued, $\mathbb{F}^{s,T}$ -progressively measurable process $(Z_r^w)_{r \in [s,T]}$ by

$$Z_r^w = \zeta(s, w) + \int_s^r \phi(u, w + W^{s,T}) du + \int_s^r \chi(u, w + W^{s,T}) du, \quad r \in [s, T].$$

Let μ be the distribution of $W^{t,s}$ on $C^d[t,s]$. Then, $Z^w \in \mathcal{A}_m^{s,\xi(w)}$ for μ -almost all $w \in C^d[t,s]$, and

$$\begin{split} 1 &=& P\left[Y_T^{t,x,y,Z} \geq g(X_T^{t,x})\right] \\ &=& \int_{C^d[t,s]} P\left[Y_T^{s,\xi(w),\psi(w),Z^w} \geq g(X_T^{s,\xi(w)}) \mid W^{t,s} = w\right] d\mu(w) \,. \end{split}$$

Hence, for μ -almost all $w \in C^d[t, s]$, the control Z^w satisfies

$$P\left[Y_T^{s,\xi(w),\psi(w),Z^w} \geq g(X_T^{s,\xi(w)})\right] = 1\,,$$

which shows that $\psi(w) = Y_s^{s,\xi(w),\psi(w),Z^w} \geq V^m(s,\xi(w))$. In view of the definition of the functions ξ and ψ , this implies that $Y_s^{t,x,y,Z} \geq V^m(s,X_s^{t,x})$.

Since we have no a priori knowledge of any regularity of the functions V^m and U_m , we introduce the semi-continuous envelopes as in Barles and Perthame (1988)

$$V_*^m(t,x) := \liminf_{(\tilde{t},\tilde{x}) \to (t,x)} V^m\left(\tilde{t},\tilde{x}\right) \text{ and } U_m^*(t,x) := \limsup_{(\tilde{t},\tilde{x}) \to (t,x)} U_m\left(\tilde{t},\tilde{x}\right),$$

for all $(t,x) \in [0,T] \times \mathbb{R}^d$, where we set by convention $V^m(t,x) = -U^m(t,x) = \infty$ for $t \notin [0,T]$. For the theory of viscosity solutions, we refer to Crandal et al. (1992) and the book of Fleming and Soner (1993).

Theorem 4.2 Let $m \geq 0$, and assume that V_*^m is \mathbb{R} -valued. Then V_*^m is a viscosity supersolution of the PDE

$$-v_t(t,x) + \sup_{\beta \in \mathcal{S}_+^d} f(t,x,v(t,x),Dv(t,x),D^2v(t,x) + \beta) = 0 \quad on [0,T) \times \mathbb{R}^d.$$

Before turning to the proof of this result, let us state the corresponding claim for the value function U_m .

Corollary 4.3 Let $m \geq 0$, and assume that U_m^* is \mathbb{R} -valued. Then U_m^* is a viscosity subsolution of the PDE

$$-u_{t}(t,x) + \inf_{\beta \in \mathcal{S}^{d}_{+}} f(t,x,u(t,x),Du(t,x),D^{2}u(t,x) - \beta) = 0 \quad on \ [0,T) \times \mathbb{R}^{d} . (4.5)$$

Proof. Observe that for all $(t, x) \in [0, T) \times \mathbb{R}^d$,

$$-U_m(t,x) = \inf \left\{ y \in \mathbb{R} : \hat{Y}_T^{t,x,y,Z} \ge -g(X_T^{t,x}) \text{ for some } Z \in \mathcal{A}_m^{t,x} \right\},$$

where for given $(y, Z) \in \mathbb{R} \times \mathcal{A}_m^{t,x}$, the process $\hat{Y}^{t,x,y,Z}$ is the unique strong solution of the SDE

$$Y_s = y + \int_t^s -f(r, X_r^{t,x}, -Y_r, -Z_r, -\Gamma_r) dr + \int_t^s (Z_r)' \circ dX_r^{t,x}, \quad s \in [t, T].$$

Hence, it follows from Theorem 4.2 that $-U_m^* = (-U_m)_*$ is a viscosity supersolution of the PDE

$$-u_t(t,x) - \inf_{\beta \in \mathcal{S}_+^d} f(t,x,-u(t,x), -Du(t,x), -D^2u(t,x) - \beta) = 0 \text{ on } [0,T) \times \mathbb{R}^d,$$

which shows that U_m^* is a viscosity subsolution of the PDE (4.5) on $[0,T)\times\mathbb{R}^d$.

Proof of Theorem 4.2 Choose $(t_0, x_0) \in [0, T) \times \mathbb{R}^d$ and $\varphi \in C^{\infty}([0, T) \times \mathbb{R}^d)$ such that

$$0 = (V_*^m - \varphi)(t_0, x_0) = \min_{(t, x) \in [0, T) \times \mathbb{R}^d} (V_*^m - \varphi)(t, x).$$

Let $(t_n, x_n) \to (t_0, x_0)$ such that $V^m(t_n, x_n) \to V^m_*(t_0, x_0)$. There exist positive numbers $\varepsilon_n \to 0$ such that for $y_n = V^m(t_n, x_n) + \varepsilon_n$, there exists $Z^n \in \mathcal{A}_m^{t_n, x_n}$ with

$$Y_T^n \geq g(X_T^n),$$

where we denote $(X^n, Y^n) = (X^{t_n, x_n}, Y^{t_n, x_n, y_n, Z^n})$ and

$$Z_s^n = z_n + \int_{t_n}^s A_r^n dr + \int_{t_n}^s \Gamma_r^n dX_r^n, \quad s \in [t_n, T].$$

Note that for all n, $\Gamma_{t_n}^n$ is almost surely constant, and $|z_n|, |\Gamma_{t_n}^n| \leq m(1+|x_n|^{p_4})$ by assumption (4.1). Hence, by passing to a subsequence, we can assume that $z_n \to z_0 \in \mathbb{R}^d$ and $\Gamma_{t_n}^n \to \gamma_0 \in \mathcal{S}^d$. Observe that $\alpha_n := y_n - \varphi(t_n, x_n) \to 0$. We choose a decreasing sequence of numbers $\delta_n \in (0, T - t_n)$ such that $\delta_n \to 0$ and $\alpha_n/\delta_n \to 0$. By Lemma 4.1,

$$Y_{t_n+\delta_n}^n \geq V^m \left(t_n+\delta_n, X_{t_n+\delta_n}^n\right),$$

and therefore,

$$Y_{t_n+\delta_n}^n - y_n + \alpha_n \geq \varphi(t_n + \delta_n, X_{t_n+\delta_n}^n) - \varphi(t_n, x_n),$$

which, after two applications of Itô's Lemma, becomes

$$\alpha_{n} + \int_{t_{n}}^{t_{n}+\delta_{n}} [f(s, X_{s}^{n}, Y_{s}^{n}, Z_{s}^{n}, \Gamma_{s}^{n}) - \varphi_{t}(s, X_{s}^{n})] ds$$

$$+ [z_{n} - D\varphi(t_{n}, x_{n})]' [X_{t_{n}+\delta_{n}}^{n} - x_{n}]$$

$$+ \int_{t_{n}}^{t_{n}+\delta_{n}} \left(\int_{t_{n}}^{s} A_{r}^{n} - \mathcal{L}D\varphi(r, X_{r}^{n}) dr \right)' \circ dX_{s}^{n}$$

$$+ \int_{t_{n}}^{t_{n}+\delta_{n}} \left(\int_{t_{n}}^{s} \Gamma_{r}^{n} - D^{2}\varphi(r, X_{r}^{n}) dX_{r}^{n} \right)' \circ dX_{s}^{n} \geq 0$$

$$(4.6)$$

It is shown in Lemma 4.4 below that the sequence of random vectors

$$\begin{pmatrix}
\delta_{n}^{-1} \int_{t_{n}}^{t_{n}+\delta_{n}} [f(s, X_{s}^{n}, Y_{s}^{n}, Z_{s}^{n}, \Gamma_{s}^{n}) - \varphi_{t}(s, X_{s}^{n})] ds \\
\delta_{n}^{-1/2} [X_{t_{n}+\delta_{n}}^{n} - x_{n}] \\
\delta_{n}^{-1} \int_{t_{n}}^{t_{n}+\delta_{n}} \left(\int_{t_{n}}^{s} A_{r}^{n} - \mathcal{L}D\varphi(r, X_{r}^{n}) dr \right)' \circ dX_{s}^{n} \\
\delta_{n}^{-1} \int_{t_{n}}^{t_{n}+\delta_{n}} \left(\int_{t_{n}}^{s} \Gamma_{r}^{n} - D^{2}\varphi(r, X_{r}^{n}) dX_{r}^{n} \right)' \circ dX_{s}^{n}
\end{pmatrix}, n \geq 1, \tag{4.7}$$

converges in distribution to

$$\begin{pmatrix} f(t_0, x_0, \varphi(t_0, x_0), z_0, \gamma_0) - \varphi_t(t_0, x_0) \\ \sigma(x_0) W_1 \\ 0 \\ \frac{1}{2} W_1' \sigma(x_0)' [\gamma_0 - D^2 \varphi(t_0, x_0)] \sigma(x_0) W_1 \end{pmatrix}$$
(4.8)

Set $\eta_n = |z_n - D\varphi(t_n, x_n)|$, and assume $\delta_n^{-1/2} \eta_n \to \infty$ along a subsequence. Then, along another subsequence, $\eta_n^{-1}(z_n - D\varphi(t_n, x_n))$ converges to some $\eta_0 \in \mathbb{R}^d$ with

$$|\eta_0| = 1. (4.9)$$

Multiplying inequality (4.6) with $\delta_n^{-1/2}\eta_n^{-1}$ and passing to the limit yields

$$\eta_0'\sigma(x_0)W_1 \ge 0\,,$$

which, since $\sigma(x_0)$ is invertible, contradicts (4.9). Hence, the sequence $(\delta_n^{-1/2}\eta_n)$ has to be bounded, and therefore, possibly after passing to a subsequence,

$$\delta_n^{-1/2}[z_n - D\varphi(t_n, x_n)]$$
 converges to some $\xi_0 \in \mathbb{R}^d$.

It follows that $z_0 = D\varphi(t_0, x_0)$. Moreover, we can divide inequality (4.6) by δ_n and pass to the limit to get

$$f(t_0, x_0, \varphi(t_0, x_0), D\varphi(t_0, x_0), \gamma_0) - \varphi_t(t_0, x_0) + \xi_0' \sigma(x_0) W_1 + \frac{1}{2} W_1' \sigma(x_0)' [\gamma_0 - D^2 \varphi(t_0, x_0)] \sigma(x_0) W_1 \ge 0.$$

$$(4.10)$$

Since the support of the random vector W_1 is \mathbb{R}^d , it follows from (4.10) that

$$f(t_0, x_0, \varphi(t_0, x_0), D\varphi(t_0, x_0), \gamma_0) - \varphi_t(t_0, x_0)$$
+ $\xi'_0 \sigma(x_0) w + \frac{1}{2} w' \sigma(x_0)' [\gamma_0 - D^2 \varphi(t_0, x_0)] \sigma(x_0) w \ge 0$,

for all $w \in \mathbb{R}^d$. This shows that

$$f(t_0, x_0, \varphi(t_0, x_0), D\varphi(t_0, x_0), \gamma_0) - \varphi_t(t_0, x_0) \ge 0$$
 and $\beta := \gamma_0 - D^2 \varphi(t_0, x_0) \ge 0$, and hence,

$$-\varphi_t(t_0, x_0) + \sup_{\beta \in \mathcal{S}^d_+} f(t_0, x_0, \varphi(t_0, x_0), D\varphi(t_0, x_0), D^2\varphi(t_0, x_0) + \beta) \ge 0.$$

Lemma 4.4 The sequence of random vectors (4.7) converges in distribution to (4.8).

Proof. With the methods used to solve Problem 5.3.15 in Karatzas and Shreve (1991) it can also be shown that for all fixed q > 0 and $m \ge 0$, there exists a constant $C \ge 0$ such that for all $0 \le t \le s \le T$, $x \in \mathbb{R}^d$, $y \in \mathbb{R}$ and $Z \in \mathcal{A}_m^{t,x}$,

$$\operatorname{E}\left[\max_{s\in[t,T]}|X_s^{t,x}|^q\right] \leq C(1+|x|^q) \tag{4.11}$$

$$E\left[\max_{r\in[t,s]}|X_r^{t,x}-x|^q\right] \le C(1+|x|^q)(s-t)^{q/2}$$
(4.12)

$$E\left[\max_{s\in[t,T]}|Y_{s}^{t,x,y,Z}|^{q}\right] \leq C(1+|y|^{q}+|x|^{\tilde{q}})$$
(4.13)

$$E\left[\max_{r\in[t,s]}|Y_r^{t,x,y,Z}-y|^q\right] \leq C(1+|y|^q+|x|^{\tilde{q}})(s-t)^{q/2}, \tag{4.14}$$

where $\tilde{q} := \max\{p_2q, p_2p_4q, (p_4+2p_1)q\}$. For every $n \geq 1$, we introduce the $\mathbb{F}^{t_n,T}$ -stopping time

$$\tau_n := \inf\{s \ge t_n : X_s^n \notin B_1(x_0)\} \wedge (t_n + \delta_n),$$

where $B_1(x_0)$ denotes the open unit ball in \mathbb{R}^d around x_0 . It follows from the fact that $x_n \to x_0$ and (4.12) that

$$P[\tau_n < t_n + \delta_n] \to 0. \tag{4.15}$$

The difference

$$(X_{t_n+\delta_n}^n - x_n) - \sigma(x_0)(W_{t_n+\delta_n} - W_{t_n})$$

can be written as

$$\int_{t_n}^{t_n+\delta_n} \mu(X_s^n) ds + \int_{t_n}^{t_n+\delta_n} [\sigma(X_s^n) - \sigma(x_n)] dW_s + (\sigma(x_n) - \sigma(x_0)) (W_{t_n+\delta_n} - W_{t_n}),$$

and obviously,

$$\frac{1}{\sqrt{\delta_n}}(\sigma(x_n) - \sigma(x_0))(W_{t_n + \delta_n} - W_{t_n}) \to 0 \quad \text{in } L^2.$$

Moreover, it can be deduced with standard arguments from (2.1), (2.2), (4.11) and (4.12) that

$$\frac{1}{\sqrt{\delta_n}} \int_{t_n}^{t_n + \delta_n} \mu(X_s^n) ds \to 0 \quad \text{and} \quad \frac{1}{\sqrt{\delta_n}} \int_{t_n}^{\tau_n} [\sigma(X_s^n) - \sigma(x_n)] dW_s \to 0 \quad \text{in } L^2.$$

This shows that

$$\frac{1}{\sqrt{\delta_n}} \left\{ X_{t_n + \delta_n}^n - x_n - \sigma(x_0) (W_{t_n + \delta_n} - W_{t_n}) \right\} \to 0 \quad \text{in probability.} \tag{4.16}$$

Similarly, it can be derived from the boundedness assumption (4.1) on A^n that

$$\frac{1}{\delta} \int_{t_n}^{t_n + \delta_n} \left(\int_{t_n}^s A_r^n - \mathcal{L}D\varphi(r, X_r^n) \, dr \right)' \circ dX_s^n \to 0 \quad \text{in probability.} \tag{4.17}$$

By the continuity assumption (4.2) on Γ^n ,

$$\frac{1}{\delta_n} \int_{t_n}^{t_n + \delta_n} \left(\int_{t_n}^s [\Gamma_r^n - \Gamma_{t_n}^n] dX_r^n \right)' \circ dX_s^n \to 0 \quad \text{in } L^2,$$

and

$$\frac{1}{\delta_n} \left\{ \int_{t_n}^{t_n + \delta_n} \left(\int_{t_n}^{s} \Gamma_{t_n}^n dX_r^n \right)' \circ dX_s^n - \frac{1}{2} (W_{t_n + \delta_n} - W_{t_n})' \sigma(x_0)' \gamma_0 \sigma(x_0) (W_{t_n + \delta_n} - W_{t_n}) \right\} \to 0 \quad \text{in probability.}$$

Hence,

$$\frac{1}{\delta_n} \left\{ \int_{t_n}^{t_n + \delta_n} \left(\int_{t_n}^{s} \Gamma_r^n dX_r^n \right)' \circ dX_s^n \right.$$

$$\left. -\frac{1}{2} (W_{t_n + \delta_n} - W_{t_n})' \sigma(x_0)' \gamma_0 \sigma(x_0) (W_{t_n + \delta_n} - W_{t_n}) \right\} \to 0 \quad \text{in probability.}$$
(4.18)

Similarly, it can be shown that

$$\frac{1}{\delta_n} \left\{ \int_{t_n}^{t_n + \delta_n} \left(\int_{t_n}^s D^2 \varphi(r, X_r^n) dX_r^n \right)' \circ dX_s^n \right.$$

$$\left. -\frac{1}{2} (W_{t_n + \delta_n} - W_{t_n})' \sigma(x_0)' D^2 \varphi(t_0, x_0) \sigma(x_0) (W_{\tau_n} - W_{t_n}) \right\} \to 0 \quad \text{in probability.}$$
(4.19)

Finally, it follows from the continuity of f and φ_t as well as (4.1), (4.2), (4.11), (4.12) and (4.14) that

$$\frac{1}{\delta_n} \int_{t_n}^{t_n + \delta_n} [f(X_s^n, Y_s^n, Z_s^n, \Gamma_s^n) - \varphi_t(s, X_s^n)] ds \to f(x_0, D\varphi(t_0, x_0), z_0, \gamma_0) - \varphi_t(t_0, x_0)$$
(4.20)

in probability. Now, the lemma follows from (4.16)–(4.20) and the simple fact that for each n, the random vector

$$\begin{pmatrix} f(x_0, \varphi(t_0, x_0), z_0, \gamma_0) - \varphi_t(t_0, x_0) \\ \delta_n^{-1/2} \sigma(x_0) (W_{t_n + \delta_n} - W_{t_n}) \\ 0 \\ \delta_n^{-1} \frac{1}{2} (W_{t_n + \delta_n} - W_{t_n})' \sigma(x_0)' [\gamma_0 - D^2 \varphi(t_0, x_0)] \sigma(x_0) (W_{t_n + \delta_n} - W_{t_n}) \end{pmatrix}$$

has the same distribution as

$$\begin{pmatrix} f(x_0, \varphi(t_0, x_0), z_0, \gamma_0) - \varphi_t(t_0, x_0) \\ \sigma(x_0)W_1 \\ 0 \\ \frac{1}{2}W_1'\sigma(x_0)'[\gamma_0 - D^2\varphi(t_0, x_0)]\sigma(x_0)W_1 \end{pmatrix}$$

We conclude this subsection by the following bounds on the growth of the value functions V^m and U_m .

Proposition 4.5 Let $p = \max \{p_2, p_3, p_2p_4, p_4 + 2p_1\}$. Then there exists for every $m \ge 0$ a constant $C_m \ge 0$ such that

$$V_*^m(t,x) \ge -C_m(1+|x|^p) \quad and$$
 (4.21)

$$U_m^*(t,x) \le C_m(1+|x|^p)$$
 (4.22)

for all $(t,x) \in [0,T] \times \mathbb{R}^d$. Moreover,

$$V_*^m(T, x) \ge g(x) \quad and \tag{4.23}$$

$$U_m^*(T, x) \le g(x) \tag{4.24}$$

for all $x \in \mathbb{R}$.

Proof. We show (4.22) and (4.24). The proofs of (4.21) and (4.23) are completely analogous. To prove (4.22) it is enough to show that for fixed $m \geq 0$, there exists a constant $C_m \geq 0$ such that for all $(t,x) \in [0,T] \times \mathbb{R}^d$ and $(y,Z) \in \mathbb{R} \times \mathcal{A}_m^{t,x}$ satisfying $Y_T^{t,x,y,Z} \leq g(X_T^{t,x})$, we have

$$y \leq C_m(1+|x|^p).$$

For $y \leq 0$ there is nothing to show. So, we assume y > 0 and introduce the stopping time

$$\tau := \inf \left\{ s \ge t \mid Y_s^{t,x,y,Z} = 0 \right\} \wedge T.$$

Then, we have for all $s \in [t, T]$,

$$\begin{split} Y_{s \wedge \tau}^{t,x,y,Z} + \int_{s}^{s \vee \tau} f(r,X_r^{t,x},Y_r^{t,x,y,Z},Z_r,\Gamma_r) dr + \int_{s}^{s \vee \tau} Z_r' \mu(X_r^{t,x}) dr \\ + \int_{s}^{s \vee \tau} Z_r' \sigma(X_r^{t,x}) dW_r + \frac{1}{2} \int_{s}^{s \vee \tau} \mathrm{Tr}[\Gamma_r \sigma(X_r^{t,x}) \sigma(X_r^{t,x})'] dr \\ = Y_{s \wedge \tau}^{t,x,y,Z} + \int_{s}^{s \vee \tau} f(r,X_r^{t,x},Y_r^{t,x,y,Z},Z_r,\Gamma_r) dr + \int_{s}^{s \vee \tau} Z_r' \circ dX_r^{t,x} \\ = Y_{\tau}^{t,x,y,Z} \\ \leq g(X_T^{t,x}) \vee 0 \,. \end{split}$$

Hence, it follows from (A2), (A3), (2.2) and (4.1) that for $\tilde{p} = \max\{p_2, p_2p_4, p_4 + 2p_1\}$,

$$\begin{split} h(s) &:= \operatorname{E}\left[Y_{s \wedge \tau}^{t,x,y,Z}\right] \\ &\leq \operatorname{E}\left[\left|g(X_T^{t,x})\right|\right] + \operatorname{E}\left[\int_s^{s \vee \tau} \left|f(r,X_r^{t,x},Y_r^{t,x,y,Z},Z_r,\Gamma_r)\right|dr\right] \\ &+ \operatorname{E}\left[\int_s^{s \vee \tau} \left|Z_r'\mu(X_r^{t,x})\right|dr\right] + \operatorname{E}\left[\frac{1}{2}\int_s^{s \vee \tau} \left|\operatorname{Tr}\left[\Gamma_r\sigma(X_r^{t,x})\sigma(X_r^{t,x})'\right]\right|dr\right] \\ &\leq G\operatorname{E}\left[1 + \left|X_T^{t,x}\right|^{p_3}\right] + F\int_s^T h(r)dr + K\int_s^T (1 + \left|X_r^{t,x}\right|^{\tilde{p}})dr \\ &\leq \tilde{K}(1 + |x|^p) + F\int_s^T h(r)dr\,, \end{split}$$

for constants K and \tilde{K} independent of t, x, y and Z. It follows from Gronwall's lemma that

$$h(s) \le \tilde{K}(1+|x|^p)e^{F(T-s)}$$
 for all $s \in [t,T]$.

In particular, $y = h(t) \le C_m(1 + |x|^p)$ for some constant C_m independent of t, x, y and Z.

To prove (4.24), we assume by way of contradiction that there exists an $x \in \mathbb{R}^d$ such that $U_m^*(T,x) \geq g(x) + 3\varepsilon$ for some $\varepsilon > 0$. Then, there exists a sequence $(t_n,x_n)_{n\geq 1}$ in $[0,T) \times \mathbb{R}^d$ converging to (T,x) such that $U_m(t_n,x_n) \geq g(x) + 2\varepsilon$ for all $n\geq 1$. Hence, for every integer $n\geq 1$, there exists a real number $y_n\in [g(x)+\varepsilon,g(x)+2\varepsilon]$ and a process $Z^n\in \mathcal{A}_m^{t_n,x_n}$ of the form $Z_s^n=z_n+\int_{t_n}^s A_s^n ds+\int_{t_n}^s \Gamma_s^n dX_s^{t_n,x_n}$ such that

$$y_n \le g(X_T^{t_n, x_n}) - \int_{t_n}^T f(s, X_s^{t_n, x_n}, Y_s^{t, x_n, y_n, Z^n}, Z_s^n, \Gamma_s^n) ds - \int_{t_n}^T (Z_s^n)' \circ dX_s^{t_n, x_n}. \tag{4.25}$$

By (4.1), (4.11), (4.12) and (4.13), the right-hand side of (4.25) converges to g(x) in probability. Therefore, it follows from (4.25) that $g(x) + \varepsilon \leq g(x)$. But this is absurd, and hence, we must have $U_m^*(T,x) \leq g(x)$ for all $x \in \mathbb{R}^d$.

4.3 Main result

For our main result, Theorem 4.9 below, we need two more assumptions on the functions f and g, the first of which is

(A4) For all $(t, x, y, z) \in [0, T] \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d$ and $\gamma, \tilde{\gamma} \in \mathcal{S}^d$,

$$f(t, x, y, z, \gamma) \ge f(t, x, y, z, \tilde{\gamma})$$
 whenever $\gamma \le \tilde{\gamma}$.

Remark 4.6 Under (A1)–(A4) it immediately follows from Theorem 4.2 that V_*^m is a viscosity supersolution of the PDE (2.8) on $[0,T)\times\mathbb{R}^d$, provided it is \mathbb{R} -valued. Analogously, if (A1)–(A4) hold and U_m^* is \mathbb{R} -valued, Corollary 4.3 implies that U_m^* is a viscosity subsolution of the PDE (2.8) on $[0,T)\times\mathbb{R}^d$.

For our last assumption and the statement of Theorem 4.9 we need the following

Definition 4.7 Let $q \geq 0$.

- **1.** We call a function $v:[0,T]\times\mathbb{R}^d\to\mathbb{R}$ a viscosity solution with growth q of the PDE (2.8) with terminal condition (2.9) if v is a viscosity solution of (2.8) on $[0,T)\times\mathbb{R}^d$ such that $v^*(T,x)=v_*(T,x)=g(x)$ for all $x\in\mathbb{R}^d$ and there exists a constant C such that $|v(t,x)|\leq C(1+|x|^q)$ for all $(t,x)\in[0,T]\times\mathbb{R}^d$.
- **2.** We say that the PDE (2.8) with terminal condition (2.9) has comparison with growth q if the following holds:

If $w:[0,T]\times\mathbb{R}^d\to\mathbb{R}$ is lower semicontinuous and a viscosity supersolution of (2.8) on $[0,T)\times\mathbb{R}^d$ and $u:[0,T]\times\mathbb{R}^d\to\mathbb{R}$ upper semicontinuous and a viscosity subsolution of (2.8) on $[0,T)\times\mathbb{R}^d$ such that

$$w(T,x) \ge g(x) \ge u(T,x)$$
 for all $x \in \mathbb{R}^d$

and there exists a constant $C \geq 0$ with

$$w(t,x) \ge -C(1+|x|^p)$$
 and $u(t,x) \le C(1+|x|^p)$ for all $(t,x) \in [0,T) \times \mathbb{R}^d$,

then $w \ge u$ on $[0,T] \times \mathbb{R}^d$.

With this definition our last assumption on f and g is

(A5) The PDE (2.8) with terminal condition (2.9) has comparison with growth $p = \max\{p_2, p_3, p_2p_4, p_4 + 2p_1\}$.

Remarks 4.8

1. The monotonicity assumption (A4) is natural from the PDE viewpoint. It implies that f is elliptic and the PDE (2.8) parabolic. If f satisfies the following stronger version of (A4): there exists a constant C > 0 such that

$$f(t, x, y, z, \gamma - B) > f(t, x, y, z, \gamma) + C \operatorname{Tr}[B]$$
(4.26)

for all $(t, x, y, z, \gamma) \in [0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{S}^d$ and $B \in \mathcal{S}^d_+$, then the PDE (2.8) is called uniformly parabolic, and there exist general results on existence, uniqueness and smoothness of solutions, see for instance Krylov (1987) or Evans (1998). When f is linear in the γ variable (in particular, for the semi- and quasi-linear equations discussed in Subsections 5.2 and 5.3 below), the condition (4.26) essentially guarantees existence, uniqueness and smoothness of solutions to the PDE (2.8)–(2.9); see for instance, Section 5.4 in Ladyzenskaya et al. (1967). Without parabolicity there are no comparison results for PDEs of the form (2.8)–(2.9).

- 2. Condition (A5) is an implicit assumption on the functions f and g. But we find it more convenient to assume comparison directly in the form (A5) instead of putting technical assumptions on f and g which guarantee that the PDE (2.8) with terminal condition (2.9) has comparison. Several comparison results for non-linear PDEs are available in the literature; see for example, Crandall et al. (1992), Fleming and Soner (1993), Cabre and Caffarelli (1995). However, most results are stated for equations in bounded domains. For equations in the whole space, the critical issue is the interplay between the growth of solutions at infinity and the growth of the non-linearity. We list some typical situations where comparison holds:
- a) Comparison with growth 1: Assume (A1)–(A4) and there exists a function $h:[0,\infty] \to [0,\infty]$ with $\lim_{x\to 0} h(x) = 0$ such that

$$|f(t, x, y, \alpha(x - \tilde{x}), A) - f(t, \tilde{x}, y, \alpha(x - \tilde{x}), B)| \le h(\alpha|x - \tilde{x}|^2 + |x - \tilde{x}|),$$

for all (t, x, \tilde{x}, y) , $\alpha > 0$ and A, B satisfying

$$-\alpha \left[\begin{array}{cc} I & 0 \\ 0 & I \end{array} \right] \leq \left[\begin{array}{cc} A & 0 \\ 0 & -B \end{array} \right] \leq \alpha \left[\begin{array}{cc} I & -I \\ -I & I \end{array} \right].$$

Then it follows from Theorem 8.2 in Crandall et al. (1992) that equations of the form (2.8), (2.9) have comparison with growth 0 if the domain is bounded. If the domain is unbounded, it follows from the modifications outlined in Section 5.D of Crandall et al. (1992) that (2.8) and (2.9) have comparison with growth 1.

b) When the equation (2.8) is a dynamic programming equation related to a stochastic optimal control problem, then a comparison theorem for bounded solutions is given in Fleming and Soner (1993), Section 5.9, Theorem V.9.1. In this case, f has the form

$$f(t,x,y,z,\gamma) = \sup_{u \in U} \left\{ \alpha(t,x,u) + \beta(t,x,u)y + b(t,x,u)'z - \text{Tr}[c(t,x,u)\gamma] \right\},\,$$

see Subsection 5.5 below. Theorem V.9.1 in Fleming and Soner (1993) is proved under the assumption that $\beta \equiv 0$, U is a compact set and that α, b, c are uniformly Lipschitz and growing at most linearly (see IV (2.1) in Fleming and Soner, 1993). This result can be extended directly to the case where β satisfies a similar condition and to equations related to differential games, that is, when

$$f(t,x,y,z,\gamma) = \sup_{u \in U} \inf_{\tilde{u} \in \tilde{U}} \left\{ \alpha(t,x,u,\tilde{u}) + \beta(t,x,u,\tilde{u})y + b(t,x,u,\tilde{u})'z - \text{Tr}[c(t,x,u,\tilde{u})\gamma] \right\} .$$

c) Many techniques in dealing with unbounded solutions were developed by Ishii (1984) for first order equations (that is, when f is independent of γ). These techniques can be extended to second order equations. Some related results can be found in Barles et al. (1997, 2003). In Barles et al (1997), in addition to comparison results for PDEs, one can also find BSDEs based on jump Markov processes.

Theorem 4.9 (Uniqueness of 2BSDE) Assume (A1)–(A5) and there exists an $x_0 \in \mathbb{R}^d$ such that the 2BSDE corresponding to (X^{0,x_0}, f, g) has a solution $(Y^{0,x_0}, Z^{0,x_0}, \Gamma^{0,x_0}, A^{0,x_0})$ with $Z^{0,x_0} \in \mathcal{A}^{0,x_0}$. Then the following hold:

- **1.** The associated PDE (2.8) with terminal condition (2.9) has a unique viscosity solution v with growth $p = \max\{p_2, p_3, p_2p_4, p_4 + 2p_1\}$, and v is continuous on $[0, T] \times \mathbb{R}^d$.
- **2.** For all $(t,x) \in [0,T) \times \mathbb{R}^d$, there exists exactly one solution $(Y^{t,x}, Z^{t,x}, \Gamma^{t,x}, A^{t,x})$ to the 2BSDE corresponding to $(X^{t,x}, f, g)$ such that $Z^{t,x} \in \mathcal{A}^{t,x}$, and the process $Y^{t,x}$ satisfies

$$Y_s^{t,x} = v(s, X_s^{t,x}), \quad s \in [t, T],$$
 (4.27)

where v is the unique continuous viscosity solution with growth p of (2.8)–(2.9).

Before turning to the proof of this result, we make some remarks.

Remark 4.10 If the assumptions of Theorem 4.9 are fulfilled, it follows from (4.27) that $v(t,x) = Y_t^{t,x}$ for all $(t,x) \in [0,T) \times \mathbb{R}^d$. Hence, v(t,x) can be approximated by backward simulation of the process $(Y_s^{t,x})_{s \in [t,T]}$. If v is C^2 , it follows from Itô's lemma that $Z_s^{t,x} = Dv(s,X_s^{t,x})$, $s \in [t,T]$. Then, Dv(t,x) can be approximated by backward simulation of $(Z_s^{t,x})_{s \in [t,T]}$. If v is C^3 , then also $\Gamma_s^{t,x} = D^2v(s,X_s^{t,x})$, $s \in [t,T]$, and $D^2v(t,x)$ can be approximated by backward simulation of $(\Gamma_s^{t,x})_{s \in [t,T]}$. A formal discussion of a potential numerical scheme for the backward simulation of the processes $Y^{t,x}$, $Z^{t,x}$ and $\Gamma^{t,x}$ is provided in Subsection 5.4 below.

Remark 4.11 Assume there exists a classical solution v of the PDE (2.8) such that

$$v_t, Dv, D^2v, \mathcal{L}Dv$$
 exist and are continuous on $[0, T] \times \mathbb{R}^d$,

and there exists a constant $m \geq 0$ such that

and

$$|D^{2}v(t,x) - D^{2}v(s,y)| \le m(1+|x|^{p_{5}} + |y|^{p_{5}})(|t-s| + |x-y|)$$
for all $0 \le t, s \le T$ and $x, y \in \mathbb{R}^{d}$.

Note that (4.29) follows, for instance, if $\frac{\partial}{\partial t}D^2v$ and D^3v exist and

$$\max_{ij} \left| \frac{\partial}{\partial t} (D^2 v(t, x))_{ij} \right| \\
\max_{ij} \left| D(D^2 v(t, x))_{ij} \right| \\$$

$$\leq \frac{m}{d} (1 + |x|^{p_5}) \quad \text{for all } 0 \leq t \leq T \text{ and } x \in \mathbb{R}^d.$$

Fix $(t,x) \in [0,T) \times \mathbb{R}^d$. By Section 3, the processes

$$Y_{s} = v(s, X_{s}^{t,x}), \quad s \in [0, T],$$

$$Z_{s} = Dv(s, X_{s}^{t,x}), \quad s \in [0, T],$$

$$\Gamma_{s} = D^{2}v(s, X_{s}^{t,x}), \quad s \in [0, T],$$

$$A_{s} = \mathcal{L}Dv(s, X_{s}^{t,x}), \quad s \in [0, T],$$

solve the 2BSDE corresponding to $(X^{t,x}, f, g)$. By (4.28) and (4.29), Z is in $\mathcal{A}_m^{t,x}$ (see (4.1) and (4.2)). Hence, if the assumptions of Theorem 4.9 are fulfilled, (Y, Z, Γ, A) is the only solution of the 2BSDE corresponding to $(X^{t,x}, f, g)$ with $Z \in \mathcal{A}^{t,x}$.

Proof of Theorem 4.9

1. Let $(X^{0,x_0}, Y^{0,x_0}, \Gamma^{0,x_0}, A^{0,x_0})$ be a solution to the 2BSDE corresponding to (X^{0,x_0}, f, g) with $Z^{0,x_0} \in \mathcal{A}_m^{0,x_0}$ for some $m \geq 0$. Then, it follows from Lemma 4.1 that

$$Y_s^{0,x_0} \ge V^m(s, X_s^{0,x_0})$$
 for all $s \in [0, T]$, (4.30)

and by symmetry,

$$Y_s^{0,x_0} \le U_m(s, X_s^{0,x_0})$$
 for all $s \in [0, T]$. (4.31)

Recall that the inequalities (4.30) and (4.31) are understood in the P-almost sure sense. But since, by assumption, σ takes values in \mathcal{M}_{inv} , X_s^{0,x_0} has full support for all $s \in (0,T]$ (see, for instance, Nualart, 1995), and we get from (4.30) and (4.31) that

$$V^m(s,x) \leq U_m(s,x)$$
 for all (s,x) in a dense subset of $[0,T] \times \mathbb{R}^d$.

It follows that

$$V_*^m \le U_m^*$$
 on $[0, T] \times \mathbb{R}^d$.

Together with Proposition 4.5, this shows that V_*^m and U_m^* are finite on $[0,T] \times \mathbb{R}^d$. By Remark 4.6, V_*^m is a viscosity supersolution and U_m^* a viscosity subsolution of the PDE (2.8) on $[0,T) \times \mathbb{R}^d$. Therefore, it follows from Proposition 4.5 and (A5) that

$$V_*^m \ge U_m^*$$
 on $[0, T] \times \mathbb{R}^d$.

Hence, the function $v = V_*^m = V^m = U_m = U_m^*$ is continuous on $[0, T] \times \mathbb{R}^d$ and a viscosity solution with growth p of the PDE (2.8)–(2.9). By (A5), v is the only viscosity solution of the PDE (2.8)–(2.9) with growth p. This shows 1.

2. Since $v = V^m = U_m$, it follows from (4.30) and (4.31) that

$$Y_s^{0,x_0} = v(s, X_s^{0,x_0})$$
 for all $s \in [0, T]$.

By conditioning $(Y^{0,x_0}, Z^{0,x_0}, \Gamma^{0,x_0}, A^{0,x_0})$ as in the proof of Lemma 4.1 and a continuity argument, it can be deduced that for all $(t,x) \in [0,T) \times \mathbb{R}^d$, there exists a solution $(Y^{t,x}, Z^{t,x}, \Gamma^{t,x}, A^{t,x})$ of the 2BSDE corresponding to $(X^{t,x}, f, g)$ such that $Z^{t,x} \in \mathcal{A}^{t,x}$ and

$$Y_s^{t,x} = v(s, X_s^{t,x}), \quad s \in [t, T].$$

If for fixed $(t,x) \in [0,T) \times \mathbb{R}^d$, (Y,Z,Γ,A) is a solution of the 2BSDE corresponding to $(X^{t,x},f,g)$ such that $Z \in \mathcal{A}^{t,x}$, then it follows as in 1. that

$$Y_s = v(s, X_s^{t,x}) = Y_s^{t,x}, \quad s \in [t, T],$$

and therefore also,

$$(Z, \Gamma, A) = (Z^{t,x}, \Gamma^{t,x}, A^{t,x})$$

because $Z^{t,x}$ is uniquely determined by

$$Y_s^{t,x} = Y_t^{t,x} + \int_t^s f(r, X_r^{t,x_t}, Y_r^{t,x}, Z_r^{t,x}, \Gamma_r^{t,x}) dr + \int_t^s (Z_r^{t,x})' \circ dX_r^{t,x}, \quad s \in [t, T],$$

and $\Gamma^{t,x}$ and $A^{t,x}$ are uniquely determined by

$$Z_s^{t,x} = Z_t^{t,x} + \int_t^s A_r^{t,x} dr + \int_t^s \Gamma_r^{t,x} dX_r^{t,x}, \quad s \in [t,T].$$

This completes the proof of 2.

5 Monte Carlo methods for the solution of parabolic PDEs

In this section, we provide a formal discussion of the numerical implications of our representation results. We start by recalling some well-known facts in the linear case. We then review some recent advances in the semi- and quasi-linear cases, and conclude with the fully non-linear case related to Theorem 4.9.

5.1 The linear case

In this subsection, we assume that the function f is of the form

$$f(t, x, y, z, \gamma) = -\alpha(t, x) - \beta(t, x)y - \mu(x)'z - \frac{1}{2}\operatorname{Tr}\left[\sigma(x)\sigma(x)'\gamma\right]$$

Then, (2.8) is a linear parabolic PDE. Under standard conditions, it has a smooth solution v, and the Feynman-Kac representation theorem states that for all $(t, x) \in [0, T] \times \mathbb{R}^d$,

$$v(t,x) = \mathbf{E}\left[\int_{t}^{T} B_{t,s} \alpha\left(s, X_{s}^{t,x}\right) ds + B_{t,T} g\left(X_{T}^{t,x}\right)\right],$$

where

$$B_{t,s} := \exp\left(\int_{t}^{s} \beta\left(r, X_{r}^{t,x}\right) dr\right)$$

(see, for instance, Theorem 5.7.6 in Karatzas and Shreve, 1991). This representation suggests a numerical approximation of the function v by means of the so-called Monte Carlo method:

(i) Given J independent copies $\{X^j, 1 \leq j \leq J\}$ of the process $X^{t,x}$, set

$$\hat{v}^{(J)}(t,x) \ := \ \frac{1}{J} \sum_{j=1}^J \int_t^T B^j_{t,s} \; \alpha\left(s,X^j_s\right) ds + B^j_{t,T} \; g\left(X^j_T\right) \; ,$$

where $B_{t,s}^j := \exp\left(\int_t^s \beta\left(r, X_r^j\right) dr\right)$. Then, it follows from the law of large numbers and the central limit theorem that

$$\hat{v}^{(J)}(t,x) \rightarrow v(t,x) \quad \text{a.s} \quad \text{and} \quad \sqrt{J} \left(\hat{v}^{(J)}(t,x) - v(t,x) \right) \rightarrow \mathbf{N} \left(0,\rho \right) \quad \text{in distribution} \,,$$

where ρ is the variance of the random variable $\int_t^T B_{t,s} \alpha\left(s, X_s^{t,x}\right) ds + B_{t,T} g\left(X_T^{t,x}\right)$. Hence, $\hat{v}^{(J)}(t,x)$ is a consistent approximation of v(t,x). Moreover, in contrast to finite differences or finite elements methods, the error estimate is of order $J^{-1/2}$, independently of the dimension d.

(ii) In practice, it is not possible to produce independent copies $\{X^j, 1 \leq j \leq J\}$ of the process $X^{t,x}$, except in trivial cases. In most cases, the above Monte Carlo approximation is performed by replacing the process $X^{t,x}$ by a suitable discrete-time approximation X^N with time step of order N^{-1} for which independent copies $\{X^{N,j}, 1 \leq j \leq J\}$ can be produced. The simplest discrete-time approximation is the following discrete Euler scheme: Set $X_t^N = x$ and for $1 \leq n \leq N$,

$$X_{t_n}^N = X_{t_{n-1}}^N + \mu(X_{t_{n-1}}^N)(t_n - t_{n-1}) + \sigma(X_{t_{n-1}}^N)(W_{t_n} - W_{t_{n-1}}),$$

where $t_n := t + n(T - t)/N$. We refer to Talay (1996) for a survey of the main results in this area.

5.2 The semi-linear case

We next consider the case where f is given by

$$f(t, x, y, z, \gamma) = \varphi(t, x, y, z) - \mu(x)'z - \frac{1}{2}\operatorname{Tr}\left[\sigma(x)\sigma(x)'\gamma\right].$$

Then the PDE (2.8) is called semi-linear. We assume that the assumptions of Theorem 4.9 are satisfied. In view of the connection between Fisk-Stratonovich and Itô integration, the 2BSDE (2.5)–(2.7) reduces to an uncoupled FBSDE of the form

$$dY_s = \varphi(s, X_s^{t,x}, Y_s, Z_s)ds + Z_s'\sigma(X_s^{t,x})dW_s, \quad s \in [t, T),$$

$$Y_T = g(X_T^{t,x}),$$

(compare to Peng, 1991, 1992b; Pardoux and Peng 1992). For $N \geq 1$, we denote $t_n := t + n(T-t)/N$, n = 0, ..., N, and we define the discrete-time approximation Y^N of Y by the backward scheme

$$Y_T^N := g(X_T^{t,x}),$$

and, for $n = 1, \ldots, N$,

$$Y_{t_{n-1}}^{N} := \mathbb{E}\left[Y_{t_{n}}^{N} \mid X_{t_{n-1}}^{t,x}\right] - \varphi\left(t_{n-1}, X_{t_{n-1}}^{t,x}, Y_{t_{n-1}}^{N}, Z_{t_{n-1}}^{N}\right) (t_{n} - t_{n-1})$$

$$(5.1)$$

$$Z_{t_{n-1}}^{N} := \frac{1}{t_n - t_{n-1}} \left(\sigma(X_{t_{n-1}}^{t,x})' \right)^{-1} \operatorname{E} \left[(W_{t_n} - W_{t_{n-1}}) Y_{t_n}^{N} \mid X_{t_{n-1}}^{t,x} \right] . \tag{5.2}$$

Then, we have

$$\limsup_{N \to \infty} \sqrt{N} |Y_t^N - v(t, x)| < \infty,$$

see for instance, Zhang (2001), Bally and Pagès (2002), Bouchard and Touzi (2004). The practical implementation of this backward scheme requires the computation of the conditional expectations appearing in (5.1) and (5.2). This suggests the use of a Monte Carlo approximation, as in the linear case. But at every time step, we need to compute conditional expectations based on J independent copies $\{X^j, 1 \leq j \leq J\}$ of the process $X^{t,x}$. Recently, several approaches to this problem have been developed. We refer to Lions and Regnier (2001), Bally and Pagès (2003), Glasserman (2003), Bouchard and Touzi (2004) and the references therein for the methodology and the analysis of such non-linear Monte Carlo methods.

5.3 The quasi-linear case

It is shown in Antonelli (1993) and Ma et al. (1994) that coupled FBSDEs of the form

$$\begin{split} dX_s &= \mu(s,X_s,Y_s,Z_s)ds + \sigma(s,X_s,Y_s)dW_s\,, \quad s \in [t,T]\,, \\ X_t &= x \\ dY_s &= \varphi(s,X_s,Y_s,Z_s)ds + Z_s'\sigma(s,X_s,Y_s)dW_s\,, \quad s \in [t,T)\,, \\ Y_T &= g(X_T)\,, \end{split}$$

are related to quasi-linear PDEs of the form (2.8)–(2.9) with

$$f(t,x,y,z,\gamma) = \varphi(t,x,y,z) - \mu(t,x,y,z)'z - \frac{1}{2}\operatorname{Tr}\left[\sigma(t,x,y)\sigma(t,x,y)'\gamma\right],$$

see also, Pardoux and Tang (1999). Delarue and Menozzi (2004) have used this relation to build a Monte Carlo scheme for the numerical solution of quasi-linear parabolic PDEs.

5.4 The fully non-linear case

We now discuss the case of a general f as in the previous section. Set

$$\varphi(t, x, y, z, \gamma) := f(t, x, y, z, \gamma) + \mu(x)'z + \frac{1}{2}\operatorname{Tr}\left[\sigma(x)\sigma(x)'\gamma\right]. \tag{5.3}$$

Then for all $(t,x) \in [0,T) \times \mathbb{R}^d$ the 2BSDE corresponding to $(X^{t,x},f,g)$ can be written as

$$dY_s = \varphi(s, X_s^{t,x}, Y_s, Z_s, \Gamma_s) ds + Z_s' \sigma(X_s^{t,x}) dW_s, \quad s \in [t, T),$$

$$(5.4)$$

$$dZ_s = A_s ds + \Gamma_s dX_s^{t,x}, \quad s \in [t, T), \tag{5.5}$$

$$Y_T = g(X_T^{t,x}). (5.6)$$

We assume that the conditions of Theorem 4.9 hold true, so that the PDE (2.8) has a unique viscosity solution v with growth $p = \max\{p_2, p_3, p_2p_4, p_4 + 2p_1\}$, and there exists a unique solution $(Y^{t,x}, Z^{t,x}, \Gamma^{t,x}, A^{t,x})$ to the 2BSDE (5.4)–(5.6) with $Z^{t,x} \in \mathcal{A}^{t,x}$.

Comparing with the backward scheme (5.1)–(5.2) in the semi-linear case, we suggest the following discrete-time approximation of the processes $Y^{t,x}$, $Z^{t,x}$ and $\Gamma^{t,x}$:

$$Y^N_T:=g(X^{t,x}_T)\,,\quad Z^N_T:=Dg(X^{t,x}_T)\,,$$

and, for $n = 1, \ldots, N$,

$$Y_{t_{n-1}}^{N} := \mathbb{E}\left[Y_{t_{n}}^{N} \mid X_{t_{n-1}}^{t,x}\right] - \varphi(t_{n-1}, X_{t_{n-1}}^{t,x}, Y_{t_{n-1}}^{N}, Z_{t_{n-1}}^{N}, \Gamma_{t_{n-1}}^{N})(t_{n} - t_{n-1})$$

$$Z_{t_{n-1}}^{N} := \frac{1}{t_{n} - t_{n-1}} \left(\sigma(X_{t_{n-1}}^{t,x})'\right)^{-1} \mathbb{E}\left[(W_{t_{n}} - W_{t_{n-1}})Y_{t_{n}}^{N} \mid X_{t_{n-1}}^{t,x}\right],$$

$$\Gamma_{t_{n-1}}^{N} := \frac{1}{t_{n} - t_{n-1}} \mathbb{E}\left[Z_{t_{n}}^{N}(W_{t_{n}} - W_{t_{n-1}})' \mid X_{t_{n-1}}^{t,x}\right] \sigma(X_{t_{n-1}}^{t,x})^{-1}.$$

A precise analysis of the latter backward scheme is left for future research. We conjecture that

$$Y_t^N \to v(t, x)$$
 as $N \to \infty$,

if v is C^2 , then

$$Z_t^N \to Dv(t,x)$$
 as $N \to \infty$,

and if v is C^3 , then

$$\Gamma_t^N \to D^2 v(t, x)$$
 as $N \to \infty$.

5.5 Connections with standard stochastic control

Let \mathcal{U}_0 be the collection of all progressively measurable processes $(\nu_t)_{t\in[0,T]}$ with values in a given bounded subset $U\subset\mathbb{R}^k$. Let $b:[0,T]\times\mathbb{R}^d\times U\to\mathbb{R}^d$ and $a:[0,T]\times\mathbb{R}^d\times U\to\mathcal{S}^d$ be continuous functions, Lipschitz in x uniformly in (t,u). We call a process $\nu\in\mathcal{U}_0$ an admissible control if

$$\mathbb{E}\left[\int_0^T (|b(s,x,\nu_s)| + |a(s,x,\nu_s)|^2) ds\right] < \infty$$

for all $x \in \mathbb{R}^d$ and denote the class of all admissible controls by \mathcal{U} . For every pair of initial conditions $(t, x) \in [0, T] \times \mathbb{R}^d$ and each admissible control process $\nu \in \mathcal{U}$, the SDE

$$dX_{s} = b(s, X_{s}, \nu_{s})ds + a(s, X_{s}, \nu_{s})dW_{s}, \quad s \in [t, T],$$

$$X_{t} = x,$$
(5.7)

has a unique strong solution, which we denote by $(X_s^{t,x,\nu})_{s\in[t,T]}$. Let $\alpha,\beta:[0,T]\times\mathbb{R}^d\times U\to\mathbb{R}$ and $g:\mathbb{R}^d\to\mathbb{R}$ be continuous functions with $\beta\leq 0$, and assume that α and g have quadratic growth in x uniformly in (t,u). Consider the stochastic control problem

$$v(t,x) := \sup_{\nu \in \mathcal{U}} \mathbf{E} \left[\int_t^T B_{t,s}^{\nu} \alpha(s, X_s^{t,x,\nu}, \nu_s) ds + B_{t,T}^{\nu} \, g(X_T^{t,x,\nu}) \right] \,,$$

where

$$B_{t,s}^{\nu} := \exp\left(\int_t^s \beta(r, X_r^{t,x,\nu}, \nu_r) dr\right), \quad 0 \le t \le s \le T.$$

By the classical method of dynamic programming, the function v can be shown to solve the Hamilton–Jacobi–Bellman equation

$$-v_t(t,x) + f(t,x,v(t,x), Dv(t,x), D^2v(t,x)) = 0$$

$$v(T,x) = g(x),$$
 (5.8)

where

$$f(t,x,y,z,\gamma) := \sup_{u \in U} \left\{ \alpha(t,x,u) + \beta(t,x,u)y + b(t,x,u)'z - \frac{1}{2} \text{Tr}[aa'(t,x,u)\gamma] \right\}.$$

This is a fully non-linear parabolic PDE covered by the class (2.8)–(2.9). Note that f is convex in the triple (y, z, γ) . The semi-linear case is obtained when there is no control on the diffusion part, that is, a(t, x, u) = a(t, x) is independent of u.

If the value function v has a stochastic representation in terms of a 2BSDE satisfying the assumptions of Theorem (4.9), then the Monte Carlo scheme of the previous subsection can be applied to approximate v.

Under suitable regularity assumptions, the optimal control at time t is known to be of the form

$$\hat{u}(t,x,v(t,x),Dv(t,x),D^2v(t,x)),$$

where \hat{u} is a maximizer of the expression

$$\sup_{u\in U} \left\{ \alpha(t,x,u) + \beta(t,x,u)v(t,x) + b(t,x,u)'Dv(t,x) - \frac{1}{2}\mathrm{Tr}[aa'(t,x,u)D^2v(t,x)] \right\}.$$

Notice that the numerical scheme suggested in the Subsection 5.4 calculates at each step in time the values of the processes $(X^{t,x}, Y^{t,x}, Z^{t,x}, \Gamma^{t,x})$. Therefore, the optimal control is also provided by this numerical scheme by

$$\hat{\nu}_s = \hat{u}(s, X_s^{t,x}, Y_s^{t,x}, Z_s^{t,x}, \Gamma_s^{t,x})$$
.

6 Boundary value problems

In this section, we give a brief outline of an extension to boundary value problems. Namely, let $O \subset \mathbb{R}^d$ be an open set. For $(t,x) \in [0,T) \times \mathbb{R}^d$, the process $X^{t,x}$ is given as in (2.3), but we stop it at the boundary of O. Then we extend the terminal condition (2.7) in the 2BSDE to a boundary condition. In other words, we introduce the exit time

$$\theta := \inf \left\{ s \ge t \mid X_s^{t,x} \notin O \right\}$$

and modify the 2BSDE (2.5)–(2.7) to

$$\begin{split} Y_{s \wedge \theta} &= g(X_{T \wedge \theta}^{t,x}) - \int_{s \wedge \theta}^{T \wedge \theta} f(r, X_r^{t,x}, Y_r, \Gamma_r, A_r) dr - \int_{s \wedge \theta}^{T \wedge \theta} Z_r' \circ dX_r^{t,x} \,, \quad s \in [t, T] \,, \\ Z_{s \wedge \theta} &= Z_{T \wedge \theta} - \int_{s \wedge \theta}^{T \wedge \theta} A_r dr - \int_{s \wedge \theta}^{T \wedge \theta} \Gamma_r dX_r^{t,x} \,, \quad s \in [t, T] \,. \end{split}$$

Then the corresponding PDE is the same as (2.8)–(2.9), but it only holds in $[0,T) \times O$. Also, the terminal condition v(T,x) = g(x) only holds in O. In addition, the following lateral boundary condition holds

$$v(t,x) = g(x)$$
, for all $(t,x) \in [0,T] \times \partial O$.

All the results of the previous sections can easily be adapted to this case. Moreover, if we assume that O is bounded, most of the technicalities related to the growth of solutions are avoided as the solutions are expected to be bounded.

References

- [1] Antonelli, F. (1993). Backward-forward stochastic differential equations. *Annals of Applied Probability* 3, 777–793.
- [2] Arkin, V. and Saksonov, M. (1979). Necessary optimality conditions for stochastic differential equations. Soviet Math. Dokl. 20, 1–5.
- [3] Bally, V., Pagès, G. (2003). Error analysis of the quantization algorithm for obstacle problems. Stochastic Processes and their Applications 106(1), 1–40.
- [4] Barles, G., Biton, S., Bourgoing, M., Ley, O. (2003). Uniqueness results for quasilinear parabolic equations through viscosity solutions' methods. *Calc. Var. PDEs*, 18 (2),159-179.
- [5] Barles, G., Buckdahn, R., and Pardoux, E. (1997). Backward stochastic differential equations and integral-partial differential equations, Stochastics Stochastics Rep., 60, no. 1-2, 57–83.
- [6] Barles, G. and Perthame, B. (1988). Exit time problems in optimal control and vanishing viscosity solutions of Hamilton–Jacobi equations, SIAM J. Cont. Opt. 26, 1133–1148.
- [7] Bismut, J.M. (1973). Conjugate convex functions in optimal stochastic control, J. Math. Anal. Appl. 44, 384–404.
- [8] Bismut, J.M. (1978). Contrôle des système linéaire quadratiques: applications de l'integrale stochastique, Sém. Probab. XII, Lect. Notes. Math. vol. 649. Springer-Velag, 180–264.

- [9] Bouchard, B., Touzi, N. (2004). Discrete-time approximation and Monte Carlo simulation of backward stochastic differential equations, Stochastic Processes and their Applications 111, 175–206.
- [10] Cabre, X., Caffarelli, L. (1995). Fully Nonlinear Elliptic Equations, AMS, Providence.
- [11] Cheridito, P., Soner, H.M., Touzi, N. (2005a). Small time path behavior of double stochastic integrals and applications to stochastic control, *Annals of Applied Probability*, to appear.
- [12] Cheridito, P., Soner, H.M., Touzi, N. (2005b). The multi-dimensional super-replication problem under gamma constraints, *Annales de l'Institute Henri Poincaré* (C) Non Linear Analysis, to appear.
- [13] Chevance, D. (1997). Numerical Methods for Backward Stochastic Differential Equations, *Publ. Newton Inst.*, Cambridge University Press.
- [14] Crandall, M.G., Ishii, H., Lions, P.L. (1992). User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc.* 27(1), 1–67.
- [15] Cvitanić, J., Ma, J. (1996). Hedging options for a large investor and forward-backward SDE's, Ann. Appl. Probab. 6, 370–398.
- [16] Cvitanić, J., Karatzas, I. (1996). Backward stochastic differential equations with reflection and Dynkin games. *Annals of Probability*, 24(4), 2024–2056.
- [17] Cvitanić, J., Karatzas, I., Soner, H.M. (1999). Backward SDEs with constraints on the gains-process, *Annals of Probability* 26, 1522–1551.
- [18] Delarue, F. (2002). On the existence and uniqueness of solutions to FBSDEs in a non-degenerate case, Stoch. Proc. Appl. 99, 209–286.
- [19] Delarue, F., Menozzi, F. (2004). A forward-backward stochastic algorithm for quasilinear PDEs. Preprint 932, University of Paris VI & VII.
- [20] Douglas, J., Ma, J., Protter, P. (1996). Numerical methods for forward-backward stochastic differential equations. Ann. Appl. Probab. 6, 940–968.
- [21] El Karoui, N., Peng, S., Quenez, M.C. (1997). Backward stochastic differential equations in finance. *Mathematical Finance* 7(1), 1–71.
- [22] Evans, L.C. (1998). Partial Differential Equations. AMS, Providence.
- [23] Fleming, W.H., Soner, H.M. (1993). Controlled Markov Processes and Viscosity Solutions. Applications of Mathematics 25. Springer-Verlag, New York.
- [24] Glasserman, P. (2003). Monte Carlo Methods in Financial Engineering, Stochastic Modelling and Applied Probability, Vol. 53. Springer.

- [25] Ikeda, N., Watanabe, S. (1989). Stochastic Differential Equations and Diffusion Processes, Second Edition. North-Holland Publishing Company.
- [26] Ishii, H. (1984). Uniqueness of unbounded viscosity solutions of Hamilton-Jacobi equations. *Indiana U. Math. J.*. 33, 721–748.
- [27] Karatzas, I., Shreve, S. (1991). Brownian Motion and Stochastic Calculus, Second Edition. Springer-Verlag.
- [28] Krylov, N. (1987). Nonlinear elliptic and Parabolic Partial Differential Equations of Second Order. Mathematics and its Applications, Reider.
- [29] Ladyzenskaya, O.A., Solonnikov, V.A., Uralseva, N.N. (1967). Linear and Quasilinear Equations of Parabolic Type, AMS, Providence.
- [30] Lions, P.L., Regnier, H. (2001). Calcul du prix et des sensibilités d'une option américaine par une méthode de Monte Carlo. Preprint.
- [31] Ma J., Protter, P., Yong, J. (1994). Solving backward stochastic differential equations explicitly A four step scheme. *Prob. Theory Rel. Fields* 98, 339–359.
- [32] Ma, J., Yong, J. (1999). Forward-Backward Stochastic Differential Equations and their Applications. LNM 1702, Springer-Verlag.
- [33] Ma, J., Protter, P., San Martin, J., Torres, S. (2002). Numerical methods for backward stochastic differential equations, *Ann. Appl. Probab.* 12, 302–316.
- [34] Nualart, D. (1995). The Malliavin Calculus and Related Topics. Springer-Verlag.
- [35] Pardoux, E., Peng, S. (1990). Adapted solution of a backward stochastic differential equation, Systems Control Lett., 14, 55–61.
- [36] Pardoux, E., Peng, S. (1992a). Backward stochastic differential equations and quasilinear parabolic partial differential equations. Lecture Notes in CIS, Vol. 176. Springer-Verlag, 200–217.
- [37] Pardoux, E., Peng, S. (1994). Backward doubly stochastic differential equations and systems of quasilinear parabolic SPDEs, *Probab. Theory Rel. Fields* 98, 209–227.
- [38] Pardoux, E., Tang, S. (1999). Forward-backward stochastic differential equations and quasilinear parabolic PDEs. *Prob. Th. Rel. Fields* 114(2), 123–150.
- [39] Peng, S. (1990). A general stochastic maximum principle for optimal control problems, SIAM J. Control Optim. 28, 966–979.
- [40] Peng, S. (1991). Probabilistic interpretation for systems of quasilinear parabolic partial differential equations, *Stochastics* 37, 61–74.
- [41] Peng, S. (1992a) Stochastic Hamilton-Jacobi-Bellman equations, SIAM J. Control Optim. 30, 284–304.

- [42] Peng, S. (1992b). A generalized dynamic programming principle and Hamilton–Jacobi–Bellman equation, *Stochastics* 38, 119–134.
- [43] Peng, S. (1992c). A nonlinear Feynman–Kac formula and applications, in Proceedings of Symposium of System Sciences and Control Theory, Ed. Chen and Yong. Singapore: World Scientific, 173–184.
- [44] Peng, S. (1993). Backward stochastic differential equation and its application in optimal control, Appl. Math. Optim. 27, 125–144.
- [45] Soner, H.M., Touzi, N. (2002). Stochastic target problems, dynamic programming and viscosity solutions, SIAM Journal on Control and Optimization 41, 404–424.
- [46] Talay, D. (1996). Probabilistic numerical methods for partial differential equations: elements of analysis, in *Probabilistic Models for Nonlinear Partial Differential Equa*tions, D. Talay and L. Tubaro, editors, Lecture Notes in Mathematics 1627, 48–196, 1996.
- [47] Zhang, J. (2001). Some fine properties of backward stochastic differential equations. PhD thesis, Purdue University.