VARIATIONAL APPROACH TO NONLINEAR GRAVITY-DRIVEN INSTABILITIES IN A MHD SETTING

HYUNG JU HWANG

ABSTRACT. We establish a variational framework for nonlinear instabilities in a setting of the ideal magnetohydrodynamic (MHD) equations. We apply a variational method to various kind of smooth steady states which are shown to be nonlinearly unstable for both incompressible and compressible ideal MHD equations. Destabilizing effect of compressibility is justified as well as stabilizing effect of magnetic field lines arising in MHD dynamics, which distinguishes from the Rayleigh-Taylor instability in the absence of magnetic field lines.

1. INTRODUCTION

Magnetohydrodynamic equations (MHD) serve as an important model for fluid and gas dynamics and hydromagnetic instability is a fundamental phenomenon in nature, for instance, oceans, atmosphere, and plasma. MHD instability possesses extensive applications in both laboratory plasmas and astrophysics such as in nuclear fusions, compression of thin foils for Xray production, and stellar dynamics. Nevertheless, there have been not many analytical results to date due to its structural complexity such as the presence of shock waves. Furthermore, nothing has been known about nonlinear instabilities for MHD equations despite its importance and variety of instabilities.

The main purpose of this article is to present a variational framework in the passage from linear to nonlinear instability in a setting of the ideal MHD system and derive nonlinear instability around different steady states for both incompressible and compressible ideal MHD equations.

We consider the equations of ideal magnetohydrodynamics(MHD) for inviscid flows:

$$\rho_t + \nabla \cdot (\rho V) = 0,$$

$$\rho (V_t + V \cdot \nabla V) = (\nabla \times B) \times B - \nabla P + \rho \vec{g},$$

$$B_t = \nabla \times (V \times B),$$

$$\nabla \cdot B = 0.$$

(1.1)

Here ρ is plasma density, V velocity, B magnetic field, P plasma pressure, and \vec{g} is the gravitational field. The x axis is taken along the gravitational field, which is assumed to be uniform:

$$\vec{g} = (g, 0, 0)$$
.

The condition for steady state (ρ_0, B_0, P_0) with $V_0 \equiv 0$ is then

$$\frac{d}{dx}\left(P_0 + \frac{1}{2}B_0^2\right) = \rho_0 g. \tag{1.2}$$

We assume for density profile that

$$\min_{x} \rho_0 \ge c > 0. \tag{1.3}$$

Our domain is $t \ge 0$ and

$$D = \{ 0 \le x \le 2\pi, 0 \le y \le 2\pi \}.$$

We assume periodic conditions at the boundary for V and B. Both incompressible and compressible fluids are considered: In the incompressible case,

$$\nabla \cdot V = 0$$

In the compressible case, γ is an adiabatic index which relates the pressure P to the density ρ by

$$P(\rho) = C\rho^{\gamma}, \qquad (1.4)$$
$$\frac{\nabla P}{\rho} = C\gamma\rho^{\gamma-2}\nabla\rho =: q(\rho)\nabla\rho.$$

Or equivalently

$$p_t + v \cdot \nabla p_0 + \gamma p_0 \nabla \cdot v = 0 \tag{1.5}$$

We now consider perturbations (σ, v, B, p) around such a steady state $(\rho_0, B_0, V_0 \equiv 0, P_0)$ of the form:

$$v_{1} = v_{1}(t, x, y), \quad v_{2} = v_{2}(t, x, y), \quad v_{3} = v_{3}(t, x, y),$$

$$B_{1} = B_{1}(t, x, y), \quad B_{2} = B_{2}(t, x, y), \quad B_{3} = B_{3}(t, x, y).$$

Equations for perturbed quantities take the form:

$$\sigma_t + \nabla \cdot [(\rho_0 + \sigma) v] = 0,$$

$$(\rho_0 + \sigma) (v_t + v \cdot \nabla v) = (\nabla \times B_0) \times B + (\nabla \times B) \times B_0$$

$$+ (\nabla \times B) \times B - \nabla p + \sigma \vec{g},$$

$$B_t = \nabla \times (v \times (B_0 + B)),$$

$$\nabla \cdot B = 0.$$
(1.6)

We also obtain its linearized system:

$$\sigma_t + \nabla \cdot (\rho_0 v) = 0, \qquad (1.7)$$

$$\rho_0 v_t = (\nabla \times B_0) \times B + (\nabla \times B) \times B_0 - \nabla p + \sigma \vec{g},$$

$$B_t = \nabla \times (v \times B_0),$$

$$\nabla \cdot B = 0.$$

In the incompressible case, v also satisfies, in both linear and nonlinear system,

$$\nabla \cdot v = 0.$$

It is crucial to make an equivalent second-order linearized system in one quantity v and use it through our variational formulation: By taking t-derivative of v-equation in (1.7) and plugging σ - and B-equation in (1.7) (also (1.5) in the compressible case) into the resulting equation, we obtain the following second-order linear hyperbolic PDE for velocity v:

$$\rho_0 v_{tt} = (Q \cdot \nabla) B_0 + (B_0 \cdot \nabla) Q - \nabla (B_0 \cdot Q + p_t) - \vec{g} \nabla \cdot (\rho_0 v) =: L(v), \qquad (1.8)$$

where $Q = \nabla \times (v \times B_0)$ and we have used (1.11).

For notational convenience,

Notation 1. For any u and $v \in L^2(0, 2\pi)$,

$$< u, v > = \int_0^{2\pi} u \cdot v \, dx, \ (u, v) = \int_0^{2\pi} \rho_0 u \cdot v \, dx,$$

For any u and $v \in L^2(D)$,

$$\langle u, v \rangle = \iint_D u \cdot v \, dx dy, \ (u, v) = \iint_D \rho_0 u \cdot v \, dx dy.$$

Notation 2. $||f|| = (f, f)^{1/2}$, $||f||_s = \left(\sum_{\alpha} ||\partial^{\alpha} f||^2\right)^{1/2}$, where α is a multi-index with $|\alpha| \leq s$.

We make the following variational formulation and obtain a discrete set of eigenvalues for the linear operator (1.8): for each wave number $k \in \mathbb{N}$,

$$\lambda_k^2 = \sup_{\substack{u = (u_1, u_2)\\u_1 \in H^{\kappa}(0, 2\pi),\\u_2 \in H^{\kappa-1}(0, 2\pi)}} \frac{\langle L(u), u \rangle}{(u, u)},$$
(1.9)

where ∂_y is replaced with multiplying by k in L(u) and $\kappa = 1$ or 2. Notice that the RHS of (1.9) is indeed a function of k.

In the incompressible case, u_2 is replaced by $-\frac{1}{k}u_{1x}$ from divergence-free condition for the normal modes and (1.9) reduces to a variational problem for u_1 alone. On the other hand, the compressible case may not be simplified to a formula for u_1 alone.

Key step of this article is to show that this discrete set $\{\lambda_k\}_{k\in\mathbb{N}}$ of eigenvalues characterizes the continuum spectral radius of the whole linearized operator by taking the limit as $k \to \infty$ to obtain the bounded least upper bound $\Lambda > 0$: Let

$$\lim_{k} \lambda_k^2 = \Lambda^2 = \sup_{v \in H^{\kappa-1}(D)} \frac{\iint_D H(v, v_x) \, dx dy}{(v, v)},\tag{1.10}$$

where coefficients of H depend on ρ_0, p_0, B_0 . Then we obtain the following theorem:

Theorem 1. Let (σ, v) be a solution to (1.7) and let $\Lambda^2 > 0$, then we have

$$\|\sigma(t), v(t), B(t)\|_{s} \leq Ce^{\Lambda t} \|\sigma(0), v(0), B(0)\|_{s+2},$$

where $C = C(\Lambda, \rho_0, s)$.

Thanks to this theorem, we can locate a dominant eigenvalue and pass to nonlinear instability.

We establish the following dynamical instability for the fully nonlinear MHD system around different steady states satisfying $\Lambda^2 > 0$ as in Section 4:

Theorem 2. Steady states $(\rho_0, \vec{0}, B_0)$ of (1.1) in (1.2) satisfying $\Lambda^2 > 0$ are indeed nonlinearly unstable: For any s large, there exists $\varepsilon_0 > 0$, such that for any small $\delta > 0$, there exists a family of classical solutions $(\rho^{\delta}(t, x, y), V^{\delta}(t, x, y), B^{\delta}(t, x, y))$ to (1.1) such that

$$\left\|\rho^{\delta}(0,\cdot) - \rho_{0}(\cdot)\right\|_{H^{s}(D)} + \left\|V^{\delta}(0,\cdot)\right\|_{H^{s}(D)} + \left\|B^{\delta}(0,\cdot) - B_{0}(\cdot)\right\|_{H^{s}(D)} \le \delta,$$

but for $T^{\delta} = O(|\ln \delta|),$

$$\sup_{0 \le t \le T^{\delta}} \left\{ \left\| \rho^{\delta}(t, \cdot) - \rho_{0}\left(\cdot\right) \right\|_{L^{1}(D)} + \left\| V^{\delta}(t, \cdot) \right\|_{L^{1}(D)} + \left\| B^{\delta}\left(t, \cdot\right) - B_{0}\left(\cdot\right) \right\|_{L^{1}(D)} \right\} \ge \varepsilon_{0}.$$

Remark 1. This indicates the Kruskal-Schwarzschild instability for incompressible ideal MHD fluids as $\gamma \to \infty$ formally.

Remark 2. The instability time T^{δ} occurs before the possible blow-up time which is shown in the proof and we measure instability in L^1 .

Rayleigh-Taylor instability is well known as gravity-driven instability in fluids when heavy fluid is on top of light one. Linear instability for an incompressible fluid was first introduced by Rayleigh in 1883 [18]. Study on linear instability for incompressible ideal MHD system in the presence of magnetic field lines could be found in [16], [4] for instance, which includes the classical Kruskal-Schwarzschild instability in the presence of a magnetic field orthogonal to the gravitational force. In this case, instability criterion (4.4) and the growth rates (4.2), (4.3) turn out to be the same as the ones in the Rayleigh-Taylor instability as in [10] without effect of magnetic field lines on the instability. On the other hand, when we consider effect of a magnetic field parallel to direction of the gravitational force, stabilizing effect of magnetic field lines appears as in (4.13) and (4.14). In the presence of a vertical magnetic field, the rigidity produced by the magnetic field lines hinders its own way to instability and makes the growth rate slower. Condition for linear instability for a compressible fluid in the absence of a magnetic field was first derived by Schwarzschild in [19], and since been discussed by many other physicists for a certain class of steady states [4], [5], [7], [11]. The full consideration of gravity, magnetic field lines and compressibility has been also largely discussed for its linear instability by many physicists such as in [1], [2], [17], which exhibit very interesting phenomena although it accelerates analytical difficulties.

Despite extensive research and interest in this subject from physical point of view, little has been done from mathematical perspective for the MHD system. In addition, the passage from linear to nonlinear instability in a conservative PDE system is quite difficult because of the following two main obstacles: (1) presence of the continuum linear spectrum and (2) severely unbounded high-order terms in PDE systems. No systematic framework has been built up for this problem although there have been works towards this subject for specific physical systems, for example [9],[3],[10]. Variational approach was first introduced by Guo and Hwang [10] in the case of dynamical Rayleigh-Taylor instability for incompressible Euler fluids. However it

is not obvious whether it can be extended to MHD instabilities for compressible as well as incompressible fluids since MHD has more complicated structure in addition to the analytical difficulties coming from compressibility.

Crucial point is whether and how to locate a dominant eigenvalue in the complex linear spectrum of the MHD system. We use the MHD energy principle in order to estimate a sharp spectral radius and make extensive use of the variational structure of the linearized MHD system, resulting in more precise and optimal estimates. The radius (1.10) of continuum spectrum is obtained as the least upper bound for a discrete set of eigenvalues (1.9) of normal growing modes and this method is explicit and constructive. We consider three different magnetic fields as steady states which give rise to different outcome in the growth rates (4.3), (4.9), (4.14) of instabilities and different ranges of admissible density profiles for instabilities. We justify stabilizing and destabilizing effects of magnetic filed lines and compressibility as expected physically. Furthermore incompressible case can also be viewed as the limiting case of compressible case as $\gamma \to \infty$. The article is organized as follows.

We formulate the variational problems (1.9) in Section 2 and show the existence of smooth maximizers satisfying the corresponding Sturm-Liouville equations. We then derive a sharp growth rate $\Lambda > 0$ for the whole linear system in Section 3 as in the Theorem 1. In Section 4, we give different examples of steady states which result in different growth rates, Sturm-Liouville equations, and different admissibility for density profiles for instability. For instance, if a steady magnetic field is parallel to the gravity, we have the fourth-order Sturm-Liouville equation (4.12). The compressible case (4.8) is more complicated and its variational problem should be treated with more delicacy.

In Section 5, we construct approximate solutions and give energy estimates for the full system. Finally, we present nonlinear instability for the ideal MHD system around our three kind steady states in Section 6.

We put some useful vector identities which are used throughout the paper. For any two vectors \vec{a} and \vec{b} ,

$$\nabla \times \left(\vec{a} \times \vec{b}\right) = \vec{a} \left(\nabla \cdot \vec{b}\right) + \left(\vec{b} \cdot \nabla\right) \vec{a} - \vec{b} \left(\nabla \cdot \vec{a}\right) - \left(\vec{a} \cdot \nabla\right) \vec{b},\tag{1.11}$$

$$\nabla\left(\vec{a}\cdot\vec{b}\right) = \left(\vec{a}\cdot\nabla\right)\vec{b} + \left(\vec{b}\cdot\nabla\right)\vec{a} + \vec{a}\times\left(\nabla\times\vec{b}\right) + \vec{b}\times\left(\nabla\times\vec{a}\right),\tag{1.12}$$

2. General variational framework

We consider the following steady magnetic fields: (Case $B_0 \perp \vec{g}$)

$$B_{0} = (0, 0, B_{0}(x)),$$

(Case $B_0 \parallel \vec{g}$)

$$B_0 = (B_0, 0, 0)$$
.

In the case of $B_0 \parallel \vec{g}$, we assume $v_3 = B_3 = 0$. By integration by parts, using (1.7),(1.5),(1.2) and completing the square with respect to $\nabla \cdot v$, we obtain the following decomposition for

< L(v), v >: for any $v = (v_1, v_2, v_3),$

$$< L(v), v >= \iint_{D} L(v) \cdot v \, dx dy$$

$$= \iint_{D} \left[-F(v_{2x}) - G(\nabla \cdot v, v_{1}, v_{1x}) + H(v_{1}, v_{1x}) \right] dx dy.$$
(2.1)

We state some important properties which are satisfied by the above functional:

(1) L is variational, i.e., for any u and v

$$< L(u), v > = < u, L(v) > .$$

(2) < L(v), v > is concave with respect to v_{1x}, v_{2x}, v_2 respectively. (3) $< L(v), v > \to -\infty$ as $v_{1x} \to \infty$ and $< L(v), v > \to -\infty$ as $v_{2x} \to \infty$ if $F \neq 0$. (4) F > 0, G > 0, H(0) = 0 when $H = H(v_1)$ alone, and F, G, H are all quadratic.

Remark 3. G = 0 in the incompressible case.

Remark 4. For the case $B_0 \perp \vec{g}$, we have F = 0, $H = H(v_1)$, and $G = G(\nabla \cdot v, v_1)$ with b = 0 while we have $F = F(v_{2x})$, $H = H(v_1, v_{1x})$, $G = G(\nabla \cdot v, v_1, v_{1x})$ in the case $B_0 \parallel \vec{g}$ as we can see in Section 4.

We will show that for any fixed wave number $k \in \mathbb{N}$, the corresponding eigenvalue $\lambda_k > 0$ for the linearized MHD system takes the variational formulation closely related to the above variational structure. A normal mode is of the form:

$$v_1(t, x, y) = \tilde{v}_1(x) \cos(ky) \exp(\lambda_k t), \qquad (2.2)$$

$$v_2(t, x, y) = \tilde{v}_2(x) \sin(ky) \exp(\lambda_k t), \qquad (2.2)$$

$$v_3(t, x, y) = \tilde{v}_3(x) \cos(ky) \exp(\lambda_k t),$$

where k is a wave number. Substituting (2.2) into (1.8) yields the following second-order ODE for \tilde{v} :

$$\lambda_k^2 \rho_0 \left(\tilde{v}_1, \tilde{v}_2, \tilde{v}_3 \right) = L \left(\tilde{v}_1, \tilde{v}_2, \tilde{v}_3 \right).$$
(2.3)

Note that $L_3(\tilde{v}_1, \tilde{v}_2, \tilde{v}_3) = 0$ and thus $\tilde{v}_3 = 0$. We now make the following variational formulations:

$$\lambda_k^2 = \sup_{\substack{u = (u_1, u_2)\\u_1 \in H^{\kappa}(0, 2\pi)\\u_2 \in H^{\kappa-1}(0, 2\pi)}} \frac{\langle L(u), u \rangle}{(u, u)},$$
(2.4)

where $\kappa = 1$ or 2, ∂_y is replaced with multiplying by k in L(u) and integrations here are with respect to x over $(0, 2\pi)$. Indeed,

Incompressible case: we use divergence-free condition to reduce to

$$\lambda_k^2 = \sup_{u \in H^{\kappa}(0,2\pi)} \frac{\int_0^{2\pi} \left[-F\left(-\frac{u_{xx}}{k}\right) + H\left(u, u_x\right) \right] dx}{\int_0^{2\pi} \left[u^2 + \frac{u_x^2}{k^2} \right] dx},$$

where $\kappa = 1$ or 2.

Compressible case:

$$\lambda_k^2 = \sup_{\substack{u_1 \in H^{\kappa}(0,2\pi)\\u_2 \in H^{\kappa-1}(0,2\pi)}} \frac{\int_0^{2\pi} \left[-F\left(u_{2x}\right) - G\left(u_{1x} + ku_2, u_1, u_{1x}\right) + H\left(u_1, u_{1x}\right)\right] dx}{\int_0^{2\pi} \left[u_1^2 + u_2^2\right] dx},$$

where $\kappa = 1$ or 2.

Let

$$\Lambda^2 = \sup_{v \in H^{\kappa-1}(D)} \frac{\iint_D H(v, v_x) \, dx dy}{(v, v)},\tag{2.5}$$

where $\kappa = 1$ or 2. Then we show Λ^2 is the least upper bound for $\{\lambda_k^2\}_{k \in \mathbb{N}}$.

Lemma 1.

$$\lim_{k \to \infty} \lambda_k^2 = \Lambda^2.$$

Proof. Since F(0) = 0, it is easy to see that our Lemma is true for incompressible case by letting $k \to \infty$. We now treat compressible case.

Note that, with the choice of $u_2 = -\frac{1}{k}(au_1 + bu_{1x})$, we have

$$\begin{split} \lambda_k^2 &= \sup_{\substack{u = (u_1, u_2) \\ u_1 \in H^{\kappa}(0, 2\pi) \\ u_2 \in H^{\kappa-1}(0, 2\pi)}} \frac{\langle L(u), u \rangle}{(u, u)} \\ &\geq \sup_{u_1 \in H^{\kappa}(0, 2\pi)} \frac{\int_0^{2\pi} \left[-F\left(-\frac{1}{k} \left(au_1 + bu_{1x}\right)_x\right) + H\left(u_1, u_{1x}\right) \right] dx}{\int_0^{2\pi} \rho_0 \left[u_1^2 + \frac{1}{k^2} \left(au_1 + bu_{1x}\right)^2\right] dx} \end{split}$$

Thus, we have the following inequality

$$\sup_{u_{1}\in H^{\kappa}(0,2\pi)}\frac{\int_{0}^{2\pi}\left[-F\left(-\frac{1}{k}\left(au_{1}+bu_{1x}\right)_{x}\right)+H\left(u_{1},u_{1x}\right)\right]dx}{\int_{0}^{2\pi}\rho_{0}\left[u_{1}^{2}+\frac{1}{k^{2}}\left(au_{1}+bu_{1x}\right)^{2}\right]dx}\leq\lambda_{k}^{2}\leq$$

$$\sup_{\substack{u_{1}\in H^{\kappa}(0,2\pi)\\u_{2}\in H^{\kappa-1}(0,2\pi)}}\frac{\int_{0}^{2\pi}\left[-F\left(u_{2x}\right)-G\left(u_{1x}+ku_{2},u_{1},u_{1x}\right)\right]dx}{\int_{0}^{2\pi}\rho_{0}\left[u_{1}^{2}+u_{2}^{2}\right]dx}+\sup_{\substack{u_{1}\in H^{\kappa}(0,2\pi)\\u_{2}\in H^{\kappa-1}(0,2\pi)}}\frac{\int_{0}^{2\pi}H\left(u_{1},u_{1x}\right)dx}{\int_{0}^{2\pi}\rho_{0}u_{1}^{2}dx}.$$

$$\leq \sup_{u_{1}\in H^{\kappa-1}(0,2\pi)}\frac{\int_{0}^{2\pi}H\left(u_{1},u_{1x}\right)dx}{\int_{0}^{2\pi}\rho_{0}u_{1}^{2}dx}.$$

Thanks to

$$\frac{1}{k}(au_1 + bu_{1x}) \to 0 \text{ as } k \to \infty, \ F(0) = 0,$$

letting $k \to \infty$ yields

$$\lim_{k \to \infty} \lambda_k^2 = \sup_{u_1 \in H^{\kappa - 1}(0, 2\pi)} \frac{\int_0^{2\pi} H(u_1, u_{1x}) \, dx}{\int_0^{2\pi} \rho_0 u_1^2 dx},$$

and

$$\lim_{k \to \infty} \lambda_k^2 = \sup_{u_1 \in H^{\kappa - 1}(D)} \frac{\iint_D H(u_1, u_{1x}) \, dx dy}{\iint_D \rho_0 u_1^2 dx dy},$$

since H^{κ} is dense in $H^{\kappa-1}$. Thus the proof is complete.

We now show the existence of maximizer for the variational problem (2.4). Assume that

$$(u,u) = \int_0^{2\pi} \rho_0 \left[u_1^2 + u_2^2 \right] \, dx = 1.$$
(2.6)

For fixed k, let

$$\lambda_k^2 = \sup_{\substack{u_1 \in H^{\kappa}(0,2\pi) \\ u_2 \in H^{\kappa-1}(0,2\pi)}} < L(u), u > .$$
(2.7)

Lemma 2. For any fixed k, there exists a smooth maximizer for the variational problem (2.7) with the constraint (2.6).

Proof. Let $\{u_1^n, u_2^n\}$ be a maximizing sequence with the constraint (2.6). Then u_1^n and u_2^n converge weakly in $L^2(0, 2\pi)$ to u_1^0 and u_2^0 respectively and we have

$$\langle L(u^n), u^n \rangle \rightarrow \lambda_k^2.$$
 (2.8)

Case 1 $B_0 \perp \vec{g}$, where $F = 0, H = H(u_1)$:

Since $\langle L(u), u \rangle \rightarrow -\infty$ as $u_{1x} \rightarrow \infty$ (Property 3 of the functional), u_{1x} is bounded in $L^2(0, 2\pi)$ uniformly in *n*. Thus there exists a weak limit $\{u_1^0, u_2^0\}$ such that

$$\begin{array}{rcl} u_{1x}^n & \rightharpoonup & u_{1x}^0 \text{ weakly in } L^2\left(0,2\pi\right), \ u_{2x}^n \rightharpoonup & u_2^0 \text{ weakly in } L^2\left(0,2\pi\right) \\ u_1^n & \to & u_1^0 \text{ strongly in } L^2\left(0,2\pi\right) \text{ and } u_1^0 \in H^1\left(0,2\pi\right). \end{array}$$

Next we show that $\{u_1^0, u_2^0\}$ is a maximizer and satisfies the constraint (2.6). Since $\{u_1^0, u_2^0\}$ is a weak limit of $\{u_1^n, u_2^n\}$, $(u^0, u^0) = \int_0^{2\pi} \rho_0 \left[(u_1^0)^2 + (u_2^0)^2 \right] dx \leq 1$ by lower semi-continuity of L^2 :

$$\int_{0}^{2\pi} \rho_0 \left[(u_1^n)^2 + (u_2^n)^2 \right] dx - \int_{0}^{2\pi} \rho_0 \left[\left(u_1^0 \right)^2 + \left(u_2^0 \right)^2 \right] dx$$

=
$$\int_{0}^{2\pi} 2\rho_0 \left[u_1^0 \left(u_1^n - u_1^0 \right) + u_2^0 \left(u_2^n - u_2^0 \right) \right] dx$$
$$+ \int_{0}^{2\pi} \rho_0 \left[\left(u_1^n - u_1^0 \right)^2 + \left(u_2^n - u_2^0 \right)^2 \right]$$

\geq
$$\int_{0}^{2\pi} 2\rho_0 \left[u_1^0 \left(u_1^n - u_1^0 \right) + u_2^0 \left(u_2^n - u_2^0 \right) \right] dx \to 0 \text{ as } n \to 0.$$

due to the fact that $\{u_1^0, u_2^0\}$ is a weak limit of $\{u_1^n, u_2^n\}$ in $L^2(0, 2\pi)$. We use concavity of the functional $\langle L(u), u \rangle$ with respect to u_{1x}, u_{2x}, u_2 (Property 2 of the functional) and the strong convergence of u_1^n to u_1^0 in $L^2(0, 2\pi)$ to deduce

$$< L\left(u^{0}\right), u^{0} > \ge \lambda_{k}^{2}.$$

$$(2.9)$$

Let

$$J(u_{2x}, u_{1x}, u_2, u_1) = < L(u), u > .$$

Then by concavity of this functional J, strong convergence of u_1^n to u_1^0 and weak convergence of u_{1x}^n, u_2^n, u_1^n to u_{1x}^0, u_2^0, u_1^0 respectively, we have

$$J\left(u_{2x}^{n}, u_{1x}^{n}, u_{2}^{n}, u_{1}^{n}\right) - J\left(u_{2x}^{0}, u_{1x}^{0}, u_{2}^{0}, u_{1}^{0}\right)$$

$$= J\left(u_{2x}^{n}, u_{1x}^{n}, u_{2}^{n}, u_{1}^{n}\right) - J\left(u_{2x}^{n}, u_{1x}^{n}, u_{2}^{n}, u_{1}^{0}\right) + J\left(u_{2x}^{n}, u_{1x}^{n}, u_{2}^{n}, u_{1}^{0}\right) - J\left(u_{2x}^{0}, u_{1x}^{0}, u_{2}^{0}, u_{1}^{0}\right)$$

$$\leq J\left(u_{2x}^{n}, u_{1x}^{n}, u_{2}^{n}, u_{1}^{n}\right) - J\left(u_{2x}^{n}, u_{1x}^{n}, u_{2}^{n}, u_{1}^{0}\right) + \nabla J\left(u_{2x}^{0}, u_{1x}^{0}, u_{2}^{0}, u_{1}^{0}\right) \cdot \left(u_{2x}^{n} - u_{2x}^{0}, u_{1x}^{n} - u_{1x}^{0}, u_{2}^{n} - u_{2}^{0}, 0\right)$$

$$\rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus we obtain (2.9). Notice that we can not have $u^0 = \{u_1^0, u_2^0\} = \{0, 0\}$ a.e. since, by strong convergence of u_1^n to u_1^0 and by (2.8), we have

$$\int_{0}^{2\pi} H(u_{1}^{0}) dx = \lim_{n \to \infty} \int_{0}^{2\pi} H(u_{1}^{n}) dx \ge \lambda_{k}^{2} > 0$$

where we have used the property 4 of the functional. Suppose now that $(u^0, u^0) = \alpha^2 < 1$. Then let $(\tilde{u}_1^0, \tilde{u}_2^0) = \frac{1}{\alpha} (u_1^0, u_2^0)$ so that $(\tilde{u}^0, \tilde{u}^0) = 1$. By the above argument (2.9), we have

$$< L\left(\tilde{u}^{0}\right), \tilde{u}^{0} > \geq \frac{\lambda_{k}^{2}}{\alpha} > \lambda_{k}^{2},$$

leading to a contradiction. Thus $\{u_1^0, u_2^0\}$ is a maximizer satisfying the constraint (2.6).

Case 2 $B_0 \parallel \vec{g}$, where $F(u_{2x}) = B_0^2 u_{2x}^2$: In this case, by property 3 of the functional, u_{2x}^n is also bounded in $L^2(0, 2\pi)$ uniformly in n and thus both u_1^n and u_2^n converge strongly in $L^2(0, 2\pi)$ to u_1^0 and u_2^0 respectively. Then we have

$$(u^0, u^0) = \int_0^{2\pi} \rho_0 \left[\left(u_1^0 \right)^2 + \left(u_2^0 \right)^2 \right] dx = 1.$$

In a similar manner, we obtain (2.9) and hence u^0 is a maximizer.

We finally show such a maximizer satisfies the generalized Sturm-Liouville equation (2.3) for both cases. For $\tau \in \mathbb{R}$ and $w = \{w_1, w_2\} \in H^{\kappa}(0, 2\pi) \times H^{\kappa-1}(0, 2\pi)$, define $u(\tau) = u^0 + \tau w$, then by (4.8), we have

$$< L(u(\tau)), u(\tau) > \leq \lambda_k^2 (u(\tau), u(\tau)).$$

Set

$$I(\tau) = \langle L(u(\tau)), u(\tau) \rangle - \lambda_k^2 (u(\tau), u(\tau)),$$

then we have $I(\tau) \leq 0$ for all $\tau \in \mathbb{R}$ and I(0) = 0. This implies

$$I'(0) = \langle L(w), u^{0} \rangle + \langle L(u^{0}), w \rangle - 2\lambda_{k}^{2}(u^{0}, w)$$

= 2 \le L(u^{0}) - \lambda_{k}^{2}\rho_{0}u^{0}, w \ge = 0 for all w,

since L is variational (Property 1 of the functional). Thus u^0 satisfies a normal mode, i.e.,

$$\lambda_k^2 \rho_0 u^0 = L\left(u^0\right)$$

Since ρ_0 , p_0 , and B_0 are smooth, u^0 is also smooth. This completes the proof.

3. Linear growth rate Λ

In this section, we show that Λ is the optimal growth rate for the linearized system and it serves as the spectral radius of the linear operator. We state a global existence of solutions to the linearized system, which can be obtained by a straightforward method.

Lemma 3. There exists a global in time solution $(\sigma, v, B) \in C([0, T]; H^s(D))$ to the linearized MHD system (1.7).

Let (σ, v) be a solution to (1.7) and let $\Lambda^2 > 0$, then we have

Theorem 3.

$$\left\|\sigma\left(t\right),v\left(t\right),B\left(t\right)\right\|_{s} \leq Ce^{\Lambda t} \left\|\sigma\left(0\right),v\left(0\right),B\left(0\right)\right\|_{s+2},$$

where $C = C(\Lambda, \rho_0, s)$ and $\Lambda > 0$.

Proof. We show by induction on \bar{s} , the number of x-derivatives. We first treat the case $\bar{s} = 0$: Multiply (4.6) by v_t and integrate over x and y, then we have

$$\frac{d}{dt}\left(v_{t}, v_{t}\right) = \frac{d}{dt} < L\left(v\right), v >$$

$$(3.1)$$

since L is variational. Notice that using (2.5) yields

$$\iint_{D} H(v_1, v_{1x}) \, dx dy \leq \Lambda^2(v_1, v_1) \leq \Lambda^2(v, v) \, .$$

By integrating (3.1) over time and by (2.1), we obtain

$$(v_{t}, v_{t}) + \iint_{D} [F(v_{2x}) + G(\nabla \cdot v, v_{1}, v_{1x})] dxdy$$

$$\leq \iint_{D} H(v_{1}, v_{1x}) dxdy + \|\sigma(0), v(0), B(0)\|_{1}^{2}.$$

$$\leq \Lambda^{2}(v, v) + \|\sigma(0), v(0), B(0)\|_{1}^{2}, \qquad (3.2)$$

and

$$\frac{d}{dt} \|v\| \le \|v_t\| \le \|\sigma(0), v(0), B(0)\|_1 + \Lambda \|v\|.$$

Thus, we have

 $\|v\| \le C e^{\Lambda t}.$

where $C = \|\sigma(0), v(0), B(0)\|_1$ and from now on, we will just use universal constant C which varies, only for notational convenience. Notice that these estimates exactly apply to the *t*- and *y*-derivatives of any order of v and σ since the variational structure of (1.8) is not destroyed by taking *t*- and *y*-derivatives.

Case 1. $B_0 \perp \vec{g}$ where F = 0, $G = (\gamma p_0 + B_0^2) \left(\nabla \cdot v + \frac{g \rho_0}{\gamma p_0 + B_0^2} v_1 \right)^2$ (G = 0 in the incompressible case), $H = H(v_1)$, $(Q \cdot \nabla) B_0 + (B_0 \cdot \nabla) Q = 0$:

By (3.2) and (1.7), we also have

$$\|\nabla \cdot v\| \le Ce^{\Lambda t}, \|\sigma\| \le Ce^{\Lambda t}, \|B\| \le Ce^{\Lambda t}.$$

Next, we consider the case $\bar{s} = 1$. By taking curl v-equation of (1.7), we have, with $\omega = \nabla \times v$,

$$\rho_0\left(\omega\right)_{tt} = -\nabla\rho_0 \times v_{tt} - g\sigma_{ty}\hat{k},\tag{3.3}$$

where \hat{k} is the unit vector in the z-direction. Since σ_{ty} and v_{tt} have no x-derivatives and $B_{1x} = -B_{2y}$ and so have the growth rate Λ as in the previous step, we have

$$\|\omega\| \le C e^{\Lambda t},$$

where $C = \|\sigma(0), v(0), B(0)\|_3$. Thanks to the identity $\Delta \zeta = -\nabla \times (\nabla \times \zeta) + \nabla (\nabla \cdot \zeta)$ for any $\zeta = (\zeta_1, \zeta_2)$, we conclude that all the first derivatives of v have the same growth rate Λ . Now for $\nabla \sigma$, we use the vector identity (1.11) to get

$$B_t = \nabla \left(v \times B_0 \right) = -B_{0x} v_1 \hat{k} - \hat{k} B_0 \nabla \cdot v.$$
(3.4)

Plugging (3.4) and (1.5) into (1.8) yields

$$\rho_0 v_{tt} = \nabla \left(B_0 B_{0x} v_1 \right) + \nabla \left(\left[\gamma p_0 + B_0^2 \right] \nabla \cdot v \right) + \sigma_t \vec{g}.$$

$$(3.5)$$

By induction hypotheses and σ - and B-equations in (1.7), we deduce

$$\|\nabla (\nabla \cdot v)\| \le Ce^{\Lambda t}, \|\nabla \sigma\| \le Ce^{\Lambda t}, \|\nabla B\| \le Ce^{\Lambda t}, \|\nabla B$$

For higher derivatives when $\bar{s} \geq 2$, we use induction hypotheses. Suppose all the derivatives of order $< \bar{s}$ have the growth rate Λ . Let α be multi-index whose order of x-derivative is $\bar{s} - 1$. Then, by taking ∂_{α} of the curl of (4.6), curl $(\partial_{\alpha} v)$ satisfies

$$\rho_0 \left(\partial_\alpha \omega\right)_{tt} = -\partial_a \left(\nabla \times (\rho_0 v)\right)_{tt} + \rho_0 \left(\partial_\alpha \omega\right)_{tt} - g\partial_\alpha \sigma_{ty} \hat{k}.$$
(3.6)

The right hand side of (3.6) contains derivatives of v and σ whose x-order $\langle \bar{s} \rangle$. Clearly the x-order of $\partial_{\alpha}\sigma_{ty}$ is still $\bar{s} - 1$. The first and the second term of the RHS together are of the form

$$\sum_{\substack{|\beta| \ge 1\\\beta+\gamma=\alpha}} C_{\beta,\gamma} \nabla \times \left[(\partial_{\beta} \rho_0) \left(\partial_{\gamma} v_{tt} \right) \right].$$
(3.7)

Notice that only purely x-derivatives $\partial_{\beta}\rho_0$ with $|\beta| \ge 1$ in (3.7) remain nonzero since $\rho_0(x)$ depends only on x. Then (3.6) yields

$$\|\nabla \times \partial_a v\| \le C e^{\Lambda t}$$

For $\nabla \cdot \partial_{\alpha} v$, we take ∂_{β} of the first component of (3.5) with $\partial_{\beta} \partial_x = \partial_{\alpha}$ to get

$$\partial_{\beta} \left(\rho_0 v_{1tt} \right) = \partial_{\alpha} \left(B_0 B_{0x} v_1 \right) + \partial_{\alpha} \left(\left[\gamma p_0 + B_0^2 \right] \nabla \cdot v \right) + \partial_{\beta} \sigma_t g$$

Using the induction hypotheses as for the case $\bar{s} = 1$, we deduce

$$\|v\|_{\bar{s}} \le Ce^{\Lambda t}, \|\sigma\|_{\bar{s}} \le Ce^{\Lambda t}, \|B\|_{\bar{s}} \le Ce^{\Lambda t},$$

where $C = \|\sigma(0), v(0), B(0)\|_{s+2}$. In the incompressible case, we use (3.3) together with the $\nabla \cdot v = 0$ to deduce the Lemma.

Case 2. $B_0 \parallel \vec{g}$ and incompressible case where $F = B_0^2 v_{2x}^2, G = 0, H = H(v_1, v_{1x})$:

From (3.2) and div v = 0 and using the induction hypotheses, we obtain

 $\|v_{2x}\| \le Ce^{\Lambda t}, \|v_{1x}\| \le Ce^{\Lambda t}.$

In this case, by (1.11), we have $Q = B_0 v_x$ and $B_0 \cdot Q = B_0^2 v_{1x}$. Then (1.8) yields

$$\rho_0 v_{tt} = B_0^2 v_{xx} - \nabla \left(B_0 \cdot Q + p_t \right) + \sigma_t \vec{g}.$$
(3.8)

Taking the curl of (3.8) yields

$$\rho_0 \omega_{tt} = B_0^2 \omega_{xx} - \nabla \rho_0 \times v_{tt} - g \sigma_{ty} \hat{k}.$$

Using the induction hypotheses and div v = 0, we can deduce the Lemma. Therefore the proof is complete.

4. Examples

In this section, we present examples of steady states which give rise to different results.

(1) Magnetic field is transverse to the gravity $B_0 \perp \vec{g}$ for incompressible fluids:

$$B_0 = (0, 0, B_0(x)). \tag{4.1}$$

Linear operator L reads

$$L(v) := -\nabla \left(p_t + B_0 \cdot Q \right) - \vec{g} \rho_{0x} v_1$$

Sturm-Liouville equation is

$$\rho_0 \lambda_k^2 \tilde{v}_1 = \frac{\lambda_k^2}{k^2} \left(\rho_0 \tilde{v}_{1x} \right)_x - g \rho_{0x} \tilde{v}_1,$$

Variational formulation for λ_k^2 is

$$\lambda_k^2 = \sup_{u \in H^1(0,2\pi)} \frac{\int_0^{2\pi} L(u) \cdot u \, dx}{\int_0^{2\pi} \rho_0 \left[u^2 + \frac{u_x^2}{k^2} \right] \, dx} > 0, \tag{4.2}$$

where

$$L\left(u\right)\cdot u = H\left(u\right) = -g\rho_{0x}u^{2}$$

Spectral radius $\Lambda > 0$ is given by

$$\Lambda^2 = \sup_{v \in L^2(D)} \frac{\iint_D H(v) \, dxdy}{\iint_D \rho_0 v^2 \, dxdy} > 0.$$

$$(4.3)$$

The sufficient condition for instability in the case (1) is given by

$$\rho_{0x}(x_0) < 0 \quad \text{for some } x_0 \in (0, 2\pi),$$
(4.4)

(2) Magnetic field is transverse to the gravity $B_0 \perp \vec{g}$ for compressible fluids:

$$B_0 = (0, 0, B_0(x)).$$
(4.5)

Linear operator L, Sturm-Liouville equation, and λ_k are as follows:

$$L(v) = -\nabla (B_0 \cdot Q) - \nabla (p_{0x}v_1 + \gamma p_0 \nabla \cdot v) - \vec{g} (\rho_{0x}v_1 + \rho_0 \nabla \cdot v)$$
(4.6)

$$\lambda_{k}^{2}\rho_{0}\left(\tilde{v}_{1},\tilde{v}_{2},\tilde{v}_{3}\right) = L\left(\tilde{v}_{1},\tilde{v}_{2},\tilde{v}_{3}\right), \qquad (4.7)$$
where $L = (L^{1},L^{2},L^{3})$ is a linear functional in $\tilde{v} = (\tilde{v}_{1},\tilde{v}_{2},\tilde{v}_{3})$ such that
$$L^{1} = \left(g\rho_{0}\tilde{v}_{1} + \left(\gamma p_{0} + B_{0}^{2}\right)\left(\tilde{v}_{1x} + k\tilde{v}_{2}\right)\right)_{x} - g\left(\rho_{0x}\tilde{v}_{1} + \rho_{0}\left(\tilde{v}_{1x} + k\tilde{v}_{2}\right)\right), \\ L^{2} = -k\left(g\rho_{0}\tilde{v}_{1} + \left(\gamma p_{0} + B_{0}^{2}\right)\left(\tilde{v}_{1x} + k\tilde{v}_{2}\right)\right), \\ L^{3} = 0,$$

and so $\tilde{v}_3 = 0$.

$$\lambda_k^2 = \sup_{\substack{u = (u_1, u_2)\\u_1 \in H^1(0, 2\pi),\\u_2 \in L^2(0, 2\pi)}} \frac{\int_0^{2\pi} L(u) \cdot u \, dx}{\int_0^{2\pi} \rho_0 \left[u_1^2 + u_2^2\right] \, dx} > 0, \tag{4.8}$$

where

$$L(u) \cdot u = -G(u_1, u_{1x}, u_2) + H(u_1)$$

$$G(u_1, u_{1x}, u_2) = -(\gamma p_0 + B_0^2) \left(u_{1x} + ku_2 + \frac{g\rho_0}{\gamma p_0 + B_0^2} u_1 \right)^2,$$

$$H(u_1) = \left(\frac{g^2 \rho_0^2}{\gamma p_0 + B_0^2} - g\rho_{0x} \right) u_1^2,$$

The spectral radius Λ is given by

$$\Lambda^2 = \sup_{v \in L^2(D)} \frac{\iint_D H(v) \, dx dy}{\iint_D \rho_0 v^2 \, dx dy} > 0. \tag{4.9}$$

The sufficient condition for nonlinear instability is given by, for some $x_0 \in (0, 2\pi)$,

$$\frac{g^2 \rho_0^2}{\gamma p_0 + B_0^2} \left(x_0 \right) > g \rho_{0x} \left(x_0 \right).$$
(4.10)

Notice that density inversion is not required for instability unlike the incompressible case. Thus, even if light fluid is on top of heavy fluid and magnetic field lines stabilizes, such steady state flows can't sustain themselves against small initial disturbances under the effect of compressibility. It means that the destabilizing effect of compressibility dominates over the stabilizing one of magnetic field lines. Furthermore, letting $\gamma \to \infty$ yields exactly (4.4) and the instability criterion for the incompressible case can be recovered from compressible ones as the limiting case.

(3) Magnetic field is parallel to the gravity $B_0 \parallel \vec{g}$ for incompressible fluids:

$$B_0 = (B_0, 0, 0). (4.11)$$

In this case we consider density profile ρ_0 whose gradient has negative average over $(0, 2\pi)$, which is stronger than (4.4):

$$\int_{0}^{2\pi} -g\rho_{0x}(x)\,dx > 0,$$

Linear operator L, Sturm-Liouville equation, and λ_k are given by:

 $L(v) := B_0^2 v_{xx} - \nabla (p_t + B_0 \cdot Q) - \vec{g} \rho_{0x} v_1,$

$$\rho_0 \lambda_k^2 \tilde{v}_1 = B_0^2 \tilde{v}_{1xx} + \frac{\lambda_k^2}{k^2} \left(\rho_0 \tilde{v}_{1x}\right)_x - \frac{B_0^2}{k^2} \tilde{v}_{1xxxx} - g\rho_{0x} \tilde{v}_1, \qquad (4.12)$$

$$\lambda_k^2 = \sup_{u \in H^2(0,2\pi)} \frac{\int_0^{2\pi} L(u) \cdot u \, dx}{\int_0^{2\pi} \rho_0 \left[u^2 + \frac{u_x^2}{k^2} \right] \, dx} > 0, \tag{4.13}$$

where

$$L(u) \cdot u = -F\left(-\frac{u_{xx}}{k}\right) + H(u, u_x),$$

$$F\left(-\frac{u_{xx}}{k}\right) = B_0^2 \frac{u_{xx}^2}{k^2}, \ H(u, u_x) = -g\rho_{0x}u^2 - B_0^2 u_x^2,$$

 Λ has the following formula:

$$\Lambda^{2} = \sup_{v \in H^{1}(D)} \frac{\iint_{D} H(v, v_{x}) \, dx dy}{\iint_{D} \rho_{0} v^{2} \, dx dy} > 0.$$
(4.14)

,

Lemma 4. Λ^2 in (4.14) is positive.

Proof. We can take a family of test periodic functions in H^1 which guarantees the positivity of Λ . Let $c = \min_x \rho_0(x)$ and

$$\phi^n(x,y) = c - \frac{x}{n} \text{ for } 0 \le x < 2\pi.$$

Since

$$\begin{split} \iint_{D} & -g\rho_{0x} \left(\phi^{n}\right)^{2} dx dy \quad \rightarrow \quad c^{2} \iint_{D} -g\rho_{0x} dx dy > 0 \quad \text{as } n \to \infty \\ & B_{0}^{2} \iint_{D} \left(\phi_{x}^{n}\right)^{2} dx dy \quad = \quad \frac{4\pi^{2} B_{0}^{2}}{n^{2}} \to 0 \quad \text{as } n \to \infty, \end{split}$$

we choose large n to get positivity of the numerator of Λ^2 .

(4) Magnetic field is parallel to the gravity $B_0 \parallel \vec{g}$ for compressible fluids: In this case, we obtain linear stability instead of instability and we have the following:

$$L(v) = B_0^2 v_{xx} - \nabla \left(B_0 \cdot Q \right) - \nabla \left(p_{0x} v_1 + \gamma p_0 \nabla \cdot v \right) - \vec{g} \left(\rho_{0x} v_1 + \rho_0 \nabla \cdot v \right),$$

$$\lambda_k^2 = \sup_{\substack{u = (u_1, u_2) \\ u_1 \in H^1(0, 2\pi), \\ u_2 \in H^1(0, 2\pi)}} \frac{\int_0^{2\pi} L\left(u\right) \cdot u \ dx}{\int_0^{2\pi} \rho_0 \left[u_1^2 + u_2^2\right] \ dx} < 0,$$

where

$$\begin{split} L\left(u\right) \cdot u &= -F\left(u_{2x}\right) - G\left(u_{1}, u_{1x}, u_{2}\right) + H\left(u_{1}, u_{1x}\right), \\ F\left(u_{2x}\right) &= B_{0}^{2}u_{2x}^{2}, \\ G\left(u_{1}, u_{1x}, u_{2}\right) &= \left(\gamma p_{0} + B_{0}^{2}\right) \left[\frac{\gamma p_{0}}{\gamma p_{0} + B_{0}^{2}}u_{1x} + \frac{g\rho_{0}u_{1}}{\gamma p_{0} + B_{0}^{2}} + ku_{2}\right]^{2}, \\ H\left(u_{1}, u_{1x}\right) &= \frac{1}{\gamma p_{0} + B_{0}^{2}}\left[g\rho_{0}u_{1} - B_{0}^{2}u_{1x}\right]^{2} - g\rho_{0x}u_{1}^{2} - B_{0}^{2}u_{1x}^{2} \\ &= -\frac{B_{0}^{2}}{\left(\gamma p_{0} + B_{0}^{2}\right)\gamma p_{0}}\left(g\rho_{0}u_{1} + \gamma p_{0}u_{1x}\right)^{2} < 0. \end{split}$$

and

$$\Lambda^{2} = \sup_{v \in H^{1}(D)} \frac{\iint_{D} H(v, v_{x}) \, dx dy}{\iint_{D} \rho_{0} v^{2} \, dx dy} < 0.$$
(4.15)

5. Energy estimate and approximate solution

In this section, we construct approximate solutions using a method originated by Grenier in [8] and we do energy estimates for the fully nonlinear MHD system in both incompressible and compressible cases.

We first construct approximate solutions. In our construction, $\delta > 0$ is an arbitrary small parameter, and θ is a small but fixed positive constant (independent of δ). We fix k_0 with $\lambda = \lambda_{k_0}$ (dominant eigenvalue) so that

 $\Lambda < 2\lambda.$

We define T^{δ} by

$$\theta = \delta \exp\left(\lambda T^{\delta}\right),\tag{5.1}$$

or equivalently,

$$T^{\delta} = \frac{1}{\lambda} \ln \frac{\theta}{\delta}.$$
 (5.2)

We may write the full system (1.6) in vector form for $w = (\sigma(t, x, y), v(t, x, y), p(t, x, y), B(t, x, y))$:

$$w + A^{1}(w) \partial_{x} w + A^{2}(w) \partial_{y} w + L(w) = F(w)$$

An approximate solution $w^{a}(t, x, y) = (\sigma^{a}(t, x, y), v^{a}(t, x, y), p^{a}(t, x, y), B^{a}(t, x, y))$ is of the form

$$w^{a}(t, x, y) = \sum_{j=1}^{N} \delta^{j} r_{j}(t, x, y), \qquad (5.3)$$

such that

$$w^{a} + A^{1}(w^{a}) \partial_{x} w^{a} + A^{2}(w^{a}) \partial_{y} w^{a} + L(w^{a}) = F(w^{a}) + R_{N}^{a}.$$
(5.4)

We show existence of such approximate solutions in the following lemma. The key point is that we can choose a dominant eigenvalue λ with $\Lambda < 2\lambda$ to make this construction work.

Lemma 5. For any fixed integer N > 0, there exists an approximate solution (5.3) satisfying (5.4). Furthermore, for every integer $s \ge 0$, there is small $\theta > 0$ such that if $0 \le t \le T^{\delta}$ as in (5.2), the *j*-th coefficient r_j and the remainder R_N^a satisfy

$$\|r_j\|_{H^s} \le C(s, N) \exp\left(j\lambda t\right), \text{ for } 1 \le j \le N,$$
(5.5)

$$\|R_N^a\|_{H^s} \le C(s, N) \,\delta^{N+1} \exp\left\{(N+1)\,\lambda t\right\}.$$
(5.6)

Proof. The construction of r_j will be made by induction on j. The idea is as follows. We split the system into linear and nonlinear part:

$$\partial_t w^a + A^1(0) \partial_x w^a + A^2(0) \partial_y w^a + L(w^a)$$

$$= \left[A^1(w^a) - A^1(0) \right] \partial_x w^a + \left[A^2(u^a) - A^2(0) \right] \partial_y w^a - F(w^a) =: h(\delta)$$
(5.7)

with the Taylor expansion of w^a in δ

$$w^a = \sum_{j=1}^N \delta^j r_j.$$

Thus, our r_j is the solution of the part of (5.7) which corresponds to the coefficient of δ^j in its Taylor expansion.

For j = 1, take for r_1 the smooth normal growing mode to the linearized system with our chosen wave number k_0 and the corresponding dominant eigenvalue $\lambda = \lambda_{k_0}$ as in (2.2). Clearly this growing mode fulfills (5.5).

Assuming that we have constructed r_i (j < N) which satisfies (5.5), we construct r_{i+1} . Let

$$u_j = \sum_{k=1}^{j} \delta^k r_k \left(t, x, y \right).$$

Define the nonlinear part of the system substituted by u_i as

$$h_{j+1}(\delta) = \left[A^{1}(u_{j}) - A^{1}(0)\right]\partial_{x}u_{j} + \left[A^{2}(u_{j}) - A^{2}(0)\right]\partial_{y}u_{j} - F(u_{j})$$

Since this is the nonlinear part of the system and the terms in δ^{j+1} come from the terms δ^k for $1 \leq k \leq j$, it is enough to consider u_j in order to collect the (j+1)-th coefficient of nonlinear part of the expansion. Then r_{j+1} is defined to be a solution of

$$\partial_t r_{j+1} + A^1(0) \,\partial_x r_{j+1} + A^2(0) \,\partial_y r_{j+1} + L(r_{j+1}) \\ = \frac{-h_{j+1}^{(j+1)}(0)}{(j+1)!}$$

with initial data $r_{j+1}(0, x, y) = 0$. Notice that

$$\frac{-h_{j+1}^{(j+1)}(0)}{(j+1)!} = \sum_{j_1m_1+j_2m_2+\ldots+j_{p+1}m_{p+1}=j+1} B_{J,i}^M r_{j_1}^{m_1} r_{j_2}^{m_2} \cdots r_{j_p}^{m_p} \partial_i r_{j_{p+1}}^{m_{p+1}},$$

where $m_k \ge 0$, $1 \le k \le j$, and $B_{J,i}^M$ depends on A^i and F. Induction hypothesis (5.5) for r_k , $1 \le k \le j$ applies to get, for all s,

$$\left\|\frac{h_{j+1}^{(j+1)}(0)}{(j+1)!}\right\|_{H^s} \le C(s,N) \exp\left[\left(j_1m_1 + j_2m_2 + \ldots + j_{p+1}m_{p+1}\right)\lambda t\right] = C(s,N) \exp\left[\left(j+1\right)\lambda t\right].$$

Thanks to our linear estimates for Λ in Lemma 4 and Duhamel principle, we have

$$\begin{aligned} \|r_{j+1}(t,\cdot)\|_{H^{s}} &\leq C \int_{0}^{t} e^{\Lambda(t-\tau)} \left\| \frac{h_{j+1}^{(j+1)}(0)}{(j+1)!}(\tau) \right\|_{H^{s+2}} \\ &\leq C(s,N) \int_{0}^{t} e^{\Lambda(t-\tau)} e^{(j+1)\lambda\tau} d\tau \\ &\leq C(s,N) e^{(j+1)\lambda t} \end{aligned}$$

since $j + 1 \ge 2$ and $\Lambda < 2\lambda$. We now define $w^a = \sum_{j=1}^N \delta^j r_j$ and it satisfies

$$\partial_t w^a + A^1(0) \,\partial_x w^a + A^2 \partial_y w^a + L(w^a) = -\sum_{j=1}^N \frac{\delta^j h_j^{(j)}(0)}{j!}.$$

Then R_N^a is defined to be the sum of all higher terms than N in nonlinear part of the δ -expansion (5.7):

$$R_{N}^{a} = h(\delta) - \sum_{j=1}^{N} \frac{\delta^{j} h_{j}^{(j)}(0)}{j!},$$

which clearly satisfies (5.6) and our proof is complete.

We state local in time existence for the incompressible ideal MHD equations:

Lemma 6. (Local existence to the full system) For all $s \geq 3$ and for any given initial data $(\sigma_0, v_0) \in H^s(D)$ such that $\rho(0) \equiv \rho_0(x) + \sigma_0(x, y) \ge m > 0$, there is a T > 0 such that there exists a unique solution $(\sigma, v, p, B) \in C([0, T] : H^s(D))$ to (1.6) with $\rho(t) = \rho_0(x) + \sigma(t, x, y) > 0$ 0.

We first treat the incompressible case and we use some vector identities to get:

$$\sigma_t + v \cdot \nabla \left(\rho_0 + \sigma\right) = 0,$$

$$(\rho_0 + \sigma) (v_t + v \cdot \nabla v) = -\nabla \left(p + \frac{1}{2} B^2 + B_0 \cdot B \right) + B \cdot \nabla B + B_0 \cdot \nabla B + B \cdot \nabla B_0 + \sigma \vec{g},$$

$$B_t = B_0 \cdot \nabla v - v \cdot \nabla B_0 + B \cdot \nabla v - v \cdot \nabla B,$$

where k is the unit vector in the z direction. Let $w(t, x, y) = (\sigma(t, x, y), v(t, x, y), p(t, x, y), B(t, x, y)) \in C([0, T]; H^s(D))$ be a local solution as constructed above.

Let $w^a(t, x, y) = (\sigma^a(t, x, y), v^a(t, x, y), p^a(t, x, y), B^a(t, x, y))$ be an approximate solution. We now estimate the difference

$$w^{d}(t, x, y) = w(t, x, y) - w^{a}(t, x, y).$$

$$\sigma_t^d + \nabla \sigma^d \cdot v + \nabla \left(\sigma^a + \rho_0\right) \cdot v^d = -R_{N,1}^a, \tag{5.8}$$
$$\left(\rho_0 + \sigma\right) \left(v_t^d + v^d \cdot \nabla v + v^a \cdot \nabla v^d\right) + \sigma^d \left(v_t^a + v^a \cdot \nabla v^a\right)$$
$$= -\nabla \left(p^d + \frac{1}{2}B^d \left(B + B^a\right) + B_0 \cdot B^d\right) + B^d \cdot \nabla B + B^a \cdot \nabla B^d \tag{5.9}$$

$$+ B_0 \cdot \nabla B^d + B^d \cdot \nabla B_0 + \sigma^d \vec{g} - R^a_{N,2},$$

$$B_t^d = B_0 \cdot \nabla v^d - v^d \cdot \nabla B_0 + B^d \cdot \nabla v + B^a \cdot \nabla v^d - v^d \cdot \nabla B - v^a \cdot \nabla B^d - R_{N,3}^a.$$
(5.10)

Lemma 7. For $s \ge 3$ and assume $\|\sigma\|_{\infty} \le \frac{1}{2} \|\rho_0\|_{\infty}$, then there exists a continuous positive function C depending only on s, ρ_0 such that

$$\frac{d}{dt} \left\| w^{d} \right\|_{s}^{2} \leq C \left(\left\| w^{d} \right\|_{s}, \left\| w^{a} \right\|_{s+1} \right) \left\| w^{d} \right\|_{s}^{2} \\ + \left\| R_{N,1}^{a} \right\|_{s}^{2} + \left\| R_{N,2}^{a} \right\|_{s}^{2} + \left\| R_{N,3}^{a} \right\|_{s}^{2}.$$

Proof. This energy estimate is straightforward and thus we give a brief sketch. We take ∂_{α} of the equations (5.8)-(5.10), multiply through ∂_{α} -derivatives, and integrate over D. For symmetric terms such as $\nabla \partial_{\alpha} \sigma^d \cdot v$ in the first equation vanish upon integration due to the divergence free condition for v, v^a, v^d, B, B^a, B^d . We now estimate nonsymmetric terms. Terms which need attention are $(B^d \cdot \nabla) \partial_{\alpha} B^d, (B^a \cdot \nabla) \partial_{\alpha} B^d, (B_0 \cdot \nabla) \partial_{\alpha} B^d$ from v-equation (5.9) and $(B^d \cdot \nabla) \partial_{\alpha} v^d, (B^a \cdot \nabla) \partial_{\alpha} v^d, (B_0 \cdot \nabla) \partial_{\alpha} B^d$ from B-equation (5.10). Upon integration, the three corresponding pairs exactly cancel out. To see this,

$$\begin{split} B_{i}^{d}\partial_{i}\partial_{\alpha}B_{j}^{d}\partial_{\alpha}v_{j}^{d} &= \partial_{i}B_{i}^{d}\partial_{\alpha}B_{j}^{d}\partial_{\alpha}v_{j}^{d} - B_{i}^{d}\partial_{i}\partial_{\alpha}v_{j}^{d}\partial_{\alpha}B_{j}^{d} = -B_{i}^{d}\partial_{i}\partial_{\alpha}v_{j}^{d}\partial_{\alpha}B_{j}^{d},\\ B_{i}^{a}\partial_{i}\partial_{\alpha}B_{j}^{d}\partial_{\alpha}v_{j}^{d} &= \partial_{i}B_{i}^{a}\partial_{\alpha}B_{j}^{d}\partial_{\alpha}v_{j}^{d} - B_{i}^{a}\partial_{i}\partial_{\alpha}v_{j}^{d}\partial_{\alpha}B_{j}^{d} = -B_{i}^{a}\partial_{i}\partial_{\alpha}v_{j}^{d}\partial_{\alpha}B_{j}^{d},\\ B_{0i}\partial_{i}\partial_{\alpha}B_{j}^{d}\partial_{\alpha}v_{j}^{d} &= \partial_{i}B_{0i}\partial_{\alpha}B_{j}^{d}\partial_{\alpha}v_{j}^{d} - B_{0i}\partial_{i}\partial_{\alpha}v_{j}^{d}\partial_{\alpha}B_{j}^{d} = -B_{0i}\partial_{i}\partial_{\alpha}v_{j}^{d}\partial_{\alpha}B_{j}^{d}, \end{split}$$

thanks to divergence free condition for v and B. Therefore we obtain the Lemma.

We now extend the energy estimates to the compressible case using symmetrizer, which is necessary in this compressible case. We write the full system (1.6) in components:

$$\sigma_{,t} + v_i \sigma_{,i} + (\rho_0 + \sigma) v_{i,i} + \rho_{0x} v_1 = 0,$$

$$(\rho_0 + \sigma) \{ v_{j,t} + v_i v_{j,i} \} + (\rho_0 + \sigma) q \sigma_{,j} + B_i B_{i,j} - B_i B_{j,i} + B_0 B_{3,j}$$

$$= [\rho_0 q (\rho_0) - (\rho_0 + \sigma) q] \rho_{0,j} + B_i B_{0,i} - B_{0,j} B_3$$

$$B_{j,t} - B_i v_{j,i} + v_{i,i} B_j + v_i B_{j,i} + v_{i,i} B_0 = -v_i B_{0,i}.$$

Here j = 1, 2, 3 and twice *i* means the sum over i = 1, 2, 3.

We rewrite the full system near the steady state $(\rho_0, \vec{0}, B)$ in vector notations for $w = (\sigma, v_1, v_2, v_3, B_1, B_2, B_3)$:

$$w_t + A^1(w) \partial_x w + A^2(w) \partial_y w + L(w) = F(w), \qquad (5.11)$$

Introducing and multiplying the symmetrizer with $q\left(\rho\right)=\gamma\rho^{\gamma-2}$

$$D = \text{diag}(q(\rho_0 + \sigma), \rho_0 + \sigma, \rho_0 + \sigma, \rho_0 + \sigma, 1, 1, 1),$$
(5.12)

leads to the following symmetric matrices

$$DL(u) = \begin{bmatrix} \rho_{0x}q(\rho_0)v_1 \\ q(\rho_0)\rho_{0x}\sigma + \rho_0q'(\rho_0)\rho_{0x}\sigma + B_{0x}B_3 \\ 0 \\ -B_{0x}B_1 \\ 0 \\ 0 \\ B_{0x}v_1 \end{bmatrix}$$

,

$$DF(u) = \begin{bmatrix} \rho_{0x}q(\rho_0)v_1 - \rho_{0x}q(\rho_0 + \sigma)v_1 \\ [(\sigma + \rho_0)q(\sigma + \rho_0) - \rho_0q(\rho_0) - q(\rho_0)\rho_{0x}\sigma - \rho_0q'(\rho_0)\rho_{0x}\sigma]\rho_{0x} \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Now that DA^1 , DA^2 and DA^3 are symmetric, we have the following local in time solution to the full system via standard energy estimates for symmetrizable hyperbolic system as in [3], [15] in the absence of magnetic fields.

Lemma 8. For all $s \ge 3$ and for any given initial data $(\sigma_0, v_0) \in H^s(D)$ such that $\rho(0) \equiv \rho_0(x) + \sigma_0(x, y) \ge m > 0$, there is a T > 0 such that there exists a unique solution $u = (\sigma, v, B) \in C([0, T] : H^s(D))$ to (1.6) with $\rho(t) = \rho_0(x) + \sigma(t, x, y) > 0$.

Since construction of approximate solutions is similar to the incompressible case as in Lemma 5 for a hyperbolic system, we omit it. We now estimate the difference of an exact solution and an approximate solution. Let $w(t, x, y) = (\sigma(t, x, y), v(t, x, y), B(t, x, y)) \in C([0, T] : H^s(D))$ be an exact solution and $w^a(t, x, y) = (\sigma^a(t, x, y), v^a(t, x, y), B^a(t, x, y))$ be an approximate solution as constructed in Lemma 5. Then their difference is

$$w^{d} = w - w^{a} = (\sigma - \sigma^{a}, v - v^{a}, B - B^{a}),$$

and it satisfies

$$w_t^d + \sum_{i=1}^3 A^i \left(w^a + w^d \right) \partial_i w^d + \sum_{i=1}^3 \left[A^1 \left(w^a + w^d \right) - A^1 \left(w^a \right) \right] \partial_x w^a$$
(5.13)
= $-L \left(w^d \right) + F \left(w \right) - F \left(w^a \right) - R_N^a$.

This symmetrizable hyperbolic system for w^d allows the following energy estimates. The proof is straightforward by classical energy methods as in [3], [8].

Lemma 9. For any $s \geq 3$, there exists a continuous function $g_s(\cdot, \cdot)$ such that

$$\frac{\partial}{\partial t} \left| \left| \left| w^{d} \right| \right| \right|_{s}^{2} \leq g_{s} \left(\left| \left| w^{d} \right| \right| \right|_{s}, \left| \left| w^{a} \right| \right| \right|_{s+1} \right) \left| \left| \left| w^{d} \right| \right| \right|_{s}^{2} + \left| \left| \left| R_{N}^{a} \right| \right| \right|_{s}^{2},$$

where $||| \cdot |||_s$ is defined by

=

$$|||v|||_{s}^{2} = \sum_{|\alpha| \le s} \partial_{\alpha} v D \left(\rho_{0} + \sigma\right) \partial_{\alpha} v.$$

Here D is the symmetrizer as in (5.12) and notice that $||| \cdot |||_s$ is related to the usual norm $|| \cdot ||_{H^s}$ by

 $\eta \|v\|_{H^s} \le |||v|||_s \le C_s \left(\|\sigma\|_{H^s}, \|\rho_0\|_{H^s} \right) \|v\|_{H^s},$

since $\rho_0 \ge c > 0$ and so $D \ge \eta Id$ for some η and $s \ge 3$.

Notice that all three norms $\|\cdot\|_s$, $\||\cdot\||_s$, and $\|\cdot\|_{H^s}$ are equivalent since ρ_0 is smooth with a positive minimum.

6. Nonlinear instability

Proof of Theorem 2. Let $w^a(t, x, y) = (\sigma^a(t, x, y), v^a(t, x, y), p^a(t, x, y), B^a(t, x, y))$ be an approximate solution with N to be determined later. For any $\delta > 0$, there exists a local in time solution $w^{\delta}(t, x, y) = (\sigma^{\delta}(t, x, y), v^{\delta}(t, x, y), p^{\delta}(t, x, y), B^{\delta}(t, x, y))$ with initial data $w^a(0)$ to the full system (1.6). By the Lemma 7 and Lemma 9, we have

$$\frac{d}{dt} \left\| w^d \right\|_s^2 \le C\left(\left\| w^d \right\|_s, \left\| w^a \right\|_{s+1} \right) \left\| w^d \right\|_s^2 + C\delta^{2(N+1)} e^{2(N+1)\lambda t}, \tag{6.1}$$

with $w^d(0) = 0$.

Let

$$T = \sup \left\{ t \mid \|w^a\|_{s+1} \le \omega/2, \|w^d\|_s \le \omega/2 \right\},\$$

where ω is a small positive number which assures local existence. Then T depending on δ is well defined since $w^d(0) = 0$ and $\|w^{\alpha}(0)\|_{s+1} = O(\delta)$. We claim that the instability time T^{δ} occurs within the existence time T, that is, $T^{\delta} \leq T$. Suppose not, i.e., $T < T^{\delta}$. Then for $t \leq T$, by our construction of approximate solution, we have

$$\|w^{a}\|_{s+1} \leq C \sum_{j=1}^{N} \delta^{j} \|r_{j}(t)\|_{H^{s}} \leq \sum_{j=1}^{N} C_{j} \delta^{j} e^{j\lambda t} \leq \sum_{j=1}^{N} C_{j} \theta^{j} < \omega/2 ,$$

if θ is small. Now we appeal to the definition of T and (6.1) to get, for $t \leq T$,

$$\frac{d}{dt} \left\| w^d \right\|_s^2 \le C \left(\omega/2, \omega/2 \right) \left\| w^d \right\|_s^2 + C \delta^{2(N+1)} e^{2(N+1)\lambda t}.$$

We choose N > 0 satisfying

$$C\left(\omega/2,\omega/2\right)<2\left(N+1\right),$$

so that $\|w^d\|_s$ has growth rate at most $(N+1)\lambda$. Then using Gronwall inequality leads to

$$\|w^d\|_s \le C\delta^{2(N+1)}e^{2(N+1)\lambda t} = C\theta^{2(N+1)} < \omega/2,$$

if ω is small. Thus we can deduce $T^{\delta} \leq T$. Now at the instability time T^{δ} ,

$$\begin{split} \left\|w^{a}\left(T^{\delta}\right)\right\|_{L^{1}} &\geq \delta \left\|r_{1}\left(T^{\delta}\right)\right\|_{L^{1}} - \sum_{j=2}^{N} \delta^{j} \left\|r_{j}\left(T^{\delta}\right)\right\|_{L^{1}} \\ &\geq C\delta e^{\lambda T^{\delta}} - \sum_{j=2}^{N} C_{j} \delta^{j} e^{j\lambda T^{\delta}} \\ &= C\theta - \sum_{j=2}^{N} C_{j} \theta^{j} \geq \frac{C}{2}\theta. \end{split}$$

We then deduce at time T^{δ} ,

$$\begin{aligned} \left\|w^{\delta}\left(T^{\delta}\right)\right\|_{L^{1}} &\geq \left\|w^{\delta}\left(T^{\delta}\right)\right\|_{L^{1}} - \left\|\left(w^{\delta} - w^{a}\right)\left(T^{\delta}\right)\right\|_{L^{1}} \\ &\geq \left\|w^{a}\left(T^{\delta}\right)\right\|_{L^{1}} - \left\|\left(w^{\delta} - w^{a}\right)\left(T^{\delta}\right)\right\|_{H^{s}} \\ &\geq \frac{C}{2}\theta - C\theta^{N+1} \geq \frac{C}{4}\theta = \varepsilon_{0} > 0. \end{aligned}$$

References

- [1] G. Bateman, MHD instabilities, MIT Press, Cambridge, MA, 1980.
- [2] I. B. Bernstein, E. A. Frieman, M. D. Kruskal, R. M. Kulsrud, Proc. Roy. Soc. (London) A244 (1958), 17
- [3] S. Cordier, E. Grenier, Y. Guo, Two-stream instabilities in plasmas, In honor of C. S. Morawetz. Methods Appl. Anal. 7 (2000), no. 2, 391-405.
- [4] S. Chandrasekhar, Hydrodynamic and hydromagnetic stability, The International Series of Monographs on Physics Clarendon Press, Oxford 1961.
- [5] S. Chandrasekhar, Proc. Natl. Acad. Sci. U. S. 23 (1937), 572
- [6] C. Cherfils-Cléouin, O. Lafitte, P-A. Raviart. Asymptotic results for the linear stage of the Rayleigh-Taylor instability, Mathematical fluid mechanics, 47–71, Adv. Math. Fluid Mech., Birkhäser, Basel, 2001.
- [7] T. G. Cowling, Monthly Notices Roy. Astro. Soc. 94 (1934), 768.
- [8] E. Grenier, Semiclassical limit of the nonlinear Schrödinger equations, Proc. AMS. 126 (1998), no. 2, 523-530.
- [9] Y. Guo, W. Strauss, Instability of periodic BGK equilibria, Comm. Pure. Appl. Math. 48 (1995), 861-894
- [10] H. J. Hwang, Y. Guo, On the dynamical Rayleigh-Taylor instability, Arch. Rational Mech. Anal. 167 (2003), 235-253.
- [11] H. Jeffreys, Phil. Mag. 2 (1926), 833.
- [12] M. D. Kruskal, M. Schwarzschild, Proc. Roy. Soc. (London) A223 (1954), 348.
- [13] P-L. Lions, Mathematical topics in fluid mechanics. Vol. 2. Compressible models. Oxford Lecture Series in Mathematics and its Applications, 3. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1996.
- [14] L. D. Landau, E. M. Litschitz, Fluid mechanics, 2nd ed., Pergamon, New York, 1987.
- [15] A. Majda, Compressible fluid flows and systems of conservation laws in several variables. Appl. Math. Sci 53, Springer, 1984.
- [16] D. Nicholson, Introduction to plasma theory, John Wiley & Sons.
- [17] E. R. Priest, Solar magntohydrodynamics, London, 1983.
- [18] L. Rayleigh, Analytic solutions of the Rayleigh equation for linear density profiles, Proc. London. Math. Soc. 14 (1883), 170-177.
- [19] K. Schwarzchild, Nachr. Kgl. Ges. Wiss. Göttingen. (1906) p.41.
- [20] Yu. A. Tserkovikov, Convective instability of a rarefied plasma. Doklady Akad. Nauk S. S. S. R. 130 (1960), 295 [translation: Soviet Phys.-Doklady 5 (1960), 87.

DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, DURHAM NC 27708, USA *E-mail address*: hjhwang@math.duke.edu