On low degree k-ordered graphs

Karola Mészáros Massachusetts Institute of Technology karola@math.mit.edu

Abstract

A simple graph G is k-ordered (respectively, k-ordered hamiltonian) if, for any sequence of k distinct vertices v_1, \ldots, v_k of G, there exists a cycle (respectively, a hamiltonian cycle) in G containing these k vertices in the specified order. In 1997 Ng and Schultz introduced these concepts of cycle orderability, and motivated by the fact that k-orderedness of a graph implies (k - 1)-connectivity, they posed the question of the existence of low degree k-ordered hamiltonian graphs. We construct an infinite family of graphs, which we call bracelet graphs, that are (k - 1)-regular and are k-ordered hamiltonian for odd k. This result provides the best possible answer to the question of the existence of low degree k-ordered hamiltonian graphs for odd k. We further show that for even k, there exist no k-ordered bracelet graphs with minimum degree k - 1 and maximum degree less than k + 2, and we exhibit an infinite family of bracelet graphs with minimum degree k - 1 and maximum degree k + 2 that are k-ordered for even k. A concept related to k-orderedness, namely that of k-edge-orderedness, is likewise strongly related to connectivity properties. We study this relation in both undirected and directed graphs, and give bounds on the connectivity necessary to imply k-(edge-)orderedness properties.

1 Introduction

The concept of k-ordered graphs was introduced in 1997 by Ng and Schultz [11]. A simple graph G is a graph without loops or multiple edges, and it is called *hamiltonian* if there exists a cycle (called a *hamiltonian cycle*) that contains all vertices of G. In this paper we consider only finite simple graphs. A simple graph G is called *k-ordered* (respectively, *k-ordered hamiltonian*) if, for any sequence of k distinct vertices v_1, v_2, \ldots, v_k of G, there exists a cycle (respectively, a hamiltonian cycle) in G containing these k vertices in the specified order. Previous results concerning cycle orderability mainly regard minimum degree and forbidden subgraph conditions that imply k-ordered hamiltonicity [3, 5, 6, 7, 9]. A comprehensive survey of results can be found in [4].

A notion related to k-orderedness, that of k-edge-orderedness, has been studied in [2]. A simple graph G is k-edge-ordered (respectively, k-edge-ordered eulerian) if, for any sequence of k distinct edges e_1, e_2, \ldots, e_k of G, there exists a tour (respectively, an eulerian tour, that is, a tour containing each edge of G) in G containing these k edges in the specified order. It is natural to explore analogous notions in directed graphs. A directed graph D is k-ordered (hamiltonian) if, for any sequence of k distinct vertices v_1, v_2, \ldots, v_k of D, there exists a directed (hamiltonian) cycle in D containing these k vertices in the specified order. Furthermore, D is k-edge-ordered (eulerian) if, for any sequence of k distinct edges e_1, e_2, \ldots, e_k of D, there exists a directed (eulerian) tour in D containing these k edges in the specified order. As k-orderedness implies (k-1)-connectivity, a natural question to pose is the existence of low degree k-ordered graphs. The question of the existence of 3-regular 4-ordered graphs was posed in [11] and answered in the affirmative in [10]. In Section 2, we answer the more general question of the existence of (k-1)-regular k-ordered graphs for odd k; in particular, we exhibit an infinite family of graphs, called *bracelet graphs*, that are (k-1)-regular and k-ordered hamiltonian. We also exhibit sufficient conditions for a bracelet graph to be k-ordered.

In Section 3 we exhibit a bound on the diameter of a k-ordered graph, and we show that the bound is almost tight for the bracelet graph that we constructed in Section 2. In Section 4 we continue investigating low degree k-ordered graphs for even k, and we show that for even k there are no k-ordered bracelet graphs with minimum degree k - 1 and maximum degree less than k + 2; however, we also exhibit an infinite family of bracelet graphs with minimum degree k - 1 and maximum degree k + 2 that are k-ordered for even k. This construction partially answers the question of the existence of low degree k-ordered graphs for even k.

In Section 5 we consider k-orderedness properties of directed graphs, exhibiting an infinite family of (k-1)-diregular graphs that are k-ordered hamiltonian. In Sections 6 and 7 we establish a relation between connectivity and k-(edge-)orderedness in undirected as well as directed graphs. We conclude our paper by posing open questions.

2 2k-regular (2k+1)-ordered hamiltonian graphs

As observed in [11], a k-ordered graph G is also (k-1)-connected, and hence has minimum degree at least k-1. The question of the existence of an infinite family of 3-regular 4-ordered graphs was raised in [11] and answered in [10] by constructing such a family. More generally, we are interested in whether there exists an infinite family of (k-1)-regular k-ordered graphs. In this section we answer this question in the case where k is odd, exhibiting an infinite family of (k-1)-regular k-ordered hamiltonian graphs for all odd $k \geq 3$.

We call a graph G a bracelet graph if its vertex set V can be partitioned into $V_1 \cup V_2 \cup \cdots \cup V_m$, $m \geq 3$, with V_i nonempty for all $i \in [m]$ (we denote the set $\{1, 2, \ldots, m\}$ by [m]), such that v is adjacent to u in G if and only if $v \in V_i$ and $u \in V_j$ and $i - j \equiv 1$ or $-1 \pmod{m}$. We call V_i , for $i \in [m]$, a part of G, and denote its cardinality by $|V_i|$. We say that two parts V_i and V_j are adjacent if $i - j \equiv 1$ or $-1 \pmod{m}$. We also say that parts V_i and V_j are at distance d if there is a path from a vertex in V_i to a vertex in V_j such that it contains d edges and there are no 2 vertices on the path from the same part. Note that as bracelet graphs are "cyclic" there are two options for the distance between two parts; in general, it will be clear from the context which of the two distances we mean.

Throughout this paper we will frequently want to construct a cycle or path through vertices in a specified order. We will refer to these specified vertices as *marked vertices*. We also use the idea of free vertices in the course of the paper; we shall define free vertices in the statement of Lemma 1, which we will use for proving Theorem 2.1.

Let $G_{k,2m}$ be a bracelet graph with parts $V_1, V_2, \ldots, V_{2m}, m \ge 2$, such that $|V_i| = k$ for $i \in [2m]$. It is clear that G is simple and 2k-regular by construction.

Lemma 1. Given 2k + 1 marked vertices $v_1, v_2, \ldots, v_{2k+1}$ in $G_{k,2m}$, there exists a set of 2m vertices, which we call free vertices, satisfying the following two properties: (i) there is exactly one free vertex in each part of $G_{k,2m}$, and (ii) there exists some i such that the marked vertices v_i and

 v_{i+1} are in the set of free vertices (indices taken modulo 2k + 1) and no other marked vertices are in the set of free vertices.

Proof. If each part of $G_{k,2m}$ contains at most k-1 marked vertices let B_1 any part that contains a marked vertex, and if there is exactly one part that has k marked vertices, let B_1 be that part. Take a vertex v_i from B_1 such that $v_{i+1} \notin B_1$. Then we can take vertices v_i, v_{i+1} , and one unmarked vertex from each of the parts not containing v_i or v_{i+1} as the free vertices.

If there are exactly two parts, B_1 and B_2 , that have k marked vertices, then it is not hard to see that there are two consecutive vertices v_i and v_{i+1} (indices taken modulo 2k + 1) such that $v_i \in B_1$ and $v_{i+1} \in B_2$. Then we can take vertices v_i, v_{i+1} , and one unmarked vertex from each of the parts not containing v_i or v_{i+1} as the free vertices. This completes the proof.

Theorem 2.1. For every $k \ge 1$, there exists an infinite family of 2k-regular graphs that are (2k + 1)-ordered hamiltonian.

Proof. We prove that the bracelet graphs $G_{k,2m}$ introduced above are (2k+1)-ordered hamiltonian for $k \geq 1$ and $m \geq 2$. In fact, we will prove more: given any 2k + 1 vertices $v_1, v_2, \ldots, v_{2k+1}$ in $G_{k,2m}$, there exists a hamiltonian cycle $\mathcal{H}_{k,2m}$ of $G_{k,2m}$ that traverses the vertices in order, and satisfies the following condition (\star): for any two adjacent parts B_1 and B_2 of $G_{k,2m}$, there exists an edge of $\mathcal{H}_{k,2m}$ with one vertex in each of B_1 and B_2 . We will proceed by induction on k.

Base case: k = 1.

It is clear that $G_{1,2m}$ for $m \ge 2$ is just a cycle, and it follows that $G_{1,2m}$ is 3-ordered hamiltonian; furthermore we can take $\mathcal{H}_{1,2m} = G_{1,2m}$.

Inductive step.

Suppose that $G_{k-1,2m}$ is (2k-1)-ordered hamiltonian for $m \ge 2$ and given 2k-1 vertices in $G_{k-1,2m}$ there is a hamiltonian cycle $\mathcal{H}_{k-1,2m}$ satisfying condition (*). Consider the 2k+1 marked vertices $v_1, v_2, \ldots, v_{2k+1}$ through which we wish to construct a hamiltonian cycle. By Lemma 1 it is possible to find 2m free vertices in $G_{k,2m}$, one in each part. Without loss of generality we can suppose that the two marked vertices among the 2m free vertices are v_{2k} and v_{2k+1} . Note that the graph induced by the 2m free vertices is a cycle, \mathcal{C} , and the graph induced on the vertices of $G_{k,2m}$ without the free vertices is (isomorphic to) $G_{k-1,2m}$. Therefore, by the induction hypothesis, there exists a hamiltonian cycle $\mathcal{H}_{k-1,2m}$ through the 2k-1 marked vertices $v_1, v_2, \ldots, v_{2k-1}$ in $G_{k-1,2m}$ satisfying condition (*).

We show that given $v_1, v_2, \ldots, v_{2k+1}$ in $G_{k,2m}$ there is a hamiltonian cycle $\mathcal{H}_{k,2m}$ containing the 2k + 1 vertices in the specified order, and such that for any two adjacent parts B_1 and B_2 of $G_{k,2m}$ there exists an edge of $\mathcal{H}_{k,2m}$ with one vertex in each of B_1 and B_2 . This will also show, in particular, that $G_{k,2m}$ is (2k+1)-ordered hamiltonian for $m \geq 2$. We will examine cases depending on the positions of the 2k + 1 specified vertices and show how to construct the desired hamiltonian cycle in each case.

Case 1. Suppose v_{2k-1} and v_{2k} are in different parts. In this case the hamiltonian cycle $\mathcal{H}_{k,2m}$ in $G_{k,2m}$ is as follows. Follow $\mathcal{H}_{k-1,2m}$ in $G_{k-1,2m}$ from v_1 until reaching v_{2k-1} . If v_{2k} is in a part adjacent to the part of v_{2k-1} , go to v_{2k} from v_{2k-1} and continue going to the free vertices in the not yet visited adjacent parts along the cycle \mathcal{C} so that we reach v_{2k+1} . If v_{2k} is not in a part adjacent to the part containing v_{2k-1} , then continue going to the free vertices in the not yet visited adjacent parts along the cycle \mathcal{C} so that we reach v_{2k} first and then v_{2k+1} . In both cases (whether or not v_{2k} is in a part adjacent to the part containing v_{2k-1}) continue along \mathcal{C} after meeting v_{2k+1} until reaching the free vertex of the part containing v_{2k-1} . After this, go to the vertex that is adjacent to v_{2k-1} in $\mathcal{H}_{k-1,2m}$ when going from v_{2k-1} to v_1 , and continue on $\mathcal{H}_{k-1,2m}$ until v_1 . It is clear that the hamiltonian cycle $\mathcal{H}_{k,2m}$ that we have constructed has the property that for any two adjacent parts of $G_{k,2m}$ there exists an edge of $\mathcal{H}_{k,2m}$ with one vertex in each of the two adjacent parts.

Case 2. Suppose v_{2k-1} and v_{2k} are in the same part.

Let the part containing v_{2k-1} and v_{2k} be B_1 , and the part containing v_{2k+1} be B_2 . Note that $B_1 \neq B_2$ since v_{2k} and v_{2k+1} are in different parts. Let u be the vertex adjacent to v_{2k-1} in the traversal of $\mathcal{H}_{k-1,2m}$ from v_{2k-1} to v_1 .

Case 2.1. Suppose $u \notin B_2$. Follow $\mathcal{H}_{k-1,2m}$ in $G_{k-1,2m}$ from v_1 to v_{2k-1} . From v_{2k-1} go to an unmarked free vertex in the part containing u. From this free vertex go to v_{2k} , and after this continue going to the free vertices in the not yet visited adjacent parts along \mathcal{C} . During this we meet v_{2k+1} , and continue along \mathcal{C} until we are at the free vertex of the part adjacent to the part containing u. Go to u and continue on $\mathcal{H}_{k-1,2m}$ until v_1 . The obtained hamiltonian cycle is $\mathcal{H}_{k,2m}$.

Case 2.2. Suppose $u \in B_2$. In this case, B_1 and B_2 are adjacent. Follow $\mathcal{H}_{k-1,2m}$ in $G_{k-1,2m}$ from v_1 to v_{2k-1} . From v_{2k-1} go to an unmarked free vertex in the adjacent part that is not B_2 . From here go to v_{2k} , then to v_{2k+1} , then to the free vertex in the next adjacent part and back to u. Continue on $\mathcal{H}_{k-1,2m}$ until reaching v_1 . When m = 2 this is a hamiltonian cycle in $G_{k,2m}$, but if m > 2, this is not a hamiltonian cycle. We now show that we can reroute the path from v_1 to v_{2k-1} so that we pick up all the missing vertices.

Indeed, note the following. If there is an edge ab in the hamiltonian cycle $\mathcal{H}_{k-1,2m}$ in $G_{k-1,2m}$ and c and d are free unmarked vertices, such that a and c are in the same part and b and d are in the same part, then replacing edge ab by edges ad, dc, and cb, preserves the ordering, and includes c and d in the cycle. Call this operation of rerouting α .

By the inductive hypothesis, $\mathcal{H}_{k-1,2m}$ in $G_{k-1,2m}$ is such that for any two adjacent parts B_1 and B_2 in $G_{k-1,2m}$ there exists an edge of $\mathcal{H}_{k-1,2m}$ with one vertex in each of B_1 and B_2 . Because the number of parts is even, we can pair up adjacent parts (without the part containing v_{2k+1} and the two parts adjacent to these) and perform the rerouting operation α as explained in the preceding paragraph for the m-2 part pairs. The hamiltonian path $\mathcal{H}_{k,2m}$ satisfies condition (\star), concluding the proof.

Corollary 2.2. The graphs $G_{k,m}$ are (2k+1)-ordered for all $m \ge 4$.

Proof. Note that in the proof of Theorem 2.1 the assumption about the even number of parts was used only in Case 2.2 where we rerouted the cycle using α . To prove (2k + 1)-orderedness, we do not require the rerouting, thus $G_{k,m}$ is (2k + 1)-ordered for all $m \ge 4$.

Theorem 2.3. Any bracelet graph with at least 4 parts, at least k vertices in each part, and at least 2k + 1 total vertices in every pair of parts at distance 2 is (2k + 1)-ordered.

Proof. Let G be a bracelet graph with parts V_1, V_2, \ldots, V_m , such that each part has at least k vertices and there are at least 2k + 1 vertices in any two parts at distance 2. Let $v_1, v_2, \ldots, v_{2k+1}$ be any 2k + 1 specified vertices. We show that there exists a cycle in G containing $v_1, v_2, \ldots, v_{2k+1}$ in this order, and therefore G is (2k + 1)-ordered.

We will prove the statement by induction on k.

Base case: k = 1. It is not hard to see that any bracelet graph with at least 1 vertex per part and at least 3 vertices in any two parts at distance 2 is 3-ordered.

Inductive step.

Case 1. Suppose each part contains at most k marked vertices. Let $V'_i \subset V_i$ for all $i \in [m]$ such that each V'_i contains k vertices, and all of the marked vertices are in $V' = V'_1 \cup V'_2 \cup \cdots \cup V'_m$. As the graph induced by V' is isomorphic to $G_{k,m}$, it follows by Corollary 2.3 that we can find the desired cycle.

Case 2. Suppose there is a part B that contains k + l marked vertices, $l \ge 1$. As there are a total of 2k + 1 marked vertices, there can be only one such part.

Case 2.1. Suppose B contains 2k + 1 marked vertices. As the two adjacent parts contain at least 2k + 1 vertices, it is not hard to see that there is a cycle containing the 2k + 1 vertices in the specified order.

Case 2.2. Suppose B contains fewer than 2k + 1 marked vertices. Then there exists a different part B' such that $v_i \in B$ and $v_{i+1} \in B'$ (indices taken modulo 2k + 1) for some i. Without loss of generality $v_{2k} \in B$ and $v_{2k+1} \in B'$. In all of the other parts choose one vertex that is not marked (note that this is possible as all of the other parts contain at most k - 1 marked vertices), and call it a free vertex. Consider the graph H induced on the vertices of G without v_{2k}, v_{2k+1} , and the free vertices. This graph is (2k - 1)-ordered by the inductive hypothesis. Let \mathcal{D} be the cycle containing vertices $v_1, v_2, \ldots, v_{2k-1}$ in this order in H. Let \mathcal{C} be the cycle induced on v_{2k}, v_{2k+1} , and the free vertices.

Now we show how to construct a cycle in G containing the vertices $v_1, v_2, \ldots, v_{2k+1}$ in this order.

Case 2.2.1. Suppose v_{2k-1} and v_{2k} are in different parts. In this case the cycle in G is as follows. Follow \mathcal{D} in H from v_1 to v_{2k-1} . From v_{2k-1} go to a free vertex f in one of the adjacent parts, choosing the adjacent part so that either $f = v_{2k}$ or f is unmarked. If $f \neq v_{2k}$, then continue going to the free vertices in the not yet visited adjacent parts along the cycle \mathcal{C} so that we reach v_{2k} first and then v_{2k+1} . Continue along \mathcal{C} after meeting v_{2k+1} until we reach the free vertex of the part containing v_{2k-1} . After this, go to the vertex that is adjacent to v_{2k-1} in \mathcal{D} when going from v_{2k-1} to v_1 , and continue along \mathcal{D} until v_1 .

Case 2.2.2. Suppose v_{2k-1} and v_{2k} are in the same part. Denote the part containing v_{2k-1} and v_{2k} by B_1 , and denote the part containing v_{2k+1} by B_2 . Let u be the end vertex of the edge incident to v_{2k-1} that is an edge in the segment of \mathcal{D} from v_{2k-1} to v_1 .

Case 2.2.2.1 Suppose $u \notin B_2$. Follow \mathcal{D} in H from v_1 to v_{2k-1} . From v_{2k-1} go to an unmarked free vertex in the part containing u. From the unmarked free vertex in the part containing u go to v_{2k} , and after this continue going to the free vertices in the not yet visited adjacent parts along \mathcal{C} . During this we meet v_{2k+1} , and we continue along \mathcal{C} until we reach the free vertex of the part adjacent to the part of u. Go to u and continue on \mathcal{D} until v_1 .

Case 2.2.2.2. Suppose $u \in B_2$. Then B_1 and B_2 are adjacent. Follow \mathcal{D} in H from v_1 to v_{2k-1} . From v_{2k-1} go to an unmarked free vertex in the adjacent part that is not B_2 . From here go to v_{2k} , then to v_{2k+1} , then to the free vertex in the adjacent part and back to u. Continue on \mathcal{D} until v_1 .

Corollary 2.4. Any bracelet graph with at least 4 parts and at least k + 1 vertices in each part is (2k + 1)-ordered.

In Corollary 2.5 we cannot replace the requirement of each part having at least k+1 vertices with each part having at least k vertices. For a subset S of vertices let N(S) denote the set of vertices adjacent to some vertex in S, not including vertices in S. If $v_1, v_2, \ldots, v_{2k+1}$ are independent vertices (there are no edges between any of them), and if there is a cycle containing $v_1, v_2, \ldots, v_{2k+1}$ in this order then $|N(\{v_1, v_2, \ldots, v_{2k+1}\})| \ge 2k+1$. Consider the example of a bracelet graph G with three adjacent parts B_1, B_2 , and B_3 such that $|B_1| = |B_3| = k$ and $|B_2| = 2k+1$. Specify $v_1, v_2, \ldots, v_{2k+1}$ to be in B_2 . Then $v_1, v_2, \ldots, v_{2k+1}$ are independent, and then $|N(\{v_1, v_2, \ldots, v_{2k+1}\})| = 2k$, which shows that such a bracelet graph cannot be (2k+1)-ordered. We also cannot weaken the condition in Theorem 2.3 that every two parts at distance 2 have at least 2k+1 vertices total.

Lemma 2. Given two vertices u and v in a bracelet graph G with an even number of parts, the parity of the number of vertices on any path from u to v is the same. Moreover, any cycle in a bracelet graph G with an even number of parts has even length.

Although Lemma 2 is an easy observation, it will be a convenient tool for proving non-existence of bracelet graphs with certain properties.

Corollary 2.5. There is no k-ordered hamiltonian bracelet graph G with an even number of parts and an odd number of vertices.

3 The diameter of a *k*-ordered graph

In this section we give an upper bound on the diameter of k-ordered graphs, and we study the tightness of this bound. It has been observed by Denis Chebikin (personal communication) that if n is the number of vertices in a 4-ordered graph, then the diameter of the graph is at most $\frac{n}{4} + 2$. Indeed, as shown in the following proposition and corollary, a k-ordered graph has diameter not more than roughly $\frac{n}{k}$.

Proposition 3.1. Given a 2k-ordered graph G on n vertices, the diameter d of G is at most $\lfloor \frac{n-3}{2k} \rfloor + 2$.

Proof. Take two vertices a and b of G at distance d. As G is 2k-ordered, it is (2k - 1)-connected, so in particular both a and b have at least 2k - 1 neighbors. Let $a_1, a_2, \ldots, a_{k-1}$ be k - 1 distinct neighbors of a, and let $b_1, b_2, \ldots, b_{k-1}$ be k - 1 distinct neighbors of b. Note that the distance between a_i and b is at least d - 1 for all $i \in [k - 1]$, and likewise the distance between b_i and a is at least d - 1 for all $i \in [k - 1]$. Furthermore, the distance between a_i and b_j is at least d - 2 for all $i, j \in [k - 1]$. Consider any cycle containing the 2k vertices $a, b_1, a_1, b_2, a_2, \ldots, b_{k-1}, a_{k-1}, b$ in the specified order. It contains at least 2k + (d - 2) + (2k - 3)(d - 3) + (d - 2) + (d - 1) vertices, so $n \ge 2kd - 4k + 4$, and $d \le \frac{n}{2k} + 2 - \frac{4}{2k}$. The result follows.

Corollary 3.2. Given a (2k + 1)-ordered graph G on n vertices, the diameter d of G does not exceed $\lfloor \frac{n-3}{2k} \rfloor + 2$.

Proof. The statement follows from Proposition 3.1, as a (2k + 1)-ordered graph is 2k-ordered as well.

Theorem 3.3. There exist infinitely many (2k + 1)-ordered graphs on n vertices with diameter $\left|\frac{n-3}{2k}\right| + 1$ for all $k \ge 2$.

Proof. Consider graphs $G_{k,2m}$ as constructed in Section 2. In this case n = 2mk, and it is not hard to see that the diameter of $G_{k,2m}$ is $m = \lfloor \frac{2mk-3}{2k} \rfloor + 1$.

4 Low degree 2k-ordered graphs

In this section we focus on low degree 2k-ordered graphs. Since 2k-orderedness of a graph G implies (2k-1)-connectivity, and thus a 2k-ordered graph G has minimum degree at least 2k-1, a question of interest is the existence of (2k-1)-regular 2k-ordered graphs, or low degree 2k-ordered graphs in general. Since (2k+1)-ordered graphs are 2k-ordered as well, it follows from Theorem 2.1 that there is an infinite family of 2k-regular graphs that are 2k-ordered. In this section we present a stronger result (Theorem 4.3), exhibiting an infinite family of 2k-ordered bracelet graphs such that the minimum degree of these graphs is 2k-1, and the maximum degree is 2k+2. We also show that this result is the best possible for bracelet graphs.

Note that the construction in Theorem 2.1 is specific to 2k-regular graphs. For example, consider an analogue of the graphs $G_{k,m}$ that are (2k - 1)-regular. Note that in this case the number of vertices in all the parts cannot be the same. Thus, consider the graph $H_{k,4m}$ to be a graph with 4m parts having a repeating pattern of two parts with k - 1 vertices followed by two parts with k vertices. Note that $H_{k,4}$ is the bipartite graph $K_{2k-1,2k-1}$, which was shown in [11] to be 2kordered hamiltonian. However, if m > 1, then $H_{2,4m}$ is not 4-ordered, as it contains a square (4-cycle), and by [10] a 3-regular 4-ordered graph on more than 6 vertices cannot contain a square. In fact, it is easy to see that $H_{k,4m}$ is not 2k-ordered for any m > 1, because 2k-orderedness implies (2k - 1)-connectivity and the deletion of two non-adjacent parts of k - 1 vertices would disconnect $H_{k,4m}$.

Lemma 3. Let G be a 2k-ordered graph. If $s \le k$, then there exists no subset V_1 of the vertices of G such that $|V_1| = s$, $|V \setminus (N(V_1) \cup V_1)| \ge s$, and $|N(V_1)| < 2s$.

Proof. Suppose that G is a graph such that there is a subset V_1 of the vertices of G with $|V_1| = s$, $|V \setminus (N(V_1) \cup V_1)| \ge s$, and $|N(V_1)| < 2s$. Specify 2s vertices v_1, \ldots, v_{2s} such that for odd i, the v_i are in V_1 , and for even i, the v_i are in $V \setminus (N(V_1) \cup V_1)$. Then it would be impossible to have a cycle containing them in the specified order, because each vertex in V_1 would have to be adjacent to 2 distinct vertices in $N(V_1)$. Thus, G could not have been 2s-ordered, nor 2k-ordered for $k \ge s$. \Box

Corollary 4.1. Given a 2k-ordered bracelet graph G with more than 5 parts, there exists no part B in G such that $|B| \le k$ and N(B) < 2|B|, and there also exists no part B' in G such that |B'| > k and N(B') < 2k.

Proof. Consider 6 consecutive parts B_1, B_2, \ldots, B_6 of a 2k-ordered bracelet graph G. Since 2k-orderedness implies (2k - 1)-connectivity, the total number of vertices in parts B_4 and B_6 is at least 2k - 1.

Suppose that there exists a part B in G such that $|B| \leq k$ and N(B) < 2|B|, and let $B = B_2$ without loss of generality. Taking V_1 as described in Lemma 3 to be the vertices in B_2 it follows that G cannot be 2k-ordered, contradicting our assumption.

Suppose that there exists a part B' in G such that |B'| > k and N(B') < 2k, and let $B' = B_2$, without loss of generality. Taking V_1 as described in Lemma 3 to be some k vertices in B_2 it follows that G cannot be 2k-ordered, contradicting our assumption.

Proposition 4.2. There is no 2k-ordered bracelet graph with more than 6 parts that has minimum degree 2k - 1 and maximum degree less than 2k + 2.

Proof. Suppose that there exists a 2k-ordered bracelet graph G with more than 6 parts that has minimum degree 2k - 1 and maximum degree less than 2k + 2.

Claim. There is no part containing fewer than k-1 vertices.

Let B be a part with the minimum number of vertices, b, in bracelet graph G. Let B, B_1 , B_2 , B_3 , and B_4 be a sequence of 5 adjacent parts. Then B_2 contains 2k + 1 - b, 2k - b or 2k - 1 - b vertices, while B_4 contains b, b+1 or b+2 vertices (since B was a part with the minimum number of vertices). Thus, because G is (2k-1)-connected and there are at least 6 parts, $b+(b+2) \ge 2k-1$, implying that $b \ge k-1$.

Therefore, since there exists a vertex in G with degree 2k-1, there exist parts B' and B'_2 distance 2 apart, with k-1 and k vertices respectively. Applying Corollary 4.1 to B'_1 , the part between B' and B'_2 , we see that B'_1 also has k-1 vertices. Also, since the graph is (2k-1)-connected, every part other than B' and B'_1 must have at least k vertices.

Suppose part C contains k vertices, and C is not B'_2 or the part adjacent to B' (that is not B'_1). Let C_1 be the part following C, in the same direction that B'_1 follows B'. Choose $v_2, v_4, \ldots, v_{2k-2}$ in B', choose $v_1, v_3, \ldots, v_{2k-1}$ in B'_2 , and choose v_{2k} in C_1 . Then it is not hard to see that there can be no cycle visiting the 2k vertices in order, as there could be no path leading from v_{2k} to v_1 once we have traversed v_1, v_2, \ldots, v_{2k} . Thus, the only parts that might have fewer than k + 1 vertices are B', B'_1, B'_2 and the parts adjacent to B'. Since there are at least 7 parts, it follows that there is a vertex with degree at least 2k + 2, contradicting our assumptions.

By arguments analogous to those in Proposition 4.2, it is not hard to see that there is no (2k - 1)-regular 2k-ordered bracelet graph on more than 4 parts, and that there is no 2k-ordered bracelet graph with minimum degree 2k - 1 and maximum degree 2k on more than 5 parts. Also, the k-orderedness of bracelet graphs on at most 6 parts can be easily studied. In the following theorem we show that Proposition 4.2 is the best possible by exhibiting an infinite family of 2k-ordered graphs with minimum degree 2k - 1 and maximum degree 2k + 2.

Theorem 4.3. There exists an infinite family of 2k-ordered graphs $P_{k,m}$ with minimum degree 2k - 1 and maximum degree 2k + 2.

Proof. For $m \ge 5$, let $P_{k,m}$ be the bracelet graph with m parts, V_1, V_2, \ldots, V_m , satisfying $|V_1| = |V_2| = k - 1$, $|V_3| = k$, and $|V_i| = k + 1$ for all i > 3. We will show by induction on k that the graphs $P_{k,m}$ are 2k-ordered for $k \ge 2$ and $m \ge 5$.

Base case: k = 2.

Case 1. Suppose one of the parts contains 3 of the 4 marked vertices. Call this part B. Without loss of generality we may assume that it contains vertices v_1, v_2 , and v_3 . As there is a part adjacent to B with 3 vertices, it is possible to go from v_1 to v_3 through v_2 , meeting two vertices from that adjacent part, and then it is possible to go from v_3 by going through all the parts to v_4 and on to v_1 .

Case 2. Suppose one of the parts contains 2 marked vertices. Call such a part B. Without loss of generality we may assume that it contains vertices v_1 and v_2 or v_1 and v_3 .

Case 2.1. Suppose B contains vertices v_1 and v_2 .

Using one vertex (that is not v_3 or v_4) from an adjacent part with 3 vertices we can go from v_1 to v_2 , and then, regardless of where v_3 and v_4 are, it is not hard to see that the desired cycle exists.

Case 2.2. Suppose B contains vertices v_1 and v_3 . We can divide this case up into cases depending on whether v_2 or v_4 are in parts with 1 or 3 vertices, or whether they are both in the part with 2 vertices. We can easily find the desired cycles in each case.

Case 3. Suppose all four vertices are in different parts. Let $v_1 \in V_{i_1}$, $v_2 \in V_{i_2}$, $v_3 \in V_{i_3}$, and $v_4 \in V_{i_4}$. Without loss of generality we can suppose that $0 < i_2 - i_1 < i_3 - i_1 < i_4 - i_1$ or $0 < i_2 - i_1 < i_4 - i_1 < i_3 - i_1$ (as by symmetry we can rotate and reflect an ordered cycle), where we consider subtraction modulo m taking results between 0 and m - 1. In the case $0 < i_2 - i_1 < i_3 - i_1 < i_4 - i_1 < i_3 - i_1$, analysis shows the existence of the desired cycle.

Inductive step. Suppose the claim is true for all numbers less than k. We shall show that $P_{k,m}$ is still 2k-ordered for $m \geq 5$.

Suppose there is no part with all vertices marked or there is exactly one part with all vertices marked. Then we can find two different parts containing two consecutive vertices, v_{2k-1} and v_{2k} without loss of generality, such that all other parts have a free vertex. By arguments analogous to those in Theorems 2.1 and 2.3 we can show how to find a cycle in $P_{k,m}$ using the cycle in $P_{k-1,m}$.

- If there are exactly two parts with all vertices marked, then one of the following cases occurs:
- (i) both parts have exactly k-1 vertices,
- (ii) one part has exactly k-1 vertices and the other has exactly k vertices, or

(iii) one of the parts has exactly k-1 vertices and the other has exactly k+1 vertices.

If either case (ii) or case (iii) occurs, then there exist marked vertices v_i and v_{i+1} , one in either part with all vertices marked. The result then follows from arguments similar to those in Theorems 2.1 and 2.3. If, however, the two parts with all vertices marked both have k-1 vertices and there are no two vertices in them adjacent (if there are, then we are done by arguments as before), then, without loss of generality, the two parts with all vertices marked contain the vertices v_2, v_3, \ldots, v_k and $v_{k+2}, v_{k+3}, \ldots, v_{2k}$ respectively. It is not hard to see that in this case there exists a cycle containing v_1, v_2, \ldots, v_{2k} in this order, regardless of where v_1 and v_{k+1} are.

5 *k*-Ordered directed graphs

In this section we address the existence of low degree k-ordered directed graphs. This inquiry is motivated by the analogous questions for undirected graphs.

Consider a directed graph D, and denote its set of vertices by V(D). A directed graph D is said to be *strongly connected* if given any two vertices u and v in D, there exists a directed path from u to v. A vertex cut of a digraph D is a set $S \subset V(D)$ such that D - S is not strongly connected.

Proposition 5.1. If a directed graph D is k-ordered, then every vertex cut has at least k - 1 vertices.

The proof is analogous to the proof of the undirected case in [11].

Corollary 5.2. If a directed graph D is k-ordered, then for every vertex v we have indeg(v), $outdeg(v) \ge k-1$.

Theorem 5.3. For every $k \ge 2$ there exists an infinite family \mathcal{F}_k of (k-1)-diregular graphs that are k-ordered hamiltonian.

Proof. Given k, consider the undirected bracelet graphs $G_{k-1,l}$, $l \ge 3$, as defined in Section 2. We define the directed graph $\overrightarrow{G_{k-1,l}}$ on the same vertex set with parts V_1, V_2, \ldots, V_l , each with k-1

vertices. Edge \overrightarrow{uv} is in $\overrightarrow{G_{k-1,l}}$ if and only if $u \in V_i$ and $v \in V_{i+1}$ where indices are taken modulo l. The graph $\overrightarrow{G_{k-1,l}}$ is (k-1)-diregular, and we now prove that it is k-ordered hamiltonian.

Consider marked vertices v_1, v_2, \ldots, v_k . We will construct a hamiltonian cycle containing these vertices in this order. It is easy to see that there have to be two consecutive vertices among v_1, v_2, \ldots, v_k , without loss of generality v_1 and v_2 , that are in different parts. We can also suppose without loss of generality that $v_1 \in V_1$. Write the vertices of $\overrightarrow{G_{k-1,l}}$ as a grid where the i^{th} column contains the vertices in the i^{th} part for $i \in [l]$, and where the first row contains the two marked vertices v_1 and v_2 , and where the *j*th row contains v_{j+1} for $j = 2, 3, \ldots, k-1$. Then a hamiltonian cycle containing v_1, v_2, \ldots, v_k in this order is as follows. Start at v_1 going from left to right across the first row, then from the rightmost element in the first row go to the leftmost element in the second row, then go from left to right across the second row, and so on until reaching the lower rightmost vertex, from which we close off the hamiltonian cycle by going back to v_1 .

6 Connectivity, linkage, and *k*-edge-orderedness

Connectivity, linkage and k-orderedness appear to be related concepts. As noted in [4], a k-linked graph G is also k-ordered. Let f(k) be the minimum connectivity of a graph G that implies k-orderedness; the existence of the function f(k) has been shown in [4]. By a result in [1], 22k-connected graphs are k-linked, and thus are k-ordered as well, leading to the upperbound $f(k) \leq 22k$ observed in [4].

It is natural to pose analogous questions for edge-orderedness and directed graphs. In this section we consider edge-orderedness, while in the following sections we consider directed graphs.

A graph G is said to be *weakly k-linked* if, given 2k vertices (not necessarily distinct) $s_1, s_2, \ldots, s_k, t_1, t_2, \ldots, t_k$, there exist edge-disjoint paths from s_i to t_i , for $i \in [k]$.

Lemma 4. If a graph G is weakly 2k-linked, then G is k-edge-ordered.

Proof. Consider distinct edges e_1, \ldots, e_k . Let v_i and u_i be the end vertices of e_i . As G is weakly 2k-linked, there are edge-disjoint paths from v_1 to u_1, u_1 to v_2, \ldots, v_k to u_k , and u_k to v_1 . If for all i the path we chose between v_i and u_i is the edge $v_i u_i$, then we are done. If the path from v_i to u_i is not the edge e_i , but the edge e_i has not been used in any path, then we can just replace the path from v_i and u_i with the edge e_i . On the other hand, if the edge e_i has been used in some other path, then it has been used by exactly one of them, say p. In this case we can replace the edge e_i in path p by the path that was between v_i and u_i and we can replace the path from v_i to u_i with the edge e_i . Repeating this process as necessary, we obtain a tour containing e_1, \ldots, e_k in this order, and thus D is k-edge-ordered.

Let g(k) be the minimum edge-connectivity of a graph G that implies k-edge-orderedness of G.

Proposition 6.1. The upper bound $g(k) \leq 2k + 2$ holds.

Proof. It is known that (k + 2)-edge-connectivity implies weakly k-linked [8]. Thus, the statement of the proposition is a corollary of Lemma 6.1 and [8].

It is easy to see that $g(k) \ge k - 1$.

Lemma 5. If a graph G is 2k-edge-ordered, then it is weakly k-linked.

Proof. Consider vertices s_1, s_2, \ldots, s_k and t_1, t_2, \ldots, t_k . Since 2k-edge-ordered implies 2k-edge-connected [2], the degree of every vertex is at least 2k, and therefore there exist edges $s_1s'_1, s_2s'_2, \ldots, s_ks'_k, t_1t'_1, t_2t'_2, \ldots$ such that

$$s'_1, s'_2, \dots, s'_k, t'_1, t'_2, \dots, t'_k\} \cap \{s_1, s_2, \dots, s_k, t_1, t_2, \dots, t_k\} = \emptyset.$$

Then, as G is 2k-edge-ordered, it readily follows that it is also weakly k-linked.

Lemmas 6.1 and 6.3 exhibit a relation between weak linkage and edge-orderedness, but tightness of these lemmas remains uncertain.

7 Connectivity, diameter and orderedness

ł

In the previous section we considered the relation between connectivity and edge-orderedness in undirected graphs. In this section we pose the question: What connectivity implies k-(edge-)orderedness in directed graphs?

A digraph D is k-connected if every vertex cut has at least k vertices. The minimum size of a vertex cut is the connectivity of D. For $S, T \subset V(D)$, let [S, T] be the set of edges from S to T. An edge cut is the set $[S, \overline{S}]$ for some nonempty $S \subset V(D)$. A digraph D is k-edge-connected if every edge cut has at least k edges. The minimum size of an edge cut is the edge-connectivity of D ([13], Section 4).

It is not immediately clear that any connectivity in directed graphs implies any orderedness property. Indeed, it has been shown by Thomassen [12] that for every natural number k there exists a strongly k-connected digraph D_k containing two vertices not lying on a cycle. This implies that there is no connectivity that would guarantee even the 2-(edge-)orderedness of a directed graph. When the diameter is small, however, we can prove some positive results.

Theorem 7.1. If a digraph D is g(k)-edge-connected with diameter d and $g(k) \ge (2k-1)\lceil \frac{d}{2}\rceil + 1$, then D is k-edge-ordered.

Proof. Suppose that a digraph D is g(k)-edge-connected and it has diameter d. Let $e_1 = v_1 u_1$, $\ldots, e_k = v_k u_k$. As $g(k) \ge 1$, there exists a directed path between v_1 and u_1 not longer than d. Delete the edges of this path from D, which decreases the edge-connectivity by at most $\lceil \frac{d}{2} \rceil$. Note that D still has edge-connectivity at least 1, and indeed the edge-connectivity will allow us to repeat the same argument 2k times for paths from v_1 to u_1, u_1 to v_2, \ldots, v_k to u_k , and u_k to v_1 . At the last step the connectivity will be greater than or equal to $g(k) - (2k-1)\lceil \frac{d}{2}\rceil \ge 1$. Therefore, if $g(k) \ge (2k-1)\lceil \frac{d}{2}\rceil + 1$, then we can obtain an oriented tour through the vertices $v_1, u_1, \ldots, v_k, u_k$ in this order.

If for all *i* the directed path we chose between v_i and u_i is the edge $\overrightarrow{v_i u_i}$, then we are done. If the path from v_i to u_i is not the edge $\overrightarrow{v_i u_i}$, but the edge $\overrightarrow{v_i u_i}$ has not been used in any path, then we can just replace the path from v_i and u_i with the edge $\overrightarrow{v_i u_i}$. On the other hand, if the edge $\overrightarrow{v_i u_i}$ has been used in some other path, then it has been used by exactly one of them, say p. In this case we can replace the edge $\overrightarrow{v_i u_i}$ in path p by the path that was between v_i and u_i and we can replace the path from v_i to u_i with the edge $\overrightarrow{v_i u_i}$. Repeating this process as necessary, we obtain a tour containing e_1, \ldots, e_k in this order, and thus D is k-edge-ordered.

Theorem 7.2. If a digraph D is g(k)-connected with diameter $d \ge 1$, where $g(k) \ge (k-1)d$, then D is k-ordered.

Proof. Suppose that a digraph D is g(k)-connected and that it has diameter d. Choose marked vertices v_1, v_2, \ldots, v_k .

Since $g(k) \ge k - 1$, removing vertices v_3, v_4, \ldots, v_k would not disconnect D, thus there exists a directed path from v_1 to v_2 not containing v_3, v_4, \ldots, v_k and furthermore this path has length at most d. Delete the vertices of the path from v_1 to v_2 , except v_1 and v_2 . The connectivity decreases by at most d-1, and we can repeat the same process k times for paths from v_1 to v_2, v_2 to v_3, \ldots, v_k to v_1 . At the last step the connectivity will be greater than or equal to $g(k) - (k-1)(d-1) \ge k-1$, as required. Therefore, if $g(k) \ge (k-1)d$, then we can obtain a cycle through the vertices v_1, v_2, \ldots, v_k in this order. This shows k-orderedness.

Using analogous methods, similar results can be obtained for undirected graphs. Indeed, we get a bound $g(k) \ge (2k-1)d+1$ for a statement analogous to Theorem 7.1 for undirected graphs, and the same bound in the analogue of Theorem 7.2.

8 Conclusion

We conclude by giving an overview of questions motivated by this paper. In Section 2 we constructed an infinite family of 2k-regular (2k + 1)-ordered hamiltonian bracelet graphs, and in Section 3 we showed that there are no 2k-ordered bracelet graphs with minimum degree 2k - 1 and maximum degree less than 2k + 2. We constructed an infinite family of 2k-ordered braceket graphs with minimum degree 2k - 1 and maximum degree 2k + 2. The following question, however, remains open.

Question 8.1. Is there an infinite family of 2k-ordered hamiltonian bracelet graphs with minimum degree 2k - 1 and maximum degree 2k + 2 for all $k \ge 2$?

In Theorem 2.3 we gave a sufficient condition for a bracelet graph to be (2k + 1)-ordered. Note that Theorem 2.3 only applies when each part has at least k vertices. It is not hard to see from connectivity properties that any (2k + 1)-ordered bracelet graph with at least 5 parts has at most two parts with fewer than k vertices, and if it has two such parts then they must be adjacent.

Question 8.2. What are the necessary and sufficient conditions for a bracelet graph to be (2k+1)-ordered?

Naturally, one can also pose this question for k-ordered graphs in general.

In Section 3 we showed that a 2k-ordered graph has diameter at most $\lfloor \frac{n-3}{2k} \rfloor + 2$, where *n* is the number of the vertices of the graph. We have also shown that there exists an infinite family of (2k + 1)-ordered hamiltonian graphs that have diameter $\lfloor \frac{n-3}{2k} \rfloor + 1$. It is natural to pose the following question.

Question 8.3. Is there a (2k+1)-ordered or 2k-ordered graph that has diameter $\lfloor \frac{n-3}{2k} \rfloor + 2?$

Note that one can ask the analogues of these questions for directed graphs.

9 Acknowledgments

This research was performed at the University of Minnesota Duluth under the supervision of Professor Joseph A. Gallian. The author would like to thank Professor Gallian for his support and encouragement as well as Denis Chebikin, Philip Matchett and Melanie Wood for many useful suggestions. Financial support was provided by the Massachusetts Institute of Technology and the University of Minnesota Duluth.

References

- [1] B. Bollobás and A. Thomason, *Highly linked graphs*, Combinatorica **16** (1996), 313-320.
- [2] D. Chebikin, On k-edge-ordered graphs, Discrete Mathematics **281** (2004), 115-128.
- [3] G. Chen, R.J. Gould, F. Pfender, New conditions for k-ordered Hamiltonian graphs, Ars Combinatoria vol. 70 (2003), 245-255.
- [4] R.J. Faundree, Survey of results on k-ordered graphs, Discrete Mathematics 229 (2001), 73-87.
- [5] J.R. Faundree, R.J. Faundree, R.J. Gould, M.S. Jacobson, L. Lesniak, On k-ordered graphs, Journal of Graph Theory 35 (2000), 69-82.
- [6] R.J. Faudree, R.J. Gould, A.V. Kostochka, L. Lesniak, I. Schiermeyer, A. Saito, Degree conditions for k-ordered Hamiltonian graphs, Journal of Graph Theory 42 (2003), 199-210.
- [7] J.R. Faundree, R.J. Gould, F. Pfender, A. Wolf, *On k-ordered bipartite graphs*, The Electronic Journal of Combinatorics **10** (2003), R11.
- [8] A. Huck, A sufficient condition for graphs to be weakly k-linked, Graphs and Combinatorics 7 (1991), 323-351.
- [9] H.A. Kierstead, G.N. Sárközy, S.M. Selkow, On k-ordered Hamiltonian graphs, Journal of Graph Theory 32 (1999), 17-25.
- [10] K. Mészáros, On 3-regular 4-ordered graphs, submitted.
- [11] L. Ng, M. Schultz, k-Ordered Hamiltonian graphs, Journal of Graph Theory 24 (1997), 45-57.
- [12] C. Thomassen, Highly connected non-2-linked digraphs, Combinatorica 11 (1991), 393-395.
- [13] D. B. West, Introduction to Graph Theory, Prentice Hall, 1996.