# A new criterion for finite non-cyclic groups \*

Wei Zhou

College of Mathematics & Finance, Southwest China Normal University , Chongqing 400715

People's Republic of China

E-mail: zh\_great@hotmail.com

and

Wujie Shi

School of Mathematics, Suzhou University, Suzhou 215006

People's Republic of China

E-mail: wjshi@suda.edu.cn

and

Zeyong Duan

College of Mathematics & Finance, Southwest China Normal University , Chongqing 400715 People's Republic of China

#### Abstract

Let H be a subgroup of a group G. We say that H satisfies the power condition with respect to G, or H is a power subgroup of G, if there exists a non-negative integer msuch that  $H = G^m = \langle g^m | g \in G \rangle$ . In this note, the following theorem is proved: Let Gbe a group and k the number of non-power subgroups of G. Then (1) k = 0 if and only if G is a cyclic group(theorem of F. Szász) ;(2)  $0 < k < \infty$  if and only if G is a finite non-cyclic group; (3)  $k = \infty$  if and only if G is a infinite non-cyclic group. Thus we get a new criterion for the finite non-cyclic groups.

Keywords: power subgroup, cyclic group, Dedekind group.

2000MR subject classification: 20E07, 20E34, 20D25

#### 1. Introduction

Let H be a subgroup of group G. We say that H satisfies the power condition with respect to G, or H is a power subgroup of G if there exists a non-negative integer m such

<sup>\*</sup>Project supported by the National Natural Science Foundation(Grant No.10171074).

that  $H = G^m = \langle g^m | g \in G \rangle$ . On the other hand, if  $H \neq G^m$  for all m, we say that H is a non-power subgroup of G. If H is a power subgroup of G, the power exponent is the least non-negative integer m such that  $H = G^m$ .

It is obvious that a nontrivial group G has at least two trivial power subgroups:  $\{1\}$  and G itself, and the power exponents of the two subgroups are 0 and 1, respectively.

The power subgroups have some properties. For example, if H is a power subgroup of group G, then H is a full-invariant subgroup, in particular H is normal in G; and if H is a non-power subgroup of G, then a conjugate subgroup  $H^x(x \in G)$  is also a non-power subgroup of G.

The number of non-trivial power subgroups affects the structure of the group. In [2], F.Szász proved that G is a cyclic group if and only if all subgroups of G are power subgroups. That is, a non-cyclic group contains at least one non-power subgroup. In this paper, we generalize the above result of F. Szász and prove the following theorem:

**Theorem.** Let G be a group and k the number of non-power subgroups of G. Then

- (1) k = 0 if and only if G is a cyclic group;
- (2)  $0 < k < \infty$  if and only if G is a finite non-cyclic group;
- (3)  $k = \infty$  if and only if G is a infinite non-cyclic group.

**Remark 1.** The conclusion (1) of this theorem is the theorem of F.Szász in [2].

**Remark 2.** Note that for some k in the theorem, the group G may not exist. In [3],[4] the author proved that the case of k = 1 or k = 2 do not occur. But there are groups having exactly 3 non-power subgroups: For example the quaternion group  $Q_8$  of order 8, and  $Z_2 \times Z_2$  are groups possessing just 3 non-power subgroups.

From the theorem the following corollary gives a new criterion for finite non-cyclic groups.

**Corollary.** Suppose G is a non-cyclic group. Then G is finite if and only if G contains only finitely many non-power subgroups.

**Problem.** For any integer  $k(k \ge 3)$ , does there exist groups possessing just k non-power subgroups?

In the proof of the Theorem, we use the structure of the Dedekind group. A group is called Dedekind group if all its subgroups are normal. From Theorem 5.3.7 in [1], we know that G is a Dedekind group if and only if G is abelian or the direct product of a quaternion group of order 8, an elementary abelian 2-group and an abelian group with all its elements of odd order. Notation is standard and may be found for instance in[1]. In particular, we denote k the number of of non-power subgroups of G.

### 2. Some lemmas

**Lemma 1.** Let A be an abelian power subgroup in a group G. Then the set of power subgroups of G contained in A coincides with the set of power subgroups of A.

**Proof:** If G = A, then there it is nothing to prove. Let  $G \neq A$  and let l be the power exponent of A. Then l > 1 and  $A = G^l$ . Suppose that  $G^m \leq A$  for  $m \geq 1$  and let dbe the greatest common divisor of l, m. Then d = lr + ms for some integers r and s, and  $g^d = (g^l)^r (g^m)^s \in A$  for every  $g \in G$ , so  $G^d \leq A$ . It is obvious that  $A = G^l \leq G^d$ , so d = land l divides m. Hence  $m = ls_0$  for some  $s_0 \geq 0$ . Note that  $A = G^l = \langle g^l | g \in G \rangle$  is abelian. We get  $A^{s_0} = (G^l)^{s_0} = \langle g^l | g \in G \rangle^{s_0} = \langle g^{ls_0} | g \in G \rangle = G^m$ . On the other hand, if  $A^n$  is a power subgroup of A, then  $A^n = G^{ln}$  is a power subgroup of G. The lemma is proved.

**Lemma 2.** If A/N is a non-power subgroup of a factor-group G/N, then A is a non-power subgroup of G.

**Proof:** Suppose the result is false. That is, A is a power subgroup of G. Let  $A = G^m = \langle g_i^m | g_i \in G, i = 1, 2, \dots, \rangle$ . Since  $g_1^m \cdots g_r^m N = (g_1 N)^m \cdots (g_r N)^m$ ,  $A/N = (G/N)^m$ , a contradiction. Hence the lemma is proved.

Now we need a lemma about the structure of the cyclic group.

**Lemma 3.** If G is an infinite abelian group all of whose proper quotient groups are finite, then G is infinite cyclic.

**Proof:** Suppose G is not infinite cyclic. We choose an element  $a_0 \in G$  and  $a_0 \neq 1$ .  $\langle a_0 \rangle$  is a proper subgroup of G and  $\langle a_0 \rangle \neq 1$ . Then  $G/\langle a_0 \rangle$  is finite. So  $G/\langle a_0 \rangle$  is finitely generated and G is finitely generated. Then G is a direct product of finitely many cyclic groups of infinite or prime-power orders. Let  $G = \langle g_1 \rangle \times \langle g_2 \rangle \cdots \times \langle g_n \rangle$ , and n > 1 by the assumption. If there exists an i, such that  $|g_i| = \infty$ . Then  $G/\langle g_1 \rangle \times \cdots \times \langle g_{i-1} \rangle \times \langle g_{i+1} \rangle \times \cdots \times \langle g_n \rangle \cong \langle g_i \rangle$  is finite by the condition. That is impossible. So  $|g_i| < \infty, \forall i$ , which makes G be a finite group, a contradiction. We prove the lemma.

**Lemma 4.** If the number of cyclic subgroups of a group G is finite, then G is finite.

**Proof:** Firstly, we have G is a torsion group, otherwise suppose  $g \in G$  and  $|g| = \infty$ , then  $\langle g^n \rangle$  will be different cyclic subgroups in G with infinitely many different n, a contradiction.

For any element g in G, it can generate a cyclic subgroup  $\langle g \rangle$ . Suppose G have n cyclic subgroups and  $g_i(i = 1, \dots, n)$  be the generated elements of all the cyclic subgroups.  $G = \bigcup_{i=1}^n \langle g_i \rangle$ . So  $|G| \leq \sum_{i=1}^n |g_i| < \infty$ , that is to say G is finite.

## 3. Proof of the theorem

**Proof of the theorem:** We need only to prove the necessity of (2).

Case 1. G is abelian.

i) Suppose G is a torsion abelian group, we prove that G is a finite group.

Let  $\pi(G)$  be the set of all primes dividing the orders of elements of G. For every prime p, let  $G_p$  be the set of all p-elements in G. Then  $G_p$  is a subgroup of G, that is, the p-component of G. We claim  $|\pi(G)|$  is finite. If  $|\pi(G)| = \infty$ , then there must exist a prime  $p \in \pi(G)$  such that  $G_p$  is a power subgroup for there are finitely many non-power subgroups. Hence there exists a positive integer m such that  $G_p = G^m$ , and then  $G/G_p = G/G^m$ , which means the exponent of  $G/G_p$  is finite. But  $G/G_p \cong \prod_{q \in \pi(G) - \{p\}} G_q$ . So the exponent of  $G/G_p$  can not be finite since  $|\pi(G)|$  is infinite and then we get the contradiction.

If  $G_p$  is infinite for some p, then it contains infinitely many finite cyclic subgroups  $C_i$  and hence we may assume that  $C_i$  is a power subgroup in G and so  $G/C_i$  is of finite exponent, hence G is of finite exponent and G is a direct product of cyclic subgroups of finite order. In particular,  $G_p$  is of finite exponent and since  $G_p \simeq G/N$  for some subgroup N,  $G_p$  has only finitely many non-power subgroups in  $G_p$  by Lemma 2. Since  $G_p$  has only finitely many power subgroups, i.e.,  $G_p, G_p^p, \dots, G_p^{p^l}$ , where  $p^l$  is the power exponent of  $G_p$ , join with its finitely many non-power subgroups,  $G_p$  has only finitely many subgroups and so  $G_p$  is finite. Then for every  $p \in \pi(G)$ ,  $G_p$  is finite. So G is finite.

ii) Suppose G is a non-torsion abelian group. Then G is an infinite abelian group with some elements of infinite order. We prove G is cyclic, which is contrary to k > 0(k is the number of non-power subgroups of G).

Firstly, we prove that the subgroup T of G consisting of all elements of finite orders in G is finite. Evidently, no subgroup of T can be a power subgroup of G, otherwise G would be torsion and so, by i), G is finite, a contradiction. Hence, since has only finitely many non-power subgroups, we have that T contains only finitely many subgroups. Thus T have only finitely many cyclic subgroups and T is finite by Lemma 4.

Now  $T \neq G$  and  $\overline{G} = G/T$  is an infinite torsion-free group. By Lemma 2,  $\overline{G}$  has only finitely many non-power subgroups. We prove  $\overline{G}$  is cyclic.

We claim that  $\overline{G}/\overline{H}$  is finite for all non-trivial subgroup  $\overline{H}$  of  $\overline{G}$ . In fact, let  $\overline{H}$  be a nontrivial subgroup of  $\overline{G}$ . If  $\overline{H}$  is a power subgroup of  $\overline{G}$ , then there exists a non-negative integer m such that  $\overline{H} = \overline{G}^m$ . Now  $\overline{G}/\overline{H}$  is torsion, and it has only finitely many non-power subgroups by Lemma 2. Then  $\overline{G}/\overline{H}$  is finite by i). On the other hand, suppose  $\overline{H}$  is a non-power subgroup. Since  $\overline{G}$  is torsion-free with finitely many non-power subgroups, there must exist  $1 \neq \overline{H}_1 < \overline{H}$ , with  $\overline{H}_1$  being a power subgroup of  $\overline{G}$ . Then  $\overline{G}/\overline{H}_1$  is finite and then  $\overline{G}/\overline{H}$  is finite.

By Lemma 3,  $\overline{G}$  is cyclic. Hence  $G = T \times \langle z \rangle$  where z is of infinite order. It is obvious

that  $G^m = T^m \times \langle z^m \rangle$ , so if s = |T| > 1 then  $T \times \langle z^{sn} \rangle$  is a non-power subgroup of G for each  $n = 2, 3, \cdots$ . Thus T = 1 and so G is cyclic, a contradiction.

Case 2. G is non-abelian.

If G contains no non-power subgroups, then G is cyclic and every subgroup of a cyclic group is a power subgroup. So, suppose that G contains some non-power subgroups, and let  $H_1, \dots, H_s$  be all of those. Since every conjugate of  $H_i(i = 1, \dots, s)$  is also a non-power subgroup of G, so  $H_i$  has only finitely many conjugate subgroups. Then the normalizer  $N_i$ of  $H_i$  has a finite index in G and hence the subgroup  $K_i = \bigcap_{g \in G} g^{-1} N_i g$  is normal in G and has a finite index in G. Let  $K = \bigcap_{i=1}^s K_i$ , then K is a normal subgroup of finite index in G and so it normalizes every  $H_i$ . Observe that every subgroup of K is normal in K. In fact, if  $L \leq K$  and L is a power subgroup of G, then  $L \trianglelefteq G$  and hence  $L \trianglelefteq K$ ; if L is a non-power subgroup of G, then  $L = H_i$  for some  $i = 1, \dots, s$ , and by construction of K,  $H_i \trianglelefteq K$ . Thus K is a Dedekind group. So K is either abelian or a direct product of an abelian group and the quaternion group of order 8. Hence the center Z = Z(K) of K is of finite index in K and evidently  $Z \triangleleft G$ . Thus Z is an abelian normal subgroup of finite index, say m, in G. In particular,  $x^m \in Z$  for every  $x \in G$ . Let l be the smallest natural number such that  $x^l \in Z$ for every  $x \in G$ . Then  $G^l \leq Z$ .

Observe that  $G^l$  is a subgroup of finite index in Z. Otherwise, then  $Z/G^l$  is infinite and contains infinitely many proper non-trivial subgroups  $R_i/G^l$ ,  $i = 1, 2, \cdots$ . By assumption, there exists t such that  $R_t$  is a power subgroup of G, so  $R_t = G^r$  for some  $r \neq l$ . It is easy to get r < l, but this contradicts the choice of l.

Thus the index of  $G^l$  in Z is finite, so we may replace Z by  $G^l$  and assume that Z is an abelian power subgroup of finite index in  $G, Z = G^l$ .

Observe that, by Lemma 1, every power subgroup of G, being contained in Z, is a power subgroup of Z, and every power subgroup of Z is a power subgroup of G. In particular, Zcontains only finitely many non-power subgroups, and hence Z is finite or an infinite cyclic group. If Z is finite, then G is finite and so the result is proved. Thus we suppose Z is an infinite cyclic group, that is,  $Z = \langle x \rangle$  with  $|x| = \infty$ . If  $C_G(Z) \neq G$ , then there exists  $y \in G$  such that  $x^y = y^{-1}xy \neq x \in \langle x \rangle$ . Obviously  $Z^y = Z$ . Since x and  $x^{-1}$  are the only generators of  $\langle x \rangle$ , we have  $x^y = x^{-1}$ .

Let  $Y_i = \langle x^{-i}yx^i \rangle$ ,  $i = 1, 2, \cdots$ . We claim there are infinite many different  $Y_i$ . Suppose it is false. Let  $Y_{i_1}, \cdots, Y_{i_s}$  be all the different such type subgroups. Then there are infinite subgroups of this type equal to  $Y_{i_t}, 1 \leq t \leq s$ . Without any loss, we suppose  $Y_{i_t} = Y_{m_1} =$  $\cdots = Y_{m_r} = \cdots$ . If  $|y| = \infty$ , by the generator properties of infinite cyclic group we have  $y^{x^{m_1}} = (y^{x^{m_r}})^{\varepsilon_r}, r = 2, 3, \cdots$ , where  $\varepsilon_r = 1$  or -1. Hence there must exist  $r_0, r_1$  such that  $\varepsilon_{r_0} = \varepsilon_{r_1}$  and  $r_0 \neq r_1$ . And then  $y^{x^{m_{r_0}}} = y^{x^{m_{r_1}}}$ . Suppose  $m_{r_0} < m_{r_1}$  without any loss. Then we have  $y = y^{x^{m_{r_1}-m_{r_p}}}$ . But since  $x^y = x^{-1}$ , we get  $y^{x^{m_{r_1}-m_{r_p}}} = x^{-2(m_{r_1}-m_{r_0})}y$ . Therefore  $y = y^{x^{m_{r_1}-m_{r_p}}} = x^{-2(m_{r_1}-m_{r_0})}y$ , and then  $x^{-2(m_{r_1}-m_{r_0})} = 1$ . We get contradiction for  $|x| = \infty$ . On the other hand if |y| = n. Similarly there are infinite subgroups such that  $Y_{m_1} = \cdots = Y_{m_r} = \cdots$ . By the generator properties of cyclic group we have  $y^{x^{m_1}} = (y^{x^{m_r}})^{\varepsilon_r}, r = 2, 3, \cdots$ , where  $1 \leq \varepsilon_r \leq n$  is coprime to n. Hence there must exist  $r_0, r_1$  such that  $\varepsilon_{r_0} = \varepsilon_{r_1}$  and  $r_0 \neq r_1$ . And then  $y^{x^{m_{r_0}}} = y^{x^{m_{r_1}}}$ . We get a contradiction in the same way as before.

So there are infinite many different  $Y_i$  which are conjugate with each other. In particular, these infinite many  $Y_i$  are non-normal in G and non-power subgroups of G, which is contrary to the assumption. Thus  $C_G(Z) = G$  and hence  $Z \leq Z(G)$ . So the center of G has a finite index in G. By Schur Theorem ([1] Theorem 10.1.3), the commutator subgroup C = [G, G] is finite. If C = 1 then G is abelian contrary to the assumption. So let y be an element of prime order p in C. Obviously  $Z \cap \langle y \rangle = 1$  and  $\langle y \rangle \langle Z, y \rangle$  by  $Z \leq Z(G)$ , so  $\langle Z, y \rangle = Z \times \langle y \rangle$  and all subgroups  $Z^r \times \langle y \rangle$  are distinct for  $r = 1, 2, \cdots$ . Thus there exists r such that  $A = Z^r \times \langle y \rangle$  is a power subgroup of G. A contains, by Lemma 1, only a finitely many non-power subgroups of A and hence A is finite or cyclic. But this is not true. The theorem is proved.

Acknowledegement The authors would like to thank Prof. V.D. Mazurov for his help.

# References

- [1] D.J.S. Robinson, An Course in the Theory of Groups, Springer-Verlag, Berlin 1982
- [2] F.Szász, On cyclic groups, Fund. Math., 43(1956), 238-240
- [3] Wei Zhou Lai ping, A characterization of non-cyclic groups, J. Southwest China Normal Univ. (Natural Sci.)(in Chinese), 2000, 25(3), 213-216.
- [4] Wei Zhou, An investigation on the least bound of the number of non-exponent subgroups of non-cyclic groups, ICM 2002, Abstracts of Short Comm. and Posters, Higher Education Press, Beijing, p.40