

INTEGRAL BASES FOR CERTAIN TQFT-MODULES OF THE TORUS

KHALED QAZAQZEH

ABSTRACT. We find two bases for the lattices of the $SU(2)$ -TQFT-theory modules of the torus over given rings of integers. We use variant of the bases defined in [GMW] for the lattices of the $SO(3)$ -TQFT-theory modules of the torus. Moreover, we discuss the quantization functors (V_p, Z_p) for $p = 1$, and $p = 2$. Then we give concrete bases for the lattices of the modules in the 2-theory. We use the above results to discuss the ideal invariant defined in [FK]. The ideal can be computed for all the 3-manifolds using the 2-theory, and for all 3-manifolds with torus boundary using the $SU(2)$ -TQFT-theory. In fact, we show that this ideal in the $SU(2)$ -TQFT-theory is contained in the product of the ideals in the 2-theory and the $SO(3)$ -TQFT-theory under a certain change of coefficients, and it is equal in the case of a torus boundary.

INTRODUCTION

We let p denote an odd prime or twice an odd prime unless mentioned otherwise. Also, we let Σ denote a surface of genus g . Gilmer defined an integral TQFT-functor \mathcal{S}_p in [G1] based on the integrality results of the $SO(3)$ - and $SU(2)$ -invariants in [MR, M]. This is a functor that associates to a closed surface Σ , a module $\mathcal{S}_p(\Sigma)$ over a certain cyclotomic ring of integers \mathcal{O}_p . Moreover, Gilmer showed that these modules are free in the case of p is an odd prime. Gilmer and Masbaum constructed basis for $\mathcal{S}_p(\Sigma)$ and gave an independent proof of freeness in this case. In addition, Gilmer showed that these modules are projective where p is twice an odd prime. In this paper, we prove that the modules $\mathcal{S}_p(S^1 \times S^1)$ are free by constructing two explicit bases in the case that p is twice an odd prime. In the 2-theory, we prove also that the modules $\mathcal{S}_2(\Sigma)$ are free by constructing an explicit basis for any surface.

Frohman and Kania-Bartoszyńska in [FK] defined an ideal invariant of 3-manifolds with boundary using the $SU(2)$ -TQFT-theory that is hard to compute. In fact, they make use of another ideal that they defined to give an estimate for this ideal. However, Gilmer and Masbaum in [GM] computed an analogous ideal invariant using the $SO(3)$ -TQFT-theory for 3-manifolds that are obtained by doing surgery along a knot in the complement of another knot. The computations depend entirely on the fact that bases are constructed for the integral lattices of the $SO(3)$ -TQFT-theory modules [GMW, GM] of the torus. Also, Gilmer and Masbaum gave a finite set of generators for this ideal in general. Based on our results in this paper, we compute this ideal for the above 3-manifolds with torus boundary using the $SU(2)$ -TQFT-theory. Also, we introduce a formula to give an estimate for the ideal using

Date: 09/23/2005.

Key words and phrases. Integral basis, TQFT, ideal invariant.

the $SU(2)$ -TQFT-theory in terms of the ideals using the 2- and $SO(3)$ -TQFT-theories. In fact, the same formula can be used to compute this ideal using the $SU(2)$ -TQFT-theory for the all the above 3-manifolds with torus boundary.

In §1, we describe the $SO(3)$ - and $SU(2)$ -TQFT-functors using the approach of [BHMV3] over a variant ring depending on p . We review the integral TQFT-functors in §2 that Gilmer defined in [G1]. The first bases for the lattices of the $SU(2)$ -TQFT-modules are given in §3. We review the Frohman Kania-Bartoszyńska ideal in §4, and then we draw some conclusions based on the results of the previous section regarding this ideal. The quantization functors for $p = 1$ and $p = 2$ are discussed in §5, again following [BHMV3]. Also in this section, we give basis for $\mathcal{S}_2(\Sigma)$, and then draw some conclusions regarding the Frohman and Kania-Bartoszyńska ideal for this theory. We reformulate some results given in [BHMV3] in §6 to serve our need. Finally, we give another bases for the lattices of the $SU(2)$ -TQFT-modules in §7. The advantage of this one over the first basis is that it allows us to prove Theorem (7.9).

1. THE $SO(3)$ - AND $SU(2)$ -TQFTs

We consider the (2+1)-dimensional TQFT constructed as the main example of [BHMV3, P. 456] with some modifications. In particular, we use the cobordism category \mathcal{C} discussed in [G1, GQ] where the 3-manifolds have banded links but surfaces do not have colored points. Hence the objects are oriented surfaces with extra structure (Lagrangian subspaces of their first real homology). The cobordisms are equivalence classes of compact oriented 3-manifolds with extra structure (an integer weight) with banded links sitting inside of them. Two cobordisms with the same weight are said to be equivalent if there is an orientation preserving diffeomorphism that fixes the boundary.

Let

$$k_p = \begin{cases} \mathbb{Z}[A_p, \frac{1}{p}], & \text{if } p \equiv -1 \pmod{4}; \\ \mathbb{Z}[\alpha_p, \frac{1}{p}], & \text{if } p \equiv 1 \text{ or } 2 \pmod{4}. \end{cases}$$

Here and elsewhere A_p, α_p are ζ_{2p} and ζ_{4p} respectively for $p \geq 3$.

Now, we consider the TQFT-functor (V_p, Z_p) from \mathcal{C} to the category of finitely generated projective k_p -modules. The functor (V_p, Z_p) is defined as follows. $V_p(\Sigma)$ is a quotient of the k_p -module generated by all cobordisms with boundary Σ , and $Z_p(M)$ is the k_p -linear map from $V_p(\Sigma)$ to $V_p(\Sigma')$ (where $\partial M = -\Sigma \amalg \Sigma'$) induced by gluing representatives of elements of $V_p(\Sigma)$ to M along Σ via the identification map of the first component of the boundary.

If M is a closed cobordism, then $Z_p[M]$ is the multiplication by the scalar $\langle M \rangle_p$ defined in [BHMV3, §.2]. This invariant is normalized in two other ways. The first normalization of this invariant is $I_p(M) = \mathcal{D}_p \langle M_b \rangle_p$. Here and elsewhere M_b is the 3-manifold M with a reassigned weight zero, and $\mathcal{D}_p = \langle S_b^3 \rangle_p^{-1}$. The second normalization is $\theta_p(M) = \mathcal{D}_p^{\beta_1(M)} \langle M_b \rangle_p$, i.e $\theta_p(M) = \mathcal{D}_p^{\beta_1(M)} I_p(M)$.

If $\partial M = \Sigma$ and M is considered as a cobordism from \emptyset to Σ , then $Z_p(M)(1) \in V_p(\Sigma)$ is denoted by $[M]_p$ and called a *vacuum state* and it is *connected* if M is connected. Finally, note that V_p is generated over k_p by all vacuum states.

The modules $V_p(\Sigma)$ are free modules over k_p , and carry a nonsingular Hermitian bilinear form

$$\langle \cdot, \cdot \rangle_\Sigma : V_p(\Sigma) \times V_p(\Sigma) \rightarrow k_p,$$

given by

$$(1.1) \quad \langle [M_1], [M_2] \rangle_\Sigma = \langle M_1 \cup_\Sigma -M_2 \rangle_p.$$

Here $-M$ is the cobordism M with the orientation reversed and multiplying the integer weight by -1 , and leaving the Lagrangian subspace on the boundary the same.

Let $d_p = \lfloor \frac{p-1}{2} \rfloor$, it is known that d_p is the dimension of $V_p(S^1 \times S^1)$. One has that $V_p(S^1 \times S^1) \cong k_p[z]/I$ where the ideal I is generated by $e_{d_p} - e_{d_p-1}$ in the case of p is an odd prime and by e_{d_p} in the case of p is twice an odd prime (See [BHMV1] for more details). Thus indeed, $V_p(S^1 \times S^1)$ has a basis $\{e_0, \dots, e_{d_p-1}\}$ of rank d_p .

2. THE INTEGRAL COBORDISM FUNCTOR

Let \mathcal{C}' be the subcategory of \mathcal{C} consisting of the nonempty connected surfaces and connected cobordisms between them. Let \mathcal{O}_p be the ring of integers of the ring k_p defined before. The ring of integers is given by

$$\mathcal{O}_p = \begin{cases} \mathbb{Z}[A_p], & \text{if } p \equiv -1 \pmod{4}; \\ \mathbb{Z}[\alpha_p], & \text{if } p \equiv 1 \text{ or } 2 \pmod{4}. \end{cases}$$

Thus the ring of integers of k_p is a Dedekind domain.

Definition 2.1. For the surface Σ , we define $\mathcal{S}_p(\Sigma)$ to be the \mathcal{O}_p -submodule of $V_p(\Sigma)$ generated by all connected vacuum states.

If $M : \Sigma \rightarrow \Sigma'$ is a cobordism of \mathcal{C} , then $Z_p(M)([N]_p) = [M \cup_\Sigma -N]_p \in \mathcal{S}_p(\Sigma')$. Hence we obtain a functor from \mathcal{C}' to the category of \mathcal{O}_p -modules. These modules are projective as they are finitely generated torsion-free over Dedekind domains [G1, Thm. 2.5]. Also, these modules carry an \mathcal{O}_p -Hermitian bilinear form

$$(\ , \)_\Sigma : \mathcal{S}_p(\Sigma) \times \mathcal{S}_p(\Sigma) \rightarrow \mathcal{O}_p,$$

given by

$$(2.1) \quad ([M_1], [M_2])_\Sigma = \mathcal{D}_p \langle [M_1], [M_2] \rangle = \mathcal{D}_p \langle M_1 \cup_\Sigma -M_2 \rangle_p,$$

The value of this form always lies in \mathcal{O}_p by the integrality results for closed 3-manifolds in [MR, M].

If $R \subseteq \mathcal{S}_p(\Sigma)$ is an \mathcal{O}_p -submodule define

$$R^\# = \{v \in V_p(\Sigma) \mid (r, v)_\Sigma \in \mathcal{O}_p, \forall r \in R\},$$

then we can conclude

$$(2.2) \quad R \subseteq \mathcal{S}_p(\Sigma) \subseteq \mathcal{S}_p^\#(\Sigma) \subseteq R^\#.$$

Definition 2.2. A Hermitian bilinear form on a projective module over a Dedekind domain is called *non-degenerate* if the adjoint map is injective, and *unimodular* if the adjoint map is an isomorphism.

For our use, if the matrix of the form has a nonzero (unit) determinant, then the form will be non-degenerate (unimodular) respectively. Note that the determinant of the form (2.1) is nonzero from the fact that the form (1.1) is non-degenerate. Hence the form (2.1) is non-degenerate. In fact, we prove that the form (2.1) is unimodular for the 2-theory (discussed in §5) for all surfaces and for $S^1 \times S^1$ in the case of p is twice an odd prime.

A standard basis $\{u_\sigma\}$ for $V_p(\Sigma)$ is given (see [BHMV3]) in terms of p -admissible colorings σ of the spine of a handlebody of genus g whose boundary is Σ where the set of colors is $\{0, 1, 2, \dots, d_p - 1\}$, and the sum of the colors at a 3-vertex even and less than $2d_p$ in the case that p is twice an odd prime.

All of the above elements u_σ lie in $\mathcal{S}_p(\Sigma)$ when p is twice an odd prime. This follows from the fact that the quantum integers (denominators of the Jones-Wenzel idempotents) are units in \mathcal{O}_p (see Corollary 6.4). An admissible colored trivalent graph [BHMV3] is to be interpreted, here and elsewhere, as an \mathcal{O}_p -linear combination of links.

We say $a \sim b$ in \mathcal{O}_p if a/b is a unit in \mathcal{O}_p . The following proposition is an elementary fact from number theory that gives us a family of units in the ring \mathcal{O}_p .

Proposition 2.3 ([W]). *Suppose n has at least two distinct prime factors. Then $1 - \zeta_n$ is a unit in $\mathbb{Z}[\zeta_n]$.*

We make use of the following lemma in giving the first basis for $\mathcal{S}_p(S^1 \times S^1)$ in §3 whose proof will be in §6.

Lemma 2.4.

$$\mathcal{D}_p \sim \begin{cases} (1 - A_p^2)^{\frac{p-3}{2}}, & \text{if } p \text{ is an odd prime;} \\ \sqrt{2}(1 + \alpha_p^4)^{\frac{\frac{p}{2}-3}{2}}, & \text{if } p \text{ is twice an odd prime.} \end{cases}$$

Proposition 2.5. *The elements $\{u_\sigma\}$ are orthogonal with respect to the form (2.1). Moreover, we have*

$$(2.3) \quad (u_\sigma, u_\sigma)_\Sigma \sim \mathcal{D}_p^g.$$

Proof. If we use Theorem (4.11) in [BHMV3], and the facts

$$\langle S^3 \rangle_p = \mathcal{D}_p^{-1}, \quad \#v - \#e = 1 - g.$$

We obtain the result from knowing that all the quantum integers are units over \mathcal{O}_p ([GMW, Lem. 4.1], Corollary(6.4)), and using the definition of $(\ , \)_\Sigma$. \square

Definition 2.6. Let $[i]_p$ denote $\frac{A_p^{2i} - A_p^{-2i}}{A_p^2 - A_p^{-2}}$. This is called the i -th quantum integer.

We can describe the modules $\mathcal{S}_p(\Sigma)$ in terms of ‘mixed graph’ notation in a fixed connected 3-manifold M whose boundary is Σ . By a mixed graph, we mean a p -admissibly trivalent graph whose simple closed curves may be colored ω_p or an integer from the set $\{0, 1, \dots, p - 2\}$ where

$$\omega_p = \mathcal{D}_p^{-1} \sum_{i=0}^{d_p-1} (-1)^i [i + 1]_p e_i.$$

Using the surgery axiom (S2) in [BHMV3], we can choose this fixed 3-manifold to be a handlebody whose boundary is Σ . Thus we have

Proposition 2.7. *A mixed graph in a connected 3-manifold with boundary Σ represents an element in $\mathcal{S}_p(\Sigma)$. Moreover, $\mathcal{S}_p(\Sigma)$ is generated over \mathcal{O}_p by all the elements given by a mixed graph in a fixed handlebody whose boundary is Σ with the same genus.*

Proof. The first statement follows from that fact that V_p satisfies the second surgery axiom. The second statement follows from the fact that every 3-manifold with boundary Σ is obtained by a sequence of 2-surgeries to a handlebody of the same boundary and the definition of $\mathcal{S}_p(\Sigma)$. \square

3. THE FIRST BASIS FOR $\mathcal{S}_p(S^1 \times S^1)$

In this section, we assume r is an odd prime and $p = 2r$. We give a standard basis for $\mathcal{S}_p(S^1 \times S^1)$. We need the following lemma before we state our basis.

Definition 3.1. Let μ_i be the eigenvalue for the eigenvector e_i of the twist map on the Kauffman skein module of the solid torus. It is known in [BHMV1] that $\mu_i = (-1)^i A^{i(i+2)}$.

Lemma 3.2. For $i \neq j$, we have $\mu_i - \mu_j$ is equivalent to one of the following three cases up to a unit in \mathcal{O}_p .

- (1) 1 if $i \not\equiv j \pmod{2}$ and $(j-i)(i+j+2) \not\equiv 0 \pmod{r}$.
- (2) $\sqrt{2}$ if $i \not\equiv j \pmod{2}$ and $(j-i)(i+j+2) \equiv 0 \pmod{r}$.
- (3) $1 + \alpha_p^4$ if $i \equiv j \pmod{2}$.

Proof. Without loss of generality we can assume $0 \leq i < j \leq d_p - 1$. We have

$$\begin{aligned} \mu_i - \mu_j &= (-1)^i \alpha_p^{2i(i+2)} - (-1)^j \alpha_p^{2j(j+2)} \\ &= (-1)^i \alpha_p^{2i(i+2)} (1 - (-1)^{j-i} \alpha_p^{2j^2+4j-2i^2-4i}) \\ &\sim 1 - (-1)^{j-i} \alpha_p^{2(j-i)(i+j+2)}. \end{aligned}$$

Now we have three cases:

- (1) The hypothesis implies,

$$\mu_i - \mu_j \sim 1 - (-\alpha_p^{2(j-i)(i+j+2)})$$

is a unit by Proposition (2.3), as $-\alpha_p^{2(j-i)(i+j+2)}$ has order divisible by two distinct primes.

- (2) The hypothesis implies,

$$\mu_i - \mu_j \sim 1 \pm i \sim \sqrt{2}.$$

- (3) Finally the hypothesis implies that for some $k \leq r-1$,

$$\mu_i - \mu_j \sim 1 - \alpha_p^{8k} = (1 - \alpha_p^8)(1 + \alpha_p^8 + \dots + \alpha_p^{8(k-1)}) \sim (1 - \alpha_p^4)(1 + \alpha_p^4) \sim (1 + \alpha_p^4).$$

As $(1 + \alpha_p^8 + \dots + \alpha_p^{8(k-1)})$, and $(1 - \alpha_p^4)$ are units by [MR, Lem 3.1(ii)], and Proposition (2.3) respectively. \square

Proposition 3.3. The number of pairs (i, j) with $0 \leq i < j \leq r-2$ such that,

- (1) The second case of the above lemma holds is $\binom{r-1}{2}$.
- (2) The third case of the above lemma holds is $\binom{r-1}{2} \binom{r-3}{2}$.

Proof. To prove the first part, we look at all pairs (i, j) with $(j-i)(i+j+2) \equiv 0 \pmod{r}$ which automatically will satisfy $i \not\equiv j \pmod{2}$. This implies that $i+j+2 = r$. So we have $\binom{r-1}{2}$ -pairs of such (i, j) . Hence the first part follows. Now for every $0 \leq i \leq r-4$, there are $(\lceil \frac{r-3-i}{2} \rceil)$ - j 's such that $i \equiv j \pmod{2}$. Hence, we have $2(1+2+\dots+\frac{r-3}{2})$ -pairs of such (i, j) . Hence the second part follows. \square

Theorem 3.4. *Let $\mathcal{B}_p = \{t^i(\omega_p) \mid 0 \leq i \leq d_p - 1\}$. Then \mathcal{B}_p is a basis for $\mathcal{S}_p(S^1 \times S^1)$.*

Proof. We have

$$t^j(\omega_p) = \mathcal{D}_p^{-1} \sum_{i=0}^{d_p-1} (-1)^i [i+1] \mu_i^j e_i.$$

Let W be the matrix which expresses \mathcal{B}_p in terms of $\{e_0, e_1, \dots, e_{d_p-1}\}$. The determinant of W is a unit (product of $(-1)^i [i+1]$, see Corollary (6.4)) times $\mathcal{D}_p^{-d_p}$ times the determinant of the Vandermonde matrix $[\mu_i^j]$, where $0 \leq i, j \leq d_p - 1 = r - 2$. By the previous lemma

$$\det[\mu_i^j] = \pm \prod_{i < j} (\mu_i - \mu_j) \sim \sqrt{2}^{\binom{r-1}{2}} (1 + \alpha_p^4)^{\binom{r-1}{2} \binom{r-3}{2}}.$$

As $\mathcal{D}_p \sim \sqrt{2}(1 + \alpha_p^4)^{\binom{r-3}{2}}$, we conclude

$$(3.1) \quad \det W \sim (\sqrt{2})^{-\binom{r-1}{2}} (1 + \alpha_p^4)^{-\binom{r-1}{2} \binom{r-3}{2}}.$$

Let \mathcal{W} denote the \mathcal{O}_p -submodule of $\mathcal{S}_p(S^1 \times S^1)$ generated by \mathcal{B}_p . From the fact that the determinant of W is non-zero, we conclude that \mathcal{B}_p is linearly independent. Now by Proposition (2.5), we know $(e_i, e_i) \sim \mathcal{D}_p$. Therefore the determinant of the form (2.1) with respect to this orthogonal basis is $(\sqrt{2}(1 + \alpha_p^4)^{\binom{r-3}{2}})^{(r-1)}$. By equation (3.1) and the fact $1 + \alpha_p^4 \sim \overline{1 + \alpha_p^4} = 1 + \alpha_p^{-4}$, the determinant of the form (2.1) with respect to \mathcal{B}_p is a unit. We can conclude that the form on \mathcal{W} is unimodular. Hence $\mathcal{W} = \mathcal{W}^\sharp$ and so the set \mathcal{B}_p forms a basis for $\mathcal{S}_p(S^1 \times S^1)$ by equation (2.2). \square

Remark 3.5. This theorem and its proof are analogous to [GMW, Thm. 6.1] and its proof.

Corollary 3.6. *$\mathcal{S}_p(\Sigma)$ is generated by 3-manifolds (with no banded links) with boundary Σ .*

Proof. We expand the graph in every element in the Proposition (2.7) in terms of linear combinations of banded links (with some simple curves are colored ω_p). Then we replace any link component (that is not colored ω_p) by a linear combination from the set $\{t^i(\omega_p) \mid 0 \leq i \leq d_p - 1\}$. Hence the result follows by doing the required surgery on all the components of the link in every summand. \square

Remark 3.7. The above result is true if we replace p by an odd prime as a corollary of [GMW, Thm. 6.1].

4. THE FROHMAN KANIA-BARTOSZYNSKA IDEAL

We can apply the results from the previous section to compute the Frohman Kania-Bartoszyńska ideal using the $SU(2)$ -theory for special family of 3-manifolds with torus boundary. Before we do so, we review this ideal.

Definition 4.1. ([FK]) Let N be a 3-manifold with boundary, we define $\mathcal{J}_p(N)$ to be the ideal generated over \mathcal{O}_p by

$$\{I_p(M) \mid \text{where } M \text{ is a closed connected 3-manifold containing } N\}.$$

The importance of this ideal is in being an invariant of 3-manifolds (with boundary) and an obstruction to embedding as stated in the following propositions.

Proposition 4.2. ([FK]) *The ideal \mathcal{J}_p is an invariant of oriented 3-manifolds with boundary.*

Proposition 4.3. ([FK]) *If N_1, N_2 are an oriented compact 3-manifolds, and N_1 embeds in N_2 , then $\mathcal{J}_p(N_2) \subset \mathcal{J}_p(N_1)$.*

Remark 4.4. Frohman and Kania-Bartoszyńska defined this ideal using the $SU(2)$ -TQFT-theory. Afterwards, Gilmer defined this ideal using the $SO(3)$ -TQFT-theory and the 2-theory.

In general, it is not easy to compute this ideal because we have infinitely many closed connected 3-manifolds that contains N . Following his work with Masbaum in the case p an odd prime, Gilmer observed that $\mathcal{J}_p(N)$ is finitely generated based on his result that $\mathcal{S}_p(\Sigma)$ is finitely generated in the case p twice an odd prime as well. We give a finite set of generators for this ideal for any oriented compact 3-manifold using the $SU(2)$ -TQFT-theory which can be obtained by the following construction.

Definition 4.5. Assume L is an ordered link of two components K, J . Let N_L be the manifold obtained by doing surgery in S^3 along K in the complement of J .

Proposition 4.6.

$$\mathcal{J}_p(N_L) = \langle I_p(M_i) \mid 0 \leq i \leq d_p - 1 \rangle,$$

where M_i is the 3-manifold obtained by doing surgery along the component K and the component J with framing i in S^3 .

Proof. If p is an odd prime this was proved in [GM]. With the help of Theorem (3.4), the case p twice an odd prime follows in the same way. \square

5. THE QUANTIZATION FUNCTORS FOR $p = 1$, AND 2

In order to understand the relation between \mathcal{J}_r and \mathcal{J}_{2r} when r is an odd prime. We consider the theories associated to $p = 1$ and $p = 2$.

We begin by reviewing the quantization functor for $p = 1$ in detail. We start by listing the ring $k_1 = \mathbb{Z}$, and the surgery element $\Omega_1 = \omega_1 = 1$ for this theory defined in [BHMV1]. We also have $\kappa_1 = \mathcal{D}_1 = \theta_1 = 1$. One has $I_1(M) = \langle M \rangle_1 = \theta_1(M) = (-2)^{\#k}$ where $\#k$ is the number of components of the banded link in a closed 3-manifold M . Then (by [BHMV3, Prop. 1.1]) there exists a unique cobordism generated quantization functor (V_1, Z_1) that extends this invariant. In fact, this quantization functor can be described explicitly for surfaces as follows. $V_1(\Sigma)$ is the quotient of the \mathbb{Z} -module generated by all 3-manifolds (with banded links) with boundary Σ by the radical of the following form

$$\langle \cdot, \cdot \rangle_\Sigma : V_1(\Sigma) \times V_1(\Sigma) \rightarrow \mathbb{Z},$$

given by

$$\langle [M_1], [M_2] \rangle_\Sigma = \langle M_1 \cup_\Sigma -M_2 \rangle_1.$$

This module is isomorphic to \mathbb{Z} with any handlebody whose boundary is Σ as a generator. Hence if $M : \Sigma \rightarrow \Sigma'$, then $Z_1(M) : V_1(\Sigma) \rightarrow V_1(\Sigma')$ is the just the multiplication by $(-2)^{\#k}$.

Now we consider the quantization functor for $p = 2$. We start by introducing the ring and its ring of integers used in this theory

$$k_2 = \mathbb{Z}[\alpha_2, \frac{1}{2}] \quad \text{and} \quad \mathcal{O}_2 = \mathbb{Z}[\alpha_2].$$

The surgery element for this theory is $\omega_2 = \frac{1}{\sqrt{2}}\Omega_2$ where $\Omega_2 = 1 + \frac{\alpha_2}{2}$ defined in [BHMV1]. One has $\mathcal{D}_2 = \sqrt{2}$, and $\kappa_2 = \zeta_8$. Therefore the invariant of a closed connected 3-manifold M , which is obtained by doing surgery on S^3 along the link L , in terms of ω_2 is given by

$$(5.1) \quad \langle M \rangle_2 = \frac{1}{\sqrt{2}} \kappa_2^{\sigma(L)} \langle L(\omega_2) \rangle, \quad \text{where } \langle \rangle \text{ denotes the Kauffman bracket.}$$

From this formula, we can easily verify that

$$(5.2) \quad \langle M_1 \sharp M_2 \rangle_2 = \sqrt{2} \langle M_1 \rangle_2 \langle M_2 \rangle_2.$$

Now this invariant $\langle M \rangle_2$ defined in [BHMV3, §2] is involutive and extended to be multiplicative, hence (by [BHMV3, Prop. 1.1]) there exists a unique cobordism generated quantization functor that extends $\langle M \rangle_2$ which is denoted by (V_2, Z_2) . The modules $V_2(\Sigma)$ carry a Hermitian bilinear form defined as follows.

$$\langle \cdot, \cdot \rangle_\Sigma : V_2(\Sigma) \times V_2(\Sigma) \rightarrow k_2,$$

given by

$$\langle [M_1], [M_2] \rangle_\Sigma = \langle M_1 \cup_\Sigma - M_2 \rangle_2.$$

By [BHMV3, 1.5 and 6.3], $V_2(S^1 \times S^1)$ is generated by two elements each of which is a solid torus where the core is colored either 0 or 1. The pairing in terms of this basis is given by

$$\langle 1, 1 \rangle_{S^1 \times S^1} = \langle S^1 \times S^2 \rangle_2 = 1.$$

$$\langle 1, z \rangle_{S^1 \times S^1} = \langle (S^1 \times S^2, z) \rangle_2 = \frac{1}{\sqrt{2}} \langle H \rangle = \frac{1}{2} [2 + (-2)] = 0 = \overline{\langle z, 1 \rangle}_{S^1 \times S^1}.$$

Here H is the Hopf link with one of the components is colored ω_2 . Finally,

$$\langle z, z \rangle_{S^1 \times S^1} = \langle (S^1 \times S^2, z \sqcup z) \rangle_2 = \frac{1}{\sqrt{2}} \langle K \rangle = \frac{1}{2} [4 + 4] = 4.$$

Here K is the 3-chain link where the middle chain is colored ω_2 .

Hence the matrix of the form $\langle \cdot, \cdot \rangle_{S^1 \times S^1}$ in terms of this basis is given by

$$\begin{pmatrix} 1 & 0 \\ 0 & 4 \end{pmatrix}.$$

If we restrict this theory to the category of nonempty connected objects and connected cobordisms between them, then we have an integral cobordism theory as before. This follows from the fact $\langle \cdot \rangle_2$ is integral as stated in the proof of [MR, Thm. 1.1].

Definition 5.1. We define $\mathcal{S}_2(\Sigma)$ to be the \mathcal{O}_2 -submodule of $V_2(\Sigma)$ generated by all connected vacuum states, and we define an \mathcal{O}_2 -Hermitian bilinear form on $\mathcal{S}_2(\Sigma)$ given by $(\cdot, \cdot)_\Sigma = \sqrt{2} \langle \cdot, \cdot \rangle_\Sigma$.

Remark 5.2. One could similarly define $\mathcal{S}'_2(\Sigma)$ based on the invariant $\langle \cdot \rangle'_2$ defined in [BHMV3, §.1.B]. In this case, the basis $\{1, z\}$ for $V'_2(S^1 \times S^1)$ over k_2 is also a basis for $\mathcal{S}'_2(S^1 \times S^1)$ over \mathcal{O}_2 . However, this theory is not useful for us in this paper.

The above basis for $V_2(S^1 \times S^1)$ does not generate $\mathcal{S}_2(S^1 \times S^1)$. The following theorem gives a basis.

Theorem 5.3. *Assume that t is the twist map defined in [BHMV1], and $\mathcal{B} = \{\omega_2, t(\omega_2)\}$. Then \mathcal{B} is a basis for $\mathcal{S}_2(S^1 \times S^1)$, and the form is unimodular on $\mathcal{S}_2(S^1 \times S^1)$. Moreover, the matrix of the form defined in the previous definition in terms of \mathcal{B} is given by $\begin{pmatrix} \sqrt{2} & \frac{1-i}{\sqrt{2}} \\ \frac{1+i}{\sqrt{2}} & \sqrt{2} \end{pmatrix}$.*

Proof. Let ω_2 and $t(\omega_2)$ stands for the elements in the Kauffman skein module of the solid torus where the core is colored ω_2 and $t(\omega_2)$ respectively. From the definition we know that these two elements lie in $\mathcal{S}_2(S^1 \times S^1)$, hence $\mathcal{W} = \text{Span}_{\mathcal{O}_2} \mathcal{B} \subseteq \mathcal{S}_2(S^1 \times S^1)$. The matrix of the form $(\ , \)_{S^1 \times S^1}$ is given by $\begin{pmatrix} \sqrt{2} & 0 \\ 0 & 4\sqrt{2} \end{pmatrix}$, and since the matrix of \mathcal{B} in terms of $\{1, z\}$ is given by $\begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{2\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{2\sqrt{2}} \end{pmatrix}$. Then the matrix B of the form in terms of \mathcal{B} is given by

$$\begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{2\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{2\sqrt{2}} \end{pmatrix} \begin{pmatrix} \sqrt{2} & 0 \\ 0 & 4\sqrt{2} \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{2\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{2\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \sqrt{2} & \frac{1-i}{\sqrt{2}} \\ \frac{1+i}{\sqrt{2}} & \sqrt{2} \end{pmatrix}.$$

So the form restricted on \mathcal{W} has a unit determinant. Hence $\mathcal{W} = \mathcal{W}^\sharp$. Using equation (2.2), we get that \mathcal{W} is all of $\mathcal{S}_2(S^1 \times S^1)$. In conclusion, $\{\omega_2, t(\omega_2)\}$ is a basis for $\mathcal{S}_2(S^1 \times S^1)$. \square

Definition 5.4. Let $H_{i_1 i_2 \dots i_g}$ be the boundary connected sum of g solid tori where the core of the m -th torus is colored $i_m = 0$ or 1. Also, let

$$\mathcal{B} = \{H_{i_1 i_2 \dots i_g} \mid (i_1, i_2, \dots, i_g) \text{ is a } g\text{-tuple over } \{0, 1\}\}.$$

This set \mathcal{B} is an orthogonal basis for $V_2(\Sigma)$, and the pairing is described as follows:

Proposition 5.5. *The above set \mathcal{B} forms an orthogonal basis with respect to the form $\langle \ , \ \rangle_2$ given by*

$$\langle H_{i_1 i_2 \dots i_g}, H_{i_1 i_2 \dots i_g} \rangle_\Sigma = 4^k (\sqrt{2})^{g-1},$$

where $k = i_1 + i_2 + \dots + i_g$.

Proof. By [BHMV3, 1.5, and 6.3] \mathcal{B} is a basis. The result now follows from equation (5.2), and the computations for $V_2(S^1 \times S^1)$ after that equation. \square

We can describe $\mathcal{S}_2(\Sigma)$ as the \mathcal{O}_2 -submodule of $V_2(\Sigma)$ generated by all 3-manifolds with boundary Σ and links sitting inside of them. As $z = 2\sqrt{2}\omega_2 - 2$, one has a similar result to Corollary (3.6) for this theory.

Definition 5.6. Let $H'_{i_1 i_2 \dots i_g}$ be the boundary connected sum of g solid tori where the core of the m -th torus is colored $t^{i_m}(\omega_2)$ for $i_m = 0$, or 1.

Also, let

$$\mathcal{B}' = \{H'_{i_1 i_2 \dots i_g} \mid (i_1, i_2, \dots, i_g) \text{ is a } g\text{-tuple over } \{0, 1\}\}.$$

Theorem 5.7. *The above set \mathcal{B}' forms a basis for $\mathcal{S}_2(\Sigma)$.*

Proof. Let $(S^1 \times S^2)_{ij}$ denote $S^1 \times S^2$ formed by gluing two solid tori whose cores are colored $t^i(\omega_2)$, and $t^j(\omega_2)$ where $i, j \in \{0, 1\}$. Let us look at the pairing

$$\begin{aligned} (H'_{i_1 i_2 \dots i_g}, H'_{j_1 j_2 \dots j_g})_\Sigma &= \sqrt{2} \langle H'_{i_1 i_2 \dots i_g}, H'_{j_1 j_2 \dots j_g} \rangle_\Sigma \\ &= \sqrt{2} \langle \#_{k=1}^g (S^1 \times S^2)_{i_k j_k} \rangle_2 \\ &= \sqrt{2}^g \prod_{k=1}^g \langle (S^1 \times S^2)_{i_k j_k} \rangle_2 \\ &= \prod_{k=1}^g ((S^1 \times S^2, t^{i_k}(\omega_2) \sqcup t^{j_k}(\omega_2))_2). \end{aligned}$$

With a natural order, the matrix of the form in terms of this set is given by $\otimes^g B$ (B is defined in the proof of the previous theorem). This implies that the determinant of this form is a unit. By a similar argument as in the proof of Theorem (5.3), the module generated by this set is all of $\mathcal{S}_2(\Sigma)$. \square

We define $I_2(M) = \sqrt{2} \langle M_b \rangle_2$ for a closed 3-manifold M where $\langle M \rangle_2$ as defined in Equation (5.1). Also we define the Frohman Kania-Bartoszynska ideal \mathcal{J}_2 just as in the previous section. Now we can compute this ideal easily for all 3-manifolds using the 2-theory by making use of above results. For example, we confirm a result of Gilmer and prove it using our basis.

Proposition 5.8. [G1, Prop.15] *Let N_L, L as defined in Definition (4.5). Also, let l be the linking number between K and J , and k is the framing of K . If l is odd then $\mathcal{J}_2(N_L) = \mathcal{O}_2$. If l is even, then we have the following:*

$$\mathcal{J}_2(N_L) = \begin{cases} (\sqrt{2}), & \text{if } k \equiv 0 \pmod{4}; \\ 0, & \text{if } k \equiv 2 \pmod{4}; \\ \mathcal{O}_2, & \text{if } k \equiv 1 \text{ or } 3 \pmod{4}. \end{cases}$$

Proof. From Theorem (5.3), we know that $\mathcal{J}_2(N_L)$ is generated by two elements. In fact, it is generated by $I_2(M_i)$ where M_i is the 3-manifold obtained by doing surgery along the component K and the component J with framing 0 or 1 in S^3 . Now, we use the formula in [BHMV2, Cor.2.4] to compute the two generators of this ideal. The computations shows that (note that κ_2 is a unit in \mathcal{O}_2)

$$\mathcal{J}_2(N_L) = \langle \frac{1}{2}(i^{k+2l} + i^k + 2), \frac{1}{2}(i^{k+2l+2} + i + i^k + 1) \rangle,$$

Now if we consider all the possibilities, we obtain the required result. \square

Also, we compute this ideal for all 3-manifolds N_K that are obtained by doing surgery on a knot K in the complement of a tubular neighborhood of an eyeglass graph:0 – 0 in S^3 .

Proposition 5.9. *Let l_1 and l_2 be the linking numbers of K with the first and the second loops in the eyeglass respectively, and k is the framing of K . Then we have*

$$\mathcal{J}_2(N_K) = \begin{cases} (\sqrt{2}), & \text{if } l_1 \equiv l_2 \equiv 0 \pmod{2} \text{ and } k \equiv 0 \pmod{4}; \\ 0, & \text{if } l_1 \equiv l_2 \equiv 0 \pmod{2} \text{ and } k \equiv 2 \pmod{4}; \\ \mathcal{O}_2, & \text{if any of } l_1, l_2, k \text{ is odd.} \end{cases}$$

Proof. Let m be the linking number between the loops. From Theorem (5.7), we know $\mathcal{J}_2(N_K)$ is generated by four elements. In fact, it is generated by $I_2(M_{i,j})$ where $M_{i,j}$ is the 3-manifold obtained by doing surgery along the component K and the loops with framing $i, j = 0$ or 1 in S^3 . As in the proof of the previous proposition, one sees

$$\begin{aligned} \mathcal{J}_2(N_K) = \langle & \frac{1}{2\sqrt{2}}(3 + i^k + i^{2m} + i^{k+2l_1} + i^{k+2l_2} + i^{k+2l_1+2l_2+2m}), \\ & \frac{1}{2\sqrt{2}}(2 + i + i^k + i^{2m+1} + i^{k+2l_1} + i^{k+2l_2+1} + i^{k+2l_1+2l_2+2m+1}), \\ & \frac{1}{2\sqrt{2}}(2 + i + i^k + i^{2m+1} + i^{k+2l_1+1} + i^{k+2l_2} + i^{k+2l_1+2l_2+2m+1}), \\ & \frac{1}{2\sqrt{2}}(1 + 2i + i^k + i^{2m+2} + i^{k+2l_1+1} + i^{k+2l_2+1} + i^{k+2l_1+2l_2+2m+2}) \rangle. \end{aligned}$$

If we take all possibilities, we get the required result. \square

6. RELATING THE r -TH AND $2r$ -TH THEORIES

From now on, we assume that r is an odd prime and $p = 2r$.

Remark 6.1. The results of this section are slight variations of results of [BHMV3, §.6] and [BHMV2, §.2]. The ring k_p is not exactly the same as the ring denoted this way in [BHMV3].

The ring k_p will be considered as a k_2 (or a k_r)-module via the homomorphisms defined below. The following is a slight variation of the maps defined in [BHMV2, §.2].

Lemma 6.2. *There are well-defined ring homomorphisms $i_r : k_2 \rightarrow k_p$, $j_r : k_r \rightarrow k_p$ given by*

$$\begin{aligned} i_r(\alpha_2) &= \alpha_p^{r^2}, \quad j_r(\alpha_r) = \alpha_p^{1+r^2} \text{ for } r \equiv 1 \pmod{4}, \text{ and} \\ j_r(A_r) &= A_p^{1+r^2} \text{ for } r \equiv -1 \pmod{4}. \end{aligned}$$

We need the following remark to prove that these maps are well-defined.

Remark 6.3. If α is a primitive n -th root of unity, then α^m is a primitive $\frac{n}{\gcd(n,m)}$ -th root of unity.

Proof. To prove that the map i_r is a well-defined ring homomorphism, we show $\alpha_p^{r^2}$ is a primitive $8r$ -th root of unity. This is true, as $\gcd(8r, r^2) = r$ and α_p is a primitive $8r$ -th root of unity. Similarly for j_r but we consider two cases:

- (1) For $r \equiv 1 \pmod{4}$, we have $\alpha_p^{1+r^2}$ is a primitive $4r$ -th root of unity, as $\gcd(8r, 1+r^2) = 2$ and α_p is a primitive $8r$ -th root of unity.
- (2) For $r \equiv -1 \pmod{4}$, we have $A_p^{1+r^2}$ is a primitive $2r$ -th root of unity, as $\gcd(4r, 1+r^2) = 2$ and A_p is a primitive $4r$ -th root of unity.

\square

Corollary 6.4. *The quantum integers $[i]_p, 1 \leq i \leq d_p$ are units in \mathcal{O}_p .*

Proof. We know that the quantum integers $[i]_r$ for $1 \leq i \leq r-1$ are units in the \mathcal{O}_r see [GMW, Lem. 4.1(iii)] and [MR, Lem. 3.1(ii)]. So we conclude that $[i]_p$ are units for $1 \leq i \leq r-1 = d_p$, as

$$j_r([i]_r) = (-1)^i [i]_p, \quad \forall 1 \leq i \leq d_p,$$

since

$$j_r(A_r^2) = -A_p^2.$$

□

Given any k_2 (or k_r)-module, we can define a k_p -module by tensoring the original module with k_p over k_2 (or k_r) respectively. We let $\widehat{V}_2(\Sigma)$ (or $\widehat{V}_r(\Sigma)$) be the k_p -module obtained in this way. We give a relation between $V_1, \widehat{V}_2, \widehat{V}_r$, and V_{2r} for any surface Σ , but before that we need the following slight reformulation of [BHMV2, Thm. 2.1].

Theorem 6.5. *For any closed 3-manifold M with possibly a banded link sitting inside of it we have,*

$$(6.1) \quad I_1(M)I_{2r}(M) = i_r(I_2(M))j_r(I_r(M)).$$

Proof. Theorem (2.1) in [BHMV2] states the following:

$$(6.2) \quad \theta_1(M)\theta_{2r}(M) = i_r(\theta_2(M))j_r(\theta_r(M)).$$

Letting $M = S^3$, we get

$$\mathcal{D}_{2r}^{-1} = i_r(\mathcal{D}_2^{-1})j_r(\mathcal{D}_r^{-1}),$$

as $\theta_1(S^3) = 1$. Now multiply both sides of equation (6.2) by $\mathcal{D}_{2r}^{-\beta_1(M)}$, and replace it by $i_r(\mathcal{D}_2^{-\beta_1(M)})j_r(\mathcal{D}_r^{-\beta_1(M)})$ in the right hand side. Then the result follows from the relation between θ and I . □

We let κ_n to be an element that plays the role of κ^3 in [BHMV3]. We define this element as follows:

$$\kappa_n = \begin{cases} \alpha_{4n}^{-6 - \frac{n(n+1)}{2}}, & \text{if } n \text{ is an odd prime;} \\ -\alpha_{4n}^{-6 - \frac{n(n+1)}{2}}, & \text{if } n \text{ is twice an odd prime.} \end{cases}$$

Changing the weight by one multiplies the invariant $\langle \rangle_n$ by κ_n .

Lemma 6.6. *For the above ring homomorphisms. We have*

$$\kappa_p = i_r(\kappa_2)j_r(\kappa_r)$$

Proof. We have

$$i_r(\kappa_2)j_r(\kappa_r) = i_r(\alpha_2)j_r(\alpha_r^{-6 - \frac{r(r+1)}{2}}) = \alpha_p^{r^2}(\alpha_p^{-6 - \frac{r(r+1)}{2}})^{1+r^2} = -\alpha_p^{-6 - \frac{2r(2r+1)}{2}} = \kappa_p,$$

as

$$r^2 + (-6 - \frac{r(r+1)}{2})(1+r^2) \equiv -6 - \frac{2r(2r+1)}{2} + 4r \pmod{8r}$$

□

We are now able to give the proof of a result used in §2.

Proof of Lemma 2.4. The first case follows from [GMW, Lem. 4.1(ii)]. The second case follows from the following facts from the proof of Corollary (6.4) and from Theorem (6.5).

- (1) $\mathcal{D}_p = \mathcal{D}_{2r} = i_r(\mathcal{D}_2)j_r(\mathcal{D}_r)$, where $\mathcal{D}_2 = \sqrt{2}$.
- (2) $j_r(A_r^2) = -A_{2r}^2 = -\alpha_{2r}^4$.

□

Theorem 6.7. *There is a natural k_p -isomorphism $F : V_1(\Sigma) \otimes V_p(\Sigma) \rightarrow \widehat{V}_2(\Sigma) \otimes \widehat{V}_r(\Sigma)$ such that*

$$(6.3) \quad F([M]_1 \otimes [M]_p) = [M]_2 \otimes [M]_r,$$

where M is a 3-manifold with banded link (but not linear combination of links) sitting inside of it.

Corollary 6.8. *The map in the previous theorem defines a k_p -isomorphism between $V_p(\Sigma)$, and $\widehat{V}_2(\Sigma) \otimes \widehat{V}_r(\Sigma)$.*

To prove this theorem, we use the following version of [BHMV3, Lemm. 6.4].

Lemma 6.9. *Let \mathcal{V}, W be free modules over an integral domain R (with involution) equipped with Hermitian sesquilinear forms $\langle \cdot, \cdot \rangle_{\mathcal{V}}, \langle \cdot, \cdot \rangle_W$, and let $F : \mathcal{V} \rightarrow W$ be a form-preserving linear map. Let $(V, \langle \cdot, \cdot \rangle_V)$ be the quotient of \mathcal{V} by the radical of $\langle \cdot, \cdot \rangle_{\mathcal{V}}$. Suppose that $\langle \cdot, \cdot \rangle_W$ is non-degenerate. Suppose V and W are free of finite rank and $\langle \cdot, \cdot \rangle_V$ is unimodular and furthermore that $\text{rank}(W) \leq \text{rank}(V)$. Then F induces an isometry $\langle \cdot, \cdot \rangle_V \rightarrow \langle \cdot, \cdot \rangle_W$.*

Proof of Theorem 6.7. It follows from Theorem (6.5) and Lemma (6.6) that formula (6.3) defines a form-preserving linear map. We know already that the form on $\widehat{V}_2(\Sigma) \otimes \widehat{V}_r(\Sigma)$ is non-degenerate. Finally, we have two cases namely,

- If $r = 1$, then F is just flipping the tensors. Hence, it is an isometry
- If $r \geq 3$, then the result follows from the fact that $\text{rank}(V_1(\Sigma) \otimes V_{2r}(\Sigma)) = \text{rank}(\widehat{V}_2(\Sigma) \otimes \widehat{V}_r(\Sigma))$, and the second part of the lemma.

□

7. THE SECOND BASIS FOR $\mathcal{S}_p(S^1 \times S^1)$

We give new basis for $\mathcal{S}_r(S^1 \times S^1)$ that will be used in constructing another basis for $\mathcal{S}_p(S^1 \times S^1)$. To do so, we need the following lemma.

Lemma 7.1. *If $0 \leq i < j \leq \frac{r-3}{2}$, then the twist coefficients satisfy $\mu_i^2 - \mu_j^2 \sim \mu_i - \mu_j \sim 1 - A_r^2$.*

Proof. Notice that $\mu_i = q_r^{i^2+2i}$ where q_r denotes the primitive r -th root of unity given by $-A_r$.

$$\begin{aligned} \mu_i^2 - \mu_j^2 &= (\mu_i + \mu_j)(\mu_i - \mu_j) = (q_r^{i^2+2i} + q_r^{j^2+2j})(\mu_i - \mu_j) \sim \\ & (1 + q_r^{(j-i)(j+i+2)})(\mu_i - \mu_j) \sim \mu_i - \mu_j \sim 1 - A_r^2. \end{aligned}$$

We used the result of the fourth part of [GMW, Lem. (4.1)] in the last equality up to a unit. Also in the one next to last, we used the fact that $1 + q_r^{(j-i)(j+i+2)}$ is a unit by [MR, Lem. (3.1)] as $\text{gcd}(r, (j-i)(j+i+2)) = 1$. □

Theorem 7.2. *Let $\mathcal{B}_{2r} = \{t^{2i}(\omega_r) \mid 0 \leq i \leq d_r - 1\}$, and $\mathcal{B}_{2r+1} = \{t^{2i+1}(\omega_r) \mid 0 \leq i \leq d_r - 1\}$. Then \mathcal{B}_{2r} and \mathcal{B}_{2r+1} form bases for $\mathcal{S}_r(S^1 \times S^1)$.*

Proof. The proof of [GMW, Thm. 4.1] now goes through with μ_i^2 playing the role of μ_i and t^{2j} playing the role of t^j . We use the previous lemma when appropriate to obtain that \mathcal{B}_{2r} is a basis. To prove that \mathcal{B}_{2r+1} is a basis, we use the fact that the twist map t is an isomorphism of the Kauffman skein module of the solid torus. \square

Notation: We use the notation $\widehat{\mathcal{S}}_i(S^1 \times S^1) = \mathcal{S}_i(S^1 \times S^1) \otimes_{k_i} k_p$ for $i = 2$ or r .

Definition 7.3. Let

$$\delta_i = \begin{cases} 0, & \text{if } i + p \equiv 2 \text{ or } 3 \pmod{4}; \\ 1, & \text{if } i + p \equiv 0 \text{ or } 1 \pmod{4}. \end{cases}$$

We defined δ_i so that the following two lemmas hold.

Lemma 7.4. *If $i + \delta_i p \equiv 0 \pmod{2}$, then $i + \delta_i p \equiv 0 \pmod{4}$.*

Proof. We know that $p \equiv 2 \pmod{4}$, now we have two cases to consider

- If $\delta_i = 0$, then $i + p \equiv 2 \pmod{4}$ as i and p are even. So we conclude $i + \delta_i p = i \equiv 0 \pmod{4}$.
- If $\delta_i = 1$, then $i + p \equiv 0 \pmod{4}$ as $i + p$ is even. So we conclude $i + \delta_i p \equiv i + p \equiv 0 \pmod{4}$.

\square

Lemma 7.5. *If $i + \delta_i p \equiv 1 \pmod{2}$, then $i + \delta_i p \equiv 1 \pmod{4}$.*

A similar proof can be given for this lemma. The following theorem gives another basis for $\mathcal{S}_p(S^1 \times S^1)$.

Theorem 7.6. *Let $\mathcal{B}_p = \{t^{i+\delta_i p}(\omega_p) \mid 0 \leq i \leq d_p - 1\}$. Then \mathcal{B}_p is a basis for $\mathcal{S}_p(S^1 \times S^1)$.*

Proof. We have that $\text{Span}_{\mathcal{O}_p} \mathcal{B}_p \subseteq \mathcal{S}_p(S^1 \times S^1)$, and $F(1 \otimes \mathcal{S}_p(S^1 \times S^1)) \subseteq \widehat{\mathcal{S}}_2(S^1 \times S^1) \otimes \widehat{\mathcal{S}}_r(S^1 \times S^1)$ where F is the map defined in formula (6.3). It is enough now to show that $F(1 \otimes \mathcal{B}_p)$ generates $\widehat{\mathcal{S}}_2(S^1 \times S^1) \otimes \widehat{\mathcal{S}}_r(S^1 \times S^1)$ which implies that $F(1 \otimes \mathcal{S}_p(S^1 \times S^1)) \subseteq \text{Span}_{\mathcal{O}_p} F(1 \otimes \mathcal{B}_p)$, i.e $\mathcal{S}_p(S^1 \times S^1) \subseteq \text{Span}_{\mathcal{O}_p} \mathcal{B}_p$. Hence, we conclude that \mathcal{B}_p is a basis for $\mathcal{S}_p(S^1 \times S^1)$ from the fact that $\text{rank}(\mathcal{S}_p(S^1 \times S^1)) = d_p$. To prove the claim, let us look at the image of \mathcal{B}_p under F .

$$t^{i+\delta_i p}(\omega_p) \rightarrow t^{i+\delta_i p}(\omega_2) \otimes t^{i+\delta_i p}(\omega_r)$$

where $i \in \{0, 1, \dots, d_p - 1\}$.

Let us consider first all the elements of \mathcal{B}_p with even number of twists, i.e $i + \delta_i p \equiv 0 \pmod{2}$. By Lemma (7.4), we get that $i + \delta_i p \equiv 0 \pmod{4}$. Hence those elements get mapped to $t^{4m}(\omega_2) = \omega_2$ for some m , as t^4 is the identity map in the 2-theory. Also they get mapped to $t^{2j}(\omega_r)$ for some $0 \leq j \leq d_p - 1$, as t^p is the identity map in the $SO(3)$ -TQFT-theory and i is even. The later elements form the basis \mathcal{B}_{2r} defined in the previous theorem. In short, the above elements get mapped to $\omega_2 \otimes \mathcal{B}_{2r}$.

Now we consider the elements of \mathcal{B}_p with odd number of twists, i.e $i + \delta_i p \equiv 1 \pmod{2}$. By Lemma (7.5), we get that $i + \delta_i p \equiv 1 \pmod{4}$. Hence those elements get mapped to $t^{4m+1}(\omega_2) = t(\omega_2)$ for some m , as t^4 is the identity map in the

2-theory. Also they get mapped to $t^{2j+1}(\omega_r)$ for some $0 \leq j \leq d_p - 1$, as t^p is the identity map in the $SO(3)$ -TQFT-theory and i is odd. The later elements form the basis \mathcal{B}_{2r+1} defined in the previous theorem. In short, the above elements get mapped to $\omega_2 \otimes \mathcal{B}_{2r+1}$.

Hence the image of \mathcal{B}_p under F is a basis for $\widehat{\mathcal{S}}_2(S^1 \times S^1) \otimes \widehat{\mathcal{S}}_r(S^1 \times S^1)$, i.e generates it as required. \square

Corollary 7.7. *From the above proof, we conclude $\mathcal{S}_p(S^1 \times S^1) \cong \widehat{\mathcal{S}}_2(S^1 \times S^1) \otimes \widehat{\mathcal{S}}_r(S^1 \times S^1)$.*

We do not know if this holds for higher genus surfaces, but it is clear that $\mathcal{S}_p(\Sigma)$ maps into $\widehat{\mathcal{S}}_2(\Sigma) \otimes \widehat{\mathcal{S}}_r(\Sigma)$ under the map F .

Proposition 7.8.

$$\mathcal{J}_p(N_L) = \langle I_p(M_{i+\delta_i p}) \mid 0 \leq i \leq d_p - 1 \rangle,$$

where M_i is the 3-manifold obtained by doing surgery along the component K and the component J with framing $i + \delta_i p$ in S^3 .

Finally, a good question would be: “Is there a relation between the Frohman Kania-Bartoszynska ideals in the $SU(2)$ - and the $SO(3)$ -TQFT-theories?” An answer is given by the following theorem.

Theorem 7.9. *Let N be an oriented compact 3-manifold with boundary. Then we have*

$$\mathcal{J}_p(N) \subseteq i_r(\mathcal{J}_2(N))j_r(\mathcal{J}_r(N)),$$

where i_r and j_r are defined as in the previous section. Moreover, we have equality if N has a torus boundary.

Proof. To prove the inclusion, we have $F(1 \otimes \mathcal{S}_p(\Sigma)) \subseteq \widehat{\mathcal{S}}_2(\Sigma) \otimes \widehat{\mathcal{S}}_r(\Sigma)$. So if $[M]_p \in \mathcal{S}_p(\Sigma)$, then $F(1 \otimes [M]_p) = [M]_2 \otimes [M]_r \in \widehat{\mathcal{S}}_2(\Sigma) \otimes \widehat{\mathcal{S}}_r(\Sigma)$. By Theorem (6.5) as $I_1(M) = 1$, we have

$$I_p(N \cup_\Sigma -M) = i_r(I_2(N \cup_\Sigma -M))j_r(I_r(N \cup_\Sigma -M)) \in i_p(\mathcal{J}_2(N))j_p(\mathcal{J}_r(N)).$$

So we can conclude that $\mathcal{J}_p(N) \subseteq i_p(\mathcal{J}_2(N))j_p(\mathcal{J}_r(N))$. Now we prove the equality in the case of a torus boundary. Let $M_{i+\delta_i p}$ be the solid torus where its core colored $t^{i+\delta_i p}(w)$. From the previous proposition, we have

$$\begin{aligned} \mathcal{J}_p(N) &= \langle I_p(N \cup_{S^1 \times S^1} M_{i+\delta_i p}) \mid 0 \leq i \leq d_p - 1 \rangle \\ &= \langle i_r(I_2(N \cup_{S^1 \times S^1} M_{i+\delta_i p}))j_r(I_r(N \cup_{S^1 \times S^1} M_{i+\delta_i p})) \mid 0 \leq i \leq d_p - 1 \rangle \\ &= i_r(\mathcal{J}_2(N))j_r(\mathcal{J}_r(N)). \end{aligned}$$

The last equality follows from the fact that $F(1 \otimes \mathcal{B}_p)$ is a basis for $\widehat{\mathcal{S}}_2(S^1 \times S^1) \otimes \widehat{\mathcal{S}}_r(S^1 \times S^1)$. \square

Acknowledgment. I like to express my deep thanks and gratitude to my Ph.D advisor Dr. P. Gilmer for all the discussions and the support that were very important to finish this paper. Also, I like to thank Dr. P. van Wamelen, and Dr. N. Stoltzfus.

REFERENCES

- [BHMV1] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel, *Three-Manifold Invariants Derived from the Kauffman Bracket*, *Topology* **31** (1992), 685-699.
- [BHMV2] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel, *Remarks on the Three-Manifold Invariants θ_p* , 'Operator Algebras, Mathematical Physics, and Low Dimensional Topology' (NATO Workshop July 1991) Edited by R. Herman and B. Tanbay, *Research Notes in Mathematics* **5** (1993), 39-59.
- [BHMV3] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel, *Topological Quantum Field Theories derived from the Kauffman bracket*, *Topology* **34** (1995), 883-927.
- [FK] C. Frohman, J. Kania-Bartoszyńska, *A quantum obstruction to embedding*, *Math. Proc. Cambridge Philso. Soc.* **131** (2001), 279-293.
- [G1] P. M. Gilmer, *Integrality For TQFTS*, *Duke Mathematical Journal* **125**(2004), 389-413.
- [G2] P. M. Gilmer, *On The Frohman Kania-Bartoszyńska Ideal*, *Math. Proc. Cambridge Philso. Soc.*, to appear.
- [GM] P. M. Gilmer, and G. Masbaum, *Integral Lattices in TQFT*, arXiv: math.GT/0411029.
- [GMW] P. M. Gilmer, G. Masbaum, and P. van Wamelen, *Integral bases for TQFT modules and unimodular representations of mapping class groups*, *Comment. Math. Helv.* **79** (2004), 260-284.
- [GQ] P. M. Gilmer, K. Qazaqzeh, *The parity of the Maslov index and the even cobordism category*, *Fund. Math.*, to appear.
- [M] H. Murakami, *Quantum $SO(3)$ -invariants dominate the $SU(2)$ -invariant of Casson and Walker*, *Math. Proc. Cambridge Philso. Soc.* **117** (1995), no.2 237-249.
- [MR] G. Masbaum, J. D. Roberts, *A simple proof of integrality of quantum invariants at prime roots of unity*, *Math. Proc. Camb. Phil. Soc.* **121** (1997), 443-454.
- [W] L. Washington, *Introduction to Cyclotomic Fields*, Springer-Verlag 83, 1980.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, 70803, USA
E-mail address: qazaqzeh@math.lsu.edu
URL: www.math.lsu.edu/~qazaqzeh/