

A NOTE ON THE DUAL COMPLEX ASSOCIATED TO A RESOLUTION OF SINGULARITIES

D. A. STEPANOV

1. CONSTRUCTION

The dual complex associated to a resolution of singularities generalizes the notion of a resolution graph of a surface singularity to any dimension. Let X be an algebraic variety over a field of characteristic 0 or an analytic space and let $S \subset X$ be the singular locus of X . Usually we consider the germ of singularity (X, S) . Take a resolution $\pi: (Y, E) \rightarrow (X, S)$ such that the exceptional set E is a divisor with simple normal crossings. This resolution exists by Hironaka theorem (see [1], [2]); following terminology of the minimal model theory, we say that π is a *log-resolution*.

The dual complex $\Gamma(E)$ associated to the resolution π is a dual CW-complex associated to the divisor E as to a reducible variety. More precisely, decompose the divisor E to its prime components E_i : $E = \sum_{i=1}^N E_i$. Then k -dimensional cells $\Delta_{i_0 \dots i_k}^j$ of the complex $\Gamma(E)$ are in one-to-one correspondence with irreducible components $E_{i_0 \dots i_k}^j$ of the intersections

$$E_{i_0} \cap \dots \cap E_{i_k} = \cup_j E_{i_0 \dots i_k}^j,$$

$$k = 0, \dots, n-1, \quad 1 \leq i_0 < i_1 < \dots < i_k \leq N.$$

Every cell $\Delta_{i_0 \dots i_k}^j$ is a standard k -dimensional simplex with vertices marked by numbers i_0, \dots, i_k . For instance, the 0-dimensional skeleton of the complex $\Gamma(E)$ is a union of the points Δ_i corresponding to the prime divisors E_i . The patching map from the boundary $\partial \Delta_{i_0 \dots i_k}^j$ of a k -dimensional cell to the $k-1$ -dimensional skeleton is simplicial. It maps the face $\Delta_{i_0 \dots \widehat{i_s} \dots i_k}^j$ onto simplex $\Delta_{i_0 \dots \widehat{i_s} \dots i_k}^{j'}$ such that

$$E_{i_0 \dots i_k}^j \cap E_{i_0 \dots \widehat{i_s} \dots i_k}^{j'} \neq \emptyset,$$

and sends the vertex i_r of the face $E_{i_0 \dots \widehat{i_s} \dots i_k}^j$ to the vertex i_r of the simplex $\Delta_{i_0 \dots \widehat{i_s} \dots i_k}^{j'}$.

The complex $\Gamma(E)$ was considered by G. L. Gordon in the paper [3].

The research was supported by RFBR grant no. 05-01-00353.

2. HOMOTOPY TYPE OF THE DUAL COMPLEX IS AN INVARIANT OF A SINGULARITY

Theorem 1. *Let $\pi': (Y', E') \rightarrow (X, o)$ and $\pi'': (Y'', E'') \rightarrow (X, o)$ be two log-resolutions of an isolated singularity (X, o) . Then the topological spaces $\Gamma(E')$ and $\Gamma(E'')$ have the same homotopy type.*

This result is based on the following theorem by Abramovich-Karu-Matsuki-Włodarczyk on factorization of birational maps (see [4]).

Theorem 2 (Weak Factorization Theorem in the Logarithmic Category). *Let (U_{X_1}, X_1) and (U_{X_2}, X_2) be complete nonsingular toroidal embeddings (this means that U_{X_i} is an open set in a nonsingular variety X_i and the boundary $X_i \setminus U_{X_i}$ is a divisor with simple normal crossings, $i = 1, 2$) over an algebraically closed field of characteristic zero. Let*

$$\varphi: (U_{X_1}, X_1) \dashrightarrow (U_{X_2}, X_2)$$

be a birational map which is an isomorphism over $U_{X_1} = U_{X_2}$. Then the map φ can be factored into a sequence of blowups and blowdowns with smooth admissible and irreducible centers disjoint from $U_{X_1} = U_{X_2}$. That is to say, there exists a sequence of birational maps between complete nonsingular toroidal embeddings

$$\begin{aligned} (U_{X_1}, X_1) &= (U_{V_1}, V_1) \xrightarrow{\psi_1} (U_{V_2}, V_2) \xrightarrow{\psi_2} \dots \\ &\dots \xrightarrow{\psi_{i-1}} (U_{V_i}, V_i) \xrightarrow{\psi_i} (U_{V_{i+1}}, V_{i+1}) \xrightarrow{\psi_{i+1}} \dots \\ &\dots \xrightarrow{\psi_{l-2}} (U_{V_{l-1}}, V_{l-1}) \xrightarrow{\psi_{l-1}} (U_{V_l}, V_l) = (U_{X_2}, X_2), \end{aligned}$$

where

- (i) $\varphi = \psi_{l-1} \circ \psi_{l-2} \circ \dots \circ \psi_1$,
- (ii) ψ_i are isomorphisms over U_{V_i} , and
- (iii) either ψ_i or ψ_i^{-1} is a morphism obtained by blowing up a smooth irreducible center C_i (or C_{i+1}) disjoint from $U_{V_i} = U_{V_{i+1}}$ and transversal to the boundary $D_{V_i} = V_i \setminus U_{V_i}$ (or $D_{V_{i+1}} = V_{i+1} \setminus U_{V_{i+1}}$), i. e., at each point $p \in V_i$ (or $p \in V_{i+1}$) there exists a regular coordinate system $\{x_1, \dots, x_n\}$ in a neighborhood $p \in U_p$ such that

$$D_{V_i} \cap U_p \text{ (or } D_{V_{i+1}} \cap U_p) = \left\{ \prod_{j \in J} x_j = 0 \right\}$$

and

$$C_i \cap U_p \text{ (or } C_{i+1} \cap U_p) = \left\{ \prod_{j \in J} x_j = 0, x_{j'} = 0 \forall j' \in J' \right\},$$

$$J, J' \subseteq \{1, \dots, n\}.$$

Remark. The same holds if X_i are complex manifolds and φ, ψ_i are bimeromorphic maps.

To apply Theorem 2 in our case, take the resolutions $(Y' \setminus E', Y')$ and $(Y'' \setminus E'', Y'')$ as toroidal embeddings and compactify Y' and Y'' to smooth varieties (here we use the fact that the given singularity (X, o) is isolated). Now Theorem 1 follows from the

Lemma. *Let $\sigma: (X' \setminus E', X') \rightarrow (X \setminus E, X)$ be a blowup of an admissible center $C \subset E$ in a nonsingular toroidal embedding $(X \setminus E, X)$, $X' \setminus E' \simeq X \setminus E$. Then the topological spaces $\Gamma(E')$ and $\Gamma(E)$ have the same homotopy type.*

Proof. Let $E = \sum_{i=1}^N E_i$ be the decomposition of E into its prime components, and let $C \subset E_i$ for $1 \leq i \leq l$ and $C \not\subset E_i$ for $l < i \leq N$. Assume that C has nonempty intersections also with E_{l+1}, \dots, E_r , $l < r \leq N$. There are two possibilities.

1) $\dim C = n - l$ ($n = \dim X$), i. e., C coincides with one of the irreducible components of the intersection $E_1 \cap \dots \cap E_l$: $C = E_{1\dots l}^1$. Then after the blowup the intersection of the proper transforms E'_1, \dots, E'_l of the divisors E_1, \dots, E_l has $J - 1$ irreducible components (if J is the number of components of $E_1 \cap \dots \cap E_l$), but all these proper transforms intersect the exceptional divisor F of the blowup σ . Furthermore, F intersects proper transforms E'_{l+1}, \dots, E'_r of the divisors E_{l+1}, \dots, E_r . Now it is clear that the complex $\Gamma(E')$ is obtained from $\Gamma(E)$ by the barycentric subdivision of the simplex $\Delta_{1\dots l}^1$ with the center at the point corresponding to the divisor F . Thus the complexes $\Gamma(E')$ and $\Gamma(E)$ are even homeomorphic.

2) $\dim C < n - l$, let $C \subset E_{1\dots l}^1$. In this case divisors E_{i_1}, \dots, E_{i_s} have nonempty intersection if and only if their proper transforms $E'_{i_1}, \dots, E'_{i_s}$ have nonempty intersection. Therefore the complex $\Gamma(E')$ is obtained from the complex $\Gamma(E)$ in the following way. Add to $\Gamma(E)$ a new vertex corresponding to the exceptional divisor F of the blowup σ and construct cones with vertex at F over all the maximal cells $\Delta_{i_1\dots i_s}^j$ of the complex $\Gamma(E)$ possessing the property

$$E_{i_1\dots i_s}^j \cap C \neq \emptyset.$$

Note that the simplex $\Delta_{F,1\dots l}$ corresponding to the intersection $F \cap E_{1\dots l}^1$ is regarded as a common simplex for all constructed cones. Now we can define the homotopy equivalence between $\Gamma(E')$ and $\Gamma(E)$ as a contraction of the constructed cones: it sends the vertex F of the complex $\Gamma(E')$ to any of the vertices E_1, \dots, E_l of the cell $\Delta_{1\dots l}^1$ of the complex $\Gamma(E)$ and it is identity on other vertices of $\Gamma(E')$ ($\Gamma(E)$). Then the induced simplicial map is our homotopy equivalence. \square

3. TORIC SINGULARITIES

As an example let us find the homotopy type of the dual complex associated to a resolution of a toric singularity.

Theorem 3. *Let $o \in X$ be an isolated singularity of a toric variety X . Then for any log-resolution $\pi: (Y, E) \rightarrow (X, o)$ the dual complex $\Gamma(E)$ is homotopy equivalent to a point.*

Proof. We can restrict ourselves to the case when the variety X is affine. Then it can be represented as a toric variety $X = U_\sigma = \text{Spec } \mathbb{C}[\sigma]$ corresponding to a convex polyhedral cone $\sigma \subset \mathbb{R}^n$. According to Theorem 1, it is enough to determine the homotopy type for one arbitrary log-resolution of the singularity (X, o) .

There exists a fan Σ such that the support $\text{supp } \Sigma = \sigma$ and the birational morphism $\pi: Y = X(\Sigma) \rightarrow X$ is a log-resolution for the variety X (see [5]). First consider the dual complex for the divisor $E' = \sum T_i + E$, where T_i are invariant divisors corresponding to the edges τ_i of the cone σ and E is the exceptional divisor of the morphism π . Recall that the prime exceptional divisors E_i , $E = \sum E_i$, correspond to 1-dimensional cones of the fan Σ different from τ_i ; the divisors E_i and E_j have a nonempty intersection if and only if the corresponding 1-dimensional cones span a 2-dimensional cone belonging to the fan Σ , and so on.

Let L be a hyperplane in the space \mathbb{R}^n such that $L \cap \sigma = K$ is a compact polyhedron. Then the fan Σ determines some triangulation of the polyhedron K . We denote by the same letter K the corresponding simplicial complex. Now it is clear that the dual complex $\Gamma(E')$ is homeomorphic to the complex K and it obviously has the homotopy type of a point.

The dual complex $\Gamma(E)$ is obtained from K by deleting simplexes which contain at least one of divisors T_i as a vertex. The triangulation Σ can be chosen sufficiently small, so it is evident that after such deleting we obtain a complex which is homotopy equivalent to the original complex K , i. e., which has the homotopy type of a point. \square

4. SOME REMARKS

The surface cusp

$$X = \{x^4 + y^4 + z^4 + xyz = 0\} \subset \mathbb{C}^3;$$

gives an example of singularity with homotopy nontrivial dual complex associated to a resolution. It is easy to see that if $\pi: Y \rightarrow X$ is a

log-resolution, E is its exceptional divisor, then the complex $\Gamma(E)$ is homotopy equivalent to the circle S^1 .

The singularity

$$\{x^8 + y^8 + z^8 + x^2y^2z^2 = 0\} \subset \mathbb{C}^3$$

gives a more complicated example (see [3]). If we blow up the origin, the exceptional divisor $E'|_{X'}$ consists of 3 lines E_i , $i = 1, 2, 3$; every 2 of them intersect at a single point. The proper transform X' is singular along these lines. A resolution of X can be obtained by blowing up the variety X' along $E_1 \cup E_2 \cup E_3$. The dual complex associated to this resolution has a nontrivial first homology group; in particular, $\dim H_1(\Gamma(E), \mathbb{Q}) = 4$.

On the other hand, it is known (see [6]) that the resolution graphs of rational surface singularities are trees, thus they have homotopy type of a point. It would be interesting to verify if this fact generalizes to higher dimensions. More precisely, if (X, o) is an isolated rational singularity and $\pi: (Y, E) \rightarrow (X, o)$ is a log-resolution, then is it right that the complex $\Gamma(E)$ has the homotopy type of a point?

REFERENCES

- [1] Hironaka, H. *Resolution of singularities of an algebraic variety over a field of characteristic zero*, Annals of Math. **79** (1964), 109–326.
- [2] Włodarczyk, J. *Simple Hironaka resolution in characteristic zero*, e-print: arXiv:math.AG/0401401.
- [3] Gordon, G.L. *On a simplicial complex associated to the monodromy*, Transactions of the AMS **261** (1980), 93–101.
- [4] Abramovich, D., Karu, K., Matsuki, K., Włodarczyk, J. *Torification and factorization of birational maps*, J. Amer. Math. Soc. **15** (2002), 531–572.
- [5] Danilov, V.I. *The geometry of toric varieties* (Russian), Uspekhi matem. nauk **33** (1978), 97–154.
- [6] Artin, M. *On isolated rational singularities of surfaces*, Amer. J. Math. **88** (1966), 129–136.

THE DEPARTMENT OF MATHEMATICAL MODELLING, BAUMAN MOSCOW STATE
TECHNICAL UNIVERSITY, MOSCOW 105005, RUSSIA

E-mail address: dstepanov@bmstu.ru