Quadrance polygons, association schemes and strongly regular graphs

Le Anh Vinh School of Mathematics University of New South Wales Email: vinh@maths.unsw.edu.au

June 27, 2018

Abstract

Quadrance between two points $A_1 = [x_1, y_1]$ and $A_2 = [x_2, y_2]$ is the number $Q(A_1, A_2) := (x_2 - x_1)^2 + (y_2 - y_1)^2$. In this paper, we present some interesting results arise from this notation. In Section 1, we will study geometry over finite fields under quadrance notations. The main purpose of Section 1 is to answer the question, for which $a_1, \ldots a_n$, we have a polygon $A_1 \ldots A_n$ such that $Q(A_i, A_{i+1}) = a_i$ for $i = 1, \ldots, n$. In Section 2, using tools developed in Section 1, we define a family of association schemes over finite field space $F_q \times F_q$ where q is a prime power. These schemes give rise to a graph V_q with vertices the points of F_q^2 , and where (X, Y) is an edge of V_q if and only if Q(X, Y)is a nonzero square number in F_q . In Section 3, we will show that V_q is a strongly regular graph and propose a conjecture about the maximal clique of V_q .

1 Universal geometry over finite fields

Suppose that q is an odd prime power, and that F_q is the finite field with q elements. To avoid lengthly calculations (but mostly repetition), we assume that q is of form q = 4l + 3 for some integer l throughout Sections 1-3. In Section 4, we will present companion results for case q = 4l+1. The following definitions follow [5] where the importance of the notation of quadrance is developed.

Definition 1 The quadrance $Q(A_1, A_2)$ between the points $A_1 = [x_1, y_1]$,

and $A_2 = [x_2, y_2]$ is the number

$$Q(A_1, A_2) := (x_2 - x_1)^2 + (y_2 - y_1)^2.$$

Definition 2 A circle in a finite field F_q with center $A_0 \in F_q \times F_q$ and quadrance $K \in F_q$ is set of all points X in $F_q \times F_q$ such that

$$Q(A_0, X) = K.$$

Recall that a (multiplicative) character of F_q is a homomorphism from F^* , the multiplicative group of the non-zero elements of F_q , to the multiplicative group of complex numbers with modulus 1. The identically 1 function is the principal character of F_q and is denoted χ_0 . Since $x^{q-1} = 1$ for every $x \in F_q^*$ we have $\chi^{q-1} = \chi_0$ for every character χ . A character χ is of order d if $\chi^d = \chi_0$ and d is the smallest positive integer with this property. By convention, we extend a character χ to the whole of F_q by putting $\chi(0) = 0$. The quadratic (residue) character is defined by $\chi(x) = x^{(q-1)/2}$. Equivalently, χ is 1 on square, 0 at 0 and -1 otherwise. It is easy to see that $\sum_{i \in F_q} \chi(i) = 0$. Let A_q, B_q, C_q and D_q be the numbers of i in F_q such that $\chi(i) = \chi(i+1) = 1$, $\chi(i) = \chi(i+1) = -1$, $\chi(i) = -\chi(i+1) = 1$ and $\chi(i) = -\chi(i+1) = -1$, respectively. We have the following lemma.

Lemma 1 Let q be an odd prime power of the form q = 4l + 3 for some integer l. Then $A_q = B_q = (q-3)/4$, $C_q = (q+1)/4$ and $D_q = (q-3)/4$.

Proof We have

$$\begin{split} \sum_{i \in F_q} \left(1 + \chi(i) \right) \left(1 + \chi(i+1) \right) &= q + \sum_{i \in F_q} \left\{ \chi(i) + \chi(i+1) + \chi(i(i+1)) \right\} \\ &= q + \sum_{i \in F_q, \neq 0, -1} \chi(i^2(1+i^{-1})) \\ &= q + \sum_{i \in F_q, \neq 0, -1} \chi(1+i^{-1}) \\ &= q - \chi(1) - \chi(0) \\ &= q - 1. \end{split}$$

But we have

$$(1 + \chi(i)) (1 + \chi(i+1)) = \begin{cases} 4 & \text{if both } i \text{ and } i+1 \text{ are square,} \\ 2 & \text{if } i=0. \\ 0 & \text{otherwise.} \end{cases}$$

Thus, $4A_q + 2 = q - 1$ or $A_q = (q - 3)/4$. Similarly, consider the sum

$$\sum_{\in F_q} \left(1 - \chi(i)\right) \left(1 - \chi(i+1)\right).$$

This gives $4B_q + 1 = q - 2$ or $B_q = (q - 3)/4$. Similarly, consider the sum

$$\sum_{i \in F_q} \left(1 + \chi(i) \right) \left(1 - \chi(i+1) \right),$$

we have $C_q = (q+1)/4$. And consider the sum

$$\sum_{i\in F_q} \left(1-\chi(i)\right) \left(1+\chi(i+1)\right),$$

we have $D_q = (q-3)/4$. This concludes the proof.

From Lemma 1, we can count the number of points in any circle in F_q^2 .

Lemma 2 Let C be a circle with nonzero quadrance in F_q^2 . Then C has exactly N = q + 1 points.

Proof Without lose of generality, we may assume that C is centered at the origin. Let K be the quadrance of C. We will prove the lemma for the case K = 1 as other cases are similar. The number of points in C is

$$N = \left| \left\{ [x, y] \in F_q^2 \mid x^2 + y^2 = 1 \right\} \right|.$$

We have $x^2+y^2 = 1$ if and only if $x^2 = (-y^2)+1$. Since $-y^2$ is not a square in F_q^2 . We have the numbers of (x^2, y^2) with $x, y \neq 0$ is $D_q = (q-3)/4$. For each (x^2, y^2) with $x, y \neq 0$, we have 4 corresponding points [x, y]. Besides, we have 4 other points [0, 1], [0, -1], [-1, 0] and [1, 0]. Hence, $N = 4D_q + 4 = q + 1$. This concludes the proof of the lemma.

Let $C_i(X)$ denote the circle centered at $X \in F_q^2$ with quadrance *i*.

Lemma 3 Let $i, j \neq 0$ in F_q and let X, Y be two distinct points in F_q^2 such that $k = Q(X, Y) \neq 0$. Then $|C_i(X) \cap C_j(Y)|$ only depends on i, j and k. Precisely, define

$$f(i, j, k) := ij - \frac{(i - j - k)^2}{4}$$

Then the number of intersection points is p_{ij}^k , where

$$p_{ij}^{k} = \begin{cases} 0 & \text{if } f(i, j, k) \text{ is non-square,} \\ 1 & \text{if } f(i, j, k) = 0, \\ 2 & \text{if } f(i, j, k) \text{ is square.} \end{cases}$$
(1)

Proof Suppose that X = [m, n] and Y = [m+x, n+y] for some $m, n, x, y \in F_q$ then $x^2 + y^2 = k$. Suppose that $Z \in C_i(X) \cap C_j(Y)$ where Z = [m + x + u, n + y + v] for some $u, v \in F_q$. Then we have $u^2 + v^2 = j$ and $(x+u)^2 + (y+v^2) = i$. This implies that xu + yv = (i-j-k)/2. But we have $(xu + yv)^2 + (xv - yu)^2 = (x^2 + y^2)(u^2 + v^2)$ so

$$(xv - yu)^2 = kj - \frac{(i - j - k)^2}{4} = ij - \frac{(k - i - j)^2}{4} = f(i, j, k).$$

If f(i, j, k) is non-square number in F_q then it is clear that there does not exist such x, y, u, v, or $p_{ij}^k = 0$. Otherwise, let $\alpha = (i - j - k)/2$ and $f(i, j, k) = \beta^2$ for $0 \leq \beta \leq (p+1)/2$ then

$$xv - yu = \pm \beta, \qquad xu + yv = \alpha.$$

Solving for (u, v) with respect to (x, y) we have

$$u = (\alpha x \mp \beta y)/k, \qquad v = (\alpha y \pm \beta x)/k.$$

If $\beta = 0$ then we have only one (u, v) for each (x, y), but if $\beta \neq 0$ then we have two pairs (u, v). This implies (1), completing the proof.

The function f(i, j, k) defined above has an interesting property.

Lemma 4 Given $i, j \neq 0$ in F_q , there exists at least $|C_i|/2$ values of k such that f(i, j, k) is square (maybe zero).

Proof There are two cases.

1. Suppose that ij is square in F_q . Then $ij = v^2$ for some $v \in F_q$. We have

$$f(i, j, k) = v^2 \left(1 - \left(\frac{k - i - j}{2v} \right)^2 \right).$$

From Lemma 1, we have $D_q = (q-3)/4$ values of $-t^2$ $(t \neq 0)$ such that $1-t^2$ is square. For each $t^2 \neq 0$, we have two values of $k = \pm 2tv + i + j$. Besides, we may choose (k - i - j) from $0, \pm 2v$ so we have three more values of k. Thus, the number of k such that f(i, j, k) is square (maybe zero) is 3 + (q-3)/2 = (q+3)/2.

2. Suppose that ij is non square in F_q . Then $ij = -v^2$ for some $v \in F_q$. We have

$$f(i,j,k) = v^2 \left(-1 - \left(\frac{k-i-j}{2v}\right)^2 \right)$$

From Lemma 1, we have $C_q = (q+1)/4$ values of $-t^2$ $(t \neq 0)$ such that $-1 - t^2$ is square. For each $t^2 \neq 0$, we have two values of $k = \pm 2tv + i + j$. These are all possibilities for k. Thus the number of k such that f(i, j, k) is square (maybe zero) is (q+1)/2.

This concludes the proof of the lemma.

The Lemmas 3 and 4 can be applied to yield an interesting result.

Theorem 1 If $q \equiv 3 \pmod{4}$ then for any $n \ge 4$ and $a_1, \ldots, a_n \in F_q^*$, there exists a polygon $A_1 \ldots A_n$ in F_q^2 such that $Q(A_i, A_{i+1}) = a_i$ for all $i \in \{1, \ldots, n\}$ (note that $A_{n+1} = A_1$).

Proof It suffices to prove for n = 4. For any $a_1, \ldots, a_4 \in F_q^*$, we need to show that there exists a quadrangle $A_1A_2A_3A_4$ such that $Q(A_i, A_{i+1}) = a_i$ for $i = 1, \ldots, 4$. From Lemma 4, there are at least (q + 1)/2 values of k such that $f(a_1, a_2, k)$ is square and at least (q + 1)/2 values of k such that $f(a_3, a_4, k)$ is square. We have only q possible values for k so by the pigeonhole principal, there must be some value k such that both $f(a_1, a_2, k)$ and $f(a_3, a_4, k)$ are square. Choose any two points $X, Z \in F_q^2$ with Q(X, Z). From Lemma 3, $f(a_1, a_2, k)$ and $f(a_3, a_4, k)$ are square in F_q so there exists Y, T such that $Q(X, Y) = a_1, Q(Y, Z) = a_2, Q(T, Z) = a_3$ and $Q(Z, X) = a_4$. Thus we can choose $A_1A_2A_3A_4 \equiv XYZT$. This concludes the proof. \Box

2 Quadrance Association Schemes

We now have enough tools to construct an association scheme on F_q^2 . Let us recall the formal definition of an association scheme from [1].

Definition 3 An association scheme with s associate classes on a finite set Ω is a partition of $\Omega \times \Omega$ into sets C_0, C_1, \ldots, C_s (called associate classes) such that

- (i) $C_0 = \text{Diag}(\Omega) = \{(\omega, \omega) : \omega \in \Omega\};\$
- (ii) C_i is symmetric for $i = 1, \ldots, s$, i.e. $C'_i := \{(\beta, \alpha) : (\alpha, \beta) \in C_i\} = C_i$;
- (iii) for all i, j, k in $\{0, ..., s\}$ there is an integer p_{ij}^k such that, for all (α, β) in C_k

$$|\{\gamma \in \Omega : (\alpha, \gamma) \in C_i, (\gamma, \beta) \in C_j\}| = p_{ij}^k.$$

The numbers p_{ij}^k are called the *intersection numbers*. The number $a_i = p_{ii}^0$ is called the *valency* of the *i*-th associate class.

Our association schemes are defined by quadrances between elements of F_q^2 . To meet the condition (i) in Definition 3, we need Q(X,Y) = 0 if and only if $X \equiv Y$ for all $X, Y \in F_q^2$, which is equivalent to $q \equiv 3 \pmod{4}$. The following theorem gives us an association scheme on F_q^2 .

Theorem 2 Let $\Omega = F_q^2$, and partition $\Omega \times \Omega$ into q subsets $C_0, C_1, \ldots, C_{q-1}$ (indexed by elements of F_q ; for example q-1 = -1) such that for X, Yin F_q^2 , $(X,Y) \in C_i$ if and only if Q(X,Y) = i. This partition gives us an association scheme on F_q^2 . Furthermore, for $i, j, k \in F_q$, let f(i, j, k) = $ij - (i - j - k)^2/4$. Then the intersection numbers of the scheme are

$$p_{ij}^{0} = \begin{cases} 0 & \text{if } i \neq j, \\ 1 & \text{if } i = j = 0, \\ q+1 & \text{otherwise}, \end{cases}$$
(2)

and

$$p_{ij}^{k} = \begin{cases} 0 & \text{if } f(i,j,k) \text{ is non-square,} \\ 1 & \text{if } f(i,j,k) = 0, \\ 2 & \text{if } f(i,j,k) \text{ is nonzero square.} \end{cases}$$
(3)

for $k \neq 0$.

Proof The conditions (i) and (ii) of Definition 3 are easy to check. We will check (iii). Suppose that we have $(X, Y) \in C_k$ where X = [m, n] and Y = [m + x, n + y] for some $m, n, x, y \in F_p$. Then $x^2 + y^2 = k$. For k = 0, it is clearly that x = y = 0, i.e. X = Y. We have

$$\{Z \in F_p : (X,Z) \in C_i, (Z,Y) \in C_j\} = \begin{cases} \emptyset & \text{if } i \neq j, \\ \{Z \in F_p : Q(X,Z) = i\} & \text{if } i = j. \end{cases}$$

This equation and Lemma 2 imply (2). Moreover, Lemma 3 implies (3), completing the proof of the theorem. \Box

We call this scheme the **quadrance association scheme**. This scheme can be used to obtain some other schemes. Precisely, we have the following theorem.

Theorem 3 For any $t \mid (q-1)$, there exists an association scheme with t+1 associate classes on F_q^2 .

Proof If $t \mid q-1$ then q-1 = tn for some positive integer n. There exists a primitive element of F_q , says g. Let $\alpha = g^t$ and let M be set of all nonzero n^{th} powers in F_q . We have |M| = t, and write $M = \{a_1^n, \ldots, a_t^n\}$. Set $x_i = a_i^n$ for $1 \leq i \leq t$. We partition F_q^2 into t+1 sets D_0, D_1, \ldots, D_t by

$$D_0 = C_0, \qquad D_h = \bigcup_{v=0}^{n-1} C_{\alpha^v x_j} \qquad \text{for} \qquad h = 1, \dots, t.$$

Here $C_0, C_1, \ldots, C_{q-1}$ are the associate classes of the quadrance association scheme. We check that this partition gives us an association scheme. The conditions (i) and (ii) of Definition 3 are easy to check. We will check (iii). For k = 0, $(X, Y) \in D_0$ iff X = Y. Similarly as in proof of Theorem 2, we have that

$$| \{ Z \in F_p : (X, Z) \in D_i, (Z, Y) \in D_j \} |$$

is the same for any $(X, Y) \in D_0$.

For $k \neq 0$, suppose that $(X, Y) \in D_k$. Then $(X, Y) \in C_{\alpha^v x_k}$ for some $0 \leq v \leq n-1$. The number of Z's in F_q^2 such that $(X, Z) \in D_i$ and $(Z, Y) \in D_j$ is

$$V_{ij}^k(v) := \sum_{c=0}^{n-1} \sum_{d=0}^{n-1} p_{\alpha^c x_i, \alpha^d x_j}^{\alpha^v x_k}.$$
(4)

From Theorem 2, p_{ij}^k depends only on whether 4f(i, j, k) is nonzero square, zero or nonsquare. Thus, we only need to look at this condition. For $0 \leq v, w \leq n-1$, we have

$$\begin{aligned} \alpha^{2(w-v)} 4f(\alpha^{c} x_{i}, \alpha^{d} x_{j}, \alpha^{v} x_{k}) &= \alpha^{2(w-v)} \left[4\alpha^{c+d} x_{i} x_{j} - (\alpha^{v} x_{k} - \alpha^{c} x_{i} - \alpha^{d} x_{j})^{2} \right] \\ &= 4\alpha^{(c+w-v)+(d+w-v)} x_{i} x_{j} - (\alpha^{w} x_{k} - \alpha^{c+w-v} x_{i} - \alpha^{d+w-v} x_{j})^{2} \\ &= 4f(\alpha^{(c+w-v)} x_{i}, \alpha^{(d+w-v)} x_{j}, \alpha^{w} x_{k}). \end{aligned}$$

So that $4f(\alpha^c x_i, \alpha^d x_j, \alpha^v x_k)$ is a square (nonzero/zero) or non square if and only if so is $f(\alpha^{(c+w-v)}x_i, \alpha^{(d+w-v)}x_j, \alpha^w x_k)$. Hence, if we replace (c, d)by (c+w-v, d+w-v) in (4) then $V_{ij}^k(v) = V_{ij}^k(w)$. This implies that $|\{Z \in F_p : (X, Z) \in D_i, (Z, Y) \in D_j\}|$ is the same for any $(X, Y) \in D_k$, completing the proof. \Box

3 Quadrance Graphs

Strongly Regular Graphs 3.1

Remind that $q \equiv 3 \pmod{4}$. In Theorem 3, we saw that for any $t \mid (q-1)$, there exists an association scheme with t+1 associate classes on F_q^2 . Now q is an odd prime power so $2 \mid (q-1)$ and we can compose the quadrance association scheme into an association scheme with three associative classes. It is not hard to see (from the construction in Theorem 3) that the association scheme with 3 associative classes is a partition $F_q^2 = \{D_0, D_1, D_2\}$ where

$$D_{0} = \{(X, X) : X \in F_{q}\},\$$

$$D_{1} = \{(X, Y) : X, Y \in F_{q}, Q(X, Y) \text{ is square}\},\$$

$$D_{2} = \{(X, Y) : X, Y \in F_{q}, Q(X, Y) \text{ is nonsquare}\}$$

Consider a graph of V_q with vertices the points of F_q^2 , and where (X, Y) is an edge of graph if and only if (X, Y) is in D_1 . For $X, Y \in F_q^2$, define

$$\delta(X, Y) = \{ Z : (X, Z), (Z, Y) \} \in D_1 \}.$$

We have the following theorem.

Theorem 4 Let X, Y be any two points in F_q^2 .

- (a) If $(X, Y) \in D_1$ then $|\delta(X, Y)| = (q^2 5)/4$.
- (b) If $(X, Y) \in D_2$ then $|\delta(X, Y)| = (q^2 1)/4$.

Proof a) Suppose that $(X, Y) \in D_1$. Then $Q(X, Y) = k^2$ for some $k \in F_q^*$. We count the number of points Z in $\delta(X, Y)$. Let Z be any point in $\delta(X,Y)$ then there exists $i, j \in F_q^*$ such that $Q(X,Z) = i^2$ and $Q(Y,Z) = j^2$. Without loss of generality, we can divide both i, j by k and assume that k = 1. We have

$$4f(i^2, j^2, 1) = 4i^2j^2 - (1 - i^2 - j^2)^2$$

= $((i+j)^2 - 1)(1 - (i-j)^2).$

There are three separate cases.

- 1. Suppose that $f(i^2, j^2, 1)$ is nonsquare. Then from Lemma 3, there does not exist Z such that $Q(X,Z) = i^2$ and $Q(Y,Z) = j^2$.
- 2. Suppose that $f(i^2, j^2, 1) = 0$. We have three subcases.

- (a) If $(i+j)^2 1 = 1 (i-j)^2 = 0$ then $i+j = \pm 1$ and $i-j = \pm 1$. But it contradicts to the condition $i, j \in F_q^*$.
- (b) If $(i+j)^2 1 = 0 \neq 1 (i-j)^2$ then $i+j = \pm 1$ and $(i-j)^2 \neq \pm 1$. Thus, we have 2(q-2) pairs for (i+j, i-j). From Lemma 3, each pair gives us one point Z.
- (c) If $1 (i j)^2 = 0 \neq (i + j)^2 1$ then $i j = \pm 1$ and $i + j \neq \pm 1$. Similarly, we have 2(q - 2) points Z.
- 3. Suppose that $f(i^2, j^2, 1)$ is a nonzero square. Then either both $(i + j)^2 1$ and $1 (i j)^2$ are nonzero square or both $(i + j)^2 1$ and $1 (i j)^2$ are nonsquare.
 - (a) Suppose that $(i + j)^2 1$ and $1 (i j)^2$ are nonzero square. From Lemma 1, there are (q - 3)/4 nonzero values for $(i + j)^2$ and (q - 3)/4 nonzero values for $(i - j)^2$. But i - j = 0 also satisfies $1 - (i - j)^2$ is nonzero square. Thus, we have (q - 3)/2values for i + j and (q - 1)/2 values for i - j. From Lemma 3, each pair (i + j, i - j) gives us two points Z and Z'. Thus, we have (q - 1)(q - 3)/2 points Z.
 - (b) Suppose that $(i+j)^2 1$ and $1 (i-j)^2$ are nonsquare. Similarly, we have (q-1)/2 values of i+j and (q-3)/2 values of i-j. Thus, we also have (q-1)(q-3)/2 points Z.

Therefore, the number of points Z such that $Q(X,Z)=i^2$ and $Q(Y,Z)=j^2$ is

$$4(q-2) + (q-1)(q-3) = q^2 - 5.$$

But any point Z is counted four times (for the four pairs (i, j), (i, -j), (-i, j)and (-i, -j)), so $|\delta(X, Y)| = (q^2 - 5)/4$.

b) Suppose that $(X, Y) \in D_2$. Then $Q(X, Y) = -k^2$ for some $k \in F_q^*$. We count the number of points Z in $\delta(X, Y)$. Let Z be any point in $\delta(X, Y)$ then there exists $i, j \in F_q^*$ such that $Q(X, Z) = i^2$ and $Q(Y, Z) = j^2$. Without loss of generality, we can divide both i, j by k and assume that k = -1. We have

$$4f(i^2, j^2, -1) = 4i^2j^2 - (-1 - i^2 - j^2)^2$$

= -((i + j)^2 + 1)(1 + (i - j)^2).

There are three separate cases.

1. Suppose that $f(i^2, j^2, -1)$ is nonsquare. Then from Lemma 3, there does not exist Z such that $Q(X, Z) = -i^2$ and $Q(Y, Z) = -j^2$.

- 2. Suppose that $f(i^2, j^2, -1) = 0$. This is impossible as -1 is not a square number in F_q .
- 3. Suppose that $f(i^2, j^2, -1)$ is a nonzero square. Again, there are two subcases.
 - (a) Suppose that $(i + j)^2 + 1$ is nonzero square and $1 + (i j)^2$ is nonsquare. From Lemma 1, there are (q-3)/4 nonzero values for $(i+j)^2$ and (q+1)/4 nonzero values for $(i-j)^2$. But i+j=0 also satisfies $1 + (i + j)^2$ is nonzero square. Thus, we have (q-1)/2values for i + j and (q + 1)/2 values for i - j. From Lemma 3, each pair (i + j, i - j) gives us two points Z and Z'. Thus, we have (q - 1)(q + 1)/2 points Z.
 - (b) Suppose that $(i+j)^2 + 1$ is nonsquare and $1 + (i-j)^2$ is nonzero square. Similarly, we have (q+1)/2 values of i+j and (p-1)/2 values of i-j. Thus, we also have (q-1)(q+1)/2 points Z.

Therefore, the number of points Z such that $Q(X,Z) = i^2$ and $Q(Y,Z) = j^2$ is $2(q-1)(q+1)/2 = q^2 - 1$. But any point Z is counted four times (for the four pairs (i, j), (i, -j), (-i, j) and (-i, -j)) so $|\delta(X, Y)| = (q^2 - 1)/4$. This completes the proof of the theorem.

From Theorem 4, we have V_q is a strongly regular graph with parameters $\{(q^2-1)/2, (q^2-5)/4, (q^2-1)/4\}$; that is V_q is $(q^2-1)/2$ -regular, any two adjacent vertices have $(q^2-5)/4$ common neighbours and any two non-adjacent vertices have $(q^2-1)/4$ common neighbours. For any two vertices a and b, there are precisely $(q^2-1)/4$ vertices $c \neq b$ joined to a and not joined to b. We call V_q quadrance graph.

3.2 Maximal complete subgraphs

Let U be the vertex set of a subgraph of V_q and e(U) be the number of edges joining vertices in U. In Section 3.1, we see that the graph V_q is a strongly regular graph with parameters $\{(q^2 - 1)/2, (q^2 - 5)/4, (q^2 - 1)/4\}$. The well-known Paley graph P_{q^2} has the same parameter. Thus, we can follow Bollobas's proof for Paley graphs (see [2], page 321-322) to show that

Theorem 5 Let U be a set of u vertices of the Quadrance graph V_q . Then

$$\left|e(U) - \frac{1}{2} \binom{u}{2}\right| \leqslant \frac{u}{4} \frac{q^2 - u}{q}$$

Proof We may assume that $0 < u < q^2$. Let n(U) be the number of unordered triples (X, Y, Z) with $X, Y \in U$ and $(X, Z), (Y, Z) \in E(V_q)$. By Theorem 4, for each pair (X, Y) of adjacent vertices of U there are $(q^2-5)/4$ such triples and for each pair (X, Y) of non-adjacent vertices of U there are $(q^2-1)/4$ such triples. Thus, we have

$$n(U) = e(U)\left(\frac{q^2 - 5}{4}\right) + \left\{\binom{u}{2} - e(U)\right\}\left(\frac{q^2 - 1}{4}\right) = \binom{u}{2}\left(\frac{q^2 - 1}{4}\right) - e(U).$$

Let d = 2e(U)/u be the average degree in the induced subgraph G[U]. There are $[\{(q^2 - 1)/2\} - d]$ edges joining U to V - U. So on average, a vertex of U is joined to d vertices in U, and a vertex in V - U is joined to $d_0 = \{u/(q^2 - u)\}[\{(q^2 - 1)/2)\} - d]$ vertices in U. Thus, we have

$$n(U) \ge u \binom{d}{2} + (q^2 - u) \binom{d_0}{2}.$$

This implies

$$\frac{ud(d-1)}{2} + \frac{u}{2}\left(\frac{q^2-1}{2} - d\right) \left\{\frac{u}{q^2-u}\left(\frac{q^2-1}{2} - d\right) - 1\right\} \leqslant \frac{u(u-1)(q^2-1)}{8} - \frac{du}{2}$$

Multiplying by $u(q^2 - u)/2q^2$ and rearranging, we have

$$\left\{\frac{du}{2} - \frac{1}{2}\binom{u}{2}\right\}^2 \leqslant \frac{u^2}{16q^2}(q^2 - u)^2.$$

Hence

$$\left| e(U) - \frac{1}{2} \binom{u}{2} \right| \leqslant \frac{u}{4} \frac{q^2 - u}{q}$$

This completes the proof.

If U spans a complete subgraphs or an empty subgraphs of V_q and |U| = u then

$$\left|e(U) - \frac{1}{2} \binom{u}{2}\right| = \frac{1}{2} \binom{u}{2} \leqslant \frac{u}{4} \frac{q^2 - u}{q}.$$

Hence $u \leq q$. It is easy to see that the equality holds when U is a set of q points in a line. Inspections for q = 3, 7 support for the converse statement. We propose the following conjecture.

Conjecture 1 If U spans a complete subgraphs or an empty subgraphs of order q of V_q then U is a line in $F_q \times F_q$.

4 Companion Results

In this section, we suppose that $q \equiv 1 \pmod{4}$. We present companion results of previous sections in this case. Some proofs of the results in this section require extra works but they are basically similar to proofs in previous sections. Therefore, we obmit the details. The following lemma is an analogue to Lemma 1.

Lemma 5 Let $q \equiv 1 \pmod{4}$ be an odd prime power. Then $A_q = (q-5)/5$, $B_q = (q-1)/4$ and $C_q = D_q = (q-1)/4$.

Using Lemma 5, we have the following result, which is an analogue to Lemma 2.

Lemma 6 Let C be a circle with nonzero quadrance in F_q^2 . Then C has exactly N = q - 1 points.

Lemma 4 still holds in this case. From Lemma 2 and Lemma 6, it is clear that the circle $C_0(X)$ is a single point when q = 4l + 3 for some integer l. But for q = 4l + 1, the circle $C_0(X)$ consists of 2q - 1 points. This property causes some troubles in calculations (see [4]) which can be resolved by redefine the circle $C_0(X)$. We redefine $C_0(X) = \{X\}$ and $C_q(X) = \{Y \in$ $F_q \times F_q \mid Q(Y, X) = 0, Y \neq X\}$. We have the following lemmas.

Lemma 7 For any $i, j \neq 0$ in F_q . Suppose that X, Y are two distinct points in F_q^2 with Q(X,Y) = 0. Then the circle $C_i(X)$ intersects $C_j(Y)$ if and only if $i \neq j$. Furthermore, if $i \neq j$ then two circles intersect at only one point.

Lemma 8 Let $i \in F_q^*$ then there exists $j \in F_q^*$ such that f(i, i, j) and f(j, j, i) are square in F_q .

Proof There are three cases.

- 1. Suppose that 3 is square in F_q . Then $4f(i, i, i) = 4i^2 (i i i)^2 = 3i^2$ is square in F_q . Thus, we can choose j = i.
- 2. Suppose that 5 is square in F_q . We have

$$4f(i, i, -i) = 4f(-i, -i, i) = -5i^{2}.$$

Since -1 is square in F_q so we can choose j = -i.

3. Suppose that neither 3 nor 5 is square in F_q . Then 15 is a square in F_q . We have

$$4f(i, i, -2i) = 4i^{2} - (-6i - i - i)^{2} = -60i^{2},$$

$$4f(-6i, -6i, i) = 144i^{2} - (i + 6i + 6i)^{2} = -25i^{2}.$$

Thus, both f(i, i, -2i) and f(-2i, -2i, i) are square in F_q . Hence we can choose j = -6i.

This concludes the proof.

Lemmas 3, 4, 7 and 8 can be used to obtain an analogue result for Theorem 1.

Theorem 6 If $q \equiv 1 \pmod{4}$ then for any $n \ge 5$ and $a_1, \ldots, a_n \in F_q^*$, there exists a polygon $A_1 \ldots A_n$ in F_q^2 such that $Q(A_i, A_{i+1}) = a_i$ for all $i \in \{1, \ldots, n\}$.

Proof Note that the condition $n \ge 5$ is necessary in this case. A simple illustration is that there does not exist a quadrangle with quadrances 1, 1, 1 and 3 in F_5^2 . Now we prove that for any $n \ge 5$ and $a_1, \ldots, a_n \in F_q^*$ then there exists a polygon $A_1 \ldots A_n$ in F_q^2 such that $Q(A_i, A_{i+1}) = a_i$ for all *i*. It is clear that we only need to prove the result for n = 5. The case q = 5 is easy to check, so we assume that q > 5 (then $q \ge 3^2 = 9$). We consider two cases.

Case 1: Suppose that $a_1 = a_2 = \ldots = a_5$. From Lemma 8, there exists $j \in F_q^*$ such that $f(a_1, a_1, j)$ and $f(j, j, a_1)$ are square in F_q . From Lemma 3, there exists a triangle $A_1A_2A_4$ with $Q(A_1A_2) = a_1$ and $Q(A_2, A_4) = Q(A_1, A_4) = j$. We have $Q(A_2, A_4) = j$ so from Lemma 3, there exists a point A_3 such that $Q(A_2A_3) = Q(A_3A_4) = a_1$. Similarly, there exists A_5 such that $Q(A_4A_5) = Q(A_1A_5) = a_1$. And we have a pentagon $A_1A_2A_3A_4A_5$ as required.

Case 2: Suppose that $a_1 \neq a_2$. Since Lemma 4 still holds, there are at least (p-1)/2 > 3 values of k such that $f(a_3, a_4, k)$ is square. Thus, we can choose some $k \neq 0, a_5$. Now we have $k \neq a_5$ and $a_1 \neq a_2$. Hence by Lemma 7, for any $X \neq Z \in F_q^2$ such that Q(X, Z) = 0 then there exists Y, V such that $Q(X, Y) = a_1, Q(Y, Z) = a_2, Q(Z, V) = k$ and $Q(V, X) = a_5$. But $f(a_3, a_4, k)$ is square so by Lemma 3, there exists T such that $Q(Z, T) = a_3$ and $Q(T, V) = a_4$. Hence we can choose the pentagon $A_1 \dots A_5 \equiv XYZTV$. This concludes the proof.

We have an open question.

Problem 1 For $q \equiv 1 \pmod{4}$ is an odd prime power. Find the necessary and sufficient condition for $i, j, k, l \in F_q$ such that there exist a quadrangle with quadrances i, j, k and l.

We can define quadrance association scheme on F_q^2 similarly as in Section 2. But note that in this case, we need to partition F_q^2 into q + 1 partitions. Precisely, we have the following theorem.

Theorem 7 Let $\Omega = F_q^2$, we partition $\Omega \times \Omega$ into q+1 subsets C_0, C_1, \ldots, C_q such that for X, Y in F_q^2 , $(X, Y) \in C_i$ if and only if $X \in C_i(Y)$. This partition gives us an association scheme on F_q^2 . Furthermore, for $i, j, k \in F_q$, let $f(i, j, k) = ij - (i - j - k)^2/4$ then the intersection numbers of the scheme are

$$p_{ij}^{0} = \begin{cases} 0 & \text{if } i \neq j, \\ 1 & \text{if } i = j = 0, \\ 2(q-1) & \text{if } i = j = q, \\ q-1 & \text{otherwise}, \end{cases}$$
$$p_{ij}^{q} = \begin{cases} 1 & \text{if } i \neq j, \\ q-2 & \text{if } i = j = q, \\ 0 & \text{otherwise}, \end{cases}$$

and

$$p_{ij}^{k} = \begin{cases} 0 & \text{if } f(i, j, k) \text{ is non-square,} \\ 1 & \text{if } f(i, j, k) = 0, \\ 2 & \text{if } f(i, j, k) \text{ is nonzero square.} \end{cases}$$

for $k \neq 0, q$.

We neither know an analogue to Theorem 3 for the association scheme above nor any strongly regular quadrance graph in this case.

References

- [1] Bailey, R. A., Association Schemes: Designed Experiments, Algebra and Combinatorics, Cambridge University Press, Cambridge, 2004.
- [2] Bollobas, B., Random graphs, Academic Press, 1985.
- [3] Burr, S. A., Diagonal Ramsey Numbers for Small Graphs, Journal of Graph Theory, 7 (1983), 57-67.

- [4] Vinh, L. A., Random walks on hypergroup of circles in a finite field, In The proceedings of Australasian Workshop on Combinatorial Algorithms, (2005), 341-351.
- [5] Wildberger, N. J., Divine Proportions: Rational trigonometry to universal geometry, WildEgg, 2005.