

LIMIT LAWS FOR DISTORTED RETURN TIME PROCESSES FOR INFINITE MEASURE PRESERVING TRANSFORMATIONS

MARC KESSEBÖHMER AND MEHDI SLASSI

ABSTRACT. We consider conservative ergodic measure preserving transformations on infinite measure spaces and investigate the asymptotic behaviour of distorted return time processes with respect to sets satisfying a type of Darling-Kac condition. We identify two critical cases for which we prove uniform distribution laws. For this we introduce the notion of uniformly returning sets and discuss some of their properties.

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

In this paper (X, T, \mathcal{A}, μ) will always denote a conservative ergodic measure preserving dynamical systems where μ is an infinite σ -finite measure. In particular, this means that the mean return time to sets of finite positive measure is infinite. Hence if the system is given by a Markov chain, this corresponds to the *null recurrent* situation. The investigation of ergodic and probabilistic properties of such dynamical systems leads to a number of interesting results which can always be interpreted within the theory of null recurrent Markov chains and which sometimes generalize classical theorems within this theory.

In this paper we present a generalization of the Thaler-Dynkin-Lamperti arc-sine law (cf. [Tha98] and (T) in Subsection 1.3) describing the asymptotic behaviour of the renewal theoretic process Z_n given by

$$Z_n(x) := \begin{cases} \max\{k \leq n : T^k(x) \in A\}, & x \in A_n := \bigcup_{k=0}^n T^{-k}A, \\ 0, & \text{else.} \end{cases}$$

For a regularly varying function F we consider the *distorted processes*

$$\frac{F(Z_n)}{F(n)} \quad \text{and} \quad \frac{F(n - Z_n)}{F(n)}.$$

In particular we introduce the processes

$$\Phi_n := \frac{\sum_{k=0}^{Z_n} \mu(A \cap \{\varphi > k\})}{\mu(A_n)} \quad \text{and} \quad \Psi_n := \frac{\sum_{k=0}^{n-Z_n} \mu(A \cap \{\varphi > k\})}{\mu(A_n)},$$

which we refer to as the *normalized Kac process* and *normalized spent time Kac process*, respectively. In here,

$$\varphi(x) = \inf\{n \geq 1 : T^n(x) \in A\}, \quad x \in X, \quad (1.1)$$

denotes the *first return time* to the set A .

Date: 23 March 2006.

2000 Mathematics Subject Classification. 37A40.

Key words and phrases. Limit laws, infinite ergodic theory, return time process.

Research supported by Zentrale Forschungsförderung Universität Bremen.

In Proposition 1.3 we give a purely probability theoretical result allowing us to derive limit laws of distorted processes if the limit law for the corresponding original process is known. This result is then applied in Corollary 1.4 to treat Φ_n and Ψ_n .

Two critical cases are identified which are not covered by Proposition 1.3 and are subject of our two main Theorems 1.5 and 1.6. More precisely this means that if the sequence $(\frac{Y_n}{n})$ converges in distribution to 0 and L is slowly varying, then in general nothing is known about the asymptotics of $\frac{L(Y_n)}{L(n)}$. In this situation we are able to show that under suitable conditions on the wandering rate of a uniform set A we have

$$\frac{L(Z_n)}{L(n)} \xrightarrow{\mathcal{L}(\mu)} U,$$

and if A is a uniformly returning set – as introduced in Subsection 1.2 – we have

$$\Psi_n \xrightarrow{\mathcal{L}(\mu)} U.$$

In here, U denotes a random variable *distributed uniformly* on $[0, 1]$.

Obviously, Theorem 1.5 can be applied to infinite measure preserving interval maps $T : [0, 1] \rightarrow [0, 1]$ with indifferent fixed points satisfying the Thaler condition stated in [Tha95], whereas Theorem 1.6 is applicable to those map satisfying the corresponding condition in [Tha00]. Other examples in the context of continued fractions – also covered by Theorem 1.6 – are treated in [KS05]. For related results we refer to [TZ] and for further interesting results concerning distributional limit theorems for ergodic sums in this context to [Zwe03].

1.1. Infinite ergodic theory. A characterization of (X, T, \mathcal{A}, μ) being a conservative ergodic measure preserving dynamical system where μ is an infinite σ -finite measure as used in this paper will be given at the end of this subsection. For further definitions and details we refer the reader to [Aar97].

Let

$$\mathcal{P}_\mu := \{\nu : \nu \text{ probability measure on } \mathcal{A} \text{ with } \nu \ll \mu\}$$

denote the set of probability measures on \mathcal{A} which are absolutely continuous with respect to μ . The measures from \mathcal{P}_μ represent the admissible initial distributions for the processes associated with the iteration of T . The symbol \mathcal{P}_μ will also be used for the set of the corresponding densities.

Let us recall the notion of the wandering rate. For a fixed set $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ we set

$$A_n := \bigcup_{k=0}^n T^{-k}A \quad \text{and} \quad W_n := W_n(A) := \mu(A_n), \quad n \geq 0,$$

and call the sequence $(W_n(A))$ the *wandering rate* of A . Note that for the wandering rate the following identities hold

$$W_n(A) = \sum_{k=0}^n \mu(A \cap \{\varphi > k\}) = \int_A \min(\varphi, n+1) d\mu, \quad n \geq 0.$$

Since T is conservative and ergodic, for all $\nu \in \mathcal{P}_\mu$,

$$\lim_{n \rightarrow \infty} \nu(A_n) = 1 \quad \text{and} \quad \nu(\{\varphi < \infty\}) = 1.$$

The key to an understanding of the stochastic properties of a nonsingular transformation of a σ -finite measure space often lies in the study of the long-term behaviour of the iterates of its *transfer operator*

$$\hat{T} : L_1(\mu) \longrightarrow L_1(\mu), f \longmapsto \hat{T}(f) := \frac{d(\nu_f \circ T^{-1})}{d\mu},$$

where ν_f denote the measure with density f with respect to μ . Clearly, \hat{T} is a positive linear operator characterized by

$$\int_B \hat{T}(f) d\mu = \int_{T^{-1}(B)} f d\mu, \quad f \in L_1(\mu), \quad B \in \mathcal{A}.$$

An approximation argument shows that equivalently for all $f \in L_1(\mu)$ and $g \in L_\infty(\mu)$

$$\int_X \hat{T}(f) \cdot g d\mu = \int_X f \cdot g \circ T d\mu.$$

The ergodic properties of (X, T, \mathcal{A}, μ) can be characterized in terms of the transfer operator in the following way (cf. [Aar97, Proposition 1.3.2]). A system is conservative and ergodic if and only if for all $f \in L_1^+(\mu) := \{f \in L_1(\mu) : f \geq 0 \text{ and } \int_X f d\mu > 0\}$ we have μ -a.e.

$$\sum_{n \geq 0} \hat{T}^n(f) = \infty. \quad (1.2)$$

Invariance of μ under T means $\hat{T}(1) = 1$.

1.2. Uniform and uniformly returning sets. The following two definitions are in many situation crucial within infinite ergodic theory.

- A set $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ is called *uniform* for $f \in \mathcal{P}_\mu$ if there exists a sequence (a_n) of positive reals such that

$$\frac{1}{a_n} \sum_{k=0}^{n-1} \hat{T}^k(f) \longrightarrow 1 \quad \mu\text{-a.e. uniformly on } A$$

(i.e. uniform convergence in $L_\infty(\mu|_{A \cap \mathcal{A}})$).

- The set A is called a *uniform set* if it is uniform for some $f \in \mathcal{P}_\mu$.

Remark. Note that from [Aar97, Proposition 3.8.7] we know, that (b_n) is regularly varying with exponent α (for the definition of this property see Section 2) if and only if (W_n) is regularly varying with exponent $(1 - \alpha)$. In this case α lies in the interval $[0, 1]$ and

$$a_n W_n \sim \frac{n}{\Gamma(1 + \alpha)\Gamma(2 - \alpha)}. \quad (1.3)$$

In here, $c_n \sim a_n$ for some sequences (c_n) and (a_n) means that $a_n \neq 0$ has only finitely many exceptions and $\lim_{n \rightarrow \infty} \frac{c_n}{a_n} = 1$.

Next, we define a new property for sets similar to that of being uniform. It will be used to state the conditions in Theorem 1.6.

Definition. A set $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ is called *uniformly returning* for $f \in \mathcal{P}_\mu$ if there exists a positive increasing sequence (b_n) diverging to ∞ such that

$$b_n \hat{T}^n(f) \longrightarrow 1 \quad \mu\text{-a.e. uniformly on } A.$$

The set A is called *uniformly returning* if it is uniformly returning for some $f \in \mathcal{P}_\mu$.

The following Example show the existence of uniformly returning sets.

Example. Let $T : [0, 1] \rightarrow [0, 1]$ be an interval map with two increasing full branches and an indifferent fixed point at 0 satisfying Thaler's conditions in [Tha00]. Then any set $A \in \mathcal{B}_{[0,1]}$ with positive distance from the indifferent fixed point 0 and $\lambda(A) > 0$ is uniformly returning. As an typical example see the Lasota–York interval map (cf. Example after Theorem 1.6).

Proposition 1.1. *Any uniformly returning set is uniform.*

Proof. Let A be a uniformly returning set for $f \in \mathcal{P}_\mu$. Then for each $\varepsilon \in (0, 1)$ there exists a positive integer n_0 such that for all $n \geq n_0$ we have

$$(1 - \varepsilon) \frac{1}{b_n} \leq \hat{T}^n(f) \leq (1 + \varepsilon) \frac{1}{b_n} \quad \mu\text{-a.e. uniformly on } A.$$

Set $\tilde{f} := \hat{T}^{n_0}(f) \in \mathcal{P}_\mu$. Since $b_n \uparrow \infty$ we deduce that $(\hat{T}^n(\tilde{f}))_{n \geq 0}$ is uniformly bounded μ -a.e. on A . Furthermore, since T is conservative and ergodic we have by (1.2)

$$\sum_{k=0}^{\infty} \hat{T}^k(\tilde{f}) = \infty \quad \mu\text{-a.e.}$$

Using the fact that

$$b_{n+n_0} \hat{T}^n(\tilde{f}) \rightarrow 1 \quad \mu\text{-a.e. uniformly on } A$$

and that $(\hat{T}^n(\tilde{f}))_{n \geq 0}$ is uniformly bounded, we get

$$\frac{1}{\sum_{k=0}^n \frac{1}{b_k}} \cdot \sum_{k=0}^n \hat{T}^k(\tilde{f}) \rightarrow 1 \quad \mu\text{-a.e. uniformly on } A.$$

This shows that A is a uniform set for \tilde{f} . □

Remark. The inverse implication of Proposition 1.1 is stated in [KS05] under some additional assumptions.

To characterize the difference of the notions of uniform and uniformly returning set we make the following considerations. Suppose A is uniformly returning for $f := \frac{1}{\mu(B)} 1_B$, $B \in \mathcal{A}$, $0 < \mu(B) < \infty$, then there exists a sequence (b_n) such that

$$\frac{b_n}{\mu(B)} \hat{T}^n(1_B) \rightarrow 1 \quad \mu\text{-a.e. uniformly on } A.$$

Integrating over A yields

$$b_n \mu(B \cap T^{-n}A) \rightarrow \mu(B) \mu(A).$$

In an analog way we deduce for a uniform set A that

$$\frac{1}{a_n} \sum_{k=1}^n \mu(B \cap T^{-k}A) \rightarrow \mu(B) \mu(A).$$

Hence, we may interpret 'uniform returning' as a version of 'strong mixing' in infinite ergodic theory, whereas 'uniform' corresponds to a version of 'ergodicity'.

Next we characterize the sequence (b_n) in this definition by the wandering rate similarly to (1.3).

Proposition 1.2. *For $\beta \in [0, 1)$ we have that (b_n) is regularly varying with exponent β if and only if (W_n) is regularly varying with the same exponent. In this case,*

$$b_n \sim W_n \Gamma(1 - \beta) \Gamma(1 + \beta) \quad (n \rightarrow \infty).$$

The proof of this proposition and of the following proposition and theorems will be postponed to Section 3.

1.3. Limit laws. An important question when studying convergence in distribution for processes defined in terms of a non-singular transformation is to what extent the limiting behaviour depends on the initial distribution. This is formalized as follows.

Let ν be a probability measure on the measurable space (X, \mathcal{A}) and $(R_n)_{n \geq 1}$ be a sequence of measurable real functions on X , distributional convergence of $(R_n)_{n \geq 1}$ w.r.t. ν to some random variable R with values in $[-\infty, \infty]$ will be denoted by $R_n \xrightarrow{\nu} R$. *Strong distributional convergence* abbreviated by $R_n \xrightarrow{\mathcal{L}(\mu)} R$ on the σ -finite measures space (X, \mathcal{A}, μ) means that $R_n \xrightarrow{\nu} R$ for all $\nu \in \mathcal{P}_\mu$. In particular for $c \in [-\infty, \infty]$,

$$R_n \xrightarrow{\mathcal{L}(\mu)} c \iff R_n \rightarrow c \text{ locally in measure,}$$

which we also denote by $R_n \xrightarrow{\mu} c$.

Now we are in the position to state the first interesting limit law for the process Z_n which is due to Thaler [Tha98].

- (T) **Thaler's Dynkin-Lamperti arc-sin Law.** Let $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ be a uniform set. If the wandering rate $(W_n(A))$ is regularly varying with exponent $1 - \alpha$ for $\alpha \in [0, 1]$, then we have

$$\frac{Z_n}{n} \xrightarrow{\mathcal{L}(\mu)} \xi_\alpha. \quad (1.4)$$

In here, for $\alpha \in (0, 1)$, ξ_α denotes the random variable on $[0, 1]$ with density

$$f_{\xi_\alpha}(x) = \frac{\sin \pi \alpha}{\pi} \frac{1}{x^{1-\alpha} (1-x)^\alpha}, \quad 0 < x < 1.$$

The distribution of ξ_α is also called *the generalized arc-sine distribution*. The continuous extension is given by $\xi_0 = 0$ and $\xi_1 = 1$.

To apply (T) to the distorted processes we need the following proposition.

Proposition 1.3. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $Y_n : \Omega \rightarrow [0, \infty]$ be measurable ($n \geq 1$), and let Y be a random variable with values in $[0, \infty]$.*

- (1) *If $\mathbb{P}(Y = 0) = 0 = \mathbb{P}(Y = \infty)$ and F is a regularly varying function with exponent $\beta \in \mathbb{R}$, then*

$$\frac{Y_n}{n} \xrightarrow{\mathbb{P}} Y \implies \frac{F(Y_n)}{F(n)} \xrightarrow{\mathbb{P}} Y^\beta.$$

- (2) *If $Y = 0$ and F is a regularly varying function with exponent $\beta \in \mathbb{R} \setminus \{0\}$ then*

$$\frac{Y_n}{n} \xrightarrow{\mathbb{P}} 0 \implies \frac{F(Y_n)}{F(n)} \xrightarrow{\mathbb{P}} \begin{cases} 0 & \text{for } \beta > 0 \\ \infty & \text{for } \beta < 0 \end{cases}.$$

The following corollary is a direct consequence of (1.4), Proposition 1.3, and the fact that $\Phi_n = \frac{F(Z_n)}{F(n)}$, $\Psi_n = \frac{F(n-Z_n)}{F(n)}$ with $F(n) := W_n$.

Corollary 1.4. *Let $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ be a uniform set. If the wandering rate (W_n) is regularly varying with exponent $1 - \alpha$, then*




FIGURE 1.1. The densities f_{X_α} of the limiting distribution of the normalized Kac process for different values of $\alpha \in (0, 1)$. The extreme distributions corresponding to $\alpha = 0$ and 1 are the Dirac measures δ_0 and δ_1 , respectively.

- (1) If $0 \leq \alpha \leq 1$, then we have

$$\Phi_n \xrightarrow{\mathcal{L}(\mu)} X_\alpha,$$

where X_α denotes the random variable on $[0, 1]$ with density

$$f_{X_\alpha}(x) = \frac{1}{1-\alpha} \frac{\sin \pi \alpha}{\pi} \frac{1}{x^{\frac{1-2\alpha}{1-\alpha}} \left(1-x^{\frac{1}{1-\alpha}}\right)^\alpha}, \quad \alpha \in (0, 1)$$

and $X_0 = 0, X_1 = 1$ (cf. Fig. 1.1).

- (2) If $0 \leq \alpha < 1$, then we have

$$\Psi_n \xrightarrow{\mathcal{L}(\mu)} Y_\alpha,$$

where Y_α denotes the random variable on $[0, 1]$ with density

$$f_{Y_\alpha}(x) = \frac{1}{1-\alpha} \frac{\sin \pi \alpha}{\pi} \frac{1}{\left(1-x^{\frac{1}{1-\alpha}}\right)^{1-\alpha}}, \quad \alpha \in (0, 1)$$

and $Y_0 = 1$ (cf. Fig. 1.2).




FIGURE 1.2. The densities f_{Y_α} of the limiting distribution of the normalized spent time Kac process for different values of $\alpha \in (0, 1)$. The extreme distribution corresponding to $\alpha = 0$ and 1 are the Dirac measure δ_1 and the uniform distribution on $[0, 1]$, respectively.

Remark. For $\alpha \in (0, 1)$ we have

$$X_\alpha \stackrel{\text{dist.}}{=} (\xi_\alpha)^{1-\alpha} \quad \text{and} \quad Y_\alpha \stackrel{\text{dist.}}{=} (1 - \xi_\alpha)^{1-\alpha}.$$

Note, that in particular both $X_{\frac{1}{2}}$ and $Y_{\frac{1}{2}}$ obey the arc-sine law, i.e. they have density

$$f_{Y_{\frac{1}{2}}}(x) = f_{X_{\frac{1}{2}}}(x) = \frac{2}{\pi} \frac{1}{(1-x^2)^{\frac{1}{2}}}, \quad 0 < x < 1.$$

Now we state as our first main result the following *uniform distribution law*. Note that this corresponds to the case $\beta = 0$ and $Y = 0$ in Proposition 1.3, which therefore is not applicable.

Theorem 1.5. *Let $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ be a uniform set. If the wandering rate (W_n) is regularly varying with exponent 1 such that $W_n \sim n \cdot \frac{1}{L(n)}$, where L is a slowly varying function. Then we have*

$$\frac{L(Z_n)}{L(n)} \xrightarrow{\mathcal{L}(\mu)} U,$$

where U denotes the uniformly distributed random variable on $[0, 1]$.

Remark. Under the assumptions of Theorem 1.5 we have by (T) and the first part of Corollary 1.4 that

$$\frac{Z_n}{n} \xrightarrow{\mathcal{L}(\mu)} 0 \quad \text{and} \quad \frac{1}{\Phi_n} \xrightarrow{\mathcal{L}(\mu)} \infty.$$

However, for the product of the two processes we have by Theorem 1.5

$$\frac{Z_n}{n\Phi_n} \xrightarrow{\mathcal{L}(\mu)} U.$$

Example. Let $f(0) = 0$, $f(x) = x + x^2 e^{-\frac{1}{x}}$, $x > 0$, and let $a \in (0, 1)$ be determined by $f(a) = 1$. Define $T : [0, 1] \rightarrow [0, 1]$ by

$$T(x) := \begin{cases} f(x), & x \in [0, a], \\ \frac{x-a}{1-a}, & x \in (a, 1]. \end{cases}$$

Then the map T satisfies Thaler's conditions (T1)–(T4) in [Tha95]. Any set $A \in \mathcal{B}_{[0,1]}$ with $\lambda(A) > 0$ which is bounded away from the indifferent fixed points is a uniform set for T . Furthermore we have

$$W_n \sim \text{const} \cdot \frac{n}{\log(n)} \quad (n \rightarrow \infty).$$

Hence,

$$\frac{\log Z_n}{\log n} \xrightarrow{\mathcal{L}(\mu)} U.$$

Next we state as our second main result the uniform distribution law for the normalized spent time Kac process. This again corresponds to the case $\beta = 0$ and $Y = 0$ in Proposition 1.3.

Theorem 1.6. *Let $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ be a uniformly returning set. If the wandering rate (W_n) is slowly varying, then we have*

$$\Psi_n \xrightarrow{\mathcal{L}(\mu)} U,$$

where the random variable U is distributed uniformly on $[0, 1]$.

Example. We consider the Lasota–Yorke map $T : [0, 1] \rightarrow [0, 1]$, defined by

$$T(x) := \begin{cases} \frac{x}{1-x}, & x \in [0, \frac{1}{2}], \\ 2x-1, & x \in (\frac{1}{2}, 1]. \end{cases}$$

This map satisfies the Thaler's conditions (i)–(iv) in [Tha00]. Any compact subset A of $(0, 1]$ with $\lambda(A) > 0$ is a uniformly returning set and we have

$$W_n \sim \log(n) \quad \text{as} \quad n \rightarrow \infty.$$

Hence,

$$\frac{\log(n - Z_n)}{\log(n)} \xrightarrow{\mathcal{L}(\mu)} U.$$

2. REGULAR VARIATION AND TAUBERIAN RESULTS

In this section we give some preparatory facts and results needed for the proofs of the main statements in the following Section 3.

We first recall the concepts of regularly varying functions and sequences (see also [BGT89] for a comprehensive account). Throughout we use the convention that for two sequences $(a_n), (b_n)$ we write $a_n = o(b_n)$ if $b_n \neq 0$ fails only for finitely many n and $\lim_{n \rightarrow \infty} \left| \frac{a_n}{b_n} \right| = 0$.

A measurable function $R : \mathbb{R}^+ \rightarrow \mathbb{R}$ with $R > 0$ on (a, ∞) for some $a > 0$ is called *regularly varying at ∞* with exponent $\rho \in \mathbb{R}$ if

$$\lim_{t \rightarrow \infty} \frac{R(\lambda t)}{R(t)} = \lambda^\rho \quad \text{for all } \lambda > 0.$$

A regularly varying function L with exponent $\rho = 0$ is called *slowly varying at ∞* , i.e.

$$\lim_{t \rightarrow \infty} \frac{L(\lambda t)}{L(t)} = 1 \quad \text{for all } \lambda > 0.$$

Clearly, a function $R : \mathbb{R}^+ \rightarrow \mathbb{R}$ is regularly varying at ∞ with exponent $\rho \in \mathbb{R}$ if and only if

$$R(t) = t^\rho L(t), \quad t \in \mathbb{R}^+,$$

for L slowly varying at ∞ .

A function R is said to be *regularly varying at 0* if $t \mapsto R\left(\frac{1}{t}\right)$ is regularly varying at ∞ .

A sequence (u_n) is *regularly varying with exponent ρ* if $u_n = R(n)$, $n \geq 1$, for $R : \mathbb{R}^+ \rightarrow \mathbb{R}$ regularly varying at ∞ with exponent ρ .

In the following list we state those Tauberian results needed in the proofs of the preparatory lemmas and propositions of this sections, as well as for the main theorems.

(KTT) **Karamata's Tauberian Theorem** ([Fel71], [Sen76]) Let $(b_n)_{n \geq 0}$ be a non-negative sequence such that for all $s > 0$, $B(s) := \sum_{n \geq 0} b_n e^{-ns} < \infty$. Let L be slowly varying at ∞ , and $\rho \in [0, \infty)$. Then

$$B(s) \sim \left(\frac{1}{s}\right)^\rho L\left(\frac{1}{s}\right) \quad \text{as } s \searrow 0,$$

if and only

$$\sum_{k=0}^{n-1} b_k \sim \frac{1}{\Gamma(\rho+1)} n^\rho L(n) \quad \text{as } n \rightarrow \infty.$$

If (b_n) is eventually monotone and $\rho > 0$, then both are equivalent to

$$b_n \sim \frac{1}{\Gamma(\rho)} n^{\rho-1} L(n) \quad \text{as } n \rightarrow \infty.$$

(KL) **Karamata's Lemma** ([Fel71, Kar33]). If (a_n) is a regularly varying sequence with exponent ρ and if $p \geq -\rho - 1$, then

$$\lim_{n \rightarrow \infty} \frac{n^{p+1} a_n}{\sum_{k \leq n} k^p a_k} = p + \rho + 1.$$

(UA) **Uniformly asymptotic** ([Sen76]) Let (p_n) and (q_n) be two positive sequences with $p_n \rightarrow \infty$ and $\frac{p_n}{q_n} \in [1/k, k]$, $k > 0$. Then we have for L a slowly varying function

$$\lim_{n \rightarrow \infty} \frac{L(p_n)}{L(q_n)} = 1.$$

(EL) **Erickson Lemma** ([Eri70]) Let $L \nearrow \infty$ be a monotone increasing continuous slowly varying function. Let $a_t(x)$ be defined by $a_t(x) := L^{-1}(xL(t))$ with $x \in (0, 1)$, where $L^{-1}(\cdot)$ denoting the inverse function of $L(\cdot)$. Then we have for every fixed $x \in (0, 1)$

$$a_t(x) = o(t) \quad \text{and} \quad a_t(x) \rightarrow \infty \quad (t \rightarrow \infty).$$

3. PROOF OF MAIN RESULTS

Proof of Proposition 1.2. Let $A \in \mathcal{A}$, $0 < \mu(A) < \infty$, be a uniformly returning set, and let the functions $U(s)$, $Q(s)$, $s > 0$, be defined as Laplace transforms

$$\begin{aligned} Q(s) &:= \sum_{n=0}^{\infty} \frac{\mu(A \cap \{\varphi > n\})}{\mu(A)} e^{-ns} \\ U(s) &:= \sum_{n=0}^{\infty} \nu(T^{-n}A) e^{-ns}, \end{aligned} \quad (3.1)$$

where ν denotes the probability measure with density $f \in \mathcal{P}_\mu$.

Since

$$A_n := \bigcup_{k=0}^n T^{-k}A = \bigcup_{k=0}^n T^{-k}(A \cap \{\varphi > n-k\}),$$

and the sets $T^{-k}(A \cap \{\varphi > n-k\})$, $0 \leq k \leq n$, are disjoint, we have

$$\nu(A_n) = \int_A \sum_{k=0}^n \hat{T}^k(f) \cdot 1_{A \cap \{\varphi > n-k\}} d\mu, \quad n \geq 0.$$

Thus,

$$\sum_{n=0}^{\infty} \nu(A_n) e^{-ns} = \int_A \left(\sum_{n=0}^{\infty} \hat{T}^n(f) e^{-ns} \right) \left(\sum_{n=0}^{\infty} 1_{A \cap \{\varphi > n\}} e^{-ns} \right) d\mu.$$

From

$$\hat{T}^n(f) \sim \frac{\nu(T^{-n}A)}{\mu(A)} \quad \text{as } n \rightarrow \infty \quad \mu\text{-a.s. uniformly on } A,$$

it follows that

$$\sum_{n=0}^{\infty} \hat{T}^n(f) e^{-ns} \sim \frac{U(s)}{\mu(A)} \quad \text{as } s \rightarrow 0 \quad \mu\text{-a.s. uniformly on } A.$$

This implies

$$\sum_{n=0}^{\infty} \nu(A_n) e^{-ns} \sim U(s) Q(s) \quad \text{as } s \rightarrow 0.$$

Hence, since $\lim_{n \rightarrow \infty} \nu(A_n) = 1$, we obtain

$$\frac{1}{s} \sim U(s) Q(s). \quad (3.2)$$

If $b_n \sim n^\beta L(n)$ for $\beta \in [0, 1)$, L denoting some slowly varying function, then due to (KL) we have

$$\sum_{k=0}^{n-1} \nu(T^{-k}A) \sim \frac{n^{1-\beta}}{(1-\beta)L(n)} \mu(A).$$

Thus, by (KTT) we obtain

$$Q(s) \sim \frac{1}{\Gamma(1-\beta)\mu(A)} \left(\frac{1}{s}\right)^\beta L\left(\frac{1}{s}\right).$$

Hence,

$$W_n \sim \frac{1}{\Gamma(1-\beta)\Gamma(1+\beta)} b_n.$$

Now let $W_n \sim n^\beta \tilde{L}(n)$ for $\beta \in [0, 1)$, \tilde{L} denoting some slowly varying function. From (3.2), it follows by (KTT)

$$\sum_{k=0}^{n-1} \mathbf{v}(T^{-k}A) \sim \frac{n^{1-\beta}}{\Gamma(2-\beta)\Gamma(1+\beta)\tilde{L}(n)} \mu(A).$$

Hence since

$$\sum_{k=0}^{n-1} \mathbf{v}(T^{-k}A) \sim \mu(A) \sum_{k=0}^{n-1} \frac{1}{b_k},$$

(b_n) is monotone, and $1 - \beta > 0$, we obtain by (KTT)

$$\frac{1}{b_n} \sim \frac{n^{-\beta}}{\Gamma(1-\beta)\Gamma(1+\beta)\tilde{L}(n)}.$$

Thus,

$$b_n \sim \Gamma(1-\beta)\Gamma(1+\beta)W_n.$$

From this the assertion follows. \square

Proof of Proposition 1.3.

ad (1) It is known that for every regularly varying function with exponent $\beta \in \mathbb{R}$ there exists a slowly varying function such that $F(x) = x^\beta L(x)$ for all $x > 0$. Therefore to prove the result in Proposition 1.3, it suffices to show

$$\frac{L(Y_n)}{L(n)} \xrightarrow{\mathbb{P}} 1.$$

We have for all $\delta > 0$ and $K > 0$ with $\delta < K$

$$\liminf \mathbb{P} \left(\delta \leq \frac{Y_n}{n} \leq K \right) \geq 1 - C_{\delta, K}.$$

Due to the uniform convergence theorem for slowly varying functions (cf. [Sen76]) we have, for all $\varepsilon > 0$ there exists $n_0 := n_0(\varepsilon)$ such that

$$n \geq n_0 \quad \Rightarrow \quad \left| \frac{L(\lambda n)}{L(n)} - 1 \right| < \varepsilon \quad \text{for all } \lambda \in [\delta, K].$$

Hence, for sufficiently large n ,

$$\mathbb{P} \left(\left| \frac{L(Y_n)}{L(n)} - 1 \right| \geq \varepsilon \right) \leq 1 - \mathbb{P} \left(\delta \leq \frac{Y_n}{n} \leq K \right).$$

This implies

$$\limsup \mathbb{P} \left(\left| \frac{L(Y_n)}{L(n)} - 1 \right| \geq \varepsilon \right) \leq C_{\delta, K}.$$

Since $C_{\delta, K} \rightarrow 0$ as $\delta \rightarrow 0$ and $K \rightarrow \infty$, the first part of the proposition follows.

ad (2) Now let $Y = 0$ and $\beta > 0$. Without loss of generality we assume that F is a positive and locally bounded function on $[0, \infty)$. Then from [BGT89, p. 28] we know that

$$F'(x) := \inf \{y \geq 0 : F(y) > x\}, \quad x \in [0, \infty),$$

defines an asymptotic inverse of F , i.e.

- (1) $F'(F(x)) \sim x$ as $x \rightarrow \infty$,
- (2) F' is regular varying with exponent $1/\beta$.

Since $F(x) > y$ implies $y \geq F'(x)$ we have for $\varepsilon > 0$

$$\frac{F(Y_n)}{F(n)} > \varepsilon \implies Y_n \geq F'(\varepsilon F(n)).$$

Hence for all $n \geq 1$ we have

$$\mathbb{P}\left(\frac{F(Y_n)}{F(n)} > \varepsilon\right) \leq \mathbb{P}\left(\frac{Y_n}{n} > \varepsilon_n\right)$$

with $\varepsilon_n := F'(\varepsilon F(n))n^{-1}$. Since $F(n) \rightarrow \infty$, $n \rightarrow \infty$, we have by the properties (1) and (2) of the asymptotic inverse that $\lim_{n \rightarrow \infty} \varepsilon_n = \varepsilon^{1/\beta}$. Thus for n sufficiently large, we have

$$\mathbb{P}\left(\frac{F(Y_n)}{F(n)} > \varepsilon\right) \leq \mathbb{P}\left(\frac{Y_n}{n} > \frac{\varepsilon^{1/\beta}}{2}\right)$$

proving the second part of the proposition for $\beta > 0$. The case $\beta < 0$ is reduced to the above case by considering $1/F$ instead of F . \square

If T is a nonsingular ergodic transformation on (X, \mathcal{A}, μ) , the compactness theorem in [Aar97, Sec. 3.6] implies that $R_n \circ T - R_n \xrightarrow{\mu} 0$ and $R_n \xrightarrow{\nu} R$ for some $\nu \in \mathcal{P}_\mu$ is sufficient for $R_n \xrightarrow{\mathcal{L}(\mu)} R$. Hence, before proving the main theorems we state and prove the following two lemmata.

Lemma 3.1. *Let $A \in \mathcal{A}$ be a set of positive finite measure $\mu(A)$ and $F(t) \rightarrow \infty$, $t \rightarrow \infty$, be a regularly varying function with exponent $\beta \geq 0$. Then we have*

$$\frac{1}{F(n)} (F(Z_n \circ T) - F(Z_n)) \xrightarrow{\mu} 0. \quad (3.3)$$

Proof. Due to the Representation Theorem of regularly varying functions (c.f. [Sen76]) we have for some $B > 0$

$$F(x) = x^\beta \psi(x) \exp\left(\int_B^x \frac{\zeta(t)}{t} dt\right) \quad \text{for all } x \geq B, \quad (3.4)$$

where ψ is a positive measurable function on $[B, \infty)$ and ζ a continuous function on $[B, \infty)$ such that

$$\psi(x) \longrightarrow C \in (0, \infty) \quad \text{und} \quad \zeta(x) \longrightarrow 0 \quad (x \rightarrow \infty).$$

Without loss of generality we assume that there exists $\delta \in (0, 1)$ such that

$$|\zeta(t)| < \delta \quad \text{for all } t \geq B.$$

By (3.4) we have

$$F(x) \sim Cx^\beta \exp\left(\int_B^x \frac{\zeta(t)}{t} dt\right) \quad (x \rightarrow \infty).$$

We define $\tilde{F}(x) := Cx^\beta \exp\left(\int_B^x \frac{\zeta(t)}{t} dt\right)$ for $x \geq B$ and show first that the claim in the lemma holds for \tilde{F} .

Let φ be the first return time to the set A defined as in (1.1). For $\varepsilon > 0$ we define

$$K_{\varepsilon, n} := \left\{ \varphi \leq n \wedge \frac{1}{\tilde{F}(n)} \left| \tilde{F}(Z_n \circ T) - \tilde{F}(Z_n) \right| \geq \varepsilon \right\} \quad (n \in \mathbb{N}).$$

Since

$$Z_n(T(x)) = \begin{cases} Z_n(x) - 1, & x \in \{\varphi \leq n\} \cap T^{-(n+1)}A^c, \\ n, & x \in T^{-(n+1)}A, \end{cases} \quad (3.5)$$

we conclude

$$\begin{aligned} K_{\varepsilon, n} \subset & \left(\{\varphi \leq n\} \cap T^{-(n+1)}A^c \cap \left\{ \frac{1}{\tilde{F}(n)} \left(\tilde{F}(Z_n) - \tilde{F}(Z_n - 1) \right) \geq \varepsilon \right\} \right) \\ & \cup \left(T^{-(n+1)}A \cap \left\{ \frac{1}{\tilde{F}(n)} \left(\tilde{F}(n) - \tilde{F}(Z_n) \right) \geq \varepsilon \right\} \right). \end{aligned}$$

Note that \tilde{F} is a monotone increasing regularly varying function and that $Z_n \rightarrow \infty$ μ -a.s. Therefore,

$$\frac{1}{\tilde{F}(n)} \left(\tilde{F}(Z_n) - \tilde{F}(Z_n - 1) \right) \leq \frac{1}{\tilde{F}(Z_n)} \left(\tilde{F}(Z_n) - \tilde{F}(Z_n - 1) \right) \rightarrow 0 \text{ } \mu\text{-a.s.}$$

This implies for all $v \in \mathcal{P}_\mu$

$$\lim_{n \rightarrow \infty} v \left(\{\varphi \leq n\} \cap T^{-(n+1)}A^c \cap \left\{ \frac{1}{\tilde{F}(n)} \left(\tilde{F}(Z_n) - \tilde{F}(Z_n - 1) \right) \geq \varepsilon \right\} \right) = 0. \quad (3.6)$$

For $n \geq B$ large enough let $\Omega_n := \{\omega : Z_n(\omega) \geq B\}$. Then for all $\omega \in \Omega_n$

$$\begin{aligned} \tilde{F}(n) - \tilde{F}(Z_n(\omega)) &= C \exp \left(\int_B^{Z_n(\omega)} \frac{\zeta(t)}{t} dt \right) \times \\ & \quad \left[n^\beta \exp \left(\int_{Z_n(\omega)}^n \frac{\zeta(t)}{t} dt \right) - (Z_n(\omega))^\beta \right]. \end{aligned}$$

Since $|\zeta(t)| < \delta$ on $[B, \infty)$, there exists a constant C_δ , such that

$$\tilde{F}(n) - \tilde{F}(Z_n(\omega)) \leq C_\delta \left(n^{\beta+\delta} - (Z_n(\omega))^{\beta+\delta} \right) =: E.$$

Now we distinguish two cases.

Case 1: For $\beta + \delta \geq 1$, by the Mean-Value Theorem, we have

$$E \leq C_\delta (\beta + \delta) n^{\beta+\delta-1} (n - Z_n(\omega)).$$

Case 2: For $\beta + \delta < 1$ we have $E \leq C_\delta (n - Z_n(\omega))$. Hence,

$$\begin{aligned} T^{-(n+1)}A \cap \left\{ \frac{\tilde{F}(n) - \tilde{F}(Z_n)}{\tilde{F}(n)} \geq \varepsilon \right\} & \cap \Omega_n \\ & \subset T^{-(n+1)}A \cap \{n - Z_n \geq c_n \varepsilon\} \\ & = \bigcup_{c_n \varepsilon \leq k \leq n-1} \{Z_n = n - k\} \cap T^{-(n+1)}A \\ & = \bigcup_{c_n \varepsilon + 1 \leq k \leq n} T^{-(n-k+1)}(A \cap \{\varphi = k\}), \end{aligned}$$

where in Case 1 we set $c_n := \frac{\tilde{F}(n)}{n^{\beta+\delta-1} C_\delta (\beta+\delta)} = \frac{n^{1-\delta} L(n)}{C_\delta (\beta+\delta)}$ for some slowly varying function L and in Case 2 $c_n := C_\delta \tilde{F}(n)$. By the choice of δ , in both cases we have $c_n \rightarrow \infty$ as $n \rightarrow \infty$.

Using the invariance of μ we obtain

$$\begin{aligned} \mu\left(T^{-(n+1)}A \cap \left\{\frac{1}{\tilde{F}(n)}\left(\tilde{F}(n) - \tilde{F}(Z_n)\right) \geq \varepsilon\right\} \cap \Omega_n\right) \\ \leq \sum_{k=\lfloor c_n \varepsilon + 1 \rfloor}^n \mu(A \cap \{\varphi = k\}) \\ \leq \mu(A \cap \{\varphi \geq \lfloor c_n \varepsilon + 1 \rfloor\}) \\ \rightarrow 0 \quad \text{für } n \rightarrow \infty. \end{aligned}$$

This gives for $\nu \in \mathcal{P}_\mu$

$$\lim_{n \rightarrow \infty} \nu\left(T^{-(n+1)}A \cap \left\{\frac{1}{\tilde{F}(n)}\left(\tilde{F}(n) - \tilde{F}(Z_n)\right) \geq \varepsilon\right\} \cap \Omega_n\right) = 0.$$

Since $\lim_{n \rightarrow \infty} \nu(\Omega_n^c) = 0$ we conclude

$$\lim_{n \rightarrow \infty} \nu\left(T^{-(n+1)}A \cap \left\{\frac{1}{\tilde{F}(n)}\left(\tilde{F}(n) - \tilde{F}(Z_n)\right) \geq \varepsilon\right\}\right) = 0.$$

Using this, the fact $\lim_{n \rightarrow \infty} \nu(\{\varphi > n\}) = 0$, and equation (3.6) we finally conclude for all $\nu \in \mathcal{P}_\mu$

$$\begin{aligned} \lim_{n \rightarrow \infty} \nu\left(\left\{\left|\frac{\tilde{F}(Z_n \circ T) - \tilde{F}(Z_n)}{\tilde{F}(n)}\right| \geq \varepsilon\right\}\right) &\leq \lim_{n \rightarrow \infty} \nu(K_{\varepsilon, n}) + \lim_{n \rightarrow \infty} \nu(\{\varphi > n\}) \\ &= 0. \end{aligned}$$

This gives (3.3) for \tilde{F} .

We are left to show that (3.3) holds for arbitrary F . For

$$Y_n := F(Z_n \circ T) - F(Z_n) \quad \text{and} \quad \tilde{Y}_n := \tilde{F}(Z_n \circ T) - \tilde{F}(Z_n)$$

we first show

$$\frac{Y_n}{\tilde{F}(n)} - \frac{\tilde{Y}_n}{\tilde{F}(n)} \xrightarrow{\mu} 0.$$

Indeed, since $F(x) \sim \tilde{F}(x)$ for $x \rightarrow \infty$ and $Z_n \rightarrow \infty$ μ -a.s. we find for a.e. $\omega \in X$ and $\varepsilon > 0$ a number $n_0 := n_0(\omega, \varepsilon)$ such that

$$\left(1 - \frac{\varepsilon}{2}\right) \tilde{F}(Z_n(\omega)) \leq F(Z_n(\omega)) \leq \left(1 + \frac{\varepsilon}{2}\right) \tilde{F}(Z_n(\omega)) \quad \forall n \geq n_0$$

and

$$\left(1 - \frac{\varepsilon}{2}\right) \tilde{F}(Z_n \circ T(\omega)) \leq F(Z_n \circ T(\omega)) \leq \left(1 + \frac{\varepsilon}{2}\right) \tilde{F}(Z_n \circ T(\omega)) \quad \forall n \geq n_0.$$

This implies for all $n \geq n_0$

$$\left|\tilde{Y}_n(\omega) - Y_n(\omega)\right| \leq \frac{\varepsilon}{2} \left(\tilde{F}(Z_n \circ T(\omega)) + \tilde{F}(Z_n(\omega))\right)$$

and by the monotonicity of \tilde{F} we have

$$\left|\frac{\tilde{Y}_n(\omega)}{\tilde{F}(n)} - \frac{Y_n(\omega)}{\tilde{F}(n)}\right| \leq \varepsilon.$$

This shows that $\left(\frac{\tilde{Y}_n(\omega)}{\tilde{F}(n)} - \frac{Y_n(\omega)}{\tilde{F}(n)}\right) \rightarrow 0$ μ -a.e. and consequently this convergence holds also locally stochastic with respect to μ . Using $F(n) \sim \tilde{F}(n)$ this gives (3.3) for F .

□

Lemma 3.2. *Let $A \in \mathcal{A}$ be a set of positive finite measure $\mu(A)$, then*

$$\Psi_n \circ T - \Psi_n \xrightarrow{\mu} 0.$$

Proof. Let φ be the first return time to the set A . Let $\varepsilon > 0$ be given, and let

$$K_{\varepsilon,n} := \{\varphi \leq n \wedge |\Psi_n \circ T - \Psi_n| \geq \varepsilon\}.$$

Choose n large enough such that $\frac{\mu(A)}{W_n} < \varepsilon$. By (3.5) we have

$$\begin{aligned} K_{\varepsilon,n} &\subset T^{-(n+1)}A \cap \left\{ n - Z_n \geq \frac{W_n}{\mu(A)}\varepsilon \right\} \\ &= \bigcup_{\frac{W_n}{\mu(A)}\varepsilon \leq k \leq n-1} \{Z_n = n - k\} \cap T^{-(n+1)}A \\ &= \bigcup_{\frac{W_n}{\mu(A)}\varepsilon + 1 \leq k \leq n} T^{-(n-k+1)}(A \cap \{\varphi = k\}). \end{aligned}$$

Using the invariance of μ we obtain

$$\mu(K_{\varepsilon,n}) \leq \sum_{k=\lceil \frac{W_n}{\mu(A)}\varepsilon + 1 \rceil}^n \mu(A \cap \{\varphi = k\}) \leq \mu\left(A \cap \left\{ \varphi \geq \left\lceil \frac{W_n}{\mu(A)}\varepsilon + 1 \right\rceil \right\}\right),$$

and therefore

$$\lim_{n \rightarrow \infty} \mu(K_{\varepsilon,n}) = 0.$$

This implies

$$\lim_{n \rightarrow \infty} \nu(K_{\varepsilon,n}) = 0 \quad \text{for all } \nu \in \mathcal{P}_\mu.$$

Since also $\lim_{n \rightarrow \infty} \nu(\{\varphi > n\}) = 0$ we have

$$\lim_{n \rightarrow \infty} \nu(\{|\Psi_n \circ T - \Psi_n| \geq \varepsilon\}) \leq \lim_{n \rightarrow \infty} \nu(K_{\varepsilon,n}) + \lim_{n \rightarrow \infty} \nu(\{\varphi > n\}) = 0.$$

□

Now we are in the position to give the proof of Theorem 1.5.

Proof of Theorem 1.5. Let A be a uniform set for some $f \in \mathcal{P}_\mu$. We suppose without loss of generality that L is strictly monoton increasing and continuous. Then for every $x \in (0, 1)$ we have

$$\begin{aligned} \nu\left(\frac{L(Z_n)}{L(n)} \leq x\right) &= \nu(Z_n \leq a_n(x)) \\ &= \int_A \sum_{k=0}^{\lfloor a_n(x) \rfloor} 1_{A \cap \{\varphi > n-k\}} \hat{T}^k(f) d\mu, \end{aligned}$$

where ν denotes the probability measure with density $f \in \mathcal{P}_\mu$ and $a_n(x) := L^{-1}(xL(n))$. By (EL) we obtain

$$a_n(x) \rightarrow \infty \quad \text{and} \quad \frac{a_n(x)}{n} \rightarrow 0 \quad \text{for } n \rightarrow \infty.$$

Using the monotonicity of the sequence $(1_{A \cap \{\varphi > n\}})$ we obtain by the asymptotic in (1.3) on the one hand that

$$\begin{aligned} \mathbf{v} \left(\frac{L(Z_n)}{L(n)} \leq x \right) &\leq \int_A 1_{A \cap \{\varphi > n - \lfloor a_n(x) \rfloor\}} \sum_{k=0}^{\lfloor a_n(x) \rfloor} \hat{T}^k(f) \, d\mu \\ &\sim \mu(A \cap \{\varphi > n - \lfloor a_n(x) \rfloor\}) \cdot L(\lfloor a_n(x) \rfloor). \end{aligned}$$

This and (UA) imply

$$\limsup \mathbf{v} \left(\frac{L(Z_n)}{L(n)} \leq x \right) \leq x.$$

On the other hand we derive in a similar way

$$\begin{aligned} \mathbf{v} \left(\frac{L(Z_n)}{L(n)} \leq x \right) &\geq \int_A 1_{A \cap \{\varphi > n\}} \sum_{k=0}^{\lfloor a_n(x) \rfloor} \hat{T}^k(f) \, d\mu \\ &\sim \mu(A \cap \{\varphi > n\}) \cdot L(\lfloor a_n(x) \rfloor). \end{aligned}$$

This gives the opposite inequality

$$\liminf \mathbf{v} \left(\frac{L(Z_n)}{L(n)} \leq x \right) \geq x.$$

Finally the theorem follows from Lemma 3.1 (for the case $\beta = 0$) by the compactness theorem. \square

For the proof of Theorem 1.6 we need the following lemma.

Lemma 3.3. *Let $A \in \mathcal{A}$ with $0 < \mu(A) < \infty$ be a uniformly returning set, $W_n \sim L(n)$, where L satisfies the properties stated in (EL), and $x \in (0, 1)$ fixed. Then for all $\varepsilon \in (0, 1)$ there exists n_0 such that for all $n \geq n_0$ and $k \in [n - a_n(x), n]$ we have uniformly on A*

$$(1 - \varepsilon) \frac{1}{W_n} \leq \hat{T}^k(f) \leq (1 + \varepsilon)^2 \frac{1}{W_n},$$

where $a_n(x)$ is defined as in (EL).

Proof. Due to Proposition 1.2 we have

$$W_n \hat{T}^n(f) \longrightarrow 1 \quad \mu - \text{a.e. uniformly on } A.$$

Thus, for all $\varepsilon \in (0, 1)$ there exists $k_0 := k_0(\varepsilon)$ such that we have uniformly on A

$$(1 - \varepsilon) \frac{1}{W_k} \leq \hat{T}^k(f) \leq (1 + \varepsilon) \frac{1}{W_k} \quad \text{for all } k \geq k_0.$$

By (EL) there exist n_1 and n_2 such that

$$n - a_n(x) \geq k_0 \quad \text{for all } n \geq n_1 \quad \text{and} \quad \frac{1}{W_{[n - a_n(x)]}} \leq (1 + \varepsilon) \frac{1}{W_n} \quad \text{for all } n \geq n_2.$$

Let us denote $n_0 := \max\{n_1, n_2\}$. Then by monotonicity of W_n we obtain uniformly on A that, for all $n \geq n_0$ and $k \in [n - a_n(x), n]$,

$$(1 - \varepsilon) \frac{1}{W_n} \leq (1 - \varepsilon) \frac{1}{W_k} \leq \hat{T}^k(f) \leq (1 + \varepsilon) \frac{1}{W_k} \leq \frac{(1 + \varepsilon)}{W_{[n - a_n(x)]}} \leq \frac{(1 + \varepsilon)^2}{W_n}.$$

\square

Proof of Theorem 1.6. Let $W_n \sim L(n)$ as $n \rightarrow \infty$, without loss of generality we may assume that L is monotone increasing and continuous. We have for every fixed $x \in (0, 1)$

$$\begin{aligned} \mathbf{v} \left(\frac{L(n - Z_n)}{L(n)} \leq x \right) &= \mathbf{v}(Z_n \geq n - a_n(x)) \\ &= \int_A \sum_{n - a_n(x) \leq k \leq n} \mathbf{1}_{A \cap \{\varphi > n - k\}} \hat{T}^k(f), \end{aligned}$$

where $a_n(x) = L^{-1}(xL(n))$. Let $\varepsilon > 0$. From Lemma 3.3 it follows that, for sufficiently large n

$$\mathbf{v} \left(\frac{L(n - Z_n)}{L(n)} \leq x \right) \leq (1 + \varepsilon)^2 \frac{1}{W_n} W_{[a_n(x)]} \sim (1 + \varepsilon)^2 x.$$

Similarly for sufficiently large n ,

$$x(1 - \varepsilon) \sim (1 - \varepsilon) \frac{1}{W_n} W_{[a_n(x)]} \leq \mathbf{v} \left(\frac{L(n - Z_n)}{L(n)} \leq x \right).$$

Both inequalities give

$$x(1 - \varepsilon) \leq \liminf \mathbf{v} \left(\frac{L(n - Z_n)}{L(n)} \leq x \right) \leq \limsup \mathbf{v} \left(\frac{L(n - Z_n)}{L(n)} \leq x \right) \leq (1 + \varepsilon)^2 x.$$

Since ε was arbitrary, we obtain

$$\mathbf{v} \left(\frac{L(n - Z_n)}{L(n)} \leq x \right) \rightarrow x \quad \text{as } n \rightarrow \infty \quad \text{for all } x \in (0, 1). \quad (3.7)$$

To show that the above convergence still holds if we replace L by the wandering rate, we firstly point out that (3.7) in particular implies

$$n - Z_n \rightarrow \infty \quad \text{in probability (w.r.t. } \mathbf{v}) \quad \text{as } n \rightarrow \infty. \quad (3.8)$$

This can be seen as follows. At first note that (3.7) is equivalent to

$$\mathbf{v}(n - Z_n \leq a_n(x)) \rightarrow x \quad \text{as } n \rightarrow \infty \quad \text{for all } x \in (0, 1). \quad (3.9)$$

Now we suppose that (3.8) fails. Then there exists $\varepsilon > 0$, a monotone increasing sequence $t_n \nearrow \infty$ and an integer N such that

$$\mathbf{v}(t_n - Z_{t_n} \leq N) \geq \varepsilon \quad \text{for all } n \in \mathbb{N}.$$

For arbitrary but fixed $x \in (0, \varepsilon)$ we have $\lim_{n \rightarrow \infty} a_n(x) = \infty$. Hence, there exists $n_0 \in \mathbb{N}$ with

$$a_{t_n}(x) \geq N \quad \text{for all } n \geq n_0.$$

Thus,

$$\mathbf{v}(t_n - Z_{t_n} \leq a_{t_n}(x)) \geq \mathbf{v}(t_n - Z_{t_n} \leq N) \geq \varepsilon \quad \text{for all } n \geq n_0.$$

This implies

$$\lim_{n \rightarrow \infty} \mathbf{v}(t_n - Z_{t_n} \leq a_{t_n}(x)) \geq \varepsilon,$$

contradicting (3.9).

Finally, since $n - Z_n \rightarrow \infty$ in probability, it is clear that the slowly varying function L may be replaced by any function L_1 with $L_1(n) \sim C \cdot L(n)$, $C > 0$, as $n \rightarrow \infty$. From this and Lemma 3.2 by the compactness result the theorem follows. \square

REFERENCES

- [Aar97] J. Aaronson. *An introduction to infinite ergodic theory*, volume 50 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [BGT89] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [Eri70] K. B. Erickson. Strong renewal theorems with infinite mean. *Trans. Amer. Math. Soc.*, 151:263–291, 1970.
- [Fel71] W. Feller. *An introduction to probability theory and its applications. Vol. II*. Second edition. John Wiley & Sons Inc., New York, 1971.
- [Kar33] J. Karamata. Sur un mode de croissance régulière. Théorèmes fondamentaux. *Bull. Soc. Math. Fr.*, 61:55–62, 1933.
- [KS05] M. Kesseböhmer and M. Slassi. A distributional limit law for continued fraction digit sums. *preprint: arXiv:math.NT/0509559*, pages 1–15, 2005.
- [Sen76] E. Seneta. *Regularly varying functions*. Springer-Verlag, Berlin, 1976. Lecture Notes in Mathematics, Vol. 508.
- [Tha95] M. Thaler. A limit theorem for the Perron-Frobenius operator of transformations on $[0, 1]$ with indifferent fixed points. *Israel J. Math.*, 91(1-3):111–127, 1995.
- [Tha98] M. Thaler. The Dynkin-Lamberti arc-sine laws for measure preserving transformation. *Trans. Amer. Math. Soc.*, 350:4593–4607, 1998.
- [Tha00] M. Thaler. The asymptotics of the Perron-Frobenius operator of a class of interval maps preserving infinite measures. *Studia Math.*, 143(2):103–119, 2000.
- [TZ] M. Thaler and R. Zweimüller. Distributional limit theorems in infinite ergodic theory. *preprint*.
- [Zwe03] R. Zweimüller. Stable limits for probability preserving maps with indifferent fixed points. *Stoch. Dyn.*, 3(1):83–99, 2003.

E-mail address: mhk@math.uni-bremen.de, slassi@math.uni-bremen.de

UNIVERSITÄT BREMEN, FACHBEREICH 3 FÜR MATHEMATIK UND INFORMATIK, BIBLIOTHEKSTRASSE 1,
D–28359 BREMEN, GERMANY.