

On the alternating sums of powers of consecutive q -integers

Taekyun Kim

Institute of Science Education,
Kongju National University, Kongju 314-701, Korea
e-mail: tkim @kongju.ac.kr

Abstract In this paper we construct q -Genocchi numbers and polynomials. By using these numbers and polynomials, we investigate the q -analogue of alternating sums of powers of consecutive integers due to Euler.

2000 Mathematics Subject Classification : 11S80, 11B68

Key words : Genocchi numbers and polynomials, q -Genocchi numbers and polynomials, alternating sums of powers

1. Introduction

The Genocchi numbers G_m are defined by the generating function:

$$F(t) = \frac{2t}{e^t + 1} = \sum_{m=0}^{\infty} G_m \frac{t^m}{m!}, (|t| < \pi), \text{ cf. [3]} \quad (1)$$

where we use the technique method notation by replacing G^m by $G_m (m \geq 0)$ symbolically. It satisfies $G_1 = 1, G_3 = G_5 = G_7 = \dots = 0$, and even coefficients are given $G_m = 2(1 - 2^{2m})B_{2m} = 2mE_{2m-1}$, where B_m are Bernoulli numbers and E_m are Euler numbers which are defined by

$$\frac{2}{e^t + 1} = \sum_{m=0}^{\infty} E_m \frac{t^m}{m!}, \text{ cf. [5, 6].}$$

For $x \in \mathbb{R}$ (= the field of real numbers), we consider the Genocchi polynomials as follows:

$$F(x, t) = F(t)e^{xt} = \frac{2t}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}. \quad (2)$$

Note that $G_m(x) = \sum_{k=0}^m \binom{m}{k} G_k x^{m-k}$. Let us also define the Genocchi polynomials of order r as follows:

$$2 \left(\frac{1}{1 + e^t} \right)^r e^{xt} = \sum_{n=0}^{\infty} G_n^{(r)}(x) \frac{t^n}{n!}, \text{ cf. [3].}$$

In the special case $x = 0$, we define $G_n^{(r)}(0) = G_n^{(r)}$. What is the value of the following sum for a given positive integer k ?

$$1^k + 2^k + 3^k + \dots + n^k.$$

Let us denote this sum by $f_k(n)$. Finding formulas for $f_k(n)$ has interested mathematicians for more than 300 years since the time of Jacob Bernoulli, cf. [1, 7, 9]. It was well known that

$$f_n(k-1) = \frac{1}{n+1} \sum_{i=0}^n \binom{n+1}{i} B_i k^{n+1-i}, \text{ cf. [9] ,} \quad (3)$$

where $\binom{n}{k}$ is binomial coefficients.

Let n, k be positive integers ($k > 1$), and let

$$T_n(k) = -1^k + 2^k - 3^k + 4^k - 5^k + \dots + (-1)^{k-1}(n-1)^k.$$

Following an idea due to Euler, it was known that

$$T_n(k) = \frac{(-1)^{k+1}}{2} \sum_{l=0}^{k-1} \binom{n}{l} E_l k^{n-l} + \frac{E_n}{2} (1 + (-1)^{k+1}), \text{ cf. [5].} \quad (4)$$

Let q be an indeterminate which can be considered in complex number field, and for any integer k define the q -integer as

$$[k]_q = \frac{q^k - 1}{q - 1} = 1 + q + \dots + q^{k-1}.$$

Throughout this paper we assume that $q \in \mathbb{C}$ with $0 < q < 1$. Recently many authors studied q -analogue of the sums of powers of consecutive integers. In [2], Garrett and Hummel gave a combinatorial proof of a q -analogue of $\sum_{k=1}^n k^3 = \binom{n+1}{2}^2$ as follows:

$$\sum_{k=1}^n q^{k-1} [k]_q^2 \left(\left[\frac{k-1}{2} \right]_{q^2} + \left[\frac{k+1}{2} \right]_{q^2} \right) = \left[\begin{matrix} n+1 \\ 2 \end{matrix} \right]_q^2,$$

where

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_q = \prod_{j=1}^k \frac{[n+1-j]_q}{[j]_q} \text{ denotes the } q\text{-binomial coefficients.}$$

Garrett and Hummel, in their paper, asked for a simpler q -analogue of the sums of cubes. As a response to Garrett and Hummel's question, Warnaar gave a simple q -analogue of the sums of cubes as follows:

$$\sum_{k=1}^n q^{2n-2k} [k]_q^2 [k]_{q^2} = \left[\begin{matrix} n+1 \\ 2 \end{matrix} \right]_q^2, \text{ cf. [10] .}$$

Let

$$f_{m,q}(n) = \sum_{k=1}^n [k]_{q^2} [k]_q^{m-1} q^{(n-k)\frac{m+1}{2}}.$$

Then we note that $\lim_{q \rightarrow 1} f_{m,q}(n) = f_m(n)$.

Warnaar [10] (for $m = 3$) and Schlosser [8] gave formulae for $m = 1, 2, 3, 4, 5$ as the meaning of the q -analogues of the sums of consecutive integers, squares, cubes, quarts and quints. Let $n, k (> 1)$ be the positive integers. In the recent paper, it was known that

$$\sum_{j=0}^{k-1} q^j [j]_q^n = \frac{1}{n+1} \sum_{j=0}^n \binom{n+1}{j} \beta_{j,q} q^{kj} [k]_q^{n+1-j} - \frac{1}{n+1} (1 - q^{(n+1)k}) \beta_{n+1,q}, \text{ see [4] ,}$$

where $\beta_{j,q}$ are called Carlitz's q -Bernoulli numbers. Originally q -Genocchi numbers and polynomials were introduced by Kim-Jang-Pak in 2001 [3], but they do not seem to be the most natural ones. In this paper we give another construction of a q -Genocchi numbers and polynomials which are different than q -Genocchi numbers and polynomials of Kim-Jang-Pak in 2001 [3]. By using these numbers and polynomials, we investigate the q -analogue of alternating sums of powers of consecutive integers.

2. q -Genocchi numbers and polynomials

Let $F_{q,k}(t)$ be the generating functions of the q -Genocchi numbers as follows:

$$F_{q,k}(t) = [2]_q t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} (-1)^{j-1} \exp\left(t[j]_{q^2} q^{\frac{k-j}{2}}\right) = \sum_{j=0}^{\infty} G_{n,q} \frac{t^n}{n!}$$

By using Taylor expansion in the above, we see that

$$\begin{aligned} \sum_{j=0}^{\infty} G_{n,k,q} \frac{t^n}{n!} &= [2]_q t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} (-1)^{j-1} \sum_{n=0}^{\infty} \frac{[j]_q^n q^{\frac{n(k-j)}{2}}}{n!} t^n \\ &= [2]_q t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} (-1)^{j-1} \sum_{n=0}^{\infty} \left\{ \frac{1}{(1-q)^n} q^{\frac{n(k-j)}{2}} \sum_{m=0}^n \binom{n}{m} (-1)^m q^{jm} \right\} \frac{t^n}{n!} \\ &= \frac{[2]_q t}{1-q^2} \sum_{n=0}^{\infty} \frac{1}{(1-q)^n} q^{k+\frac{nk}{2}} \sum_{m=0}^n \binom{n}{m} (-1)^m \left(\sum_{j=0}^{\infty} (-1)^{j-1} q^{mj-j-\frac{nj}{2}} (1-q^{2j}) \right) \frac{t^n}{n!} \\ &= \frac{[2]_q t}{1-q^2} \sum_{n=0}^{\infty} \frac{1}{(1-q)^n} \sum_{m=0}^n \binom{n}{m} (-1)^m \frac{q^{m-1-\frac{n}{2}+k+\frac{nk}{2}} (1-q^2)}{(1+q^{m-1-\frac{n}{2}})(1+q^{m+1-\frac{n}{2}})} \frac{t^n}{n!}. \end{aligned}$$

Note that $G_{0,k,q} = 0$. Hence, we have

$$\begin{aligned} \sum_{n=1}^{\infty} G_{n,k,q} \frac{t^n}{n!} &= t \sum_{n=1}^{\infty} \frac{1}{(1-q)^n} \sum_{m=1}^n \binom{n-1}{m-1} \frac{(-1)^{m-1} q^{m+k+\frac{(n-1)k}{2}-\frac{n-1}{2}-2}}{(1+q^{-2+m-\frac{n-1}{2}})(1+q^{m-\frac{n-1}{2}})} \frac{t^{n-1}}{(n-1)!} \\ &= \sum_{n=1}^{\infty} \frac{1}{(1-q)^n} \sum_{m=1}^n \binom{n}{m} \frac{(-1)^{m-1} m q^{m+k+\frac{(n-1)(k-1)}{2}-2}}{(1+q^{-2+m-\frac{n-1}{2}})(1+q^{m-\frac{n-1}{2}})} \frac{t^n}{n!} \end{aligned}$$

By comparing the coefficients of $\frac{t^n}{n!}$ on both sides of the above equation, we obtain the below:

Theorem 1. Let $k, n (n \geq 1)$ be positive integers. Then we have

$$G_{n,k,q} = \left(\frac{1}{1-q} \right)^n \sum_{m=1}^n \binom{n}{m} \frac{(-1)^{m-1} m q^{m+k+\frac{(n-1)(k-1)}{2}-2}}{(1+q^{-2+m-\frac{n-1}{2}})(1+q^{m-\frac{n-1}{2}})}.$$

We also define the generating function.

$$F_{q,k}(t, k) = [2]_q t \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} (-1)^{j+k-1} \exp\left(t[j+k]_q q^{-\frac{j}{2}}\right) = \sum_{n=0}^{\infty} G_{n,k,q}(k) \frac{t^n}{n!}$$

By using the binomial theorem and some elementary calculations in the above equation, we

have

$$\begin{aligned}
\sum_{n=0}^{\infty} G_{n,k,q}(k) \frac{t^n}{n!} &= \frac{[2]_q t}{1-q^2} \sum_{j=0}^{\infty} q^{-j} (1-q^{2j+2k}) (-1)^{j+k-1} \sum_{n=0}^{\infty} [j+k]_q^n q^{-\frac{j}{2}n} \frac{t^n}{n!} \\
&= \frac{[2]_q t}{1-q^2} \sum_{j=0}^{\infty} q^{-j} (1-q^{2j+2k}) (-1)^{j+k} \sum_{n=0}^{\infty} \left\{ \left(\frac{1}{1-q} \right)^n \sum_{m=0}^n \binom{n}{m} (-1)^m q^{j m + k m} q^{-\frac{j}{2}n} \right\} \frac{t^n}{n!} \\
&= \frac{[2]_q t}{1-q^2} \sum_{n=0}^{\infty} \left(\frac{1}{1-q} \right)^n \sum_{m=0}^n \binom{n}{m} (-1)^{m+k} q^{m k} \sum_{j=0}^{\infty} (1-q^{2j+2k}) (-1)^j q^{-j+jm-\frac{j}{2}n} \frac{t^n}{n!} \\
&= \frac{[2]_q t}{1-q^2} \sum_{n=0}^{\infty} \left(\frac{1}{1-q} \right)^n \sum_{m=0}^n \binom{n}{m} (-1)^{m+k} q^{m k} \left\{ \frac{1}{1+q^{m-\frac{n}{2}-1}} - \frac{q^{2k}}{1+q^{1+m-\frac{n}{2}}} \right\} \frac{t^n}{n!} \\
&= t \sum_{n=0}^{\infty} \left(\frac{1}{1-q} \right)^{n+1} \sum_{m=0}^n \binom{n}{m} (-1)^{m+k} \left(\frac{q^{m k}}{1+q^{m-\frac{n}{2}-1}} - \frac{q^{(m+2)k}}{1+q^{1+m-\frac{n}{2}}} \right) \frac{t^n}{n!} \\
&= t \sum_{n=1}^{\infty} \left(\frac{1}{1-q} \right)^n \sum_{m=1}^n \binom{n-1}{m-1} (-1)^{m-1+k} \left(\frac{q^{(m-1)k}}{1+q^{m-2-\frac{n-1}{2}}} - \frac{q^{(m+1)k}}{1+q^{m-\frac{n-1}{2}}} \right) \frac{t^{n-1}}{(n-1)!} \\
&= \sum_{n=1}^{\infty} \left(\frac{1}{1-q} \right)^n \sum_{m=1}^n \binom{n}{m} (-1)^{m-1+k} \left(\frac{m q^{(m-1)k}}{1+q^{m-2-\frac{n-1}{2}}} - \frac{m q^{(m+1)k}}{1+q^{m-\frac{n-1}{2}}} \right) \frac{t^n}{n!}.
\end{aligned}$$

Note that $G_{0,k,q}(k) = 0$. Therefore we obtain the following theorem.

Theorem 2. Let $k, n (n \geq 1)$ be positive integers. Then we have

$$G_{n,k,q}(k) = \left(\frac{1}{1-q} \right)^n \sum_{m=1}^n \binom{n}{m} (-1)^{m-1+k} \left(\frac{m q^{(m-1)k}}{1+q^{m-2-\frac{n-1}{2}}} - \frac{m q^{(m+1)k}}{1+q^{m-\frac{n-1}{2}}} \right).$$

Remark 3. Note that

$$(1) \lim_{q \rightarrow 1} G_{n,k,q} = G_n^{(2)}, \quad (2) \lim_{q \rightarrow 1} G_{n,k,q}(k) \neq G_n^{(2)}(k).$$

It is easy to see that

$$\begin{aligned}
&[2]_q t \sum_{j=0}^{\infty} q^{k-j} [j]_{q^2} (-1)^{j-1} \exp\left(t [j]_{q^2} q^{\frac{k-j}{2}}\right) - [2]_q t \sum_{j=0}^{\infty} q^{-j} [j+k]_{q^2} (-1)^{j-1+k} \exp\left(t [j+k]_q q^{-\frac{j}{2}}\right) \\
&= [2]_q t \sum_{j=0}^{k-1} (-1)^{j-1} [j]_{q^2} q^{k-j} \exp\left(t [j]_q q^{\frac{k-j}{2}}\right).
\end{aligned}$$

Thus, we easily see that

$$[2]_q \sum_{j=0}^{k-1} [j]_{q^2} (-1)^{j-1} [j]_q^{n-1} q^{\frac{(k-j)(n+1)}{2}} = \frac{G_{n,k,q} - G_{n,k,q}(k)}{n}.$$

Therefore we obtain the following theorem.

Theorem 4. Let $k, n (n \geq 1)$ be positive integers. Then we have

$$\sum_{j=0}^{k-1} [j]_{q^2} (-1)^{j-1} [j]_q^{n-1} q^{\frac{(k-j)(n+1)}{2}} = \frac{G_{n,k,q} - G_{n,k,q}(k)}{n [2]_q}$$

References

- [1] J. FAULHABER, *Academia Algebrae, Darinnen die miraculosische inventiones zu den höchsten Cossen weiters continuirt und profitiert werden*, Augspurg, bey Johann Ulrich Schönigs, 1631.
- [2] K.C. GARRETT, K. HUMMEL, A combinatorical proof of the sum of q -cubes, *Electro. J. Comb.*, **11** #R 9, (2004).
- [3] T. KIM, L.-C. JANG, H. K. PAK, A note on q -Euler and Genocchi numbers , *Proc. Japan Acad.* , **77 A**, 139-141 (2001).
- [4] T. KIM, Sums of powers of consecutive q -integers, *Advan. Stud. Contemp. Math.*, **9**, 15-18 (2004).
- [5] T. KIM, A note on the alternating sums of powers of consecutive integers, *arXiv.org:math/0508233* , **1**, 1-4 (2005).
- [6] T. KIM, Analytic continuation of multiple q -Zeta functions and their values at negative integers , *Russ. J. Math. Phys.*, **11 (1)**, 71-76 (2004).
- [7] D. E. KNUTH, Johann Faulhaber and sums of powers, *Math. Comput.*, **61**, 277-294 (1993).
- [8] M. SCHLOSSER, q -analogues of the sums of consecutive integers squares, cubes, quarts, quints, *Electro. J. Comb.*, **11**, # R 71 (2004).
- [9] Y.-Y. SHEN, A note on the sums of powers of consecutive integers, *Tunghai Science*, **5**,101-106 (2003).
- [10] S. O. WARNAAR, On the q -analogues of the sums of cubes, *Electro. J. Comb.*, **11**,#N 13 (2004).