

FUNCTION SPACES ASSOCIATED WITH SCHRÖDINGER OPERATORS: THE PÖSCHL-TELLER POTENTIAL

GESTUR ÓLAFSSON AND SHIJUN ZHENG

ABSTRACT. We address the function space theory associated with the Schrödinger operator $H = -d^2/dx^2 + V$. The discussion is featured with potential $V(x) = -\lambda(\lambda-1)\operatorname{sech}^2 x$, which is called in quantum physics the Pöschl-Teller potential. Using biorthogonal dyadic system, we introduce Besov spaces and Triebel-Lizorkin spaces (including Sobolev spaces) associated with H . We then use interpolation method to identify these spaces with the classical ones for a certain range of $p, q > 1$. A physical implication is that the corresponding wave function $\psi(t, x) = e^{-itH}f(x)$ admits appropriate time decay in the Besov space scale.

INTRODUCTION

Let $H = -d^2/dx^2 + V$ be a Schrödinger operator on \mathbb{R} with real valued potential function V . In quantum physics, H is the energy operator of a particle with one degree of freedom with potential $V(x)$. The Schrödinger equation

$$i \partial_t \psi = H\psi, \quad \psi(0, x) = f(x)$$

has solution given by $\psi(t, x) = e^{-itH}f(x)$. If the potential has certain decay at ∞ , then one expects that asymptotically, as time tends to infinity, the motion of the associated perturbed quantum system resembles the free evolution. In fact, if V is in $L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ or if $\int_{\mathbb{R}} (1 + |x|^2)|V(x)|dx < \infty$, then it is known, c.f. [CK01], [GH98], [Z04a], that the essential spectrum of H and $H_0 = -d^2/dx^2$ coincide and there is no singular continuous spectrum; the wave operators $W_{\pm} = s - \lim_{t \rightarrow \pm\infty} e^{itH}e^{-itH_0}$ exists and are complete.

Recently, several authors have considered function spaces associated with Schrödinger operators, cf. [JN94, E95, E96, DZ98, DZ02, BZ04]. One of the goals has been to develop the associated Littlewood-Paley theory, in order to give a unified approach. Motivated by the treatment in [BZ04, E95] we study the negative potential

$$(0.1) \quad V(x) = -\lambda(\lambda - 1)\operatorname{sech}^2 x$$

Date: July 19, 2021.

2000 Mathematics Subject Classification. Primary: 42B25; Secondary: 35P25.

Key words and phrases. spectral multiplier, Schrödinger operator, Littlewood-Paley theory.

G. Ólafsson was supported by NSF grants DMS-0139473 and DMS-0402068. S. Zheng was partially supported by DARPA grant MDA 972-01-1-0033. He also gratefully thanks NSF for support during his visit at Louisiana State University in 2003.

called Pöschl-Teller potential. It is a fundamental model in soliton theory. In this article, we will mainly consider the case where $\lambda = n + 1$, $n \in \mathbb{N}_0$. In that case we obtain a compact expression for eigenfunctions of H . It turns out that for the absolute continuous part of H , the analysis is simpler than the barrier potential, although H has a nonempty pure point spectrum.

Denote by $C_0(\mathbb{R})$ the space of compactly supported functions on the line. Suppose $\Phi, \varphi, \Psi, \psi \in C_0^\infty(\mathbb{R})$ satisfy the ‘‘biorthogonal’’ condition (see Section ??)

$$(0.2) \quad \Phi(x)\Psi(x) + \sum_{j=1}^{\infty} \varphi_j(x)\psi_j(x) = 1,$$

where $\varphi_j(x) = \varphi(2^{-j}x)$. Let $\alpha \in \mathbb{R}$, $0 < p, q < \infty$. The *Triebel-Lizorkin spaces* associated with H , denoted by $F_p^{\alpha,q} := F_p^{\alpha,q}(H)$, is defined to be the completion of the subspace $L_0^2(\mathbb{R}) := \{f \in L^2(\mathbb{R}) : \|f\|_{F_p^{\alpha,q}} < \infty\}$, where the quasi-norm $\|\cdot\|_{F_p^{\alpha,q}}$ is initially defined for $f \in L^2(\mathbb{R})$ by:

$$(0.3) \quad \|f\|_{F_p^{\alpha,q}} = \left\| \left(\sum_{j=0}^{\infty} 2^{j\alpha q} |\varphi_j(H)f|^q \right)^{1/q} \right\|_p.$$

The main result is an equivalence theorem for $F_p^{\alpha,q}(H)$ using Peetre’s maximal function, which suggests that that $\|f\|_{F_p^{\alpha,q}}^\phi$ and $\|f\|_{F_p^{\alpha,q}}^\psi$ are equivalent quasi-norms on $F_p^{\alpha,q}(H)$, for given two systems of smooth dyadic systems $\{\phi_j\}, \{\psi_j\}$. The following is proved in Theorem 2.7:

Theorem 0.1. *Let $\alpha \in \mathbb{R}$, $0 < p, q \leq \infty$. If $\varphi_j^* f$ is defined for $j \geq 0$ with $s > 1/\min(p, q)$, we have for $f \in L^2(\mathbb{R})$*

$$(0.4) \quad \|f\|_{F_p^{\alpha,q}} \sim \|\Phi^* f\|_p + \left\| \left(\sum_{j=1}^{\infty} (2^{j\alpha} \varphi_j^* f)^q \right)^{1/q} \right\|_p.$$

Furthermore, $F_p^{\alpha,q}$ is a quasi-Banach space (Banach space if $p \geq 1, q \geq 1$) and it is independent of the choice of $\{\Phi, \varphi_j\}_{j \geq 1}$.

Here $\varphi_j^* f = \varphi_{j,s}^* f$ denotes the Peetre type maximal function, see Section ?? for details.

As in [BZ04] the essential part is to obtain the decay estimates of the kernel of the operator $\phi_\lambda(H)$ as well as of its derivatives. This requires an examination of the differential properties of the eigenfunctions $e_k(x) := e(x, k)$. For $k = i, 2i, \dots, ni$, where as usually $i = \sqrt{-1}$, the bounded states are Schwartz functions. In Section 1 we solve the Lippmann-Schwinger equation (1.5) for $k \in \mathbb{R} \cup \{i, \dots, ni\}$ based on a general expression suggested in [Lam80, p.40, p.54]. Furthermore, we determine precisely the transmission and reflection coefficients.

We mention that our results are also true for translation and regular scaling (i.e., $V \rightarrow \alpha^2 V(\alpha)$) of the potential V .

A natural question is: what is the relation between the new function spaces and classical ones, namely, $F_p^{\alpha,q}(\mathbb{R})$ and $B_p^{\alpha,q}(\mathbb{R})$. Concerning identification of the F and B spaces, it can be shown that the following holds:

Theorem 5.2 If $\alpha > 0$, $1 < p < \infty$ and $2p/(p+1) < q < 2p$, then

$$F_p^{\alpha,q}(H) = F_p^{2\alpha,q}(\mathbb{R})$$

and if $\alpha > 0$, $1 \leq p < \infty$, $1 \leq q \leq \infty$, then

$$B_p^{\alpha,q}(H) = B_p^{2\alpha,q}(\mathbb{R}).$$

For $0 < p \leq 1$, it would be interesting to see whether it holds that $F_p^{0,1} = \mathcal{H}^p$, where \mathcal{H}^p is the Hardy spaces associated with H defined by atomic decomposition or the maximal function associated to the heat kernel, cf. [DZ98, DZ02].

In a sequel to this paper we will study the spectral multiplier problem on the F and B spaces.

1. THE EIGENFUNCTIONS OF H

In this section we derive a simple expression for the eigenfunctions on the Schrödinger operator $H = -d^2/dx^2 - \lambda(\lambda-1)\text{sech}^2(x)$ in the case where $\lambda = n+1$ is an integer.

1.1. The Lippmann-Schwinger equation. Consider the eigenvalue problem

$$(1.1) \quad He(x, k) = k^2 e(x, k),$$

with asymptotic behavior

$$(1.2) \quad e_{\pm}(x, k) \sim \begin{cases} T_{\pm}(k)e^{ikx} & \text{for } x \rightarrow \pm\infty \\ e^{ikx} + R_{\pm}(k)e^{-ikx} & \text{for } x \rightarrow \mp\infty \end{cases}$$

We will often use the notation

$$(1.3) \quad e(x, k) = \begin{cases} e_+(x, k) & \text{for } k > 0 \\ e_-(x, k) & \text{for } k < 0 \end{cases}$$

The coefficients $T_{\pm}(k)$ and $R_{\pm}(k)$ in (1.2) are called the *transmission coefficients* and *reflection coefficients*, resp. They satisfy the relation

$$(1.4) \quad |T_{\pm}(k)|^2 + |R_{\pm}(k)|^2 = 1.$$

The equations (1.1) and (1.2) are equivalent to the Lippmann-Schwinger equation

$$(1.5) \quad e_{\pm}(x, k) = e^{ikx} + \frac{1}{2i|k|} \int e^{i|k||x-y|} V(y) e_{\pm}(y, k) dy,$$

where \pm indicates the sign of k . For $V(x) = -\lambda(\lambda-1)\text{sech}^2 x$, we seek a solution to the integral equation (1.5) for each k .

1.2. Inductive construction of the solution. Let us recall how to obtain the solution for the eigenvalue problem (1.1) for integer values of $\lambda = n + 1$. By regularity, a weak solution u solving $Hu = k^2u$ is smooth on the domain where u is smooth. For $n \in \mathbb{N}_0$ denote by T_n the differential operator

$$(1.6) \quad T_n = \frac{d}{dx} - n \tanh x.$$

With $y_n = e(x, k)$ the equation (1.1) can then be written as

$$(1.7) \quad y_n'' + n(n+1)\operatorname{sech}^2 x y_n = -k^2 y_n.$$

If $n = 0$, then the general solution is given by $y_0 = Ae^{ikx} + Be^{-ikx}$. For $n \geq 1$ we obtain

$$(1.8) \quad y_n(x) = T_n y_{n-1}(x) = T_n \cdots T_1 y_0(x) = A(k)e^{ikx} + B(k)e^{-ikx}.$$

Lemma 1.3. *Assume that $\lambda = n + 1$, with $n \in \mathbb{N}_0$. Then there exists a polynomial $p_n(t, ik)$ of degree n in both t and k , such that*

$$e_{\pm}(x, k) = A_n^{\pm}(k)p_n^{\pm}(\tanh x, ik)e^{ikx}.$$

Furthermore the following holds:

(1) The constant $A_n^{\pm}(k)$ is given by

$$A_n^+(k) = \prod_{j=1}^n \frac{1}{j + ik} \quad \text{and} \quad A_n^-(k) = (-1)^n \prod_{j=1}^n \frac{1}{j - ik}.$$

(2) The transmission coefficient $T_{\pm}(k)$ are

$$T_+(k) = (-1)^n \prod_{j=1}^n \frac{j - ik}{j + ik} \quad \text{and} \quad T_-(k) = (-1)^n \prod_{j=1}^n \frac{j + ik}{j - ik}.$$

(3) The reflection coefficients $R_{\pm}(k)$ are all zero.

Proof. As n is fixed here, we will omit the subscript n in the following except in obvious induction steps. As $\frac{d}{dx}(\tanh x) = 1 - \tanh^2 x$ it follows from (1.8) that

$$(1.9) \quad e_{\pm}(x, k) = A^{\pm}(k)p_n^{\pm}(x, ik)e^{ikx} + B(k)q_n^{\pm}(x, ik)e^{-ikx},$$

where $p^{\pm}(x, k) := p^{\pm}(\tanh x, ik)$ and $q^{\pm}(x, k) = q^{\pm}(\tanh x, ik)$ are polynomials of degree n in k and $\tanh x$.

Assume for the moment, that $k > 0$. We will therefore also omit the superscript \pm for the moment. Substituting (1.9) into the asymptotics for $k > 0$,

$$e(x, k) \longrightarrow \begin{cases} T_+(k)e^{ikx}, & x \rightarrow \infty, \\ e^{ikx} + R_+(k)e^{-ikx}, & x \rightarrow -\infty. \end{cases}$$

we obtain $B(k) = 0 = R_+(k)$,

$$(1.10) \quad A(k)p(-1, ik) = 1$$

and

$$(1.11) \quad T_+(k) = A(k)p(1, ik).$$

This implies easily the recurrence formula

$$(1.12) \quad p_n(\tanh x, ik) = \frac{d}{dx}p_{n-1}(\tanh x, ik) + (ik - n \tanh x)p_{n-1}(\tanh x, ik).$$

Note that $\frac{d}{dx}p_{n-1}(\tanh x, ik) = p'_{n-1}(\tanh x, ik)(1 - \tanh^2(x))$. As $p'_{n-1}(t, ik)$ is a polynomial in t it follows that

$$\lim_{x \rightarrow \pm\infty} p'_{n-1}(\tanh x, ik) = p'_{n-1}(\pm 1, ik)$$

is bounded, and hence

$$(1.13) \quad \lim_{x \rightarrow \pm\infty} \frac{d}{dx}p_{n-1}(\tanh x, ik) = 0.$$

By the recursion formula (1.12) we therefore get

$$(1.14) \quad p_n(\pm 1, ik) = (ik \mp n)p_{n-1}(\pm 1, ik)$$

and by induction

$$(1.15) \quad p(1, ik) = (-1)^n \prod_{j=1}^n j - ik$$

and

$$(1.16) \quad p(-1, ik) = \prod_{j=1}^n j + ik = (-1)^n \overline{p(1, k)}$$

By (1.10) we now get:

$$(1.17) \quad A_n(k) = \frac{1}{p_n(-1, ik)} = \prod_{j=1}^n \frac{1}{j + ik}.$$

By the recursion formula (1.12) and (1.14) we get $p(1, ik) = \prod_{j=1}^n (j - ik)^{-1}$ and hence (1.10) and (1.19) imply the the statement for $T_+(k)$.

For k negative we notice the following changes have to be made. Instead of (1.10) we now have

$$(1.18) \quad A_n(k)p(1, ik) = 1$$

and instead of (1.19) we have

$$(1.19) \quad T_+(k) = A(k)p(-1, ik).$$

The claim follows now from (1.15) and (1.16). □

The proof shows that $p_n^+(\tanh x, ik) = p_n^-(\tanh x, ik)$. Furthermore, simple induction shows that

$$(1.20) \quad p_n(\tanh x, -k) = (-1)^n p(\tanh(-x), ik).$$

We therefore get the following explicit formula for the eigenfunctions $e(x, k)$:

Theorem 1.4. *Assume that $k \neq 0$. Then*

$$e(x, k) = (\text{sign}(k))^n \left(\prod_{j=1}^n \frac{1}{j + i|k|} \right) P_n(x, ik) e^{ikx}$$

where $P_n(x, k) = p_n(\tanh(x), ik)$ is defined by the recursion formula

$$p_n(\tanh x, ik) = \frac{d}{dx} p_{n-1}(\tanh x, ik) + (ik - n \tanh x) p_{n-1}(\tanh x, ik).$$

In particular the function

$$\mathbb{R} \times (\mathbb{R} \setminus \{0\}) \ni (x, k) \mapsto e(x, k) \in \mathbb{C}$$

is analytic, and $e(x, -k) = e(-x, k)$. Finally, the function

$$(x, y, k) \mapsto e(x, k) \overline{e(y, k)} = \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, ik) P_n(-y, ik) e^{ik(x-y)}$$

is real analytic on \mathbb{R}^3 .

1.5. The point spectrum. In this subsection we discuss briefly the point spectrum of H . We start with:

Lemma 1.6. *If $\lambda = n(n+1)$, then the point spectrum of H is given by*

$$\sigma_p = \{-1, -4, \dots, -n^2\}.$$

Proof. By [GH98, Theorem 6.1, Theorem 7.3] or [Ti58] the point spectrum is given by the points μ^2 , such that $\text{Im}\mu > 0$ and μ and $T(k)$ has a pole at μ . The claim now follows from the explicit description of $T(k)$ in Lemma 1.3. \square

In (1.7) replace k by ji , where $j \in \{1, 2, \dots, n\}$, and denote the solution by $y_{n,j}$. We have two particular solutions: $y_{n,n} = \cosh^n x$ and $y_{n,n+1} = \text{sech}^n x$. Induction shows that

$$y_{n,n+2} = T_{n+1} \text{sech}^n x$$

and

$$y_{n,n+3} = T_{n+2} y_{n,n+2}.$$

For $n = 1$ the particular solution is $y_{1,3} = -3 \tanh x \text{sech} x$. For $n = 2$ the point spectrum is $\sigma_p = \{-1, -4\}$, and the corresponding eigenfunctions (bound states) are $\text{sech} x \tanh x$ and $\text{sech}^2 x$.

Finally we note that ([Lam80, p.46]) the bound-state solution is orthogonal to the propagating solutions, that is,

$$\int \operatorname{sech}(kz)(i\eta - k \tanh(kz))e^{i\eta z} dz = 0$$

1.7. Examples. Assume first that $\lambda = n + 1 = 2$. In this case $n = 1$, the continuous spectrum is $\sigma_c = [0, \infty)$, and the point spectrum is $\sigma_p = -1$.

The boundary condition $u(\pm\infty) = 0$ leads to the single eigenvalue $E = -1$ with the associated eigenfunction $u = \operatorname{sech}(x)$. In quantum theory, the interpretation of this result is that a particle is confined by a potential well having a shape proportional to $\operatorname{sech}^2(x)$ while the single value of E is proportional to the energy that the particle confined by this well can possess. When the sign of the potential is reversed so that $V(x) = 2\operatorname{sech}^2(x)$, the potential is repulsive and no bound state occurs. The solutions for the eigenfunctions in the continuous spectrum are given by

$$e(x, k) = \operatorname{sign}(k) \frac{ik - \tanh(x)}{1 + i|k|} e^{ikx}.$$

If $\lambda = 3$. Thus $n = 2$. Then $\sigma_c = [0, \infty)$, and $\sigma_p = \{-1, -4\}$. The eigenfunctions in the continuous spectrum for $k \neq 0$ are give by

$$e(x, k) = \frac{3 \tanh^2(x) - 3ik \tanh(x) - k^2 - 1}{-k^2 + 3i|k| + 2} e^{-kx}.$$

For point spectrum $\sigma_p = \{-1, -4\}$, the corresponding eigenfunctions (bounded states) are $\operatorname{sech}x \tanh x$ and sech^2x respectively.

1.8. The continuous parameter $\lambda > 1$. For $\alpha > 0$ set $a = \frac{1}{2}(\lambda + i\frac{k}{\alpha})$, $b = \frac{1}{2}(\lambda - i\frac{k}{\alpha})$. Then the fundamental system (1.1) of two real standard solutions u_e and u_o , even and odd in x , are given by [Flu74],

$$u_e(x) = \cosh^\lambda(\alpha x) F(a, b, \frac{1}{2}; -\sinh^2 \alpha x)$$

and

$$u_o(x) = \cosh^\lambda(\alpha x) \sinh(\alpha x) F(a + \frac{1}{2}, b + \frac{1}{2}, \frac{3}{2}; -\sinh^2 \alpha x).$$

Since $-\sinh^2 \alpha x \sim -\frac{e^{2\alpha|x|}}{4}$ and $|x| \rightarrow \infty$, we obtain by the asymptotic of hypergeometric functions:

$$u_e(x) \sim \frac{e^{\lambda\alpha|x|}\Gamma(\frac{1}{2})}{2^\lambda} \left\{ \frac{\Gamma(b-a)}{\Gamma(b)\Gamma(\frac{1}{2}-a)} 2^{2a} e^{-2a\alpha|x|} + \frac{\Gamma(a-b)}{\Gamma(a)\Gamma(\frac{1}{2}-b)} 2^{2b} e^{-2b\alpha|x|} \right\}$$

and

$$u_o(x) \sim \pm \frac{e^{(\lambda+1)\alpha|x|}\Gamma(\frac{3}{2})}{2^{\lambda+1}} \left\{ \frac{\Gamma(b-a)}{\Gamma(b+\frac{1}{2})\Gamma(1-a)} 2^{2a+1} e^{(-2a+1)\alpha|x|} \right. \\ \left. + \frac{\Gamma(a-b)}{\Gamma(a+\frac{1}{2})\Gamma(1-b)} 2^{2b+1} e^{-(2b+1)\alpha|x|} \right\}$$

(\pm sign corresponding to $x > 1$ or $x < 1$), [Le75, p.249].

If the energy $E = k^2$ is non-zero, which corresponds to the scattering solution, we have the following asymptotic expressions for $x > 0$ or $x < 0$:

$$u_e(x) \simeq C_e \cos(k|x| + \phi_e)$$

and

$$u_o(x) \simeq \pm C_o \cos(k|x| + \phi_o)$$

with the abbreviations,

$$(1.21) \quad \phi_e(x) = \arg \frac{\Gamma(ik/\alpha) e^{-i\frac{k}{\alpha} \log 2}}{\Gamma(\frac{\lambda}{2} + i\frac{k}{2\alpha}) \Gamma(\frac{1-\lambda}{2} + i\frac{k}{2\alpha})}$$

$$(1.22) \quad \phi_o(x) = \arg \frac{\Gamma(ik/\alpha) e^{-i\frac{k}{\alpha} \log 2}}{\Gamma(\frac{\lambda-1}{2} + i\frac{k}{2\alpha}) \Gamma(1 - \frac{\lambda}{2} + i\frac{k}{2\alpha})}.$$

Seeking a solution of the asymptotic form

$$u = \begin{cases} T e^{ikx} & x > 0 \\ e^{ikx} + R e^{-ikx} & x < 0 \end{cases}$$

we compute T, R as

$$T = (e^{2i\phi_e} - e^{2i\phi_o})/2, R = (e^{2i\phi_e} + e^{2i\phi_o})/2.$$

Hence,

$$|T|^2 = \sin^2(\phi_e - \phi_o), |R|^2 = \cos^2(\phi_e - \phi_o),$$

which satisfy the conservation law $|T|^2 + |R|^2 = 1$.

Using (1.21),(1.22) we can write

$$\phi_e - \phi_o = \tan^{-1} \left(\frac{\sinh \pi k}{\sin \pi \lambda} \right).$$

For the discrete spectrum let $k = ij$, $j \in \mathbb{N}_0$ yields the eigenvalues $-(\lambda - j - 1)^2$, $j \leq \lambda - 1$, $j \in \mathbb{N}_0$. Hence for the eigenstate equation

$$-\frac{\hbar^2}{2m} \frac{d^2 u}{dx^2} + V = Eu,$$

the corresponding energies are, for $j \leq \lambda - 1$

$$E_n = -\frac{\hbar^2}{2m} (\lambda - 1 - j)^2,$$

where the zero energy is a resonance (as $k \rightarrow 0$) rather than an eigenvalue.

1.9. Projection of the spectral operator $\phi(H)$. Let $\mathcal{H} = L^2$. Then $\mathcal{H} = \mathcal{H}_{ac} \oplus \mathcal{H}_{pp}$, where \mathcal{H}_{ac} denotes the absolute continuous subspace of H and \mathcal{H}_{pp} the pure point subspace. The corresponding orthogonal projections will be denoted by E_{ac} respectively E_{pp} . Note that $E_{ac} = E_{[0,\infty)}$ and $E_{pp} = E_{\sigma_p}$, where E stand for the spectral projection. Hence, if $f \in L^2$, then $f = E_{ac}f + E_{pp}f$. It follows that

$$\phi(H)f = \phi(H)E_{ac}f + \phi(H)E_{pp}f = \phi(H)|_{\mathcal{H}_{ac}}f + \phi(H)|_{\mathcal{H}_{pp}}f.$$

For $\phi \in C_0$ we have

$$(1.23) \quad \phi(H)f(x) = \int K(x, y)f(y)dy + \sum_{k^2 \in \sigma_p} \phi(k^2)(f, e_k)e_k$$

where

$$(1.24) \quad K(x, y) = (2\pi)^{-1} \int \phi(k^2)e(x, k)\bar{e}(y, k)dk$$

is the kernel of $\phi(H)|_{\mathcal{H}_{ac}}$. As $e(\cdot, k)$ is smooth for $k \neq 0$, it follows from (1.24) that $K(x, y)$ is smooth if ϕ is compactly supported. Hence, if σ_p is finite, which is the case when $\int x|V(x)|dx < \infty$, and if $e(\cdot, k) \in L^2(\mathbb{R}) \cap L^p(\mathbb{R})$, as in our case, we have $\|\phi(H)|_{\mathcal{H}_p}f\|_{L^p} \leq C\|f\|_{L^p}$, $f \in L^2 \cap L^p$.

2. DECAY ESTIMATES OF THE SPECTRAL OPERATORS $\phi_j(H)$

In this section we introduce the Triebel-Lizorkin and Besov spaces associated with the Schrödinger operator H and show that they are well-defined, i.e., different dyadic systems give rise to equivalent quasi-norms. In doing so, we show that the quasi-norm can be characterized by using Peetre type maximal function φ_j^*f .

Let $\Phi, \varphi, \Psi, \psi \in C_0^\infty(\mathbb{R})$ satisfy [FJW, E95]

- i) $\text{supp } \Phi, \text{supp } \Psi \subset \{|\xi| \leq 1\}$; $|\Phi(\xi)|, |\Psi(\xi)| \geq c > 0$ if $|\xi| \leq \frac{1}{2}$;
- ii) $\text{supp } \varphi, \text{supp } \psi \subset \{\frac{1}{4} \leq |\xi| \leq 1\}$; $|\varphi(\xi)|, |\psi(\xi)| \geq c > 0$ if $\frac{3}{8} \leq |\xi| \leq \frac{7}{8}$;
- iii) $\Phi(\xi)\Psi(\xi) + \sum_{j=1}^{\infty} \varphi(2^{-j}\xi)\psi(2^{-j}\xi) = 1$, $\forall \xi \in \mathbb{R}$.

Definition 2.1. The Triebel-Lizorkin space $F_p^{\alpha, q} := F_p^{\alpha, q}(H)$ associated with H is the completion of the set $\{f \in L^2 : \|f\|_{F_p^{\alpha, q}} < \infty\}$ with respect to the norm $\|\cdot\|_{F_p^{\alpha, q}}$, where

$$\|f\|_{F_p^{\alpha, q}} = \left\| \left(\sum_{j=0}^{\infty} 2^{j\alpha q} |\varphi_j(H)f|^q \right)^{1/q} \right\|_p.$$

Note, that $1 < p < \infty, q = 2$, $F_p^{\alpha,q}(H)$ corresponds to Sobolev type spaces [Tr92, p.15]. For $j \in \mathbb{Z}$ let $\phi_j(x) = \phi(2^{-j}x)$ and $K_j(x, y) = \phi(2^{-j}H)(x, y)$.

As in [BZ04], the following lemma is essential to establish the Peetre type maximal inequalities Lemma 2.4 and Lemma 2.5. We postpone the proof till Section 3. To simplify the notation we set

$$(2.1) \quad w_j(x) := (1 + 2^{j/2}|x|).$$

Lemma 2.2. *For each $n \in \mathbb{N}_0$ there exist constants $C_n, D_n > 0$ such that*

$$(1) \quad |K_j(x, y)| \leq C_n 2^{j/2} w_j(x - y)^{-n};$$

$$(2) \quad \left| \frac{\partial}{\partial x} K_j(x, y) \right| \leq D_n 2^j w_j(x - y)^{-n}.$$

Note that the statement here is *simpler* than the corresponding statements in Lemmas 3.1-3.3 in [BZ04].

Remark 2.3. *If $H_\alpha = -\Delta + V_\alpha$ on \mathbb{R}^d , where $V_\alpha(x) = \alpha^2 V(\alpha x)$ is the regular scaling, then for $\phi \in L^\infty$ we have:*

$$(2.2) \quad \phi(H_\alpha)(x, y) = \alpha^d \phi(H)(\alpha x, \alpha y).$$

To see this, we note that $\delta_\alpha \circ H \circ \delta_{\alpha^{-1}} = \alpha^{-2} H_\alpha$, hence $\delta_\alpha \phi(H) \delta_{\alpha^{-1}} = \phi(\alpha^{-2} H_\alpha)$. From this we obtain

$$\phi(H)(x, y) = \alpha^{-d} \phi(H_\alpha)(\alpha^{-1}x, \alpha^{-1}y).$$

If $H_h = -\Delta + V_h, V_h = V(\cdot - h)$ is a translation of $V, h \in \mathbb{R}^d$, we have for $\phi \in L^\infty$,

$$(2.3) \quad \phi(H_h)(x, y) = \phi(H)(x - h, y - h),$$

where we note that $H_h = \tau_h \circ H \circ \tau_{-h}$, hence $\phi(H_h) = \tau_h \circ \phi(H) \circ \tau_{-h}$.

The above identities mean that the decay estimates for the kernels, hence the results in this paper, still hold true for the case when our V is replaced by its regular scaling or translation.

For $s > 0$ define the *Peetre maximal function* for H by:

$$(2.4) \quad \phi_j^* f(x) := \phi_{j,s}^* f(x) = \sup_{t \in \mathbb{R}} \frac{|\phi_j(H)f(t)|}{w_j(x - t)^s}$$

and

$$\phi_j^{**} f(x) := \phi_{j,s}^{**} f(x) = \sup_{t \in \mathbb{R}} \frac{|(\phi_j(H)f)'(t)|}{w_j(x - t)^s}.$$

Lemma 2.4. *Let $\phi \in C_0^\infty$ and $s > 0$. Then there exists a constant $c_s > 0$ such that*

$$\phi_j^{**} f(x) \leq C_s 2^{j/2} \phi_j^* f(x).$$

Proof. Using the diadic partition of unity from the beginning of the section and noting that the sum contains at most three non-zero terms we write using the convention that $\psi_{-1} = 0$:

$$\begin{aligned} \frac{d}{dt}(\phi_j(H)f)(t) &= \frac{d}{dt} \sum_{\nu=-1}^1 (\varphi\psi)_{j+\nu}(H)\phi_j(H)f(t) \\ &= \sum_{\nu=-1}^1 \int_{\mathbb{R}} \frac{\partial}{\partial t} K_{j+\nu}(t, y)\phi_j(H)f(y) dy, \end{aligned}$$

where $K_{j+\nu}(t, y)$ denotes the kernel of $(\varphi\psi)_{j+\nu}(H)$.

Apply Lemma 2.2 to obtain

$$\left| \frac{d}{dt}(\phi_j(H)f)(t) \right| \leq C_n \sum_{\nu=-1}^1 2^{j+\nu} \frac{|\phi_j(H)f(y)|}{w_{j+\nu}(t-y)^n w_j(x-t)^s} dy.$$

We now claim that

$$(2.5) \quad \frac{|\phi_j(H)f(y)|}{w_j(x-t)^s} \leq \max_k \phi_j^* f(x) w_j(t-y)^s.$$

To confirm the claim we note that the left hand side of (2.5) is bounded by

$$\begin{aligned} \frac{|\phi_j(H)f(y)|}{w_j(x-y)^s} \cdot \frac{w_j(x-y)^s}{w_j(x-t)^s} &\leq \phi_j^* f(x) \frac{w_j(x-t)^s w_j(y-t)^s}{w_j(x-t)^s} \\ &= \phi_j^* f(x) w_j(t-y)^s, \end{aligned}$$

It follows that

$$\begin{aligned} \left| \frac{d}{dt}(\phi_j(H)f)(t) \right| &\leq C_n \sum_{\nu=-1}^1 2^{j+\nu} \phi_j^* f(x) \int_{\mathbb{R}} \frac{w_j(t-y)^s}{w_{j+\nu}(t-y)^s} dy \\ &\leq C_n 2^{(j+n)/2} \phi_j^* f(x) \int_{\mathbb{R}} \frac{1}{w_j(t-y)^{n-s}} dy \\ &\leq C_{n,s} 2^{j/2} \phi_j^* f(x) \end{aligned}$$

provided $n - s > 1 \leftrightarrow n = [s + 1] + 1$. This proves Lemma 2.4. \square

We are ready to show Peetre maximal inequality.

Lemma 2.5. *Denote by M the Hardy-Littlewood maximal operator. If $0 < r < \infty$ there exists a constant $C > 0$ such that*

$$\phi_j^* f(x) C_{1/r} [M(|\phi_j(H)f|^r)]^{1/r}(x).$$

Proof. The same as those found in [BZ04, E95], which is a modification of the proof in the Fourier case [Tr83, p.16]. \square

Remark 2.6. *It is known that the Hardy-Littlewood maximal operator M is bounded on L^p , $1 < p < \infty$. Lemma 2.5 implies that*

$$(2.6) \quad \|\phi_j^* f\|_p \leq c \|\phi_j(H)f\|_p, \quad 0 < p \leq \infty$$

Moreover, if $1 < p, q < \infty$ and $\{f_j\}$ is a sequence of functions, then

$$(2.7) \quad \left\| \left(\sum_j |Mf_j|^q \right)^{1/q} \right\|_{L^p} \leq C_{p,q} \left\| \left(\sum_j |f_j|^q \right)^{1/q} \right\|_{L^p},$$

by the Fefferman-Stein vector-valued H-L maximal function inequality.

We are ready to state the main theorem on characterization of $F_p^{\alpha,q}(H)$ using the Peetre maximal function.

Theorem 2.7. *Let $\alpha \in \mathbb{R}$, $0 < p, q \leq \infty$. If $\varphi_j^* f$ is defined for $j \geq 0$ with $s > 1/\min(p, q)$, we have for $f \in L^2(\mathbb{R})$*

$$(2.8) \quad \|f\|_{F_p^{\alpha,q}} \sim \|\Phi^* f\|_p + \left\| \left(\sum_{j=1}^{\infty} (2^{j\alpha} \varphi_j^* f)^q \right)^{1/q} \right\|_p.$$

Furthermore, $F_p^{\alpha,q}$ is a quasi-Banach space (Banach space if $p \geq 1, q \geq 1$) and it is independent of the choice of $\{\Phi, \varphi_j\}_{j \geq 1}$.

Proof. Noting that $\phi_j^* f(x) \geq |\phi_j(H)f(x)|$, it is sufficient to show that

$$\|\Phi^* f\|_p + \left\| \left(\sum_{j=1}^{\infty} (2^{j\alpha} \varphi_j^* f)^q \right)^{1/q} \right\|_p \leq C \|f\|_{F_p^{\alpha,q}}$$

but this follows from (2.7) in the following way:

$$\begin{aligned} \|\{2^{j\alpha} \varphi_j^* f\}\|_{L^p(\ell^q)} &\leq C_{\epsilon,s} \|\{2^{j\alpha} [M(|\varphi_j(H)f|^r)]^{1/r}\}\|_{L^p(\ell^q)} \\ &= C_s \|\left\{ \sum_0^{\infty} [M(2^{j\alpha r} |\varphi_j(H)f|^r)]^{q/r} \right\}^{r/q}\|_{L^{p/r}}^{1/r} \\ &\leq C_{s,p,q} \|\{2^{j\alpha} \varphi_j(H)f\}\|_{L^p(\ell^q)} \\ &= C_{s,p,q} \|f\|_{F_p^{\alpha,q}}. \end{aligned}$$

Next we show that $F_p^{\alpha,q}$ is independent of the generating function ϕ , i.e. if $\tilde{\phi}$ is another function given a dyadic partition of unity then for $f \in L^2$, $\|f\|_{F_p^{\alpha,q}}^{\phi}$ and $\|f\|_{F_p^{\alpha,q}}^{\tilde{\phi}}$ are equivalent quasi-norms on $F_p^{\alpha,q}$.

Write $\phi_j(H) = \sum_{\nu=-1}^1 \phi_j(H) \tilde{\phi}_{j+\nu}(H)$ by the identity $\phi_j(x) = \phi_j(x) \sum_{\nu=-1}^1 \tilde{\phi}_{j+\nu}(x)$, for $x \in \mathbb{R}, j \geq 0$, where as before we use the convention that $\psi_{-1} \equiv 0$.

We have by Lemma 2.2 (1),

$$\begin{aligned} |\phi_j(H)f(x)| &\leq 2^{j/2} \sum_{\nu=-1}^1 \int_{\mathbb{R}} \frac{|\tilde{\phi}_{j+\nu}(H)f(y)|}{w_j(x-y)^n} dy \\ &\leq C \sum_{\nu=-1}^1 2^{j/2} \tilde{\phi}_{j+\nu}^* f(x) \int_{\mathbb{R}} \frac{w_j(x-y)^s}{w_j(x-y)^n} dy \\ &= C_N \sum_{\nu=-1}^1 \tilde{\phi}_{j+\nu}^* f(x), \end{aligned}$$

provided $n - s > 1$. Thus, for $f \in L^2$

$$\|\{2^{j\alpha} \phi_j(H)f\}\|_{L^p(\ell^q)} \leq C_{s,p,q} \|\{2^{j\alpha} \tilde{\phi}_j^* f\}\|_{L^p(\ell^q)} \sim \|f\|_{\dot{F}_p^{\alpha,q}}^\psi,$$

This concludes the proof of Theorem 2.7. \square

As expected from Lemma 2.5 we can define the homogeneous Triebel-Lizorkin spaces $\dot{F}_p^{\alpha,q}$ and obtain a maximal function characterization as well.

Definition 2.8. *The homogeneous Triebel-Lizorkin space $\dot{F}_p^{\alpha,q} := \dot{F}_p^{\alpha,q}(H)$ associated with H is the completion of the set $\{f \in L^2 : \|f\|_{\dot{F}_p^{\alpha,q}} < \infty\}$ with respect to the norm $\|\cdot\|_{\dot{F}_p^{\alpha,q}}$, where*

$$\|f\|_{\dot{F}_p^{\alpha,q}} = \left\| \left(\sum_{j=-\infty}^{\infty} (2^{j\alpha} \phi_j(H)f)^q \right)^{1/q} \right\|_p.$$

Theorem 2.9. *Let $\alpha \in \mathbb{R}, 0 < p, q \leq \infty$. If $\phi_j^* f$ is defined for $j \in \mathbb{Z}$ with $s > 1/\min(p, q)$, then for $f \in L^2$*

$$\|f\|_{\dot{F}_p^{\alpha,q}} \sim \left\| \left(\sum_{j=-\infty}^{\infty} (2^{j\alpha} \phi_j^* f)^q \right)^{1/q} \right\|_p.$$

Furthermore, $\|\cdot\|_{\dot{F}_p^{\alpha,q}}^\varphi$ and $\|\cdot\|_{\dot{F}_p^{\alpha,q}}^\psi$ are equivalent norms on the quasi-Banach space $\dot{B}_p^{\alpha,q}$.

The proof is completely implicit in that of Theorem 2.7 and hence omitted.

Moreover, like in the Fourier case and [Tr83], Peetre maximal inequality allows us to define and characterize H -Besov spaces.

Let $\alpha \in \mathbb{R}, 0 < p < \infty, q \leq \infty$. We define the Besov space $B_p^{\alpha,q} := B_p^{\alpha,q}(H)$ associated with H to be the completion of the set $\{f \in L^2 : \|f\|_{B_p^{\alpha,q}} < \infty\}$ with respect to the norm $\|\cdot\|_{B_p^{\alpha,q}}$, where

$$\|f\|_{B_p^{\alpha,q}} = \|\Phi(H)f\|_p + \left\{ \sum_{j=1}^{\infty} 2^{j\alpha q} \|\varphi_j(H)f\|_{L^p}^q \right\}^{1/q}.$$

Thus $B_p^{\alpha,q}$ is a quasi Banach space (Banach space if $p, q \geq 1$). The proof of the following Theorem is analogous to that of Theorem 2.7 and thus left to the reader.

Theorem 2.10. *Let $\alpha \in \mathbb{R}$, $0 < p < \infty$, $0 < q \leq \infty$. If $\Phi^* f$, $\phi_j^* f$ is defined for $j > 0$ with $s > 1/p$, then for $f \in L^2$*

$$\|f\|_{B_p^{\alpha,q}} \sim \|\Phi^* f\|_p + \left(\sum_{j=1}^{\infty} 2^{j\alpha q} \|\phi_j^* f\|_{L^p}^q \right)^{1/q}.$$

Furthermore, $B_p^{\alpha,q}$ is well defined and independent of the choice of generating functions $\varphi = \varphi_0$.

Choosing ϵ small enough in Lemma 2.5 we obtain by taking $r < p (\leftrightarrow s > 1/p)$

$$\begin{aligned} \|\{2^{j\alpha} \phi_j^* f\}\|_{B_p^{\alpha,q}} &\leq C_s \left(\sum_j 2^{j\alpha q} \| [M(|\phi_j(H)f|^r)]^{1/r} \|_{L^p}^q \right)^{1/q} \\ &= C_s \left(\sum_j 2^{j\alpha q} \| M(|\phi_j(H)f|^r) \|_{L^{p/r}}^{q/r} \right)^{1/q} \\ &\leq C \left(\sum_j 2^{j\alpha q} \| |\phi_j(H)f|^r \|_{L^{p/r}}^{q/r} \right)^{1/q} \\ &= C \left(\sum_j 2^{j\alpha q} \|\phi_j(H)f\|_p^q \right)^{1/q} \\ &= C_s \|f\|_{B_p^{\alpha,q}}, \end{aligned}$$

where we used the Hardy-Littlewood maximal function inequality.

The definition and characterization for the homogeneous Besov spaces $\dot{B}_p^{\alpha,q}$ can be formulated similarly.

3. HIGH ENERGY ESTIMATES

In this section we proof Lemma 2.2. We divide the estimates into high and low energy cases.

Recall from Theorem 1.4 that

$$e(x, k) = (\text{sign}(k))^n \left(\prod_{j=1}^n \frac{1}{j + i|k|} \right) P_n(x, ik) e^{ikx}$$

where $P_n(x, k) = p_n(\tanh(x), ik)$ is defined by the recursion formula

$$p_n(\tanh x, ik) = \frac{d}{dx} p_{n-1}(\tanh x, ik) + (ik - n \tanh x) p_{n-1}(\tanh x, ik).$$

By the same Theorem the function

$$e(x, k) \overline{e(y, k)} = \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, ik) P_n(-y, ik) e^{ik(x-y)}$$

is analytic in $k \in \mathbb{R}$ and $e(x, -k) = e(-x, k)$. Furthermore we will also use the obviously relation $\phi(\lambda^2 H)(x, y) = \overline{\phi(\lambda^2 H)(y, x)}$. For simplicity we write

$$(3.1) \quad Q(x, y, k) = P(x, ik)P(y, -ik) = P(x, ik)\overline{P(y, ik)}.$$

In this section we set $\lambda = 2^{-j/2}$ and $\psi(x) = \phi(x^2)$, where $\text{supp}\phi \subset [\frac{1}{4}, 1]$ is fixed. Then

$$K_j(x, y) = \frac{1}{2\pi} \int \psi(\lambda\xi)e(x, k)\overline{e(y, k)} dk.$$

3.1. High and low energy estimates. Using the above explicit formula for $e(x, k)\overline{e(x, k)}$ we set

$$I^+(x, y) = \int_0^\infty \phi_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) Q(x, y, k) e^{ik(x-y)} dk$$

and

$$I^-(x, y) = \int_{-\infty}^0 \phi_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) Q(x, y, k) e^{ik(x-y)} dk.$$

Then we have

$$(3.2) \quad 2\pi\phi_j(H)|_{H_c}(x, y) = I^+(x, y) + I^-(x, y).$$

It is enough to deal $I^+(x, y)$ since $I^-(x, y) = I^+(-x, -y)$ by a change of variable $k \rightarrow -k$.

We note, that

$$k \mapsto R(x, y, k) = R(k) = \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) Q(x, y, k)$$

is a rational function with numerator and denominator both of degree $2n$, and uniformly bounded in x and y . It follows that for $|k| > \epsilon > 0$, we have $|\frac{d^j}{dk^j} R(x, y, k)| \leq C_j |k|^{-j}$ where $C_j > 0$. Furthermore $|\frac{d^j}{dk^j} \psi(\lambda k)| \leq \|\psi^{(n)}\|_\infty \lambda^j$. Note, that the left hand is only non-zero for $|k| \in [2^{j/2-1}, 2^{j/2}]$. This gives first of all an uniform estimate, independent of x and y :

$$|I^+(x, y)| \leq \|R\|_\infty \int_{2^{j/2-1}}^{2^{j/2}} |\psi(\lambda k)| dk \leq 2^{-1} \|R\|_\infty \|\psi\|_\infty \lambda^{-1}.$$

Let $I_j = [2^{j/2-1}, 2^{j/2}]$ and note that $|I_j| = 2^{j/2-1}$, where $|\cdot|$ stand for the standard Lebesgue measure. Then integration by parts gives:

$$\begin{aligned}
|I^+(x, y)| &= \left| \frac{(-1)^n}{i^n(x-y)^n} \int_0^\infty \frac{d^n}{dk^n} [\psi(\lambda k) R(k)] e^{ik(x-y)} dk \right| \\
&\leq |x-y|^{-n} \int_0^\infty \left| \sum_{s=0}^n \binom{n}{s} \frac{d^s}{dk^s} (\psi(\lambda k)) \frac{d^{n-s}}{dk^{n-s}} R(k) \right| dk \\
&\leq C_n |x-y|^{-n} \sum_{s=0}^n \lambda^s \binom{n}{s} \int_{|k| \in I_j} \left| \frac{d^{n-s}}{dk^{n-s}} R(k) \right| dk \\
&= C'_n \lambda^{n-1} / |x-y|^n,
\end{aligned}$$

where we used that

$$\lambda^s \int_{|k| \in I_j} \left| \frac{d^{n-s}}{dk^{n-s}} R(k) \right| dk \leq C \lambda^{n-1}$$

for all $s = 0, 1, \dots, n$. It follows, that there exists a constant C_n such that

$$(3.3) \quad |I^+(x, y)| \leq C_n 2^{j/2} / (1 + 2^{j/2} |x-y|)^n = C_n 2^{j/2} w_j (x-y)^{-n}.$$

The same estimate holds also for for $I^-(x, y)$. Hence it it follows that

$$(3.4) \quad |\phi_j(H)|_{\mathcal{H}_c}(x, y)| \leq C_n \lambda^{-1} / (1 + \lambda^{-1} |x-y|)^n \quad j > 0.$$

The low energy estimates follows in the same way by setting $j = 0$ using that

$$2\pi \Phi(H)|_{\mathcal{H}_c}(x, y) = \int_{-1}^1 \Phi(k^2) |R(x, y, k)| e^{ik(x-y)} dk.$$

Hence

$$|\Phi(H)|_{\mathcal{H}_c}(x, y)| \leq C_n (1 + |x-y|)^{-n}.$$

3.2. Proof of Lemma 2.2 part 2. In the following the subscript \mathcal{H}_c is suppressed, and we denote by $\phi_j(H)(x, y)$ the kernel of the continuous part of the operator $\phi_j(H)(x, y)$.

According to equation 1.24, we know the kernel $\phi_j(H)(x, y) = K_j(x, y)$ ($j \geq 0$) has a unified expression:

$$\phi_j(H)(x, y) = \frac{1}{2\pi} \int \phi_j(k^2) R(x, y, k) e^{ik(x-y)} dk.$$

As $\phi_j(k^2)$ has compact support, and $x \mapsto \frac{d}{dx}\phi_j(k^2)R(x, y, k)e^{ik(x-y)}$ is smooth and hence bounded on the support of $\phi_j(k^2)$, we can interchange the order of integration and differentiation to get:

$$\begin{aligned} 2\pi \frac{\partial}{\partial x} \phi_j(H)(x, y) &= \frac{\partial}{\partial x} \int \phi_j(k^2)R(x, y, k)e^{ik(x-y)} dk \\ &= \int \phi_j(k^2) \frac{\partial}{\partial x} [R(x, y, k)e^{ik(x-y)}] dk, \\ &= \int \phi_j(k^2) |A(k)|^2 [ikP(x, ik) + \frac{\partial}{\partial x} P(x, ik)] P(y, -ik) e^{ik(x-y)} dk \end{aligned}$$

The function $[\partial/\partial x]P(x, ik)$ is a polynomial of $\tanh x$ and ik having degree n in k . Note also, that

$$\left| \frac{d^i}{dk^i} [k\phi_j(k^2)] \right| = \lambda^i C(k) \|\phi_j^{(i)}\|_\infty = O(\lambda^{i-1})$$

where $C(k)$ is a polynomial in k , $i \geq 0$. By the same arguments as in the beginning of this section, we get

$$\left| \frac{\partial}{\partial x} \phi_j(H)(x, y) \right| \leq C_n \lambda^{n-2} (1 + |x - y|)^{-n}, \quad n \geq 1.$$

Together with the fact that

$$\left| \frac{\partial}{\partial x} \phi_j(H)(x, y) \right| \leq O(\lambda^{-2})$$

this implies that

$$\left| \frac{\partial}{\partial y} \phi_j(H)(x, y) \right| \leq C_n \lambda^{-2} (1 + \lambda^{-1}|x - y|)^{-n}$$

completing the proof of Lemma 2.2 (2). \square

4. INTERPOLATION OF $F_p^{0,2} \sim L^p$, $1 < p < \infty$

Let ϕ_j, ψ_j be as in §3. We may assume that $\|\Phi\|_\infty, \|\phi\|_\infty, \|\Psi\|_\infty, \|\psi\|_\infty$ are all ≤ 1 . Let $\phi_0 = \Phi$. Define

$$Q : L^2 \rightarrow L^2(\ell^2) \quad Qf = \{\phi_j(H)f\}_0^\infty.$$

and

$$R : L^2(\ell^2) \rightarrow L^2 \quad R(\{g_j\}_0^\infty) = \sum_{j=0}^\infty R_j g_j = \sum_{j=0}^\infty \psi_j(H) g_j.$$

Note that $\|f\|_{F_p^{0,2}(H)} = \|Qf\|_{L^p(\ell^2)}$

It is easy to see that $RQ = Id : L^2 \rightarrow L^2$ and $QR \leq 3Id : L^2(\ell^2) \rightarrow L^2(\ell^2)$. We will use Q and R to identify $F_p^{0,2}(H)$ with L^p .

Theorem 4.1. *Let $1 < p < \infty$. Then $F_p^{0,2}$ and L^p are isomorphic and have equivalent norms.*

To prove the theorem, we first show that $Q: L^p \rightarrow L^p(\ell^2)$ and $R: L^p(\ell^2) \rightarrow L^p$, $1 < p < \infty$. First we note that

$$(4.1) \quad \|f\|_{F_p^{0,2}(H)} \lesssim \|f\|_p$$

and

$$(4.2) \quad \|g\|_p \lesssim \|g\|_{F_p^{0,2}(H)}$$

for $f, g \in L^2 \cap L^p \cap F_p^{0,2}(H)$. Here (4.1) follows directly from the definition of Q (and $F_p^{0,2}(H)$) and equation (4.2) follows from the identity $R \circ Q = id_{L^2}$, i.e., $\sum \varphi(H)\psi_j(H) = id$. By continuity extension of Q, R , (4.1) and (4.2) hold for all of $F_p^{0,2}(H)$ and L^p . So we obtain the continuous embedding

$$(4.3) \quad L^p \subset F_p^{0,2}(H)$$

$$(4.4) \quad F_p^{0,2}(H) \subset L^p,$$

which proves Theorem 4.1.

The remaining part of this section will be devoted to showing the boundedness of Q and R . In the following, Lemma 4.2 and Lemma 4.4 imply that Q is bounded from L^p to $L^p(\ell^2)$, and, Lemma 4.2 and Lemma 4.6 imply that R is bounded from $L^p(\ell^2)$ to L^p by interpolation and duality.

Lemma 4.2. $Q : L^2 \rightarrow L^2(\ell^2)$ and $R : L^2(\ell^2) \rightarrow L^2$ are well-defined bounded operators.

Proof. Let $\{g_j\} \in L^2(\ell^2)$. Note that $R_j = \psi_j(H)$ is bounded on L^2 : $\|R_j g\|_2 \leq \|\psi\|_\infty \|g\|_2$. Thus

$$\begin{aligned} \left(\sum_{j=0}^{\infty} R_j g_j, \sum_{j=0}^{\infty} R_j g_j \right) &= \sum_{\nu=-1}^1 \sum_{j=0}^{\infty} (R_j g_j, R_{j+\nu} g_j) \\ &\leq \sum_{\nu=-1}^1 \sum_j \|R_j g_j\|_2 \|R_{j+\nu} g_j\|_2 \\ &\leq 3 \sum_j \|g_j\|_2^2 \\ &= 3 \|g_j\|_{L^2(\ell^2)}^2. \end{aligned}$$

□

We now derive some needed estimates for the kernel of $Q_j = \phi_j(H)$. As before, $Q_j(x, y)$ denotes the kernel of $\phi_j(H)E_{ac} = \phi_j(H_{ac})$.

Define

$$\tilde{Q}_j(x, y) = \begin{cases} \int_{I_k} Q_j(x, y) b_k(y) dy & \text{if } 2^{j/2} |I_k| \geq 1 \\ \int_{I_k} (Q_j(x, y) - Q_j(x, \bar{y})) b_k(y) dy & \text{if } 2^{j/2} |I_k| < 1. \end{cases}$$

Lemma 4.3. Let $y \in I = (\bar{y} - \frac{t}{2}, \bar{y} + \frac{t}{2})$, $t = |I|$ and $I^* = (\bar{y} - t, \bar{y} + t)$. Then the following holds:

(1) If $2^{j/2}|I| \geq 1$, then

$$\sup_{y \in I} \int_{R \setminus I^*} |Q_j(x, y)| dx \leq C(2^{j/2}|I|)^{-1}$$

(2) If $2^{j/2}|I| < 1$, then

$$\sup_{y \in I} \int_{R \setminus I^*} |Q_j(x, y) - Q_j(x, \bar{y})| dx \leq C2^{j/2}|I|.$$

In particular we have

$$(4.5) \quad \int_{R \setminus I^*} \sum_j |\tilde{Q}_j(x, y)| \leq 3C.$$

Proof. For the first inequality, let $2^{j/2}|I| \geq 1$.

$$\begin{aligned} \int_{\mathbb{R} \setminus I^*} |Q_j(x, y)| dx &\leq C_n \int_{|x-y|>t/2} \frac{2^{j/2}}{(1+2^{j/2}|x-y|)^n} dx \\ &\leq C(2^{j/2}|I|)^{-1}, \quad (n=2). \end{aligned}$$

If $2^{j/2}|I| < 1$, $y \in I$, \bar{y} = center of $I = (\bar{y} - \frac{t}{2}, \bar{y} + \frac{t}{2})$.

$$\begin{aligned} \int_{\mathbb{R} \setminus I^*} |Q_j(x, y) - Q_j(x, \bar{y})| dx &= \int_{\mathbb{R} \setminus I^*} \left| \int_{\bar{y}}^y \frac{\partial}{\partial z} \psi_j(H)(x, z) dz \right| dx \\ &\leq C_n |y - \bar{y}| \int_{|x-\bar{y}|>t} \frac{2^j}{(1+2^{j/2-1}|x-\bar{y}|)^n} dx \\ &\leq C_n 2^{j/2} t \int_{|u|>(2^{-j/2+1})^{-1}t} \frac{du}{(1+|u|)^n} \quad (n=2) \\ &\leq C2^{j/2}|I|. \end{aligned}$$

Now (4.5) follows easily from the above two inequalities. \square

Lemma 4.4. Q is bounded from L^1 to weak $L^1(\ell^2)$, i.e.,

$$|\{x : (\sum_0^\infty |Q_j f(x)|^2)^{1/2} > \lambda\}| \leq C\lambda^{-1} \|f\|_1, \quad \lambda > 0.$$

Proof. Let $f \in L^1 \cap L^2$. By the Calderón-Zygmund decomposition, there exists a sequence of intervals $\{I_k\}$ and functions $\{b_k\}$ with $\text{supp } b_k \subset I_k$ such that with $g = \sum_k b_k$ we have $f = g + b$, for some $g \in L^2$. Furthermore:

- i) $|g(x)| \leq C\lambda$ a.e.
- ii) $b_k(x) = f(x) - |I_k|^{-1} \int_{I_k} f dx$
- iii) $\lambda \leq |I_k|^{-1} \int_{I_k} |f| dx \leq 2\lambda$.
- iv) $|D_\lambda| = |\cup_k I_k| = \sum_k |I_k| \leq \lambda^{-1} \|f\|_1$.

Let $\{e_n\}_m$ be an orthonormal basis in \mathcal{H}_{pp} and λ_m the corresponding eigenvalues in σ_p . Note that this set is finite.

$$\begin{aligned}
\int \sum_j |Q_j g(x)|^2 dx &= \sum_j \|\phi_j(H)g\|_2^2 \\
&= \sum_j (\|\mathcal{F}^* \phi_j(k^2) \mathcal{F}g\|_2^2 + \sum_{\sigma_p} \|\phi_j(\lambda_n)(g, e_m)e_m\|_2^2) \\
&= \sum_j (\|\phi_j(k^2) \mathcal{F}g(k)\|_2^2 + \sum_{\sigma_p} |\phi_j(\lambda_n)|^2 |(g, e_m)|^2) \\
&\leq \int (\sum_j |\phi_j(k^2)|^2) |\mathcal{F}g(k)|^2 dk + 3\|\phi\|_\infty^2 \sum_{\sigma_p} |(g, e_m)|^2 \\
&\leq 3\|\phi\|_\infty^2 \int |\mathcal{F}g(k)|^2 dk + 3\|\phi\|_\infty^2 \|E_{pp}g\|^2 \\
&\leq 3\|\phi\|_\infty^2 \|E_{ac}g\|^2 + 3\|\phi\|_\infty^2 \|E_{pp}g\|^2 \\
&= 3\|\phi\|_\infty^2 \|g\|^2,
\end{aligned}$$

where we note that $\mathcal{F}: L^2 \rightarrow L^2$ surjection and $\mathcal{F}: \mathcal{H}_{ac} \rightarrow L^2$ isometry. By Chebychev inequality we have

$$|\{x : (\sum_0^\infty |Q_j g(x)|^2)^{1/2} > \lambda/2\}| \leq C\lambda^{-2} \|g\|_2^2 \leq C\lambda^{-1} \|f\|_1.$$

We now only need to show

$$|\{x \notin \cup I_k^* : (\sum_j |Q_j b(x)|^2)^{1/2} > \lambda/2\}| \leq C\lambda^{-1} \|f\|_1.$$

The left hand side is bounded by

$$\frac{2}{\lambda} \sum_k \int_{\mathbb{R} \setminus \cup I_k^*} (\sum_j |Q_j b_k(x)|^2)^{1/2} dx.$$

For each k , since $\int b_k = 0$,

$$\begin{aligned}
(\sum_j |Q_j b_k(x)|^2)^{1/2} &\leq \sum_j |Q_j(E_{ac} + E_{pp})b_k(x)| \\
&\leq \sum_j \left| \int \tilde{Q}_j(x, y) b_k(y) dy \right| + \sum_j \left| \sum_{\sigma_{pp}} \phi_j(\lambda_m)(b_k, e_m)e_m(x) \right| \\
&\leq \int \sum_j |\tilde{Q}_j(x, y) b_k(y)| dy + \sum_{\sigma_{pp}} (\sum_j |\phi_j(\lambda_m)|) \|b_k\|_1 |e_m(x)| \\
&\leq \int \sum_j |\tilde{Q}_j(x, y) b_k(y)| dy + 3\|\phi\|_\infty \|b_k\|_1 \sum_{\sigma_{pp}} |e_m(x)|
\end{aligned}$$

where we note that $e_m \in L^1 \cap L^\infty$ and σ_{pp} is finite, having n eigenvalues.

Equation (4.5) gives now the following estimates for the left hand side

$$\begin{aligned}
l.h.s. &\leq \frac{2}{\lambda} \sum_k \int_{\mathbb{R} \setminus \cup I_k^*} \sum_j \int |\tilde{Q}_j(x, y)| |b_k(y)| dy dx + \\
&\quad 3 \|\phi\|_\infty \frac{2}{\lambda} \sum_k \int_{\mathbb{R} \setminus \cup I_k^*} \|b_k\|_1 \left(\sum_{\sigma_{pp}} |e_m(x)| \right) dx \\
&\leq \frac{2}{\lambda} \sum_k \int_{y \in I_k} |b_k(y)| dy \int_{\mathbb{R} \setminus I_k^*} \sum_j |\tilde{Q}_j(x, y)| dx + \frac{C}{\lambda} \sum_k \|b_k\|_1 \sum_{\sigma_{pp}} \|e_m\|_1 \\
&\leq \frac{C'}{\lambda} \sum_k \int_{y \in I_k} |b_k(y)| dy \leq C \lambda^{-1} \|f\|_1.
\end{aligned}$$

□

By Lemma 4.3, we have the following estimates which will be used in the proof of Lemma 4.6:

If $2^{j/2}|I_k| \geq 1$, then

$$\sum_{j: 2^{j/2}|I_k| \geq 1} \int_{\mathbb{R} - I^*} |R_j(x, y)| \leq C |I_k|^{-1} \sum_{2^{j/2}|I_k| \geq 1} 2^{-j/2} \leq C.$$

If $2^{j/2}|I_k| < 1$, then

$$\sum_{j: 2^{j/2}|I_k| < 1} \int_{\mathbb{R} - I^*} |R_j(x, y) - R_j(x, \bar{y})| \leq C |I_k| \sum_{2^{j/2}|I_k| < 1} 2^{j/2} \leq 2C.$$

Remark 4.5. $R_j f(x) = \int \psi_j(H)(x, y) f(y) dy + \sum_{\sigma_{pp}} \psi_j(\lambda_m)(f, e_m) e_m(x)$ is well defined for $f \in L^1 \cap L^2$ and for $f \in L^1$ by all appropriate means, since $\psi_j(H)(x, \cdot) \in L^1 \cap L^\infty$.

Lemma 4.6. Let $R_j = \psi_j(H)$. Then $R = \{R_j\}$ are bounded from $L^1(\ell^2)$ to weak- L^1 .

Proof. We will show that there exists a positive constant C such that

$$(4.6) \quad \left\{ x : \left| \sum_0^N R_j f_j(x) \right| > \lambda \right\} \leq C \lambda^{-1} \|\{f_j\}\|_{L^1(\ell^2)},$$

for all $N \in \mathbb{N}$, $\{f_j\} \in L^1(\ell^2)$ and $\lambda > 0$. Then, the inequality holds for $N = \infty$ and $\{f_j\} \in L^1(\ell^2) \cap L^2(\ell^2)$ by passing to the limit, noting that $\sum_0^\infty R_j f_j(x) \in L^2$ if $\{f_j\} \in L^2(\ell^2)$. Finally, the lemma follows by the fact that $L^1(\ell^2) \cap L^2(\ell^2)$ is dense in $L^1(\ell^2)$.

Alternatively, suppose $\{f_j\} \in L^1(\ell^2)$ only. Observe that (4.6) suggests that the partial sum $R_N f = \sum_0^N R_j f_j$ is a Cauchy sequence in measure, therefore there exists a unique function, which we define to be Rf , so that $R_N f$ converges to Rf in measure and $Rf = \sum_{j=0}^\infty R_j f_j$ satisfies (4.6).

So, it is sufficient to show (4.6). Calderón-Zygmund decomposition: Let $F(x) = (\sum_{j=0}^{\infty} |f_j(x)|^2)^{1/2} \in L^1$. There exists a collection of countable disjoint open intervals $\{I_k\}$ such that

- a) $|F(x)| \leq \lambda$, a.e. $x \in \mathbb{R} \setminus \cup_k I_k$,
- b) $\sum_k |I_k| \leq \lambda^{-1} \|F\|_1$,
- c) $\lambda \leq |I_k|^{-1} \int_{I_k} |F(x)| dx \leq 2\lambda$, $\forall k$.

Define

$$g_j(x) = \begin{cases} |I_k|^{-1} \int_{I_k} f_j dy, & x \in I_k \\ f_j(x) & \text{otherwise} \end{cases}$$

$$b_j(x) = \begin{cases} f_j - g_j, & x \in I_k \\ 0 & \text{otherwise} \end{cases}$$

Then, if $x \in \mathbb{R} \setminus \cup_k I_k$, $(\sum_{j=0}^{\infty} |g_j(x)|^2)^{1/2} = (\sum_{j=0}^{\infty} |f_j(x)|^2)^{1/2}$. And, if $x \in I_k$

$$\begin{aligned} (\sum_{j=0}^{\infty} |g_j(x)|^2)^{1/2} &\leq (\sum_{j=0}^{\infty} |I_k|^{-2} |\int_{I_k} f_j(y) dy|^2)^{1/2} \\ &\leq |I_k|^{-1} \int_{I_k} (\sum_{j=0}^{\infty} |f_j(y)|^2)^{1/2} dy \\ &\leq 2\lambda \end{aligned}$$

by Minkowski inequality.

We also have

$$\begin{aligned} \|\{g_j(x)\}\|_{L^2(\ell^2)}^2 &= \sum_{k=0}^{\infty} \int_{I_k} (\sum_j |g_j(x)|^2) dx + \int_{\mathbb{R} \setminus \cup_k I_k} (\sum_j |g_j(x)|^2) dx \\ &\leq (2\lambda)^2 \sum_k |I_k| + \lambda \int_{\mathbb{R} \setminus \cup_k I_k} (\sum_j |f_j|^2)^{1/2} dx \\ &\leq C\lambda \|F\|_1. \end{aligned}$$

Finally, by Chebychev inequality and Lemma 4.2, we get:

$$\begin{aligned} |\{x : |\sum_0^N R_j g_j(x)| > \lambda/2\}| &\leq C\lambda^{-2} \|\sum_0^N R_j g_j\|^2 \\ &\leq C'\lambda^{-2} \|\{g_j\}\|_{L^1(\ell^2)}^2 \leq C\lambda^{-1} \|F\|_1. \end{aligned}$$

It remains to show

$$|\{x \notin \cup_k I_k^* : |\sum_0^N R_j b_j(x)| > \lambda/2\}| \leq C\lambda^{-1} \|F\|_1.$$

The left hand side is not exceeding $\frac{2}{\lambda} \sum_k \int_{\mathbb{R} \setminus \cup I_k^*} |\sum_{j=0}^N R_j b_{j,k}(x)| dx$, where $b_{j,k} = b_j \chi_{I_k}$. For each k , define

$$\tilde{R}_j^k(x, y) = \begin{cases} R_j(x, y) & \text{if } 2^{j/2}|I_k| \geq 1 \\ R_j(x, y) - R_j(x, \bar{y}_k) & \text{if } 2^{j/2}|I_k| < 1 \end{cases}$$

Note that the domain of R_j has a natural extension from $L^1 \cap L^2$ to L^1 . Then, since $\int b_{j,k} = 0$ and by Jensen inequality $(\sum_{j=0}^N |\tilde{R}_j^k(x, y)|^2)^{\frac{1}{2}} \leq \sum_{j=0}^N |\tilde{R}_j^k(x, y)|$ we obtain

$$\begin{aligned} \int_{\mathbb{R} \setminus I_k^*} |\sum_{j=0}^N R_j b_{j,k}(x)| dx &= \int_{\mathbb{R} \setminus I_k^*} |\sum_{j=0}^N \int_{I_k} \tilde{R}_j^k(x, y) b_{j,k}(y) dy| dx \\ &\leq \int_{I_k} (\sum_{j=0}^N |b_{j,k}|^2(y))^{\frac{1}{2}} dy \int_{\mathbb{R} \setminus I_k^*} (\sum_{j=0}^N |\tilde{R}_j^k(x, y)|^2)^{\frac{1}{2}} dx \\ &\leq C \int_{I_k} (\sum_{j=0}^N |b_{j,k}(y)|^2)^{\frac{1}{2}} dy \\ &\leq C \int_{I_k} (\sum_{j=0}^N |f_j|^2)^{\frac{1}{2}} dy + C \int_{I_k} (\sum_{j=0}^N |g_j|^2)^{\frac{1}{2}} dy \\ &\leq 2C \int_{I_k} (\sum_{j=0}^{\infty} |f_j|^2)^{\frac{1}{2}} dy, \end{aligned}$$

where we used the pointwise estimate by Minkowski inequality

$$(\sum_0^N |g_j(y)|^2)^{\frac{1}{2}} \leq (\sum_0^{\infty} (|I_k|^{-1} \int_{I_k} |f_j(y)|^2)^{\frac{1}{2}}) \leq |I_k|^{-1} \int_{I_k} (\sum_0^N |f_j(y)|^2)^{\frac{1}{2}} dy.$$

Hence

$$|\{x \in \mathbb{R} \setminus \cup I_k^* : |\sum_0^N R_j b_j(x)| > \lambda/2\}| \leq \frac{4C}{\lambda} \sum_k \int_{I_k} (\sum_j |f_j|^2)^{\frac{1}{2}} dy \leq \frac{4C}{\lambda} \|(\sum_j |f_j|^2)^{\frac{1}{2}}\|_1.$$

as desired.

By now, we have shown that

$$\begin{aligned} |\{x : |\sum_0^N R_j f_j(x)| > \lambda\}| &\leq |\{x : |\sum_0^N R_j g_j(x)| > \frac{\lambda}{2}\}| + |\{x : |\sum_0^N R_j b_j(x)| > \frac{\lambda}{2}\}| \\ &\leq C\lambda^{-1} \|F\|_1, \end{aligned}$$

where C is independent of N . Passing to limit, since convergence a.e. implies convergence in measure, we obtain

$$|\{x : |\sum_0^\infty |R_j f_j(x)| > \lambda\}| \leq C\lambda^{-1} \|F\|_1.$$

□

5. REMARK ON TIME DECAY OF THE WAVE FUNCTION

We conclude the paper with a time decay result on the wave function $\psi(t, x) = e^{-itH} f$ based on the H -Besov space method. Generalized Besov space method has been considered in e.g. [J92], [JN94], [JP85], [BZ04] in the study of PDEs related to perturbation for Schrödinger operators.

By [BZ04, Theorem 7.1] or [JN94, Theorem 5.1] we know that if V is in the Kato class \mathcal{K} and if $\mathcal{D}(H^m) = W_p^{2m}$ for some $m \in \mathbb{N}$, $1 \leq p < \infty$, then for $1 \leq q \leq \infty$, $0 < \alpha < m$, $B_p^{\alpha, q}(H) = B_p^{2\alpha, q}(\mathbb{R}^d)$. It is easy to see that if V is C^∞ with all derivatives bounded, then the domain condition on H is verified for all $m \in \mathbb{N}$. Since $V \sim \text{sech}^2 x$ has this property, $B_p^{\alpha, q}(H) = B_p^{2\alpha, q}(\mathbb{R}^d)$ for all $\alpha > 0$. Combining this with [JN94, Theorem 5.2], we obtain the time decay of $\psi(t, x)$ on ordinary Besov spaces. Let $\langle t \rangle = (1 + t^2)^{1/2}$.

Theorem 5.1. *Suppose $1 \leq p < \infty$, $1 \leq q \leq \infty$ and $\alpha > 0$. Let $\beta = |\frac{1}{2} - \frac{1}{p}|$. Then e^{-itH} maps $B_p^{\alpha+2\beta, q}(\mathbb{R})$ continuously to $B_p^{\alpha, q}(\mathbb{R})$. Moreover, e^{-itH} maps $B_p^{2\beta, q}(\mathbb{R})$ continuously to L^p . In both cases the operator norm is less than or equal to $C\langle t \rangle^\beta$. That is,*

$$\|\psi(t, x)\|_{B_p^{\alpha, q}(\mathbb{R})} \lesssim \langle t \rangle^\beta \|f\|_{B_p^{\alpha+2\beta, q}(\mathbb{R})}$$

and

$$\|\psi(t, x)\|_{L^p} \lesssim \langle t \rangle^\beta \|f\|_{B_p^{2\beta, q}(\mathbb{R})}.$$

Compare Proposition 7.3 in [BZ04].

Finally we give an interpolation result. We know that $B_p^{\alpha, q}(H) = B_p^{2\alpha, q}(\mathbb{R})$, $\alpha > 0$. In particular, $F_p^{\alpha, p}(H) = F_p^{2\alpha, p}(\mathbb{R})$. On the other hand, by Theorem 4.1, $F_p^{0, p}(H) = L^p = F_p^{0, p}(\mathbb{R})$. Thus we obtain Theorem 5.2 using complex interpolation method.

Theorem 5.2. *If $\alpha > 0$, $1 < p < \infty$ and $2p/(p+1) < q < 2p$, then*

$$F_p^{\alpha, q}(H) = F_p^{2\alpha, q}(\mathbb{R})$$

and if $\alpha > 0$, $1 \leq p < \infty$, $1 \leq q \leq \infty$, then

$$B_p^{\alpha, q}(H) = B_p^{2\alpha, q}(\mathbb{R}).$$

In general e^{-itH} is not bounded on L^p unless $p = 2$. [JN94, RS04]. However one can obtain the boundedness on F and B spaces.

Let S_p be a set of numbers q such that

$$\begin{aligned} p \leq q < 2 & \text{ if } p < 2; \\ 2 & \text{ if } p = 2; \\ 2 < q \leq p & \text{ if } p > 2. \end{aligned}$$

Based on the mapping property: if $\alpha > 0$, $1 \leq p < \infty$, $1 \leq q \leq \infty$, $\beta = |\frac{1}{p} - \frac{1}{2}|$, $e^{-itH}: F_p^{\alpha+2\beta,p} \rightarrow F_p^{\alpha,p}$ and $F_p^{2\beta,p} \rightarrow L^p$, and the interpolation $(F_p^{\alpha+2\beta,p}, F_p^{2\beta,p})_\theta \rightarrow (F_p^{\alpha,p}, F_p^{0,2})_\theta$, we obtain

Proposition 5.3. *If $\alpha_1 > 0$, $1 < p < \infty$, $q_1 \in S_p$, $\beta := \beta(p) = |\frac{1}{p} - \frac{1}{2}|$, then*

$$(5.1) \quad e^{-itH} : F_p^{\alpha_1+2\beta,p}(H) \rightarrow F_p^{\alpha_1,q_1}(H).$$

If $\alpha > 0$, $1 \leq p < \infty$, $1 \leq q_1 \leq \infty$,

$$B_p^{\alpha+2\beta,q}(H) \rightarrow B_p^{\alpha,q}(H)$$

with operator bound $C\langle t \rangle^\beta$.

If we try to use interpolation coupled with $L^2 \rightarrow L^2$, we proceed

$$(F_p^{\alpha_1+2\beta,p}, F_2^{0,2})_\theta \rightarrow (F_p^{\alpha_1,q_1}, F_2^{0,2})_\theta.$$

We find that for each $t \in (0, 1)$

$$F_{p_t}^{\alpha_2+2\beta_t,p_t}(H) \rightarrow F_{p_t}^{\alpha_2,q_2}(H) \quad \forall \alpha_2 > 0, 1 < p_t < \infty$$

where $1/p_t = (1-t)/p + t/2$, $q_2 \in S_{p_t}$. It means the interpolation with $F_2^{0,2} \rightarrow F_2^{0,2}$ does not help improve (5.1) at all. This can also be observed from the two-triangle shaped region bounded by $\{(\frac{1}{p}, \frac{1}{q_1}) : q_1 \in S_p\}$.

The result for B spaces follows from Theorem 4.6 and Remark 4.7 in [JN94].

REFERENCES

- [AGHH] S. Albeverio, F. Gesztesy, R. Hoegh-Krohn, H. Holden, *Solvable Models in Quantum Mechanics*, Springer-Verlag, 1988.
- [BZ04] J.J. Benedetto, S. Zheng, Besov spaces for the Schrödinger operator with barrier potential (submitted). <http://lanl.arXiv.org/math.CA/0411348>, (2004).
- [BL] J. Bergh, J. Löfström, *Interpolation Spaces*, Springer-Verlag, 1976.
- [B99] H.R. Beyer, On the completeness of the quasinormal modes of the Pöschl-Teller potential, *Comm. Math. Phys.* **204** (1999), no. 2, 397-423.
- [Ca64] A.P. Calderón, Intermediate spaces and interpolation, the complex method, *Stud. Math.*, **24** (1964), 113-190.
- [CaT] A.P. Calderón, A. Torchinsky, Parabolic maximal functions associated with a distribution, I,II, *Advances in Math.* **16** (1975), 1-64, and **24** (1977), 101-171.
- [CK01] M. Christ, A. Kiselev, One-Dimensional Schrödinger operators with slowly decaying potentials: spectra and asymptotics, Notes for IPAM tutorial, 2001 Workshop on Oscillatory Integrals and Dispersive Equations.
- [CK02] ———, Scattering and wave operators for one-dimensional Schrödinger operators with slowly decaying nonsmooth potentials, *Geom. Funct. Anal.* **12** (2002), no. 6, 1174-1234.

- [CMS86] M. Cwikel, M. Milman, Y. Sagher, Complex interpolation of some quasi-Banach spaces, *J. Funct. Anal.* **65** (1986), 339-347.
- [CS88] M. Cwikel, Y. Sagher, Analytic families of operators on some quasi-Banach spaces, *Proc. Amer. Math. Soc.* **102** (1988), 979-984.
- [DF05] Piero D'Ancona, Luca Fanelli, Decay estimates for the wave and Dirac equations with a magnetic potential, <http://arXiv.org/abs/math.AP/0504370>.
- [DP05] Piero D'Ancona, Vittoria Pierfelice, On the wave equation with a large rough potential, <http://arXiv.org/math.AP/0310199>.
- [DPV05] Piero D'Ancona, Vittoria Pierfelice, Nicola Visciglia, Some remarks on the Schrödinger equation with a potential in $L_t^r L_x^s$, to appear in *Math. Annalen*, <http://arXiv.org/math.AP/0501125>.
- [DOS] X.T. Duong, E.M. Ouhabaz and A. Sikora, Plancherel type estimates and sharp spectral multipliers, *J. Funct. Anal.* (2002).
- [DZ98] J. Dziubański, Atomic decomposition of H^p spaces associated with some Schrödinger operators, *Indiana Univ. Math. J.* **47**(1998), 75-98.
- [DZ02] J. Dziubański and J. Zienkiewicz, H^p spaces for Schrödinger operators, in: *Fourier Analysis and Related Topics*, Banach Center Publ. **56**, Inst. Math., Polish Acad. Sci., 2002, 45-53.
- [E95] J. Epperson, Triebel-Lizorkin Spaces for Hermite expansions, *Studia Math.* **114**(1), 1995.
- [E96] ———, Hermite multipliers and pseudo-multipliers, *Proc. Amer. Math. Soc.* **124** (1996), no. 7, 2061–2068.
- [FGr] Y.F. Fang, M.G. Grillakis, On the Dirac-Klein-Gordon system in 2+1 dimensions, *preprint*.
- [FM84] V. Ferrari and B. Mashoon, New approach to the quasinormal modes of a black hole. *Phys. Rev. D.* **30** (1984), 295-304.
- [Flu74] S. Flügge, *Practical Quantum Mechanics*, Springer-Verlag, 1974.
- [FJW] M. Frazier, B. Jawerth, G. Weiss, *Littlewood-Paley Theory and the Study of Function Spaces*, Conference Board of the Math. Sci. **79**, 1991.
- [FMV05a] Giulia Furioli, Camillo Melzi, Alessandro Veneruso, Littlewood-Paley decompositions and Besov spaces on Lie groups of polynomial growth, <http://arXiv.org/math.CA/0502406>.
- [FMV05b] ———, Strichartz inequalities for the wave equation with the full Laplacian on the Heisenberg group, <http://arXiv.org/math.AP/0502497>.
- [GH98] C.-A. Guerin, M. Holschneider, Time-dependent scattering on fractal measures, *J. Math. Physics* **39**(8), 1998.
- [He90a] W. Hebisch: A multiplier theorem for Schrödinger operators. *Colloq. Math.* **60/61** (1990), no. 2, 659-664.
- [J92] A. Jensen, High energy resolvent estimates for Schrödinger operators in Besov spaces. Festschrift on the occasion of the 70th birthday of Shmuel Agmon, *J. Anal. Math.* **59** (1992), 45-50.
- [JN94] A. Jensen, S. Nakamura, Mapping properties of functions of Schrödinger operators between L^p spaces and Besov spaces, in *Spectral and Scattering Theory and Applications*, Advanced Studies in Pure Math. **23**, 1994.
- [JP85] A. Jensen, P. Perry, Commutator methods and Besov space estimates for Schrödinger operators, *J. Operator Theory*, **14** (1985), 181-188.
- [JSS] J.-L. Journé, A. Soffer, C.D. Sogge, Decay estimates for Schrödinger operators, *Comm. Pure Appl. Math.*, vol. XLIV (1991), 573-604.
- [Kee] J.P. Keener, *Principles of applied mathematics*, Transformation and approximation, Addison-Wesley Publishing Company, 1988.
- [KMS] S. Klainerman, M. Machedon and J. Stalker, Decay of solutions to the wave equation on a spherically symmetric static background, *preprint*.
- [KO02] B. Krötz, G. Olafsson, The c -function for non-compactly causal symmetric spaces, *Invent. Math.* **149** (2002), no. 3, 647-659.

- [Lam80] G.L. Lamb, Jr., *Elements of Soliton Theory*, Pure & Applied Mathematics, Wiley-Interscience, 1980.
- [Le75] N.N. Lebedev, *Special functions and their applications*, Translation by R.A. Silverman, New York, Dover Publications, 1972.
- [Ob90] F. Oberhettinger, *Tables of Fourier Transforms and Fourier transforms of Distributions*, Springer-Verlag, 1990.
- [Ob73] ———, *Fourier Expansions :A Collection of Formulas*, Academic Press, New York-London, 1973.
- [O87] G. Ólafsson, Fourier and Poisson transformation associated to a semisimple symmetric space, *Invent. Math.* **90** (1987), no. 3, 605–629.
- [Pa80] L. Päiväranta, On the spaces $L_p^A(\ell_q)$: Maximal inequalities and complex interpolation, *Ann. Acad. Sci. Fenn. Ser. AI Math. Dissertationes* **25**, Helsinki 1980.
- [PT33] G. Pöschl, E. Teller, Bemerkungen zur Quantenmechanik des harmonischen Oszillators, *Z. Phys.* **83** (1933), 143-151.
- [RS] M. Reed, B. Simon, *Methods of Modern Mathematical Physics III: Scattering Theory, IV: Analysis of Operators*, Academic Press, New York, 1978.
- [RVVW] R.Rochberg, A. Tabacco Vignati, M.Vignati, G.Weiss, Interpolation of quasinormed spaces by the complex method. In: “Function Spaces and Applications”, Proc. Conf. Lund 1986, Lect. Notes Math. 1302. Berlin, Springer 1988, 91-98.
- [RoTao] I. Rodnianski, T. Tao, Long-time decay estimates for the Schrödinger equation on Manifolds, *preprint*. <http://lanl.arXiv.org/math.AP/0412416>, (2004).
- [Sch05] W. Schlag, Dispersive estimates for Schrödinger operators: A survey. <http://lanl.arXiv.org/math.AP/0501037>, (2005).
- [Sh95] Z. Shen, L^p estimates for Schrödinger operators with certain potentials, *Ann. Inst. Fourier (Grenoble)* **45** (1995), no. 2, 513-546.
- [St93] E.M. Stein, *Harmonic Analysis*, Princeton Univ. Press, Princeton, 1993.
- [Tay96] M. Taylor, *Partial Differential Equations II, Qualitative Studies of Linear Equations*, Applied Mathematical Sciences **116**, Springer Verlag, 1996.
- [Ti58] E.C. Titchmarsh, *Eigenfunction Expansions Associated with Second-Order Differential Equations. (Part I, II)*, Oxford Univ. Press, 1958.
- [Tr83] H. Triebel, *Theory of Function Spaces*, Birkhäuser Verlag, 1983.
- [Tr92] ———, *Theory of Function Spaces II*, Monographs Math. **84**, Birkhäuser, Basel, 1992.
- [Tr78] ———, *Interpolation Theory, Function Spaces, Differential Operators*, Berlin, VEB Deutsch. Verl. Wissenschaften 1978, Amsterdam, North-Holland 1978.
- [V88] A. Tabacco Vignati, Complex interpolation for families of quasi-Banach spaces, *Indiana Univ. Math. J.* **37** (1988), 1-21.
- [Zh01] Q. Zhang, Global bounds of Schrödinger heat kernels with negative potentials, *J. Func. Anal.* **182** (2001), no.2, 344-370.
- [Z04a] S. Zheng, A representation formula related to Schrödinger operators, *Anal. Theo. Appl.* **20** (2004), no.3. <http://lanl.arXiv.org/math.SP/0412314>.
- [Z04b] ———, Perturbed Fourier transform associated to Schrödinger operators, *preprint* (2005).
- [Z03] ———, *Besov Spaces for Schrödinger Operators*, Dissertation, University of Maryland, College Park, 2003.
- [Z05] ———, *Littlewood-Paley theory, atomic decomposition and Schrödinger equations on \mathbb{R}^d* , (In preparation). Presented at the 111th AMS Conference, Atlanta, 2005.

(Gestur Ólafsson) DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE,
LA 70803

E-mail address: olafsson@math.lsu.edu

URL: <http://www.math.lsu.edu/~olafsson>

(Shijun Zheng) DEPARTMENT OF MATHEMATICS, INDUSTRIAL MATHEMATICS INSTITUTE, UNIVER-
SITY OF SOUTH CAROLINA, COLUMBIA, SC 29208

AND

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803

E-mail address: shijun@math.sc.edu

URL: <http://www.math.sc.edu/~shijun>