Periodicities in Linear Fractional Recurrences: Degree growth of birational surface maps

Eric Bedford and Kyounghee Kim

§0. Introduction

Given complex numbers $\alpha_0, \ldots, \alpha_p$ and β_0, \ldots, β_p , we consider the recurrence relation

$$x_{n+p+1} = \frac{\alpha_0 + \alpha_1 x_{n+1} + \dots + \alpha_p x_{n+p}}{\beta_0 + \beta_1 x_{n+1} + \dots + \beta_p x_{n+p}}.$$
 (0.1)

Thus a *p*-tuple (x_1, \ldots, x_p) generates an infinite sequence (x_n) . We consider two equivalent reformulations in terms of rational mappings: we may consider the mapping $f : \mathbf{C}^p \to \mathbf{C}^p$ given by

$$f(x_1, \dots, x_p) = \left(x_2, \dots, x_p, \frac{\alpha_0 + \alpha_1 x_1 + \dots + \alpha_p x_p}{\beta_0 + \beta_1 x_1 + \dots + \beta_p x_p}\right).$$
(0.2)

Or we may use the imbedding $(x_1, \ldots, x_p) \mapsto [1 : x_1 : \ldots : x_p] \in \mathbf{P}^p$ into projective space and consider the induced map $f : \mathbf{P}^p \to \mathbf{P}^p$ given by

$$f_{\alpha,\beta}[x_0:x_1:\ldots:x_p] = [x_0\beta \cdot x:x_2\beta \cdot x:\ldots:x_p\beta \cdot x:x_0\alpha \cdot x], \qquad (0.3)$$

where we write $\alpha \cdot x = \alpha_0 x_0 + \cdots + \alpha_p x_p$.

Here we will study the degree growth of the iterates $f^k = f \circ \cdots \circ f$ of f. In particular, we are interested in the quantity

$$\delta(\alpha,\beta) := \lim_{k \to \infty} \left(\operatorname{degree}(f_{\alpha,\beta}^k) \right)^{1/k}$$

A natural question is: for what values of α and β can (0.1) generate a periodic recurrence? In other words, when does (0.1) generate a periodic sequence (x_n) for all choices of x_1, \ldots, x_p ? This is equivalent to asking when there is an N such that $f_{\alpha,\beta}^N$ is the identity map. Periodicities in recurrences of the form (0.1) have been studied in [L, KG, KoL, GL, CL]. The question of determining the parameter values α and β for which $f_{\alpha,\beta}$ is periodic has been known for some time and is posed explicitly in [GKP] and [GL, p. 161]. Recent progress in this direction was obtained in [CL]. The connection with our work here is that if $\delta(\alpha, \beta) > 1$, then the degrees of the iterates of $f_{\alpha,\beta}$ grow exponentially, and $f_{\alpha,\beta}$ is far from periodic.

In the case p = 1, f is a linear (fractional) map of \mathbf{P}^1 . The question of periodicity for f is equivalent to determining when a 2×2 matrix is a root of the identity. In this paper we address these questions in the case p = 2. In fact, our principal efforts will be devoted to determining $\delta(\alpha, \beta)$ for all of the mappings in the family above. In order to remove trivial cases, we will assume throughout this paper that

$$(\alpha_0, \alpha_1, \alpha_2)$$
 is not a multiple of $(\beta_0, \beta_1, \beta_2)$,
 $(\alpha_1, \beta_1) \neq (0, 0)$, $(\alpha_2, \beta_2) \neq (0, 0)$, and
 $(\beta_1, \beta_2) \neq (0, 0)$.
(0.4)

Note that if the first condition in (0.4) is not satisfied, then the right hand side of (0.1) is constant. If the left hand part of the second condition (0.4) is not satisfied, then f does not depend on x_1 thus has rank 1, which cannot be periodic. If the right hand part of the second condition (0.4) is not satisfied, then f^2 is essentially the 1-dimensional mapping $\zeta \mapsto \frac{\alpha_0 + \alpha_1 \zeta}{\beta_0 + \beta_1 \zeta}$. If the third condition in (0.4) is not satisfied, then f is linear. In this case, the periodicity of f is a question of linear algebra.

Since we consider all parameters satisfying (0.4), we must treat a number of separate cases. By V_n we denote the variety of parameters (α, β) such that

$$\beta_2 = 0, \text{ and } f^n_{\alpha,\beta}(q) = p,$$

where $p = [\beta_1 \alpha_2 - \beta_2 \alpha_1 : -\beta_0 \alpha_2 + \alpha_0 \beta_2 : \alpha_1 \beta_0 - \alpha_0 \beta_1],$
and $q = [\beta_1(\beta_1 \alpha_2) : \beta_1(\alpha_1 \beta_0 - \alpha_0 \beta_1) : \alpha_1(\beta_1 \alpha_2 - \alpha_1 \beta_2)].$ (0.5)

The following two numbers are of special importance here:

 ϕ (~ 1.61803 golden mean) is the largest root of $x^2 - x - 1$ (0.6)

$$\delta_{\star} \ (\sim 1.32472)$$
 is the largest root of $x^3 - x - 1$ (0.7)

Theorem 1. If $(\alpha, \beta) \notin \bigcup_{n \ge 0} V_n$, then $\phi \ge \delta(\alpha, \beta) \ge \delta_* > 1$. For generic (α, β) , the dynamic degree is $\delta(\alpha, \beta) = \phi$.

In particular, we see that $f_{\alpha,\beta}$ has exponential degree growth in all of these cases. The remaining possibilities are:

Theorem 2. If $(\alpha, \beta) \in V_n$ for some $n \ge 0$, then there is a complex manifold $X = X_{\alpha,\beta}$ obtained by blowing up \mathbf{P}^2 at finitely many points, and $f_{\alpha,\beta}$ induces a biholomorphic map $f_{\alpha,\beta} : X \to X$. Further:

If n = 0, $f_{\alpha,\beta}$ is periodic of period 6. If n = 1, $f_{\alpha,\beta}$ is periodic of period 5. If n = 2, $f_{\alpha,\beta}$ is periodic of period 8. If n = 3, $f_{\alpha,\beta}$ is periodic of period 12. If n = 4, $f_{\alpha,\beta}$ is periodic of period 18. If n = 5, $f_{\alpha,\beta}$ is periodic of period 30. If n = 6, the degree of $f_{\alpha,\beta}^n$ is asymptotically quadratic in n.

If $n \geq 7$, $f_{\alpha,\beta}$ has exponential degree growth rate $\delta(\alpha,\beta) = \delta_n > 1$, which is given by the largest root of the polynomial $x^{n+1}(x^3 - x - 1) + x^3 + x^2 - 1$. Further, δ_n increases to δ_{\star} as $n \to \infty$.

The family of maps

$$(x,y) \mapsto (y, \frac{a+y}{x})$$

has been studied by several authors (cf. [L, KoL, KLR, GBM, CL]). Within this family, the case a = 0 corresponds to V_0 , a = 1 corresponds to V_1 , and all the rest belong to the case V_6 (see §6).

In the cases $n \ge 7$, the entropy of $f_{\alpha,\beta}$ is equal to $\log \delta_n$ by Cantat [C]. The number δ_{\star} is known (see [BDGPS, Chap. 7]) to be the infimum of all Pisot numbers. Diller and Favre [DF] showed that if g is a birational surface map which is not birationally conjugate to a holomorphic automorphism, then $\delta(g)$ is a Pisot number. So the maps f in the cases $n \ge 7$ have smaller degree growth than any such g. Note that projective surfaces which have automorphisms of positive entropy are relatively rare: Cantat [C] shows that, except for nonminimal rational surfaces (like X in Theorem 2), the only possibilities are complex tori, K3 surfaces, or Enriques surfaces.



Figure 0.1. A map with (maximal) degree growth ϕ .

Determining the dynamical degree for this family of mappings may be seen as a first step towards the dynamical study of these maps. Figure 0.1 portrays the stable/unstable laminations of a mapping of maximal degree growth within the family $f_{\alpha,\beta}$. This paper is organized as follows. In §1 we give the general properties of the family $f_{\alpha,\beta}$. In §2 we show that $\delta(f_{\alpha,\beta}) = \phi$ if $f_{\alpha,\beta}$ has only two exceptional curves. Next we determine $\delta(f_{\alpha,\beta})$ in the (generic) case where it has three exceptional curves. This determination, however, threatens to involve a large case-by-case analysis. We avoid this by adopting a more general approach. In §3 we show how $\delta(f_{\alpha,\beta})$ may be derived from the set of numbers in open and closed orbit lists. We do this by showing that results of [BK] may be extended from the "elementary" case to the general case. We use this in §4 to determine $\delta(\alpha, \beta)$ when the critical triangle is nondegenerate. In §5 we handle the periodic cases in Theorem 2. In §6, we discuss parameter space and the varieties V_n for $0 \le n \le 6$. We explain the computer pictures in the Appendix.

We wish to thank Curt McMullen for helpful comments on this paper.

$\S1$. Setting and Basic Properties

In this section we review some basic properties of the map

$$f(x) = [x_0\beta \cdot x : x_2\beta \cdot x : x_0\alpha \cdot x],$$

which is the map (0.3) in the case p = 2. (We refer to [GBM] for a description of f as a real map.) The indeterminacy locus is

$$\mathcal{I} = \{ x \in \mathbf{P}^2 : x_0(\beta \cdot x) = x_2(\beta \cdot x) = x_0(\alpha \cdot x) = 0 \}$$

= $\{ e_1, p_0, p_\gamma \},$

where we set $p_0 = [0 : -\beta_2 : \beta_1]$ and $p_{\gamma} = [\beta_1 \alpha_2 - \beta_2 \alpha_1 : -\beta_0 \alpha_2 + \alpha_0 \beta_2 : \alpha_1 \beta_0 - \alpha_0 \beta_1]$. Thus f is holomophic on $\mathbf{P}^2 - \mathcal{I}$, and its Jacobian is $2x_0(\beta \cdot x)[\beta_1(\alpha \cdot x) - \alpha_1(\beta \cdot x)]$. Let us set

$$\gamma = (\beta_1 \alpha_0 - \alpha_1 \beta_0, 0, \beta_1 \alpha_2 - \alpha_1 \beta_2) \in \mathbf{C}^{\natural}$$

and note that the Jacobian vanishes on the curves

$$\Sigma_0 = \{x_0 = 0\}, \quad \Sigma_\beta = \{\beta \cdot x = 0\}, \text{ and } \Sigma_\gamma = \{\gamma \cdot x = 0\}.$$

These curves are exceptional in the sense that they are mapped to points:

$$f(\Sigma_0 - \mathcal{I}) = e_1 := [0:1:0], \quad f(\Sigma_\beta - \mathcal{I}) = e_2 := [0:0:1], \quad f(\Sigma_\gamma - \mathcal{I}) = q, \tag{1.1}$$

where q is defined in (0.5). We write the set of exceptional curves as $\mathcal{E}(f) = \{\Sigma_0, \Sigma_\beta, \Sigma_\gamma\}$.

Lemma 1.1.

$$f(\mathbf{P}^2 - \Sigma_0 \cup \Sigma_\beta) \cap \Sigma_0 = \emptyset.$$

Further, if $\beta_2 \neq 0$,

$$f(\mathbf{P}^2 - \Sigma_0 \cup \{p_\gamma\}) \cap \{p_0\} = \emptyset.$$

Proof. In $\mathbf{P}^2 - \mathcal{E}(f) \cup \mathcal{I}(f)$, f is holomorphic. It follows that for $[x_0 : x_1 : x_2] \in \mathbf{P}^2 - \mathcal{E}(f) \cup \mathcal{I}(f)$, $f([x_0 : x_1 : x_2]) \notin \Sigma_0$ since $x_0(\beta \cdot x) \neq 0$. If $\beta_1 = 0$ or $\beta_1 \alpha_2 - \alpha_1 \beta_2 = 0$ then either $\Sigma_{\gamma} = \Sigma_{\beta}$ or $\Sigma_{\gamma} = \Sigma_0$. If both β_1 and $\beta_1 \alpha_2 - \alpha_1 \beta_2$ are non-zero, we have $f(\Sigma_{\gamma}) = q \notin \Sigma_0$. In case $\beta_2 \neq 0$, for $[x_0 : x_1 : x_2] \in \Sigma_{\beta}$, we have seen that $f([x_0 : x_1 : x_2]) = e_2 \neq p_0$, which completes the proof.

The inverse of f is given by the map

$$f^{-1}(x) = [x_0 B \cdot x : x_0 A \cdot x - \beta_2 x_1 x_2 : x_1 B \cdot x],$$

where we set $A = (\alpha_0, \alpha_2, -\beta_0)$ and $B = (-\alpha_1, 0, \beta_1)$. In the special case $\beta_2 = 0$, the form of f^{-1} is similar to that of f. The indeterminacy locus $\mathcal{I}(f^{-1}) = \{e_1, e_2, q\}$ consists of the three points which are the f-images of the exceptional lines for f. The Jacobian of f^{-1} is

$$-2x_0B \cdot x(\alpha_1\beta_0x_0 - \alpha_0\beta_1x_0 - \alpha_2\beta_1x_1 + \alpha_1\beta_2x_1).$$

Let us set $C = (\alpha_1\beta_0 - \alpha_0\beta_1, \alpha_1\beta_2 - \alpha_2\beta_1, 0)$, and $\Sigma_B = \{x \cdot B = 0\}$, $\Sigma_C = \{x \cdot C = 0\}$. In fact, $\mathcal{E}(f^{-1}) = \{\Sigma_0, \Sigma_B, \Sigma_C\}$, and f^{-1} acts as: $\Sigma_0 \mapsto p_0, \Sigma_B \mapsto e_1$, and $\Sigma_C \mapsto p_\gamma$.

To understand the behavior of f at \mathcal{I} , we define the cluster set $Cl_f(a)$ of a point $a \in \mathbf{P}^2$ by

$$Cl_f(a) = \{ x \in \mathbf{P}^2 : x = \lim_{a' \to a} f(a'), a' \in \mathbf{P}^2 - \mathcal{I}(f) \}.$$

In general, a cluster set is connected and compact. In our case, we see that the cluster set is a single point when $a \notin \mathcal{I}$, i.e., when f is holomorpic. And the cluster sets of the points of indeterminacy are found by applying f^{-1} : i.e., $e_1 \mapsto Cl_f(e_1) = \Sigma_B$, $p_0 \mapsto Cl_f(p_0) = \Sigma_0$, and $p_\gamma \mapsto Cl_f(p_\gamma) = \Sigma_C$. Thus f acts as in Figure 1.1: the lines on the left hand triangle are exceptional and are mapped to the vertices of the right hand triangle, and the vertices of the left hand triangle are blown up to the sides of the right hand triangle.

Let

$$\pi: Y \to \mathbf{P}^2 \tag{1.2}$$

be the complex manifold obtained by blowing up \mathbf{P}^2 at e_1 . We will discuss the induced birational map $f_Y: Y \to Y$. We let $E_1 := \pi^{-1}e_1$ denote the exceptional blow-up fiber. The projection gives a biholomorphic map $\pi: Y - E_1 \to \mathbf{P}^2 - e_1$. For a complex curve $\Gamma \subset \mathbf{P}^2$, we use the notation $\Gamma \subset Y$ to denote the *strict transform* of Γ in Y. Namely, Γ denotes the closure of $\pi^{-1}(\Gamma - e_1)$ inside Y. Thus Γ is a proper subset of $\pi^{-1}\Gamma = \Gamma \cup E_1$.

We identify E_1 with \mathbf{P}^1 in the following way. For $[\xi_0 : \xi_2] \in \mathbf{P}^1$, we associate the point

$$[\xi_0:\xi_2]_{E_1} := \lim_{t \to 0} \pi^{-1}[t\xi_0:1:t\xi_2] \in E_1.$$

We may now determine the map f_Y on Σ_0 . For $x = [0: x_1: x_2] = \lim_{t\to 0} [t: x_1: x_2] \in \Sigma_0$, we assign $f_Y x := \lim_{t\to 0} f[t: x_1: x_2] \in Y$. That is, $f[t: x_1: x_2] = [t\beta \cdot x: x_2\beta \cdot x: t\alpha \cdot x]$, and so taking the limit as $t \to 0$, we obtain

$$f_Y[0:x_1:x_2] = [\beta \cdot x:\alpha \cdot x]_{E_1}.$$
(1.3)



Figure 1.1. Blowing-up/blowing-down behavior of f.

Now we make a similar computation for a point $[\xi_0 : \xi_2]_{E_1}$ in the fiber E_1 over the point of indeterminacy e_1 . We set $x = [t\xi_0 : 1 : t\xi_2]$ so that

$$fx = [t\xi_0\beta \cdot x : t\xi_2\beta \cdot x : t\xi_0\alpha \cdot x].$$

Taking the limit as $t \to 0$, we find

$$f_Y([\xi_0:\xi_2]_{E_1}) = [\xi_0\beta_1:\xi_2\beta_1:\xi_0\alpha_1] \in \Sigma_A$$

Thus we have:

Lemma 1.2. The map f_Y has the properties:

(i) f_Y is a local diffeomorphism at points of Σ_0 if and only if $\beta_1 \alpha_2 - \alpha_1 \beta_2 \neq 0$.

(ii) f_Y is a local diffeomorphism at points of E_1 if and only if $\beta_1 \neq 0$.

§2. Degenerate Critical Triangle

We will refer to the set $\{\Sigma_0, \Sigma_\beta, \Sigma_\gamma\}$ of exceptional curves as the *critical triangle*; we say that the critical triangle is *nondegenerate* if these three curves are distinct. Since $(\beta_1, \beta_2) \neq (0, 0)$, we have $\Sigma_0 \neq \Sigma_\beta$. Thus there are only two possibilities for a degenerate triangle. The first of these is the case $\Sigma_\gamma = \Sigma_\beta$, which occurs when $\beta_1 = 0$. The second is $\Sigma_\gamma = \Sigma_0$, which occurs when $\beta_1 \alpha_2 - \alpha_1 \beta_2 = 0$. (And since $\Sigma_0 \neq \Sigma_\beta$ we have $\beta_1 \neq 0$ in this case.) We will show that $\delta(\alpha, \beta) = \phi$ when the critical triangle is degenerate. This is different from the general case (and easier), and we treat it in this section.

In order to determine the degree growth rate of f, we will consider the induced pullback f^* on $H^{1,1}$. We will be working on compact, complex surfaces X for which $H^{1,1}(X)$ is generated by the classes of divisors. If [D] is the divisor of a curve $D \subset X$, then we define $f^*[D]$ to be the class of the divisor $f^{-1}D$. We say that f is 1-regular if $(f^n)^* = (f^*)^n$ for all $n \ge 0$. Fornaess and Sibony showed in [FS] that if

for every exceptional curve C and all
$$n \ge 0, f^n C \notin \mathcal{I}$$
 (2.1)

then f is 1-regular. We will use this criterion in the following:

Proposition 2.1. If the critical triangle is degenerate, then the map $f_Y: Y \to Y$ is 1-regular.

Proof. We treat the two possibilities separately. The first case is $\Sigma_{\gamma} = \Sigma_{\beta}$; see Figure 2.1. In this case f has two exceptional lines Σ_0 and Σ_{β} and two points of indeterminacy $\mathcal{I} = \{e_1, p_{\lambda}\}$. After we blow up e_1 to obtain Y, the line Σ_0 is no longer exceptional. (Our drawing convention in this and subsequent Figures is that exceptional curves are thick, and points of indeterminacy

are circled.) By (1.3), we see that f_Y maps E_1 to $e_2 = q$, and thus the exceptional set becomes $\mathcal{E}(f_Y) = \{E_1, \Sigma_\beta = \Sigma_\gamma\}$. Now in order to check condition (2.1), we need to follow the orbit of e_2 . By (1.3) we see that e_2 is part of a 2-cycle $\{e_2, [\beta_2 : \alpha_2]_{E_1}\}$. On the other hand, the points of indeterminacy for f_Y are p_γ and $[0:1]_{E_1} = E_1 \cap \Sigma_0$. Since $\beta_1 = 0$ in this case, we have $\beta_2 \neq 0$, so (2.1) holds.



Figure 2.1. The case $\Sigma_{\beta} = \Sigma_{\gamma}$.

The second case is $\Sigma_{\gamma} = \Sigma_0$. Again, $\mathcal{I} = \{e_1, p_{\gamma}\}$, but $\mathcal{E}(f) = \{\Sigma_0, \Sigma_\beta\}$, and the arrangement of exceptional curves and points of indeterminacy are as in Figure 2.2. In this case, we have $\beta_1 \neq 0$, so by Lemma 1.2, we hve $\mathcal{I}(f_y) = \{p_0 = p_{\gamma}\}$ and $\mathcal{E}(f_Y) = \{\Sigma_\beta\}$. As before, we need to track the orbit of e_2 . But by Lemma 1.1, we see that we can never have $f^j e_2 = p_0$ for $j \geq 1$. Thus (2.1) holds in this case, too, and the proof is complete.



Figure 2.2. The case $\Sigma_0 = \Sigma_{\gamma}$.

Now let us determine f_Y^* . The cohomology group $H^{1,1}(\mathbf{P}^2; \mathbf{Z})$ is one-dimensional and is generated by the class of a complex line. We denote this generator by L. Let $L_Y := \pi^* L \in H^{1,1}(Y; \mathbf{Z})$ be the class induced by the map (1.2). It follows that $\{L_Y, E_1\}$ is a basis for $H^{1,1}(Y; \mathbf{Z})$. Now $\Sigma_0 = L \in H^{1,1}(\mathbf{P}^2; \mathbf{Z})$. Pulling this back by π , we have

$$L_Y = \pi^* \Sigma_0 = \Sigma_0 + E_1.$$

Now f_Y^* acts by taking pre-images:

$$f_Y^* E_1 = [f^{-1} E_1] = \Sigma_0 = L_Y - E_1,$$

where the last equality follows from the equation above.

Now e_1 is indeterminate, and $fe_1 = \Sigma_A$. Since Σ_A intersects any line L, it follows that $e_1 \in f^{-1}L$. Thus

$$\pi^*[f^{-1}L] = [f^{-1}L] + E_1 \in H^{1,1}(Y; \mathbf{Z}).$$

On the other hand, $f^{-1}L = 2L \in H^{1,1}(\mathbf{P}^2; \mathbf{Z})$. Thus

$$\pi^*[f^{-1}L] = \pi^* 2L = 2L_Y.$$

Putting these last two equations together, we have $f_Y^*L_Y = 2L_Y - E_1$. Thus

$$f_Y^* = \begin{pmatrix} 2 & 1\\ -1 & -1 \end{pmatrix},$$

which is a matrix with spectral radius equal to ϕ . This yields the following:

Proposition 2.2. If the critical triangle is degenerate, then $\delta(\alpha, \beta) = \phi$.

§3. Regularization and Degree Growth

In this Section we discuss a different, but more general, family of maps. By $J: \mathbf{P}^2 \to \mathbf{P}^2$ we denote the involution

$$J[x_0:x_1:x_2] = [x_0^{-1}:x_1^{-1}:x_2^{-1}] = [x_1x_2:x_0x_2:x_0x_1].$$

For an invertible linear map L of \mathbf{P}^2 we consider the map $f := L \circ J$. The exceptional curves are $\mathcal{E} = \{ \mathbf{\Sigma}_0, \mathbf{\Sigma}_1, \mathbf{\Sigma}_2 \}$, where $\mathbf{\Sigma}_j := \{ x_j = 0 \}, j = 0, 1, 2$, and the points of indeterminacy are $\mathcal{I} = \{ \epsilon_0, \epsilon_1, \epsilon_2 \}$, where $\epsilon_i = \mathbf{\Sigma}_j \cap \mathbf{\Sigma}_k$, with $\{ i, j, k \} = \{ 0, 1, 2 \}$. We define $\mathbf{a}_j := f(\mathbf{\Sigma}_j - \mathcal{I}) = L\epsilon_j$ for j = 0, 1, 2.

For $p \in \mathbf{P}^2$ we define the orbit $\mathcal{O}(p)$ as follows. If $p \in \mathcal{E} \cup \mathcal{I}$, then $\mathcal{O}(p) = \{p\}$. If there exists an $N \geq 1$ such that $f^j p \notin \mathcal{E} \cup \mathcal{I}$ for $0 \leq j \leq N-1$ and $f^N p \in \mathcal{E} \cup \mathcal{I}$, then we set $\mathcal{O}(p) = \{p, fp, \dots, f^N p\}$. Otherwise we have $f^j p \notin \mathcal{E} \cup \mathcal{I}$ for all $j \geq 0$, and we set $\mathcal{O}(p) = \{p, fp, f^2 p, \dots\}$. We say the orbit $\mathcal{O}(p)$ is singular if it is finite; otherwise, it is non-singular. We say an orbit $\mathcal{O}(p)$ is elementary if it is either non-singular, or if it ends at a point of indeterminacy. In other words, a non-elementary orbit ends in a point of $\mathcal{E} - \mathcal{I}$.

Lemma 3.1. If f has at least one singular orbit, then it has a singular orbit that is elementary.

Proof. Suppose for all $i \in S_0$, \mathcal{O}_i is non-singular. If follows that every orbit $\mathcal{O}_j, j \notin S_0$ ends at a point in $\Sigma_i, i \in S_0$. Since all $\mathcal{O}_i, i \in S_0$ are non-singular, $\Sigma_j, j \notin S_0$ cannot end at a point of indeterminacy. This means that f is 1-regular.

Henceforth, we will assume that f has singular orbits. Let us write $\mathcal{O}_i = \mathcal{O}(\mathbf{a}_i) = \mathcal{O}(f(\mathbf{\Sigma}_i - \mathcal{I}))$ for the orbit of an exceptional curve. We set

$$S = \{i \in \{0, 1, 2\} : \mathcal{O}_i \text{ is singular}\},\$$

and

 $S_0 = \{i \in \{0, 1, 2\} : \mathcal{O}_i \text{ is singular and elementary} \}.$

Let $\mathcal{O}_{S_0} = \bigcup_{i \in S_0} \mathcal{O}_i$. We write $X_0 = \mathbf{P}^2$, and let $\pi : X_1 \to X_0$ be the complex manifold obtained by blowing up the points of \mathcal{O}_{S_0} . We let $f_1 : X_1 \to X_1$ denote the induced birational mapping. By Lemma 1.2, we see that the curves Σ_i , $i \in S_0$, are not exceptional for f_1 , and the blowing up operation constructed no new points of indeterminacy for f_1 . Thus the exceptional curves for f_1 are Σ_i for $i \notin S_0$. If S_0 is a proper subset of S, then for $i \in S - S_0$ we redefine \mathcal{O}_i to be the f_1 orbit of \mathbf{a}_i inside X_1 . Let us define $S_1 = \{i \in S - S_0 : \mathcal{O}_i \text{ is elementary}\}$. We may apply Lemma 3.1 to conclude that if $S - S_0 \neq \emptyset$, then $S_1 \neq \emptyset$. As before, we may define $\mathcal{O}_{S_1} = \bigcup_{i \in S_1} \mathcal{O}_i$, and we construct the complex manifold $\pi : X_2 \to X_1$ by blowing up all the points of \mathcal{O}_{S_1} . Doing this, we reach the situation where every singular orbit \mathcal{O}_i has the property that it is elementary in some X_i , and thus it has the form $\mathcal{O}_i = \{\mathbf{a}_i, \ldots, \epsilon_{\tau(i)}\}$ for some $\tau(i) \in \{0, 1, 2\}$.

Next we organize the singular orbits \mathcal{O}_i into lists, as follows (modulo permutation of the indices $\{0, 1, 2\}$). If there is only one singular orbit, we have the list $\mathcal{L} = \{\mathcal{O}_i = \{\mathbf{a}_i, \ldots, \epsilon_{\tau(i)}\}\}$. If $\tau(i) = i$, we say that \mathcal{L} is a *closed list*; otherwise it is an *open list*. If there are two singular orbits, we can have two closed lists:

$$\mathcal{L}_1 = \{ \mathcal{O}_0 = \{ \mathbf{a}_0, \dots, \epsilon_0 \} \}, \ \mathcal{L}_2 = \{ \mathcal{O}_1 = \{ \mathbf{a}_1, \dots, \epsilon_1 \} \}$$

or a closed list and an open list:

$$\mathcal{L}_1 = \{\mathcal{O}_0 = \{\mathbf{a}_0, \dots, \epsilon_0\}\}, \ \mathcal{L}_2 = \{\mathcal{O}_1 = \{\mathbf{a}_1, \dots, \epsilon_2\}\}.$$

We cannot have two open lists since there are only 3 orbits \mathcal{O}_i . We can also have a single list :

$$\mathcal{L} = \{\mathcal{O}_0 = \{\mathbf{a}_0, \dots, \epsilon_1\}, \mathcal{O}_1 = \{\mathbf{a}_1, \dots, \epsilon_{\tau(1)}\}\},\$$

which is a closed list if $\tau(1) = 0$ and an open list otherwise. If there are three singular orbits, then the possibilities are

$$\mathcal{L} = \{\mathcal{O}_0 = \{\mathbf{a}_0, \dots, \epsilon_1\}, \mathcal{O}_1 = \{\mathbf{a}_1, \dots, \epsilon_2\}, \mathcal{O}_2 = \{\mathbf{a}_2, \dots, \epsilon_0\}\},\$$
$$\mathcal{L}_1 = \{\mathcal{O}_0 = \{\mathbf{a}_0, \dots, \epsilon_0\}\}, \ \mathcal{L}_2 = \{\mathcal{O}_1 = \{\mathbf{a}_1, \dots, \epsilon_2\}, \mathcal{O}_2 = \{\mathbf{a}_2, \dots, \epsilon_1\}\},\$$

or

$$\mathcal{L}_1 = \{\mathcal{O}_0 = \{\mathbf{a}_0, \dots, \epsilon_0\}\}, \mathcal{L}_2 = \{\mathcal{O}_1 = \{\mathbf{a}_1, \dots, \epsilon_1\}\}, \mathcal{L}_3 = \{\mathcal{O}_2 = \{\mathbf{a}_2, \dots, \epsilon_2\}\},\$$

where all the lists are closed.

For an orbit \mathcal{O}_i , we let $n_i = |\mathcal{O}_i|$ denote its length, and for an orbit list $\mathcal{L} = \{\mathcal{O}_a, \dots, \mathcal{O}_{a+\mu}\}$, we denote the set of orbit lengths by $|\mathcal{L}| = \{n_a, \dots, n_{a+\mu}\}$. We set $\#\mathcal{L}^c = \{|\mathcal{L}_j| : \mathcal{L}_j \text{ is closed}\}$ and $\#\mathcal{L}^o = \{|\mathcal{L}_j| : \mathcal{L}_j \text{ is open}\}$. The set $\#\mathcal{L}^c$ and $\#\mathcal{L}^o$ determine $\delta(f)$, as is shown in the following:

Theorem 3.2. If $f = L \circ J$, then the dynamic degree $\delta(f)$ is the largest real zero of the polynomial

$$\chi(x) = (x-2) \prod_{\mathcal{L} \in \mathcal{L}^c \cup \mathcal{L}^o} T_{\mathcal{L}}(x) + (x-1) \sum_{\mathcal{L} \in \mathcal{L}^c \cup \mathcal{L}^o} S_{\mathcal{L}}(x) \prod_{\mathcal{L}' \neq \mathcal{L}} T_{\mathcal{L}'}(x).$$
(3.1)

Here \mathcal{L} runs over all orbit lists. For each orbit list \mathcal{L} , we let N denote the sum of all the length of the orbits in \mathcal{L} . If \mathcal{L} is closed $T_{\mathcal{L}}(x) = x^N - 1$, and if \mathcal{L} is open $T_{\mathcal{L}}(x) = x^N$. The polynomial $S_{\mathcal{L}}$ is defined by

$$S_{\mathcal{L}}(x) = 1 \quad \text{if } |\mathcal{L}| = \{n_1\} \\ = x^{n_1} + x^{n_2} + 2 \quad \text{if } \mathcal{L} \text{ is closed and } |\mathcal{L}| = \{n_1, n_2\} \\ = x^{n_1} + x^{n_2} + 1 \quad \text{if } \mathcal{L} \text{ is open and } |\mathcal{L}| = \{n_1, n_2\} \\ = \sum_{i=1}^{3} [x^{N-n_i} + x^{n_i}] + 3 \quad \text{if } \mathcal{L} \text{ is closed and } |\mathcal{L}| = \{n_1, n_2, n_3\} \\ = \sum_{i=1}^{3} x^{N-n_i} + \sum_{i \neq 2} x^{n_i} + 1 \quad \text{if } \mathcal{L} \text{ is open and } |\mathcal{L}| = \{n_1, n_2, n_3\}.$$

The rest of this section is devoted to the proof of Theorem 3.2. We start by considering the case where f is elementary. In this case we have $S = S_0$. We set $X := X_1$. It follows from (2.1) that $f_X : X \to X$ is 1-regular, and thus $\delta(f)$ is spectral radius of f_X^* . The computation given in the Appendix of [BK] then shows that (3.1) is the characteristic polynomial of f_X^* .

For $p \in \mathcal{O}_S - \mathcal{I}$ we let $\mathcal{F}_p = \pi^{-1}p$ denote the exceptional fiber over p. If $\epsilon_i \in \mathcal{O}_S \cap \mathcal{I}$, we let E_i denote the exceptional fiber over ϵ_i . We will feel free to identify curves with the classes they

generate in $H^{1,1}(X)$. Let $H \in H^{1,1}(\mathbf{P}^2)$ denote the class of a line, and let $H_X = \pi^* H$ denote the induced class in $H^{1,1}(X)$. For $i \in S$, we have

$$\Sigma_i \to \mathbf{a}_i \to \cdots \to f^{n_i-1} \mathbf{a}_i = f^{n_i} (\Sigma_i - \mathcal{I}) = \epsilon_{\tau(i)}$$

for some $\tau(i) \in \{0, 1, 2\}$. At each points $f^j \mathbf{a}_i$, $0 \le j \le n_i - 1$, f is locally biholomorphic, so f_X induces a biholomorphic map

$$f_X : \mathcal{F}_{f^j \mathbf{a}_i} \to \mathcal{F}_{f^{j+1} \mathbf{a}_i} \quad 0 \le j \le n_i - 2, \text{ and}$$

 $f_X : \mathcal{F}_{f^{n_i - 1} \mathbf{a}_i} \to E_{\tau(i)}.$

It follows that

$$f_X^* \mathcal{F}_{f^{j+1}\mathbf{a}_i} = \mathcal{F}_{f^j\mathbf{a}_i} \quad \text{for } 0 \le j \le n_i - 2, \ i \in S$$

$$f_X^* E_{\tau(i)} = \mathcal{F}_{f^{n_i - 1}\mathbf{a}_i} \tag{3.2}$$

and

$$f_X^* \mathcal{F}_{\mathbf{a}_i} = \{ \mathbf{\Sigma}_i \} \quad \text{for } i \in S \tag{3.3}$$

where $\{\Sigma_i\}$ is the induced class by Σ_i in $H^{1,1}(X)$. Let $\Omega = \mathcal{I} \cap \{\epsilon_{\tau(i)} = f^{n_i - 1}\mathbf{a}_i, i \in S\}$, the set of blow-up centers which belongs to \mathcal{I} . Let us denote \mathcal{A} the set of indices i such that \mathcal{O}_i is singular orbit and is the first orbit in an open orbit list. For each i, Σ_i contains blow-up centers in the set $\Omega - \{\epsilon_i\}$. Notice that if $i \in \mathcal{A}, \epsilon_i \notin \Omega$, otherwise $\epsilon_i \in \Omega$. Using the identigy $\pi^* \{\Sigma_i\} = \{\pi^{-1}\Sigma_i\}$, we have

$$\{\boldsymbol{\Sigma}_i\} = H_X - E_\Omega + E_i \quad i \notin \mathcal{A} \{\boldsymbol{\Sigma}_i\} = H_X - E_\Omega \quad i \in \mathcal{A}$$
(3.4)

where $E_{\Omega} := \sum_{\epsilon_t \in \Omega} E_t$. A generic hyperplane \mathcal{H} in \mathbf{P}^2 does not contain any blow-up centers and may be considered to be subset of X. Let us restrict the map to $X - \mathcal{I}$. A generic hyperplane \mathcal{H} does intersect with any line in \mathbf{P}^2 . It follows that $\epsilon_i \in f_X^{-1}\mathcal{H}$, $i \in \Omega$ and we have

$$2H_X = \pi^*(f^*H) = \pi^*\{f^{-1}\mathcal{H}\} = f_X^* + E_\Omega$$

Therefore under f_X^* , we have

$$f_X^* H_X = 2H_X - E_\Omega. (3.5)$$

Now let us suppose that f is not elementary. Let $S = S_0 \cup S_1 \cup S_2$ and the manifolds $\pi_{i+1} : X_{i+1} \to X_i$ be as above. Let us set $X := X_3$. By (2.1) again, the induced map $f_X : X \to X$ is 1-regular. If $p \in X_i$ is a center of blow-up, we let \mathcal{F}_p denote the exceptional fiber inside X_{i+1} , and we use the same notation for the divisor in X given by the strict transform of \mathcal{F}_p ; in particular, \mathcal{F}_p is irreducible. Thus for $i \in S_k$, f_X induces a biholomorphic map

$$f_X: \mathcal{F}_{f_k^j \mathbf{a}_i} \to \mathcal{F}_{f_k^{j+1} \mathbf{a}_i} \quad 0 \le j \le n_i - 1.$$

It follows that

$$\begin{aligned}
f_X^* \mathcal{F}_{f_k^{j+1} \mathbf{a}_i} &= \mathcal{F}_{f_k^j \mathbf{a}_i} \quad \text{for } 0 \le j \le n_i - 1, \ i \in S_k \\
f_X^* \mathcal{F}_{\mathbf{a}_i} &= \{ \mathbf{\Sigma}_i \} \text{ for } i \in S
\end{aligned}$$
(3.6)

where $\{\Sigma_i\}$ is the induced class by Σ_i in $H^{1,1}(X)$. Let $\Omega_i := \{p \in \mathcal{O}_S : \pi(p) = \epsilon_i\}$ the set of blow-up centers whose image of π is $\epsilon_i \in \mathcal{I}$ and let $\Omega := \bigcup_{i \in S} \Omega_i$. For each $i \in S$, we denote

 $\Xi_i := \{p \in \mathcal{O}_S : \pi(p) \in \Sigma_i - \mathcal{I}\}$ the set of blow-up centers which belongs to exceptional line $\Sigma_i - \mathcal{I}$ and we let $\Xi = \bigcup_{i \in S} \Xi_i$. We also use the notation \mathcal{A} for the set of indices *i* such that \mathcal{O}_i is singular orbit and is the first orbit in an open orbit list. For each $i \in S$, Σ_i contains blow-up centers in the set $(\Omega - \Omega_i) \cup \Xi_i$. with $H_X = \pi^* H$ the induced class in $H^{1,1}(X)$, we have

$$\{\boldsymbol{\Sigma}_i\} = H_X - E_{\Omega} + E_i - \mathcal{F}_{\Xi_i} \quad i \notin \mathcal{A} \{\boldsymbol{\Sigma}_i\} = H_X - E_{\Omega} - \mathcal{F}_{\Xi_i} \quad i \in \mathcal{A}$$

$$(3.7)$$

where $E_{\Omega} = \sum_{p \in \Omega} \mathcal{F}_p$, $E_i = \sum_{p \in \Omega_i} \mathcal{F}_p$, and $\mathcal{F}_{\Xi_i} = \sum_{p \in \Xi_i} \mathcal{F}_p$. We also have

$$2H_X = \pi^*(f^*H) = f_X^* + E_\Omega.$$
(3.8)

To finish the proof, let us suppose that g is an elementary map, and f is not elementary, but both have the same orbit list structure given by $\#\mathcal{L}^c, \#\mathcal{L}^o$. We have shown that g^* is represented by the transformation (3.2–5), and f^* is represented by the transformation (3.6–8). To finish the proof, we show that these two linear transformations have the same characteristic polynomials. We illustrate this computation with an example which appears later in the paper. (The matrix computation for the other cases are similar.) We consider the case where the list structures of fand g are both given by

$$#\mathcal{L}^o = \emptyset, \quad #\mathcal{L}^c = \{\{1, 6\}\}.$$

We may also assume that $1 \in S_0, 2 \in S_1$ and

$$\mathcal{O}_1 = \{\mathbf{a}_1 = \epsilon_2\}, \quad \mathcal{O}_2 = \{\mathbf{a}_2, f_1\mathbf{a}_2 \in \mathbf{\Sigma}_1, f_1^2\mathbf{a}_2 \in \Omega_2, f_1^3\mathbf{a}_2, \dots, f_1^5\mathbf{a}_2 = \epsilon_1\}.$$

Combining (3.2-5) and (3.6-8), we have the matrix representations for g and f:

$$M_{g} = \begin{pmatrix} 2 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 & -1 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ \end{pmatrix}, M_{f} = \begin{pmatrix} 2 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

To show they have the same characteristic polynomial we are going to look at the matrices $M_g - xI$ and $M_f - xI$ where I is the identity matrix and will show that after row and column operations to $M_f - xI$ we get the same matrix as $M_g - xI$.

$$M_f - xI = \begin{pmatrix} 2-x & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & -1-x & 0 & 0 & 0 & 0 & 0 & -1 \\ -1 & -1 & -x & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & -x & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -x & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 1 & -x & 0 & -1 \\ 0 & -1 & 0 & 0 & 0 & 1 & -x & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -x \end{pmatrix}$$

First we subtract the second row from the 6-th row. For the general situation, we subtract rows for the lower generation chain from the rows for the corresponding part of the higher generation chain. Then we add 6-th column to the second column to remove the additional part and to obtain the matrix $M_g - xI$. For the general case, we add columns for the orbit collision part of the higher generation chain to the corresponding lower generation chain to remove extra elements. It is clear that $M_f - xI$ and $M_g - xI$ have the same determinant. This gives us the desired result.

§4. Non-degenerate Critical Triangle

In this section we will determine the degree growth rate of f with non-degenerate critical triangle. As we noted at the beginning of §2, it is equivalent to assume that

$$\beta_1 \neq 0$$
, and $\beta_1 \alpha_2 - \alpha_1 \beta_2 \neq 0$. (4.1)

In particular, the curves Σ_{γ} , Σ_{β} and Σ_0 are distinct, as well as $\{e_1, e_2, q\}$, the points of indeterminacy of f^{-1} . Let us choose invertible linear maps M_1 and M_2 of \mathbf{P}^2 such that

$$M_1 \Sigma_0 = \Sigma_0, \ M_1 \Sigma_1 = \Sigma_\beta, \ M_1 \Sigma_2 = \Sigma_\gamma,$$

and

$$M_2e_1 = \epsilon_0, \ M_2e_2 = \epsilon_1, \ M_2q = \epsilon_2.$$

It follows that $M_2 \circ f_{\alpha,\beta} \circ M_1$ is a quadratic map with $\Sigma_j \leftrightarrow e_j$ and so is equal to the map J. Thus $f_{\alpha,\beta}$ is linearly conjugate to a mapping of the form $L \circ J$. We will determine $\delta(\alpha,\beta)$ by finding the possibilities for $\#\mathcal{L}^{c/o}$ and then applying Theorem 3.2. When we treat $f_{\alpha,\beta}$ as a mapping $L \circ J$, we make the identifications

$$\Sigma_0 = \Sigma_0, \quad \Sigma_1 = \Sigma_\beta, \quad \Sigma_2 = \Sigma_\gamma,$$

 $\epsilon_0 = p_\gamma, \quad \epsilon_1 = e_1, \quad \epsilon_2 = p_0,$

and

$$\mathbf{a}_0 = f(\mathbf{\Sigma}_0 - \mathcal{I}(f)) = e_1, \ \mathbf{a}_1 = f(\mathbf{\Sigma}_1 - \mathcal{I}(f)) = e_2, \ \mathbf{a}_2 = f(\mathbf{\Sigma}_2 - \mathcal{I}(f)) = q$$

Thus we have $f(\Sigma_0 - \mathcal{I}) = \mathbf{a}_0 = \epsilon_1$, so the orbit $\mathcal{O}_0 = {\mathbf{a}_0 = \epsilon_1}$ is singular and has length one. There are two possibilities for the exceptional component Σ_1 ; the first is that $\mathbf{a}_1 \in \Sigma_0 - \mathcal{I}(f)$, which occurs when $\beta_2 \neq 0$. (See Figure 4.1.) The second possibility is $\mathbf{a}_1 = \epsilon_2 \in \mathcal{I}$, which occurs when $\beta_2 = 0$. (See Figure 4.2.) An analysis of the possibilities for \mathcal{O}_1 and \mathcal{O}_2 will yield the candidates for $|\mathcal{O}_1|$, $|\mathcal{O}_2|$ and $\#\mathcal{L}^{c/o}$, and thus give the possibilities for $\delta(\alpha, \beta)$.

Theorem 4.1. If the critical triangle is non-degenerate and $\beta_2 \neq 0$, then $\delta_{\star} \leq \delta(\alpha, \beta) \leq \phi$.

Proof. Let $f_Y: Y \to Y$ be as in (1.2). Since $\mathbf{a}_1 = e_2 \neq \epsilon_i$ for i = 0, 1, 2, we have

$$f_Y: \Sigma_1 - \mathcal{I} \to \mathbf{a}_1 \to [\beta_2: \alpha_2]_{E_1} \to [\beta_1 \beta_2: \beta_1 \alpha_2: \alpha_1 \beta_2] \in \Sigma_B - \Sigma_0.$$

$$(4.2)$$

If $f^2 \mathbf{a}_1 = f_Y^2 \mathbf{a}_1 = \epsilon_0$, then both lines Σ_2 and Σ_B contain ϵ_1, ϵ_0 . Since $\mathbf{a}_2 = \Sigma_B \cap \Sigma_1$ and $\epsilon_0 = \Sigma_2 \cap \Sigma_1$, we have $\mathbf{a}_2 = \epsilon_0$. By the second statement of Lemma 1.1, we see that the end points of both orbits \mathcal{O}_1 and \mathcal{O}_2 can not be ϵ_2 . It follows that we have at most two singular orbits including \mathcal{O}_0 . We have three cases.



Figure 4.1. Nondegenerate critical triangle: case $\beta_2 \neq 0$.

The first case is where neither \mathcal{O}_1 nor \mathcal{O}_2 is singular. In this case the orbit list structure is $\#\mathcal{L}^c = \emptyset, \#\mathcal{L}^o = \{1\}$. By Theorem 3.2, $\delta(\alpha, \beta)$ is the largest real root of the polynomial

$$\chi(x) = (x-2)x + (x-1) = x^2 - x - 1 \tag{4.3}$$

and is thus equal to ϕ .

In the second case both \mathcal{O}_0 and \mathcal{O}_1 are singular. In this case the orbit \mathcal{O}_2 can not be singular and therefore $f^2 \mathbf{a}_1 \neq \epsilon_0$. By the equation (4.2) with above argument, we have $n_1 = |\mathcal{O}_1| \geq 4$ and $\mathcal{O}_1 = {\mathbf{a}_1, \ldots, \epsilon_0}$. It follows that $\#\mathcal{L}^o = \emptyset, \#\mathcal{L}^c = {1, n_1}$. The dynamic degree $\delta(\alpha, \beta)$ is the largest root of the polynomial

$$\chi(x) = (x-2)(x^{1+n_1}-1) + (x-1)(x+x^{n_1}+2) = x^{n_1}(x^2-x-1) + x^2.$$
(4.4)

When $n_1 = 4$, the characteristic polynomial is given by $x^6 - x^5 - x^4 + 2 = x^2(x-1)(x^3 - x - 1)$. Thus $\delta = \delta_{\star}$ in this case. Let us observe that the Comparison Principle [BK, Theorem 5.1] concerns the modulus of the largest zero of the characteristic polynomial of f^* . In §3 we showed that the characteristic polynomials are the same in the elementary and the non-elementary cases. Thus we may apply the Comparison Principle to conclude that $\delta(\alpha, \beta) \geq \delta_{\star}$ if $n_1 \geq 4$.



Figure 4.2. Nondegenerate critical triangle: case $\beta_2 = 0$.

The last case is where both \mathcal{O}_0 and \mathcal{O}_2 are singular. We have $n_2 = |\mathcal{O}_2| \ge 1$ and $\mathcal{O}_2 = \{\mathbf{a}_2, \ldots, \epsilon_0\}$. Therefore the orbit list structure is $\#\mathcal{L}^c = \emptyset, \#\mathcal{L}^o = \{n_2, 1\}$. By Theorem 3.2, the dynamic degree $\delta(\alpha, \beta)$ is the largest root of the polynomial

$$\chi(x) = (x-2)x^{1+n_2} + (x-1)(x+x^{n_2}+1) = x^{n_2}(x^2-x-1) + x^2 - 1.$$
(4.5)

If $n_2 = 1$, we have $\chi(x) = x^3 - x - 1$.

Theorem 4.2. Assume that the critical triangle is non-degenerate. If $\beta_2 = 0$ and $n_2 = |\mathcal{O}_2| \ge 8$, then $1 < \delta(\alpha, \beta) \le \delta_{\star}$. If $\beta_2 = 0$ and $n_2 = |\mathcal{O}_2| \le 7$, then $\delta(\alpha, \beta) = 1$.

Proof. If $\beta_2 = 0$, we have $\mathbf{a}_1 = \epsilon_2$ and therefore we have

$$\mathcal{O}_0 = \{\mathbf{a}_0 = \epsilon_1\}, \text{ and } \mathcal{O}_1 = \{\mathbf{a}_1 = \epsilon_2\}.$$

If the orbit \mathcal{O}_2 is non-singular, we have the orbit list structure $\#\mathcal{L}^o = \{1,1\}, \#\mathcal{L}^c = \emptyset$. By Theorem 3.2, the degree growth rate $\delta(\alpha, \beta)$ is the largest root of the polynomial

$$\chi(x) = (x-2)x^2 + (x-1)(x+x+1) = x^3 - x - 1.$$
(4.6)

If the orbit \mathcal{O}_2 is singular, the end point of the orbit has to be the remaining point of indeterminacy, ϵ_0 . Thus we have $n_2 = |\mathcal{O}_2| \ge 1$ and $\mathcal{O}_2 = \{\mathbf{a}_2, \ldots, \epsilon_0\}$. It follows that the orbit list structure $\#\mathcal{L}^c = \{1, 1, n_2\}, \#\mathcal{L}^o = \emptyset$. Using the Lemma 2 and Proposition 7 in §3, the dynamic degree is the largest root of the polynomial

$$\chi(x) = (x-2)(x^{2+n_2}-1) + (x-1)(2x^{1+n_2}+x^2+x^{n_2}+2x+3)$$

= $x^{n_2}(x^3-x-1) + x^3 + x^2 - 1.$ (4.7)

It follows that $1 \leq \delta(\alpha, \beta) \leq \delta_{\star}$. For $n_2 = 7$, we have $\chi(x) = (x^2 - 1)(x^3 - 1)(x^5 - 1)$ and so the $\delta(\alpha, \beta) = 1$. For $n_2 = 8$ we have $\chi(x) = (x - 1)(x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1)$ and $\chi'(1) < 0$ and therefore the largest real root is strictly bigger than 1. It follows, then, from the comparison principle ([BK, Theorem 5.1]) that $\delta(\alpha, \beta) > 1$ if $n_2 \geq 8$.

Let us note that when the orbit of q lands on p, and we blow up the orbit of q, then we have removed the last exceptional curves for f and f^{-1} . Thus we have:

Proposition 4.3. If $(\alpha, \beta) \in V_n$, then the induced map $f_X : X \to X$ is biholomorphic.

Figure 4.3 shows the arrangement of the exceptional varieties in X in the case where the orbit of q does not enter Σ_{β} .



Figure 4.3. Nondegenerate critical triangle; elementary case $(\alpha, \beta) \in V_n$.

§5. Periodic Mappings

Here we determine the precise degree growth rate when $|\mathcal{O}_2| \leq 7$. In particular, we show that the degree grows quadratically when $|\mathcal{O}_2| = 7$, and we show that f is periodic when $|\mathcal{O}_2| \leq 6$. We do this by showing that f^* is periodic in this case, and then we show that the periodicity of f^* implies the periodicity of f.

Notice that if $|\mathcal{O}_2| = n$, then $f^n(\Sigma_{\gamma}) = f^{n-1}(q) = p$, and therefore $(\alpha, \beta) \in V_{n-1}$. To show the periodicity of f_X^* it suffices to show that all roots of (4.7) with $n \leq 6$ are roots of unity and are simple. For $n \leq 6$ we list the characteristic polynomials, together with the smallest polynomials of the form $x^m - 1$ that they divide:

$$V_0 (n = 1): (x - 1)(x + 1)(x^2 + x + 1)|(x^6 - 1)$$

$$V_1 (n = 2): (x - 1)(x^4 + x^3 + x^2 + x + 1)|(x^5 - 1)$$

$$V_2 (n = 3): (x - 1)(x + 1)(x^4 + 1)|(x^8 - 1)$$

$$V_3 (n = 4): (x - 1)(x^2 + x + 1)(x^4 - x^2 + 1)|(x^{12} - 1)$$

$$V_4 (n = 5): (x - 1)(x + 1)(x^6 - x^3 + 1)|(x^{18} - 1)$$

$$V_5 (n = 6): (x - 1)(x^8 + x^7 - x^5 - x^4 - x^3 + x + 1)|(x^{30} - 1)$$

Thus we have:

Lemma 5.1. Assume that the critical triangle is non-degenerate. If $\beta_2 = 0$ and $n = |\mathcal{O}_2| \leq 6$, then f_X^* is periodic, with period κ_n , where $\kappa_n = 6, 5, 8, 12, 18, 30$ (respectively).

When $|\mathcal{O}_2| = 7$, the largest root of equation (4.7) is 1 and has multiplicity 3. Whether f is elementary or not, the matrix representation from §3 has a 3×3 Jordan block with eigenvalue 1. This means that f_X^* has quadratric growth, and we have:

Lemma 5.2. Assume that the critical triangle is non-degenerate. If $\beta_2 = 0$ and $|\mathcal{O}_2| = 7$, then f_X^* has quadratic growth.

Notice that $|\mathcal{O}_2| = 1$ if and only if $q = p_{\gamma}$, which means that the parameters in V_0 satisfy $\alpha_1\beta_0 - \alpha_0\beta_1 = -\alpha_2\beta_0 = \alpha_1\alpha_2$. With these conditions on α and β , it is not hard to check that the map f is indeed periodic with period 6. We could also see this by observing that f has a period 6 cycle $\Sigma_\beta \mapsto e_2 \mapsto \Sigma_0 \mapsto e_1 \mapsto \Sigma_\gamma \mapsto p_\gamma \mapsto \Sigma_\beta$.

Theorem 5.3. Assume that the critical triangle is non-degenerate. if $\beta_2 = 0$ and $|\mathcal{O}_2| \leq 6$, then f is periodic with period κ_n .

To prove Theorem 5.3, we use the following lemma:

Lemma 5.4. If $f : \mathbf{P}^2 \to \mathbf{P}^2$ is a linear map with five invariant lines which are in general position, then f is the identity.

Proof. Let l_i , i = 0, 1, 2, 3, 4, denote the lines fixed by f. Since they are in general position, we may assume that they have the form $\Sigma_i = \{x_i = 0\}$ for $i = 0, 1, 2, \Sigma_* = \{x_0 + w_1x_1 + w_2x_2 = 0, w_1 \neq 0\}$, and $\Sigma_{**} = \{x_0 + v_1x_1 + v_2x_2 = 0, v_2 \neq 0\}$. A computation then shows that f must be the identity.

Proof of Theorem 5.3. It suffices to show that f^{κ_n} has at least five invariant lines for $n = 2, \ldots, 6$. Consider the basis elements E_1, E_2, \mathcal{F}_q , and $\mathcal{F}_{p_{\gamma}}$. Since $(f_X^*)^{\kappa_n}$ is the identity, it fixes these basis elements. Thus f^{κ_n} fixes the base points in \mathbf{P}^2 . Since f^{κ_n} is linear, it leaves invariant every line through two of these base points.

§6. Parameter Regions

There is a natural group action on parameter space. Namely, for $(\lambda, c, \mu) \in \mathbf{C}_* \times \mathbf{C}_* \times \mathbf{C}$ we have actions

$$(\alpha,\beta) \mapsto (\lambda\alpha,\lambda\beta) \tag{6.1}$$

$$(\alpha,\beta) \mapsto (\alpha_0, c\alpha_1, c\alpha_2, c\beta_0, c^2\beta_1, c^2\beta_2) \tag{6.2}$$

$$\begin{array}{l} (\alpha,\beta) \mapsto (\alpha_0 + \mu(\alpha_1 + \alpha_2) - \mu(\beta_0 + \mu(\beta_1 + \beta_2)), \\ \alpha_1 - \mu\beta_1, \alpha_2 - \mu\beta_2, \beta_0 + \mu(\beta_1 + \beta_2), \beta_1, \beta_2). \end{array}$$
(6.3)

The first action corresponds to the homogeneity of $f_{\alpha,\beta}$. The other two are given by linear conjugacies of $f_{\alpha,\beta}$. To see them, we write f in affine coordinates, as in (0.2). Action (6.2) is given by conjugating by the scaling map $(x_1, x_2) \mapsto (cx_1, cx_2)$, and (6.3) is given by conjugating by the translation $(x_1, x_2) \mapsto (x_1 + \mu, x_2 + \mu)$.

Now consider maps of the form

$$f:(x,y)\mapsto (y,\frac{y}{b+x+cy}), \quad b\neq 0.$$
(6.4)

In this case we have $\alpha = (0, 0, 1)$, $\beta = (b, 1, c)$ and $\gamma = (0, 0, 1)$. Let Y be as in (1.2), and let $f_Y: Y \to Y$ be the induced map. Repeating the computation of (1.3), we see that

$$\Sigma_{\beta} \mapsto E_2 \mapsto [c:0:1]_{e_1} \in E_1 \mapsto [c:1:0] \in \Sigma_{\gamma}.$$
(6.5)

We conclude that the sub-family (6.4) is critically finite the following sense that all exceptional curves have finite orbits:

Proposition 6.1. If f be as in (6.4), then q = (0,0) is a fixed point, and the exceptional curves are mapped to q. In particular, f_Y is 1-regular.

Proof. If c = 0, then exceptional locus is Σ_{γ} ; if $c \neq 0$, then both Σ_{β} and Σ_{γ} are exceptional. We see from (6.5) that in either case the exceptional curves are mapped to the fixed point.

The variety $V_n \subset \{\beta_2 = 0\}$ corresponds to a dynamical property: an exceptional line is mapped to a point of indeterminacy. Thus V_n is invariant under the actions (6.1–3). For $(\alpha, \beta) \in V_n$, we have $\beta_2 = 0$, and we may apply (6.3) to obtain $\alpha_1 = 0$. Since by (0.4) we must have $\alpha_2 \neq 0$ and $\beta_1 \neq 0$, we apply (6.1) and (6.2) to obtain $\alpha_2 = \beta_1 = 1$. Thus each orbit within V_n is represented by a map which may be written in affine coordinates as

$$(x,y) \mapsto (y, \frac{a+y}{b+x}). \tag{6.6}$$

If f is of the form (6.6), then f^{-1} is conjugate via the involution $\sigma : x \leftrightarrow y$ and a transformation (6.3) to the map

$$(x,y) \mapsto (y, \frac{a-b+y}{-b+x}). \tag{6.7}$$

Such a mapping is conjugate to its inverse if b = 0.

Now we suppose that f is given by (6.6). Thus q = (-a, 0) and p = (-b, -a), and V_n is defined by the condition $f^n q = p$. The coefficients of the equations defining V_n are positive integers, and V_n is preserved under complex conjugation. An inspection of the equations defining V_n produces the first few:

- V_0 : the orbit under (6.1–3) of (a, b) = (0, 0)
- V_1 : the orbit of (a, b) = (1, 0)
- V_2 : the orbits of (a, b) = ((1 + i)/2, i) and its conjugate.
- V_3 : the orbits of $(a,b) \in \{(2+i-\sqrt{3})/2,i), (2+i+\sqrt{3})/2,i)\}$ and their conjugates.

We solve for V_4 , V_5 and V_6 by using the resultant polynomials of the defining equations, and we find:

 V_4 : the orbits of $(a, b) = (0.8711 + 0.7309i, 1.4619i), (0.6974 + 0.2538i, 0.5077i), (-0.06857 + 0.3889i, 0.7778i), and their conjugates. The exact values are roots of <math>1 - 3 a + 9a^2 - 24a^3 + 36a^4 - 27a^5 + 9a^6$ and $1 + 6b^2 + 9b^4 + 3b^6$.

 V_5 : the orbits of $(a, b) = (3.7007 + 1.2024, 2.4048i), (1.0353 + 0.3364i, 0.6728i), (0.4465 + 0.6146i, 1.2293i), (-0.1826 + 0.2513i, 0.5027i), or their conjugates. The exact values are roots of <math>1 + 3a^2 - 20a^3 + 49a^4 - 60a^5 + 37a^6 - 10\ a^7 + a^8$ and $1 + 7b^2 + 14b^4 + 8b^6 + b^8$.

 V_6 : The defining equations for V_6 are divisible by b^2 , so all points of the form (a, 0), $a \neq 0, 1$, belong to V_6 . By (6.7), these parameters correspond to maps which are conjugate to their inverses. In addition, V_6 contains the orbits of

$$a = (3 \pm \sqrt{5} + 2b)/4, \quad b = i\sqrt{(5 \pm \sqrt{5})/2}$$

and their conjugates.

By Theorem 2, mappings in V_6 have quadratic degree growth, and by [G] such mappings have invariant fibrations by elliptic curves. Let us show how our approach yields these invariant fibrations.



Figure 6.1. Points $f^j q = j', 0 \le j \le 6$, for V_6 . Case b = 0 on left; $b \ne 0$ on right.

Let us first consider parameters (a, 0). In this case, the fibration was obtained classically in [L] and [KoL]. In the space Y of (1.2), the f-orbit $\{q_j = f^j q : j = 0, 1, \dots, 6\}$ is:

$$q_0 = (-a, 0)_{\mathbf{C}^2} = [1 : -a : 0], \ q_1 = (0, -1)_{\mathbf{C}^2} = [1 : 0 : -1], \ q_2 = [0 : 0 : 1] = e_2, q_3 = [0 : 1 : -1], \ q_4 = [1 : 0 : -1]_{e_1}, \ q_5 = (-1, 0), \ q_6 = (0, -a) = p,$$

as is shown in Figure 6.1. Here we use 'j' to denote 'q_j'. The construction of X is shown in Figure 6.2, where ' $f^{j}Q$ ' denotes the blowup fiber over q_{j} . In contrast, the case corresponding to $(a, b) \in V_{6}, b \neq 0$ corresponds to Figure 4.3. Consulting Figure 6.2, we see that the cohomology class $3H_{X} - E_{1} - E_{2} - Q_{2} - Q_{4} - \sum Q_{j}$ is fixed under f^{*} . We will find polynomials which correspond as closely as possible to this class. These will be cubics which vanish on e_{i} and q_{j} . Looking for lines that contain as many of the q_{j} as possible, we see $L_{1} = \{x + y + a = 0\}$ contains 0,3,6. Mapping forward by f, we have

$$L_1 \mapsto L_2 = \{y + 1 = 0\} \mapsto L_3 = \{x + 1 = 0\} \mapsto L_1.$$

In addition, the points q_j , j = 2, 3, 4 are contained in the line at infinity $M_1 = \Sigma_0$. This maps forward as:

$$M_1 \mapsto e_1 \mapsto M_2 = \{y = 0\} \mapsto M_3 = \{x = 0\} \mapsto e_2 \mapsto M_1.$$

The cubic $c_1 = (x + y + at)(x + t)(y + t)$ defines $L_1 + L_2 + L_3$ in \mathbf{P}^2 , and $c_2 = xyt$ defines $M_1 + M_2 + M_3$. Setting t = 1 and taking the quotient, we find the classical invariant $h(x, y) = c_1/c_2$.



Figure 6.2. Space X for V_6 , b = 0.

Now we consider the other four parameters (a, b) in V_6 . Inspecting the defining equations of V_6 , we find that a and b satisfy $-2a + a^2 + b - ab = 0$ and $-b^2 - 1 + b - 2a = 0$. Using these relations, we see that the *f*-orbit of *q* is:

$$q_0 = (-a, 0), \ q_1 = (0, 1-a), \ q_2 = (1-a, 1/b), \ q_3 = (1/b, a(1+ab)/(ab-b^2)), \ q_4 = (a(1+ab)/(ab-b^2), 1-a), \ q_5 = (1-a, -b), \ q_6 = (-b, -a).$$

Looking again at the points q_j , j = 0, 3, 6, we see that they are contained in a line $L_1 = \{x + (1 - \frac{b}{a})y + a = 0\}$. Mapping L_1 forward under f, we find:

$$L_1 \mapsto L_2 = \{y + a - 1 = 0\} \mapsto L_3 = \{x + a - 1 = 0\} \mapsto L_1$$

We multiply these linear functions together to obtain a cubic c_1 which defines $\sum L_i$. We see, too, that the points q_j , j = 1, 3, 5 are contained in the line $M_1 = \{(a-b-1)x + (a-1)y + (a-1)^2 = 0\}$. Mapping forward, we find:

$$M_1 \mapsto M_2 = \{(a-1)xy + (b^2+1)y + (a-b-1)x + (a-b) = 0\} \mapsto M_1.$$

Multiplying the defining functions, we obtain a cubic c_2 which defines $M_1 + M_2$. Now we define $k(x, y) = c_1/c_2$. And inspection shows that $k \circ f = \omega k$, where ω is a 5th root of unity. Thus f is a period 5 mapping of the set of cubics $\{k = \text{const}\}$ to itself.

Appendix: Explanation of the Computer Graphics

It is useful to have visual representations for rational mappings. A number of interesting computer graphic representations of the behavior of rational mappings of the real plane have been given in various works by Bischi, Gardini and Mira; we cite [BGM] as an example. The pictures here have a somewhat different origin and are made following a scheme used earlier by one of the authors and Jeff Diller (see [BD1,3]). They are motivated by the theory of dynamics of complex surface maps. Let f be a birational map of a Kähler surface. If $\delta(f) > 1$, then there are positive, closed, (1,1)-currents T^{\pm} such that $f^*T^+ = \delta(f)T^+$ and $f^*T^- = \delta(f)^{-1}T^-$ (see Diller-Favre [DF]). These currents have the additional property that for any complex curve Γ there is a number c > 0 such that

$$cT^{+} = \lim_{n \to \infty} \frac{1}{\delta^{n}} f^{n*}[\Gamma], \qquad (A.1)$$

and similarly for T^- . By work of Dujardin [D1] these currents have the structure of a generalized lamination. We let $\mathcal{L}^{s/u}$ denote the generalized laminations corresponding to T^{\pm} . It was shown in [BD2] that the wedge product $T^+ \wedge T^-$ defines an invariant measure in many cases, and Dujardin [D2] showed that this invariant measure may be found by taking the "geometric intersection" of the measured laminations \mathcal{L}^s and \mathcal{L}^u .

When one of our mappings f has real coefficients, it defines a birational map of the real plane, and we can hope that there might be real analogues for the results of the theory of complex surfaces. This was proved to be the case for certain maps in [BD1,3] but is not known to hold for the maps studied in the present paper.

Figure 0.1 was drawn as follows. We work in the affine coordinate chart (x, y) on \mathbb{R}^2 given by $x_0 = 1$, $x = x_1/x_0 = x_1$, $y = x_2/x_0 = x_2$. We start with a long segment $L \subset \mathbb{R}^2$ and map it forward several times. The resulting curve is colored black and "represents" \mathcal{L}^u . After the first few iterates, the computer picture seems to "stabilize," and further iteration serves to "fill out" the lamination. The appearance of the computer picture obtained in this manner is independent of the choice of initial line L. To represent \mathcal{L}^s , we repeat this procedure for f^{-1} and color the resulting picture gray. In Figure 0.1 we present \mathcal{L}^s in gray in the left hand frame. Then we present \mathcal{L}^s and \mathcal{L}^u together in the right hand frame in order to show the set where they intersect.



Figure A.1. Explanation of Figure 0.1 (left); a mapping from V_7 (right).

We also want the graphic to have the appearance of a subset of \mathbf{P}^2 , so we rescale the distance to the origin. The resulting "disk" is a compactification of \mathbf{R}^2 . In fact, this is real projective space, since antipodal points of the circle are identified. The circle forming the boundary of this disk is the line at infinity Σ_0 .

Figure 0.1 is obtained using the map of the form (6.4):

$$(x,y)\mapsto (y,\frac{y}{.1+x+.3y}).$$

By Proposition 6.1, f is critically finite, so $\delta(f) = \phi$ by Theorem 4.1. On the left half of Figure A.1, we have re-drawn \mathcal{L}^s , together with the points of indeterminacy of f and f^{-1} . Pictured, for instance, are $e_1, e_2, p_0 = [0: -.3: 1], p_{\gamma} = (-.1, 0)$, and q = (0, 0). The exceptional curves are lines connecting certain pairs of these points and may be found easily using Figure 1.1 as a guide. As we expect, \mathcal{L}^s is "bunched" at the points of indeterminacy of f, i.e., p_0, e_1 , and p_{γ} . Let us track the backward orbits of these points. First, $p_0 = f^{-1}p_0$ is fixed under f^{-1} , and $f^{-1}p_{\gamma} = e_1$.

Now let Y and f_Y be as in (1.2). Repeating the calculations at equation (1.3), we see that f_Y^{-1} takes p_γ to the fiber point $[1:0:-..1]_{E_1}$ over e_1 . Then this fiber point is mapped under f^{-1} to the point $s = [0:1.03:-..1] \in \Sigma_0$. The next preimage is $f^{-1}s = p_0$, so f^{-1} is critically finite in the sense that the exceptional curves all have finite orbits. This explains why \mathcal{L}^s is "bunched" at only four points.

To explain the points where \mathcal{L}^u is "bunched," we have plotted the point $r := f^3 \Sigma_\beta = (10/3, 0)$ from (6.5). If we superimpose the picture of \mathcal{L}^u on the left panel of Figure A.1, we find that \mathcal{L}^u is "bunched" exactly on the set e_1, e_2, q , and r. The "eye" which appears in the first quadrant is due to an attracting fixed point.

The right hand side of Figure A.1 is obtained using the map

$$(x,y) \mapsto (y, \frac{-.499497 + y}{-.415761 + x}),$$

which corresponds to a real parameter $(a, b) \in V_7$. By "j", j = 0, ..., 7, we denote the point $f^j q$. Thus "7" is the point of indeterminacy $p = f^7 q$. We let $\pi : X \to \mathbf{P}^2$ be the manifold obtained by blowing up e_1, e_2 , and "j" for j = 0, ..., 7. The lamina of \mathcal{L}^u are then separated in X, and the apparent intersections may be viewed as artifacts of the projection π .

References

- [BD1] E. Bedford and J. Diller, Real and complex dynamics of a family of birational maps of the plane: the golden mean subshift, American J. of Math., to appear.
- [BD2] E. Bedford and J. Diller, Energy and invariant measures for birational surface maps, Duke Math. J., to appear.
- [BD3] E. Bedford and J. Diller, Dynamics of a two parameter family of plane birational mappings: Maximal entropy. arxiv.org/math.DS/0505062
- [BK] E. Bedford and KH. Kim, On the degree growth of birational mappings in higher dimension, J. Geom. Anal. 14 (2004), 567-596.
- [BDGPS] M.J. Bertin, A. Decomps-Guilloux, M. Grandet-Hugot, M. Pathiaux-Delefosse, and J.P. Schreiber, *Pisot and Salem Numbers*, Birkhäuser Verlag (1992)
 - [BGM] G-I. Bischi, L. Gardini, and C. Mira, Plane maps with denominator. I. Some generic properties, International J. of Bifurcation and Chaos, 9 (1999), 119–153.
 - [C] S. Cantat, Dynamique des automorphismes des surfaces projectives complexes, C.R. Acad. Sci. Paris, t. 328, p. 901–906, 1999.
 - [CL] M. Csörnyei and M. Laczkovich, Some periodic and non-periodic recursions, Monatshefte für Mathematik 132 (2001), 215-236.
 - [DF] J. Diller and C. Favre, Dynamics of bimeromorphic maps of surfaces, Amer. J. of Math., 123 (2001), 1135–1169.
 - [D1] R. Dujardin, Laminar currents in \mathbf{P}^2 , Math. Ann., 325, 2003, 745–765.
 - [D2] R. Dujardin, Laminar currents and birational dynamics. arxiv.org/math.DS/0409557
 - [FS] J-E Fornæss and N. Sibony, Complex dynamics in higher dimension, II. Modern Methods in Complex Analysis, Ann. of Math. Studies, vol. 137, Princeton Univ. Press, 1995, pp. 135–182.
 - [GBM] L. Gardini, G.I. Bischi, and C. Mira, Invariant curves and focal points in a Lyness iterative process, Int. J. Bifurcation and Chaos 13 (2003), 1841-1852.
 - [G] M. Gizatullin, Rational G-surfaces. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 44 (1980), no. 1, 110–144, 239.
 - [GKP] R.L. Graham, D.E. Knuth, and O. Patashnik, *Concrete Mathematics*, 1989.

- [GL] E.A. Grove and G. Ladas, *Periodicities in Nonlinear Difference Equations*, Kluwer Academic Publishers, 2005.
- [KoL] V.I. Kocic and G. Ladas, Global Behaviour of Nonlinear Difference Equations of Higher Order with Applications, Kluwer Academic Publishers 1993.
- [KLR] V.I. Kocic, G. Ladas, and I.W. Rodrigues, On rational recursive sequences, J. Math. Anal. Appl 173 (1993), 127-157.
- [KuL] M. Kulenovic and G. Ladas, Dynamics of Second Order Rational Difference Equations, CRC Press, 2002.
- [KG] R.P. Kurshan and B. Gopinath, Recursively generated periodic sequences, Canad. J. Math. 26 (1974), 1356–1371.
 - [L] R.C. Lyness, Notes 1581,1847, and 2952, Math. Gazette 26 (1942), 62, 29 (1945), 231, and 45 (1961), 201.

Indiana University Bloomington, IN 47405 bedford@indiana.edu

> Syracuse University Syracuse, NY 13244 kkim26@syr.edu