### A Quantum Deformation of Invariants of Higher Binary Forms

Frank Leitenberger

### FRANK LEITENBERGER

Fachbereich Mathematik, Universität Rostock, Rostock, D-18051, Germany. e-mail: frank.leitenberger@mathematik.uni-rostock.de

#### Abstract

We use the theory of the quantum group  $U_q(gl(2, \mathbb{R}))$  in order to develop a quantum theory of invariants and show a decomposition of invariants into a Gordan-Capelli series. Higher binary forms are introduced on the basis of braided algebras. We define quantised invariants and give basic examples. We show that the symbolic method of Clebsch and Gordan works also in the quantised case. We discuss the deformed discriminant of the quadratic and the cubic form, the deformed invariants  $I_1$ ,  $I_2$  of the quartic form and further invariants without a classical analog.

## 1. INTRODUCTION

In this article we develop a theory of a quantum deformation of the invariant theory of higher binary forms. The invariant theory was developed by Cayley, Sylvester, Clebsch, Gordan, Capelli etc. (cf. [1, 2, 3]). The invariant theory is one of the historical roots of the modern representation theory, which has a well developed quantum deformation. For the relation between both theories we refer to [4, 5].

To find quantum deformations of some basic notions of algebraic geometry (e.g. curves of higher genus) is an open problem. Plane curves with genus  $g \ge 1$  can be described by homogeneous ternary forms of degree  $k \ge 3$ . These ternary forms are related to binary forms by the "Übertragungsprinzip", cf. [6]. This is a motivation to consider a quantum deformation of an invariant theory of the higher binary forms.

We will demonstrate that the basic concepts of the theory of the quantum group  $U_q(sl(2, \mathbb{R}))$  can be used, to build a quantum deformation of the invariant theory of the higher binary forms. Our approach is based on replacing the commutative algebra of classical homogeneous coordinates  $x_i, y_i$  of n points of the real line by a braided algebra. If the quantum parameter q is not a root of unity, the noncommutative subalgebra of invariant elements is generated by a deformation of the bracket symbols (ij) of Clebsch. We show that there is a decomposition of invariants into Gordan-Capelli series.

Using the braided algebra we define the notions of n-forms and invariants of n-forms. The n-forms are related to quantised polynomials. We define quantum analogs of symmetric functions and power sums and demonstrate Newton relations.

It turns out that the Clebsch-Gordan symbolic method also works in our situation. For this purpose we introduce the notions of universal forms and universal invariants, and we consider quantum n-forms and invariants as realisations of these forms.

The Clebsch-Gordan method works even in the case when q is a root of unity. But in this case we cannot represent all invariant elements of the algebra of noncommutative coordinates by symbols (ij). We find that certain coefficients play the role of invariants.

By computer calculations we derive the simplest invariants of linear, quadratic, cubic and quartic forms. It turns out that the ring of invariants has a richer structure than in the classical case. For example, we obtain a nonvanishing invariant for the linear form, an invariant of degree three for the quadratic form and an invariant of degree two for the cubic form, which have no analogs in the classical case.

Furthermore we consider deformations of discriminants. For the quadratic form the situation is similar to the classical case. For the cubic form we obtain a four-dimensional space of invariants with properties of the discriminant. One of them plays a special role. For the quartic form we encounter an obstruction for a representation of the discriminant by the basic invariants of degree two and three.

We used the computer algebra programs Mathematica 3.0 and FELIX forming the computations.

# 2. PRELIMINARIES

In this section we introduce some basic concepts about the quantum group  $U_q(gl(2, \mathbb{R}))$ and the braided module algebras  $H_I$ .

We will use the q-numbers

$$[i]_{q^2} = [i] := q^{i-1} + q^{i-3} + \ldots + q^{-i+1} = \frac{q^i - q^{-i}}{q - q^{-1}}$$

and

$$(i)_{q^2} = (i) := 1 + q^2 + q^4 + \dots + q^{2i-2} = \frac{q^{2i} - 1}{q^2 - 1}$$

for  $i \in \mathbb{Z}$ , i.e.  $(i) = q^{i-1}[i]$ .

Let  $U_q(gl(2,\mathbb{R})), |q| = 1, q \neq \pm 1, \pm i$  be the unital \*-algebra, determined by generators  $E, F, K, L, K^{-1}, L^{-1}$ , where  $L, L^{-1}$  are central and relations

$$LL^{-1} = L^{-1}L = 1, \quad KK^{-1} = K^{-1}K = 1,$$
  
 $KE = qEK, \quad KF = q^{-1}FK,$   
 $[E, F] = \frac{K^2 - K^{-2}}{q - q^{-1}}.$ 

We endow  $U_q(gl(2, \mathbb{R}))$  with the involution

$$E^* = F, \qquad F^* = E, \qquad K^* = K^{-1}, \qquad L^* = L^{-1}.$$

For an arbitrary ordered index set I we consider the unital  $U_q(gl(2, \mathbb{R}))$ -module algebra  $A_I$ , which is freely generated by the variables  $x_i, y_i, i \in I$ . To determine the action of  $U_q(gl(2, \mathbb{R}))$  on  $A_I$ , we set

$$L1 = 1, Lx_i = q^{\frac{1}{2}}x_i, Ly_i = q^{\frac{1}{2}}y_i, K1 = 1, Kx_i = q^{-\frac{1}{2}}x_i, Ky_i = q^{\frac{1}{2}}y_i, E1 = 0, Ex_i = q^{\frac{1}{2}}y_i, Ey_i = 0, F1 = 0, Fx_i = 0, Fy_i = q^{-\frac{1}{2}}x_i$$
(1)

and require that

$$\begin{aligned}
K(ab) &= K(a)K(b), \\
E(ab) &= E(a)K(b) + K^{-1}(a)E(b), \\
F(ab) &= F(a)K(b) + K^{-1}(a)F(b)
\end{aligned}$$
(2)

for  $a, b \in A_I$ . A proof can be given along the lines of [8], p. 19, where a similar module occurs.

We note that

$$E(x_i^m y_i^n) = q^{\frac{2-m+n}{2}} [m] x_i^{m-1} y_i^{n+1},$$
  

$$F(x_i^m y_i^n) = q^{\frac{m-n}{2}} [n] x_i^{m+1} y_i^{n-1}.$$
(3)

Furthermore we consider the ideal J of  $A_I$  which is generated by the elements

$$x_{i}y_{i} - qy_{i}x_{i}, \qquad x_{j}x_{i} - q^{2}x_{i}x_{j}, \qquad y_{j}y_{i} - q^{2}y_{i}y_{j},$$
$$x_{j}y_{i} - qy_{i}x_{j} - (q^{2} - 1)x_{i}y_{j}, \qquad y_{j}x_{i} - qx_{i}y_{j}$$
(4)

for i < j.

One can show that the ideal J is  $U_q(gl(2, \mathbb{R}))$ -invariant (i.e. E(J) = F(J) = 0,  $K(J) \subseteq J$  and  $L(J) \subseteq J$ ) (cf. [9]).

Therefore the action of  $U_q(gl(2, \mathbb{R}))$  on  $A_I$  induces an action on  $H_I := A_I/J$ . In the following we identify  $x_i$  with its image under the quotient map  $A_I \to H_I$ .

I.e.  $H_I$  is the unital  $U_q(gl(2, \mathbb{R}))$ -module algebra with generators  $x_i, y_i \ i \in I$  and relations

$$\begin{aligned}
x_i y_i &= q y_i x_i, \\
x_j x_i &= q^2 x_i x_j, \\
y_j y_i &= q^2 y_i y_j, \\
x_j y_i &= q y_i x_j + (q^2 - 1) x_i y_j, \\
y_j x_i &= q x_i y_j
\end{aligned} (5)$$

for i < j. Since the relations (5) are \*-invariant for the involution  $x_i^* = x_i, y_i^* = y_i$  we consider  $H_I$  as a \*-module algebra. If  $I = \{1, 2, ..., n\}$  we denote the algebra by  $H_n$ . From (5) we derive the relations

From (5) we derive the relations

$$\begin{aligned} x_j y_i^k &= q^k y_i^k x_j + (q^{2k} - 1) x_i y_i^{k-1} y_j, \\ x_j^k y_i &= q^k y_i x_j^k + (q^{2k} - 1) x_i x_j^{k-1} y_j. \end{aligned}$$
(6)

We say that that an element a of the module  $H_I$  is homogeneous of degree k if  $La = q^{\frac{k}{2}}a$ .

Remark 1.1  $H_I$  is a braided module algebra (i.e.  $(x_i + x_j)(y_i + y_j) = q(y_i + y_j)(x_i + x_j)$ for  $i \neq j$ ), (cf. [9]). Furthermore the algebra  $H_I$  has the PBW property for an arbitrary order of the variables  $x_i, y_i, i \in I$ . We note that  $H_I$  is up to reordering the points the only module algebra with quadratic relations which is braided and has the PBW property.

Remark 1.2 We can consider the Eqs. (5) as rules in order to express an element  $a \in H_I$  by the PBW basis. These rules do not change the type of homogenity with respect to the indices  $i \in I$ .

At the end of this section we extend  $H_I$  to a division algebra  $Q_I$ . Let R be an algebra,  $\sigma$  an endomorphism of R and  $\delta$  a  $\sigma$ -derivation with

$$\delta(ab) = \delta(a)b + \sigma(a)\delta(b).$$

We say that  $S = R[x; \sigma, \delta]$  is an *Ore extension of* R if S is freely generated over R by an element x subject only to the relation

$$xa = \sigma(a)x + \delta(a)$$

for  $a \in R$  (cf. [10]).

LEMMA 1.1 The algebra  $H_I$  is an iterated Ore extension.

*Proof.* We restrict the consideration to the finite algebras  $H_n$ . In the case of an infinite index set I one can proceed by transfinite induction. We consider the tower of subalgebras

$$H_0 = \mathbb{C} \subset H'_1 \subset H_1 \subset \cdots \subset H'_n \subset H_n$$

where  $H'_i$  is the subalgebra generated by  $H_{i-1}$  and  $y_i$ .

We have  $H'_i \cong H_{i-1}[y_{n+1}, \sigma, \delta]$  where  $\delta = 0$  and  $\sigma$  is determined by

$$\sigma(x_i) = qx_i, \qquad \sigma(y_i) = q^2 y_i.$$

Furthermore we have  $H_i \cong H'_i[x_{n+1}, \sigma, \delta]$  where  $\sigma$  and  $\delta$  are determined by

$$\sigma(x_i) = q^2 x_i, \qquad \sigma(y_i) = q y_i, \qquad \sigma(y_{n+1}) = q y_{n+1},$$
  
$$\delta(x_i) = 0, \qquad \delta(y_i) = (q^2 - 1) x_i y_{n+1}, \qquad \delta(y_{n+1}) = 0$$

The verification of the isomorphies is similar to [11], p. 81. The assertion follows.

PROPOSITION 1.2 The algebra  $H_I$  has no zero divisors and  $H_I$  has an uniquely determined right quotient division ring  $Q_I$ , whose elements are the right quotients  $ab^{-1}$  for  $a, b \in R, b \neq 0$ .

*Proof.* Ore proved that a noncommutative ring without zero divisors has an uniquely determined quotient division ring if

$$aR \cap bR \neq \{0\} \qquad \forall a, b \in R, \qquad a, b \neq 0, \tag{7}$$

i.e., two nonvanishing elements a, b have a common right multiple (cf. [12]). Curtis proved that an iterated Ore extension of a ring without zero divisors with the property (7) is again a ring without zero divisors with the property (7) (cf. [13]). The assertion follows from both results together with Lemma 2.1.  $\bullet$ 

Example 1.1 For  $x_2$  and  $y_1$  we find the common right multiple

$$x_2(x_2y_1 - (q^2 - q^{-2})x_1y_2) = y_1(q^2x_2^2).$$

Therefore we can represent the element  $y_1^{-1}x_2$  by the right quotient  $q^2x_2^2(x_2y_1 - (q^2 - q^{-2})x_1y_2)^{-1}$ . Note, that in general it is not possible to find a common right multiple of a and b of degree  $\leq deg(a) + deg(b)$ .

Remark 1.3 We can interpret the variables  $x_i, y_i$  as noncommutative homogeneous coordinates of different points of the projective real line (cf. [14]). The noncommutative real coordinate is given by

$$v_i := q^{\frac{1}{2}} x_i y_i^{-1} \in Q_I.$$

We have  $v_i^* = v_i$  and

$$v_j v_i = q^2 v_i v_j + (1 - q^2) v_i^2$$
 for  $i < j$ .

For q = -1 we obtain a noncommutative algebra of homogeneous coordinates for the classical commutative algebra of real coordinates.

# 3. THE INVARIANTS OF THE ALGEBRA $H_I$

DEFINITION 3.1 We say that the element a of  $H_I$  is an invariant element of degree k, if Ea = Fa = 0, Ka = a and  $La = q^{\frac{k}{2}}a$ .

By the rules (2) sums and products of invariant elements are again invariant elements. Therefore they form a subalgebra  $H_I^{Inv} \subset H_I$ .

Example 3.1 The simplest invariant element of  $H_I$  is given by

$$(ij) := q^{-\frac{1}{2}} x_i y_j - q^{\frac{1}{2}} y_i x_j, \qquad i, j \in I.$$

We have

$$(ji) = -(ij), \quad \forall i, j,$$
  
 $(ij)^* = (ij), \quad \forall i, j$ 

and

 $(ii) = 0, \forall i.$ 

The condition (ji) = -(ij) is equivalent to the braiding condition  $(x_i + x_j)(y_i + y_j) = q(y_i + y_j)(x_i + x_j)$  for  $i \neq j$ , cf. Remark 2.1.

LEMMA 3.1 Let i < j < k be three ordered indices. Then the following commutation relations hold:

PROPOSITION 3.2 (i) Let i < j < k < l be four ordered indices. Then the following commutation relations hold:

$$\begin{aligned} (kl)(ij) &= q^{6}(ij)(kl), \\ (jk)(il) &= (il)(jk), \\ (jl)(ik) &= q^{4}(ik)(jl) + (q^{4} - q^{6})(ij)(kl), \\ (jk)(ij) &= q^{4}(ij)(jk), \\ (ik)(ij) &= q^{2}(ij)(ik), \\ (jk)(ik) &= q^{2}(ik)(jk). \end{aligned}$$

$$(9)$$

(ii) Let i, j, k, l be four different indices. Then the three invariant elements

commute.

(iii) Let i < j < k < l be four ordered indices. Then we have the identities

$$q^{4}(ij)(kl) + q^{2}(ik)(lj) + (il)(jk) = 0$$
(10)

(Grassmann-Plücker relations).

A way to check the identities of Lemma 3.1 and Proposition 2.2 is indicated in [14], p. 816.

Remark 3.1 We can rewrite Proposition 3.2 (ii), (iii) in a more symmetric formulation. Let

$$Z_1 := q^3(ij)(kl),$$
  

$$Z_2 := q^2(ik)(lj) + (q^4 - q^3)(ij)(kl),$$
  

$$Z_3 := (il)(jk).$$

According to Proposition 3.2 the elements  $Z_1$ ,  $Z_2$ ,  $Z_3$  commute, are \*-invariant and we have

$$Z_1 + Z_2 + Z_3 = 0.$$

Remark 3.2 If we interprete  $x_i, y_i$  as noncommutative projective coordinates, then the six elements  $-Z_1Z_2^{-1}, -Z_1Z_3^{-1}, -Z_2Z_1^{-1}, -Z_2Z_3^{-1}, -Z_3Z_1^{-1}, -Z_3Z_2^{-1} \in Q_I$  get the meaning of the six possible cross ratios of the four zeros  $v_i := q^{\frac{1}{2}}x_iy_i^{-1}$  of the form f. For example, we have

$$-Z_1 Z_3^{-1} = \frac{1}{q} (v_i - v_j) (v_j - v_k)^{-1} (v_k - v_l) (v_i - v_l)^{-1}$$

(cf. [14]).

LEMMA 3.3 Let i < j < k. The following formulas hold: (i)  $(ij)(ik) = x_i^2 y_j y_k - x_i y_i (qx_j y_k + y_j x_k) + q^2 y_i^2 x_j x_k,$ 

$$(ik)(jk) = x_i x_j y_k^2 - (q x_i y_j + y_i x_j) x_k y_k + q^2 y_i y_j x_k^2$$

$$(ij)(jk) = \frac{1}{q^2} x_i x_j y_j y_k - (x_i y_j y_j x_k + y_i x_j x_j y_k) + q y_i x_j y_j x_k,$$

$$(ij)(ij) = x_i^2 y_j^2 - [2] x_i y_i x_j y_j + q^2 y_i^2 x_j^2$$
$$= \frac{1}{q^2} x_j^2 y_i^2 - \frac{[2]}{q^2} x_j y_j x_i y_i + y_j^2 x_i^2,$$

(ii)

$$(ij)^{n} = \sum_{k=0}^{n} \begin{bmatrix} n\\k \end{bmatrix} (-1)^{k} q^{\frac{1}{2}n^{2} + (k-n)(1+k)} x_{i}^{n-k} y_{i}^{k} x_{j}^{k} y_{j}^{n-k},$$

(iii)

$$(ij)(ik)(jk) = q^{-\frac{1}{2}}x_ix_jy_jy_ky_k - q^{\frac{3}{2}}x_ix_jy_jy_jx_ky_k - q^{\frac{3}{2}}x_iy_ix_jx_jy_ky_k - q^{-\frac{3}{2}}x_iy_ix_jy_jx_ky_k + q^{\frac{5}{2}}x_iy_ix_jy_jx_ky_k + q^{\frac{3}{2}}x_iy_iy_jy_jx_kx_k + q^{\frac{3}{2}}y_iy_ix_jx_jx_ky_k - q^{\frac{7}{2}}y_iy_ix_jy_jx_kx_k.$$

(iv) Let 
$$u_i^{(0)} := x_i$$
 and  $u_i^{(1)} := y_i$ . We have  
 $(1, 2n)(2, 2n - 1)(3, 2n - 2) \cdots (n, n + 1)$   
 $= \sum_{i_1, i_2, \cdots, i_n = 0}^{1} q^{\frac{3}{2}n(n-1)}(-q)^{i_1 + i_2 + \cdots + i_n} u_1^{(i_1)} u_2^{(i_2)} \cdots u_n^{(i_n)} u_{n+1}^{(1-i_n)} u_{n+2}^{(1-i_n-1)} \cdots u_{2n}^{(1-i_1)}.$ 

Here

$$\left[\begin{array}{c}n\\i\end{array}\right] := \frac{[n]!}{[n-i]![i]!}$$

denotes the q-binomial coefficient, where

$$[n]! := [1][2]...[n],$$
  
 $i, n \in \mathbb{N}.$  (If  $q^{2d} = 1$  for all  $d \in \mathbb{N}$ , but  $q^2 \neq 1$ , we define  $\begin{bmatrix} n\\ i \end{bmatrix}$  by a limit.)

A proof of Lemma 3.3 can be given by an explicite calculation or by induction.

### 3.1 Polarisation operators

We introduce the polarisation operators  $\Delta_{kl} : H_I \to H_I, k \neq l$ . We fix a PBW basis, which consists of the elements

$$x_{a_1}^{i_1}y_{a_1}^{j_1}x_{a_2}^{i_2}y_{a_2}^{j_2}...x_{a_k}^{i_k}y_{a_k}^{j_k}$$

with  $a_1 < a_2 < \dots < a_k$ . We set

$$\Delta_{kl}(x_k) = x_l, \qquad \Delta_{kl}(y_k) = y_l$$

and

$$\Delta_{kl}(x_i) = 0, \qquad \Delta_{kl}(y_i) = 0$$

for  $i \neq k$  and require

$$\Delta_{kl}(ab) = \Delta_{kl}(a) \ b + a \ \Delta_{kl}(b),$$

if the expressions a, b and ab are elements of the PBW basis.

If f is a polynomial, which is homogeneous with respect to  $k \in I$  of degree  $n_k$  we have the *Euler identity* 

$$\Delta_{kk}f = n_k f.$$

Remark 3.3 One obtains different linear operators  $\Delta_{kl}$  for different PBW bases. For our purposes it is sufficient to work with the above fixed basis.

Furthermore, if  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ , we define a linear operator  $P_{kl}$  by

$$P_{kl}a := \frac{1}{(n_k)_{q^2}} \Delta_{kl}a,$$

if a is a monomial of  $H_I$ , which is homogeneous with respect to the index k of degree  $n_k$ .

PROPOSITION 3.4 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ . (i) We have  $P_{ij}(ik) = (jk)$ , i.e.  $P_{ij}(ij) = 0$ . (ii) For k < l and  $\beta \in H_{\{i \in I \mid i > l\}}$  we have

$$P_{kl}\left(x_k^{m-i}y_k^iy_l^{n-j}x_l^j\beta\right) =$$

$$=\frac{1}{[m]}([m-i]x_k^{m-i-1}y_k^ix_l^{n-j+1}y_l^j\beta+q^{m-i-n+j}[i]x_k^{m-i}y_k^{i-1}x_l^{n-j}y_l^{j+1}\beta)$$

and

$$P_{lk}\left(x_k^{m-i}y_k^i x_l^{n-j}y_l^j\beta\right) =$$
  
=  $\frac{1}{[n]}(q^{j-i}[n-j]x_k^{m-i+1}y_k^i x_l^{n-j-1}y_l^j\beta + [j]x_k^{m-i}y_k^{i+1}x_l^{n-j}y_l^{j-1}\beta).$ 

(iii)  $P_{kl}$  commutes with the action of  $U_q(gl(2, \mathbb{R}), i.e., if f \in H_I^{Inv}, then P_{kl}(f) \in H_I^{Inv}$ .

*Proof.* (i) For i < k we have

$$P_{ij}(ik) = \Delta_{ij}q^{-\frac{1}{2}}(x_iy_k - qy_ix_k) = q^{-\frac{1}{2}}(x_jy_k - qy_jx_k) = (jk).$$

The case i > k is analogous.

(ii) From the definition of  $\Delta_{kl}$  we obtain  $\Delta_{kl}(x_k^m) = (m)x_k^{m-1}x_l$  and  $\Delta_{kl}(y_k^m) = (m)y_k^{m-1}y_l$ . It follows

$$\Delta_{kl}(x_k^{m-i}y_k^i x_l^{n-j}y_l^j) = (m-i)x_k^{m-i-1}x_l y_k^i x_l^{n-j}y_l^j + (i)x_k^{m-i}y_k^{i-1}y_l x_l^{n-j}y_l^j.$$

Using (5) and (6) we find

$$= (m-i)q^{i}x_{k}^{m-i-1}y_{k}^{i}x_{l}^{n-j+1}y_{l}^{j} + (i)q^{2m-2i-n+j}x_{k}^{m-i}y_{k}^{i-1}x_{l}^{n-j}y_{l}^{j+1}.$$

Since  $(m) = q^{m-1}[m]$  we have

$$= [m-i]q^{m-1}x_k^{m-i-1}y_k^i x_l^{n-j+1}y_l^j + [i]q^{2m-i-n+j-1}x_k^{m-i}y_k^{i-1}x_l^{n-j}y_l^{j+1}$$

The formula for  $P_{kl}$  follows. The proof for  $P_{lk}$  is analogous.

(iii) By the definition of K, L and  $\Delta_{kl}$  we have  $K\Delta_{kl}f = \Delta_{kl}Kf$  and  $L\Delta_{kl}f = \Delta_{kl}Lf$ .

From (ii) and (3) it follows that

$$\Delta_{kl} E(x_k^m y_k^n) = q^{\frac{2-m+n}{2}} [m] \Delta_{kl}(x_k^{m-1} y_k^{n+1})$$
$$= q^{\frac{2-m+n}{2}} [m] ([m-1] x_k^{m-2} y_k^{n+1} x_l + q^{m-1} [n+1] x_k^{m-1} y_k^n y_l)$$

and

$$E\Delta_{kl}(x_k^m y_k^n) = [m]E(x_k^{m-1} y_k^n)K(x_l) + [m]K^{-1}(x_k^{m-1} y_k^n)E(x_l) + q^m[n]E(x_k^m y_k^{n-1})K(y_l)$$

$$=q^{\frac{2-m+n}{2}}[m][m-1]x_k^{m-2}y_k^{n+1}x_l + q^{\frac{m-n}{2}}[m]x_k^{m-1}y_k^ny_l + q^{\frac{2+m+n}{2}}[n][m]x_k^{m-1}y_k^ny_l$$
$$=q^{\frac{2-m+n}{2}}[m]([m-1]x_k^{m-2}y_k^{n+1}x_l + q^{m-1}[n+1]x_k^{m-1}y_k^ny_l)$$

(since  $q^{-n}[1] + q[n] = [n+1]$ ). Therefore we have  $E\Delta_{kl}f_k = \Delta_{kl}Ef_k$  for  $f_k = x_k^m y_k^n$ .

Now let f be a PBW basis element. We consider the decomposition  $f = f_- f_k f_+$  with PBW basis elements  $f_+ \in H_{\{i \in I | i < k\}}$ ,  $f_- \in H_{\{i \in I | i > k\}}$ . We have

$$E\Delta_{kl}f = E\Delta_{kl}(f_-f_kf_+) = E(f_-\Delta_{kl}(f_k)f_+)$$

$$= E(f_{-})(\Delta_{kl}(K(f_{k})))K(f_{+}) + K^{-1}(f_{-})(\Delta_{kl}(E(f_{k})))K(f_{+}) + K^{-1}(f_{-})(\Delta_{kl}(K^{-1}(f_{k})))E(f_{+})$$
  
=  $\Delta_{kl}E(f_{-}f_{k}f_{+}) = \Delta_{kl}Ef.$ 

Similarly we find  $F\Delta_{kl}f = \Delta_{kl}Ff$  for all elements f of the PBW basis. I.e., E, F, K commute with  $\Delta_{kl}$  and therefore with  $P_{kl}$ . The proposition follows.•

PROPOSITION 3.5 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$  and let  $f \in H_I$  be a linear combination of monomials

$$x_k^{m-i} y_k^i x_l^{n-j} y_l^j \beta_{ij}$$

with  $k < l, m, n \ge 1$  and  $\beta_{ij} \in H_{\{i \in I | i > l\}}$ . (i) Then we can represent f in the unique form

$$f = P_{lk}P_{kl}f + (kl)a$$

$$f = P_{kl}P_{lk}f + (kl)b$$

with  $a, b \in H_I$  and  $b = \frac{[m][n+1]}{[m+1][n]}a$ . (ii) We have [m][m+1]R

$$[m][n+1]P_{lk}P_{kl}f - [n][m+1]P_{kl}P_{lk}f = [m-n]f$$

(iii) If  $f \in H_I^{Inv}$ , then  $a \in H_I^{Inv}$ .

*Proof.* (i) It is sufficient to consider the case, where f is the monomial  $x_k^{m-i}y_k^i x_l^{n-j}y_l^j$ . With Proposition 3.4 (ii) we find

$$P_{lk}P_{kl}f = \frac{1}{[m]}P_{lk}([m-i]x_k^{m-i-1}y_k^ix_l^{n-j+1}y_l^j + q^{m-i-n+j}[i]x_k^{m-i}y_k^{i-1}x_l^{n-j}y_l^{j+1})$$
  
$$= \frac{1}{[m][n+1]}([m-i][n-j+1]q^{j-i}x_k^{m-i}y_k^ix_l^{n-j}y_l^j + [m-i][j]x_k^{m-i-1}y_k^{i+1}x_l^{n-j+1}y_l^{j-1} + [i][n-j]q^{m-n-i+j+2}x_k^{m-i+1}y_k^{i-1}x_l^{n-j-1}y_l^{j+1} + [i][j+1]q^{m-n-i+j}x_k^{m-i}y_k^ix_l^{n-j}y_l^j).$$

For  $m, n \geq 2$  it follows that

$$P_{lk}P_{kl}f = \frac{1}{[m][n+1]} x_k^{m-i-1} y_k^{i-1} (q^{m+n+2}[i][n-j]x_k^2 y_l^2 + (q^{n-2}[m-i][n-j+1] + q^{m-2}[i][j+1]) x_k y_k x_l y_l + [m-i][j] y_k^2 x_l^2) x_l^{n-j-1} y_l^{j-1} = \frac{1}{[m][n+1]} x_k^{m-i-1} y_k^{i-1} (q^{m+n-2+\frac{1}{2}}[i][n-j]x_k (kl) y_l - q^{-\frac{1}{2}}[m-i][j] y_k (kl) x_l + q^{n-1}[i][n-i] + q^{n-2}[m-i][i] + q^{n-2}[m-i][n-i] + q^{n-2}[i][i+1]) x_l y_l x_l y_l x_l y_l + q^{n-j-1} y_l^{j-1}$$

 $\begin{aligned} (q^{m+n-1}[i][n-j]+q^{-3}[m-i][j]+q^{n-2}[m-i][n-j+1]+q^{m-2}[i][j+1])x_ky_kx_ly_l)x_l^{n-j-1}y_l^{j-1}. \\ \text{Because of } [i+j] = q^j[i]+q^{-i}[j] = q^{-j}[i]+q^i[j] \text{ we obtain} \end{aligned}$ 

$$= \frac{1}{[m][n+1]} x_k^{m-i-1} y_k^{i-1} (q^{m+n-2+\frac{1}{2}}[i][n-j]x_k(kl)y_l - q^{-\frac{1}{2}}[m-i][j]y_k(kl)x_l + q^{n-j+i-2}[m][n+1]x_1y_1x_2y_2) x_l^{n-j-1}y_l^{j-1}.$$

Applying Lemma 3.1 we find

$$=\frac{1}{[m][n+1]}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i}y_k^{i-1}x_l^{n-j-1}y_l^j-q^{-m+\frac{1}{2}}[m-i][j]x_k^{m-i-1}y_k^ix_l^{n-j}y_l^{j-1})+f.$$

We encounter the same formula in the cases m < 2 and n < 2. The first formula follows. Similarly we find

$$P_{lk}P_{kl}f = \frac{1}{[n][m+1]}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i}y_k^{i-1}x_l^{n-j-1}y_l^j - q^{-m+\frac{1}{2}}[m-i][j]x_k^{m-i-1}y_k^ix_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i}y_k^{i-1}x_l^{n-j-1}y_l^j - q^{-m+\frac{1}{2}}[m-i][j]x_k^{m-i-1}y_k^ix_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i}y_k^{i-1}x_l^{n-j-1}y_l^j - q^{-m+\frac{1}{2}}[m-i][j]x_k^{m-i-1}y_k^ix_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i}y_k^{i-1}x_l^{n-j-1}y_l^j - q^{-m+\frac{1}{2}}[m-i][j]x_k^{m-i-1}y_k^ix_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i}y_k^{i-1}x_l^{n-j-1}y_l^j - q^{-m+\frac{1}{2}}[m-i][j]x_k^{m-i-1}y_k^ix_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i-1}y_k^ix_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i-1}y_k^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_k^{m-i-1}y_k^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_k^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_k^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_k^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_k^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_l^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_l^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_l^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_l^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_l^jx_l^{n-j}y_l^{j-1}) + f_{lk}(kl)(q^{j-i+\frac{3}{2}}[i][n-j]x_l^{m-i-1}y_l^jx_l^{n-j}y_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x_l^{j-j}x$$

The second formula and the relation  $b = \frac{[m][n+1]}{[m+1][n]}a$  follow. The representation is unique, since  $H_I$  has no zero divisors (c.f. Proposition 2.2).

- (ii) The assertion is a consequence of (i) and of the identity
- [m][n+1] [n][m+1] = [m-n].

(iii) By Proposition 3.4 (iii) we have

$$0 = Ef = P_{lk}P_{kl}E(f) + E((kl))K(a) + K^{-1}((kl))E(a) = (kl)E(a).$$

Since  $H_I$  has no zero divisors, we have E(a) = 0. Similarly we derive K(a) = a and F(a) = 0.

### 3.2 The main Theorem

THEOREM 3.6 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ . The subalgebra of invariant elements  $H_I^{Inv}$  is generated by the elements  $(ij), i, j \in I$ .

*Proof.* Since the  $U_q(gl(2, \mathbb{R}))$ -action on  $H_I$  does not change the type of homogenity with respect to the indices  $i \in I$ , it is sufficient to consider invariants which are homogeneous with respect to all indices.

Let  $f \in H_I^{Inv}$  be an invariant element of degree n with k series of variables  $x_{a_j}, y_{a_j}, j = 1, ..., k, a_1 < a_2 < ... < a_k$ , i.e.

$$f = \sum_{i_1,\dots,i_k} c_{i_1,i_2,\dots,i_k} x_{a_1}^{n_1-i_1} y_{a_1}^{i_1} x_{a_2}^{n_2-i_2} y_{a_2}^{i_2} \dots x_{a_k}^{n_k-i_k} y_{a_k}^{i_k}$$

with  $n_j \ge 1$ ,  $i_j \ge 0$  and  $n_1 + \ldots + n_k = n$ . We suppose  $k \ge 2$  (i.e.  $n \ge 2$ ). For simplicity we set  $a_1 = 1$  and  $a_2 = 2$ .

Using Proposition 3.4 (iii) we can form the  $n_1$  invariants

$$f, P_{12}f, P_{12}^2f, \cdots, P_{12}^{n_1-1}f.$$

By Proposition 3.5 (i) we have

$$f = P_{21}P_{12}f + (12)a_0,$$
  

$$P_{12}f = P_{21}P_{12}^2f + (12)a_1,$$
  
....,  

$$P_{12}^{n_1-1}f = P_{21}P_{12}^{n_1}f + (12)a_{n_1-1}$$

with  $a_i \in H_I^{Inv}$ . It follows that

$$f = P_{21}P_{12}f + (12)a_0$$
  
=  $P_{21}(P_{21}P_{12}^2f + (12)a_1) + (12)a_0$   
=  $P_{21}^2P_{12}^2f + (12)(a_0 + P_{21}a_1)$   
.....  
=  $P_{21}^{n_1}P_{12}^{n_1}f + (12)(a_0 + P_{21}a_1 + P_{21}^2a_2 + \dots + P_{21}^{n_1-1}a_{n_1-1})$ 

Therefore we can reduce the problem for the invariant element f of degree n with k variables to the problems for the invariant element  $P_{12}^{n_1}f$  of degree n with k-1 variables and the invariant element

$$a_0 + P_{21}a_1 + P_{21}^2a_2 + \dots + P_{21}^{n_1 - 1}a_{n_1 - 1}$$

of degree n-2 with k variables. Consequently the problem reduces to the cases n = 0, 1 or k = 1 by a finite number of steps.

For n = 0 we have only the invariant element 1.

For n = 1 there are no invariant elements, since we infer from

$$K(\sum_{i}(\alpha_{i}x_{i}+\beta_{i}y_{i}))=\sum_{i}(q^{-\frac{1}{2}}\alpha_{i}x_{i}+q^{\frac{1}{2}}\beta_{i}y_{i})=\sum_{i}(\alpha_{i}x_{i}+\beta_{i}y_{i})$$

that  $\alpha_i = \beta_i = 0$ .

Finally we consider the case k = 1. The only elements  $a \in H_I$  with k = 1 and K(a) = a are  $a = \sum_{i=0}^{n} \alpha_i x_i^m y_i^m$ . From the condition

$$E(\sum_{i} \alpha_{i} x_{i}^{m} y_{i}^{m}) = \sum_{i} \alpha_{i} q^{\frac{2-m}{2}} [m] x_{i}^{m-1} y_{i}^{m+1} = 0$$

it follows that  $\alpha_i = 0$  for  $i \ge 1$ , since we supposed  $q^{2d} \ne 1$ , i.e.  $a = \alpha_0 1$ . Therefore we have no invariant elements for k = 1. The Theorem follows. •

### 3.3 The Gordan-Capelli series

In this section we consider  $H_I$  with  $I = \{1, 2, \dots, p\}$ . We use Proposition 3.5 (i), (ii) in order to introduce the quantum  $\Omega$ -process for the two lowest indices.

DEFINITION 3.2 Let  $f = P_{21}P_{12}f + (12)a$ . Then we define  $\Omega_{12}(f) := \frac{[n+1]}{[n]}a$ .

By Proposition 3.5 (i) a is unique and we have

$$\Omega_{12}(x_1^{m-i}y_1^i x_2^{n-j}y_2^j \beta)$$

$$=\frac{1}{[m][n]}(q^{j-i+\frac{3}{2}}[i][n-j]x_1^{m-i}y_1^{i-1}x_2^{n-j-1}y_2^j\beta-q^{-m+\frac{1}{2}}[m-i][j]x_1^{m-i-1}y_1^ix_2^{n-j}y_2^{j-1}\beta)$$

for  $m, n \ge 1$  and  $\beta \in H_{\{i \in I \mid i > 2\}}$  (cf. the proof of Proposition 3.5 (i)).

Therefore we have

$$f = P_{21}P_{12}f + \frac{[n]}{[n+1]}(12)\Omega_{12}(f)$$
(11)

and

$$f = P_{12}P_{21}f + \frac{[m]}{[m+1]}(12)\Omega_{12}(f).$$
(12)

Remark 3.4 For q = 1 we have

$$\Omega_{12} = \frac{1}{mn} \begin{vmatrix} \partial_{x_1} & \partial_{y_1} \\ \partial_{x_2} & \partial_{y_2} \end{vmatrix}$$

and Eq. (11) takes the form of the *Capelli identity*:

$$\begin{vmatrix} \Delta_{11} & \Delta_{12} \\ \Delta_{21} & \Delta_{22} + 1 \end{vmatrix} (f) = (12) \begin{vmatrix} \partial_{x_1} & \partial_{y_1} \\ \partial_{x_2} & \partial_{y_2} \end{vmatrix} (f)$$

(cf. [15, 16]). We have not introduced quantum deformations of the partial derivatives  $\partial_{x_i}$ ,  $\partial_{y_i}$ . Therefore we use Eq. (11) not as an identity, but only for the definition of the operator  $\Omega_{12}$ . This will be sufficient for the proof of Theorem 3.8. For further applications of the Capelli identity we refer to [5, 17]. A Capelli identity on quantum groups was considered in [18] for  $GL_2(\mathbb{R})$  and in [19] for  $GL_n(\mathbb{R})$ .

The operators  $P_{12}$ ,  $\Omega_{12}$  have the following commutation relations

PROPOSITION 3.7 (i) We have

$$P_{12}((12)f) = q \frac{[m]}{[m+1]}(12)P_{12}f,$$
$$P_{21}((12)f) = q \frac{[n]}{[n+1]}(12)P_{21}f$$

for  $f \in H_I$  and *(ii)* 

$$\Omega_{12}P_{12} = \frac{1}{q} \frac{[n]}{[n+1]} P_{12}\Omega_{12},$$
$$\Omega_{12}P_{21} = \frac{q[m]}{[m+1]} P_{21}\Omega_{12}.$$

*Proof.* (i) A proof can be given by an explicite calculation using Proposition 3.4 (ii). (ii) By (11) and Proposition 3.5 (ii) we have

$$P_{12}((12)\Omega_{12}(f)) = \frac{[m]}{[m+1]}P_{12}(f - P_{12}P_{21}f)$$
  
$$= \frac{[m]}{[m+1]} \left( P_{12}f - \frac{[n][m+1]}{[m][n+1]}P_{12}P_{21}P_{12}f - \frac{[m-n]}{[m][n+1]}P_{12}f \right)$$
  
$$= \frac{[n+1][m] - [m-n]}{[m+1][n+1]}P_{12}f - \frac{[n]}{[n+1]}P_{12}P_{21}P_{12}f$$
  
$$= \frac{[n]}{[n+1]}(1 - P_{12}P_{21})P_{12}f = \frac{[m][n]}{[m+1][n+1]}(12)\Omega_{12}P_{12}(f).$$

Using (i), we find

$$q(12)P_{12}\Omega_{12}(f)) = \frac{[n]}{[n+1]}(12)\Omega_{12}P_{12}(f).$$

The first equation follows, since  $H_I$  has no zero divisors. The proof of the second equation is analogous.

Now we demonstrate that we can represent every invariant with n variables by powers of symbols  $(12)^k$  and polars of invariants with n-1 variables.

THEOREM 3.8 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$  and let f be an invariant element of degree n with k series of variables  $x_i, y_i, i = 1, 2, ..., p$ , homogeneous of degree  $n_i$  with respect to i. Then we have

$$f = \sum_{k=0}^{n_1} \frac{\begin{bmatrix} n_1 \\ k \end{bmatrix} \begin{bmatrix} n_2 \\ k \end{bmatrix}}{\begin{bmatrix} n_1 + n_2 - k + 1 \\ k \end{bmatrix}} (12)^k P_{21}^{n_1 - k} P_{12}^{n_1 - k} \Omega_{12}^k f.$$

*Proof.* For  $0 \leq l \leq n_1$  we can represent f in the form

$$f = \sum_{k=0}^{l} \alpha_k^{(l)} (12)^k P_{21}^{l-k} P_{12}^{l-k} \Omega_{12}^k f$$
(13)

with

$$\alpha_k^{(l+1)} = \alpha_k^{(l)} + \alpha_{k-1}^{(l)} \frac{(n_2 - k + 1)^2}{(n_2 + l - 2k + 2)(n_2 + l - 2k + 3)}$$
(14)

 $(\alpha_{-1} := 0)$ . Since  $\alpha_0^{(0)} = \alpha_0^{(1)} = 1$ , the statement is true for l = 0. We suppose the statement for l, i.e. we have

$$f = \sum_{k=0}^{l} \alpha_k^{(l)} (12)^k P_{21}^{l-k} P_{12}^{l-k} \Omega_{12}^k f.$$

Applying (13) to the elements

$$P_{12}^{l-k}\Omega_{12}^k f$$

we find

$$f = \sum_{k=0}^{l} \alpha_{k}^{(l)} (12)^{k} P_{21}^{l-k} P_{21} P_{12} P_{12}^{l-k} \Omega_{12}^{k} f + \sum_{k=0}^{l} \alpha_{k}^{(l)} \frac{[n_{2}+l-2k]}{[n_{2}+l-2k+1]} (12)^{k} P_{21}^{l-k} (12) \Omega_{12} P_{12}^{l-k} \Omega_{12}^{k} f.$$

By Proposition 3.7 it follows that

Therefore the statement is true for l + 1 with (14).

The coefficients are uniquely determined by the relations (14). Therefore it is sufficient to check the relations (14) for

$$\alpha_k^{(l)} := \frac{ \begin{bmatrix} l \\ k \end{bmatrix} \begin{bmatrix} n_2 \\ k \end{bmatrix}}{ \begin{bmatrix} l+n_2-k+1 \\ k \end{bmatrix}}$$

It follows from the identity

$$[a][b-c] + [b][c-a] + [c][a-b] = 0$$

with a = l + 1, b = k,  $c = l + n_2 - k + 2$  that

$$[l+1][l+n_2-2k+2] = [k][n_2-k+1] + [l+n_2-k+2][l+1-k].$$

Furthermore

$$[l+1][l+n_2-2k+2] = [l+n_2-k+2][l+1-k] + \frac{[k][n_2+l-2k+2][n_2+l-2k+3]}{[n_2-k+1]} \frac{[n_2-k+1]^2}{[n_2+l-2k+2][n_2+l-2k+3]}$$

Therefore

$$\frac{\left[\begin{array}{c}l+1\\k\end{array}\right]\left[\begin{array}{c}n_2\\k\end{array}\right]}{\left[\begin{array}{c}l+n_2-k+2\\k\end{array}\right]} = \frac{\left[\begin{array}{c}l\\k\end{array}\right]\left[\begin{array}{c}n_2\\k\end{array}\right]}{\left[\begin{array}{c}l+n_2-k+1\\k\end{array}\right]} + \frac{\left[\begin{array}{c}l\\k-1\end{array}\right]\left[\begin{array}{c}n_2\\k-1\end{array}\right]}{\left[\begin{array}{c}l+n_2-k+1\\k-1\end{array}\right]}\left(\frac{[n_2-k+1]^2}{[n_2+l-2k+2][n_2+l-2k+3]}\right).$$

The assertion follows.

Remark 3.5 In the classical case we can consider the vector space generated by the monomials  $x_1^{m-i}y_1^ix_2^{n-j}y_2^j$ ,  $0 \le i \le m$ ,  $0 \le j \le n$  as the representation module  $V_m \otimes V_n$  of the Lie algebra sl(2). The elements  $P_{12}^{n_1-k}\Omega_{12}^kx_1^{m-i}y_1^ix_2^{n-j}y_2^j$ ,  $0 \le k \le \min(m, n)$  depend only on  $x_2, y_2$ . One can show that the polars

$$(12)^k P_{21}^{n_1-k} P_{12}^{n_1-k} \Omega_{12}^k x_1^{m-i} y_1^i x_2^{n-j} y_2^j$$

span an irreducible subrepresentation  $V_{m+n-2k}$ . Therefore we can use the quantised Gordan-Capelli series for an explicite decomposition of the braided tensor product of  $V_m$  and  $V_n$  (cf. [9]) into irreducible representations of  $U_q(sl(2, \mathbb{R}), \text{ i.e., for } n = 2$  the Gordan-Capelli series determines the structure of the representation ring of  $U_q(sl(2, \mathbb{R}), \mathbb{R})$ .

## 4. *n*-FORMS

#### 4.1 Definition and Examples

In this section we use the algebra  $H_I$  in order to introduce noncommutative binary homogeneous *n*-forms.

For the following we suppose for the algebra  $H_I$  that  $0 \in I$ . We will use the short notations

$$x := x_0, \qquad y := y_0.$$

Furthermore let  $H_{I\setminus\{0\}}$  be the subalgebra of  $H_I$  which is generated by the elements  $x_i, y_i, i \in I\setminus\{0\}$ .

DEFINITION 4.1 We say that the nonvanishing element

$$f := x^{n} A_{0} + \begin{bmatrix} n \\ 1 \end{bmatrix} x^{n-1} y A_{1} + \begin{bmatrix} n \\ 2 \end{bmatrix} x^{n-2} y^{2} A_{2} + \dots + y^{n} A_{n}$$
(15)

of  $H_I$  with  $A_i \in H_{I \setminus \{0\}}$  is a *n*-form, if  $f \in H_I^{Inv}$ .

Remark 4.1 The representation of f with coefficients  $A_i \in H_{I \setminus \{0\}}$  is unique. This is a consequence of the PBW theorem. We do not suppose anything about the commutation behaviour of the elements  $A_i$  (cf. Remark 4.2 below).

Example 4.1 For the linear form

$$f = xA + yB := (0i)$$

 $(i \neq 0)$  we find  $A = q^{-\frac{1}{2}}y_i, B = -q^{\frac{1}{2}}x_i.$ 

Example 4.2 For the linear form

$$f = xA + yB := (01)(23) - c(03)(12)$$

a calculation yields

$$A = cq^{2}x_{1}y_{2}y_{3} + (q^{-1} - cq^{3})y_{1}x_{2}y_{3} - y_{1}y_{2}x_{3},$$
  
$$B = -x_{1}x_{2}y_{3} + (q - cq^{3})x_{1}y_{2}x_{3} + cq^{4}y_{1}x_{2}x_{3}.$$

Example 4.3 For the quadratic form "with two different real zeros"

$$f = x^{2}A + xy[2]_{q^{2}}B + y^{2}C := q(01)(02)$$

we find  $A = qy_1y_2$ ,  $B = \frac{-q^2x_1y_2-qy_1x_2}{q+q^{-1}}$  and  $C = q^3x_1x_2$ , (cf. Lemma 3.3 (i)). (Later we will be able to present examples of quantised real forms with complex zeros.)

Example 4.4 For the quadratic form "with two equal real zeros"

$$f = x^2 A + [2]_{q^2} xy B + y^2 C := (01)^2$$

we find  $A = y_1^2$ ,  $B = -x_1y_1$  and  $C = q^2x_1^2$ .

Example 4.5 Consider n-forms of the type

$$(0i_1)(0i_2)\cdots(0i_n)$$

which correspond to polynomials with real zeros. Because of  $(0j)(0i) = q^2(0i)(0j)$ ,  $(0 < i < j), (0j)(0i) = q^4(0i)(0j), (i < 0 < j)$  and  $(0j)(0i) = q^2(0i)(0j), (i < j < 0)$  a permutation of the indices yields only a multiplication of the form by a nonzero constant factor (cf. Proposition 3.2 (i)). Therefore it is sufficient to consider forms  $(0i_1)(0i_2)\cdots(0i_n)$ with  $i_1 \leq i_2 \leq \ldots \leq i_n$ . We will endow each form of this kind with a certain factor  $q^c$ ,  $c \in \mathbb{N}$  such that the form becomes a real (i.e. \*-invariant) algebra element. For example let  $0 < i_1 < i_2 < \cdots < i_n$ . Then

$$f_{i_1,i_2,\ldots,i_n} := q^{\frac{n(n-1)}{2}}(0i_1)(0i_2)\cdots(0i_n)$$

is a real form.

Remark 4.2 The commutation behaviour of the coefficients of a *n*-form depends on the concrete type of the form. In Example 4.1 the coefficients A, B of the linear form f = xA + yB = (0i) have the commutation relations

$$BA = q^2 AB.$$

However for the linear form f = xA + yB of Example 4.2 we have

$$BA = qAB + cq^{\frac{7}{2}}(q^2 - 1)(12)(13)(23).$$
(16)

One can show, that there are no closed quadratic commutation relations between A and B.

PROPOSITION 4.1 We have

$$L[A_i] = q^{\frac{n}{2}} A_i, K[A_i] = q^{\frac{n-2i}{2}} A_i, F[A_i] = -q^{\frac{2i-n+2}{2}} [i] A_{i-1}, F[A_i] = -q^{\frac{-n-2i}{2}} [n-i] A_{i+1} (17)$$

 $(A_{-1} = A_{n+1} := 0).$ 

*Proof.* We have

$$K(f) = \sum_{i=0}^{n} \begin{bmatrix} n\\i \end{bmatrix} q^{\frac{-n+2i}{2}} x^{n-i} y^{i} K(A_{i}).$$

By K(f) = f and by the PBW theorem the formula for K follows. Furthermore

$$E(f) = \sum_{i=0}^{n} \begin{bmatrix} n\\i \end{bmatrix} (E(x^{n-i})K(y^{i}A_{i}) + K^{-1}(x^{n-i})E(y^{i})K(A_{i}) + K^{-1}(x^{n-i}y^{i})E(A_{i})).$$

By  $E(x^m) = q^{\frac{2-m}{2}}[m]x^{m-1}y$  and  $E(y^m) = 0$  it follows that

$$=\sum_{i=0}^{n} \begin{bmatrix} n\\i \end{bmatrix} \left(q^{\frac{2-n+i}{2}}[n-i]x^{n-i-1}yq^{\frac{i}{2}}y^{i}q^{\frac{n-2i}{2}}A_{i} + q^{\frac{n-2i}{2}}x^{n-i}y^{i}E(A_{i})\right)$$
$$=\sum_{i=0}^{n} \begin{bmatrix} n\\i-1 \end{bmatrix} q[n-i+1]x^{n-i}y^{i}A_{i-1} + \begin{bmatrix} n\\i \end{bmatrix} q^{\frac{n-2i}{2}}x^{n-i}y^{i}E(A_{i}).$$

By E(f) = 0 and by the PBW theorem the formula for E follows. The proof for F and L is analogous. •

Therefore the action of  $U_q(gl(2, \mathbb{R}))$  on the coefficients is independent from the concrete realisation of the *n*-form.

### 4.2 Geometric interpretation of $H_I$

In Remark 2.3 we considered the generators  $x_i, y_i, i \in I$  as homogeneous coordinates of points of the real line. In this section we use these points in order to construct further points.

DEFINITION 4.2 We say that the ordered pair (X, Y) of nonvanishing elements  $X, Y \in H_{I\setminus\{0\}}^{Inv}$  is a *point* if

(i)  $q^{-1/2}x_0Y - q^{1/2}y_0X \in H_I^{Inv}$  and

(ii) X, Y are homogeneous with respect to all indices  $i \in I$  of the same degree. We say, that the two points  $(X_1, Y_1)$ ,  $(X_2, Y_2)$  are equal if  $X_1Y_1^{-1} = X_2Y_2^{-1}$  in  $Q_I$ . Remark 4.3 Le q be not a root of unity. Then X, Y are elements of odd degree, since  $H_I^{Inv}$  is generated by the elements (ij).

Example 4.6 Pairs of generators  $(x_i, y_i)$  are points. Furthermore let  $f = x_0A + y_0B$  be a linear form. Then  $(-q^{-1/2}B, q^{1/2}A)$  is a point. For the form f = (01)(23) - c(03)(12)of Example 4.2 we have

$$X = q^{\frac{-1}{2}} x_1 x_2 y_3 - (q^{\frac{1}{2}} - cq^{\frac{5}{2}}) x_1 y_2 x_3 - cq^{\frac{7}{2}} y_1 x_2 x_3$$

$$Y = cq^{\frac{5}{2}}x_1y_2y_3 + (q^{\frac{-1}{2}} - cq^{\frac{7}{2}})y_1x_2y_3 - q^{\frac{1}{2}}y_1y_2x_3$$

Example 4.7 Let (X, Y) be a point and  $a \in H_I^{Inv}$ . Then we have

$$(X,Y) = (Xa,Ya).$$

### 4.3 Polynomials, Elementary symmetric functions, Newton relations

In Remark 2.3 we have introduced the elements  $v_i = q^{\frac{1}{2}} x_i y_i^{-1} \in Q_I$  which obey the commutation relations

$$v_j v_i = q^2 v_i v_j - (1 - q^2) v_i^2$$

for i < j.

We define bracket symbols [ij] by

 $[ij] := v_i - v_j.$ 

They are related to the bracket symbols (ij) by

 $(ij) = y_i[ij]y_j.$ 

If we consider  $v_0$  as an independent variable, we can define *polynomials with zeros*  $v_{i_1}$ ,  $v_{i_2}$ , ...,  $v_{i_n}$  by

$$p_{i_1,i_2,\ldots,i_n} := [0i_1][0i_2] \cdots [0i_n].$$

Remark 4.4 The definition of  $p_{i_1,i_2,...,i_n}$  depends on the relative position of the brackets  $[0i_j]$ . By Example 4.5 a permutation of the indices of the form  $(0i_1)(0i_2)\cdots(0i_n)$  yields only a multiplication by a constant factor. We have not the same situation for polynomials  $p_{i_1,i_2,...,i_n}$ . For example we have

$$p_{1,2} = [01][02] = q^{-2}v_0^2 - q^{-4}v_0(q^2v_1 + v_2) + q^{-4}v_1v_2$$

and

$$p_{2,1} = [02][01] = q^{-2}v_0^2 - q^{-4}v_0(v_1 + q^2v_2) + q^{-2}v_1v_2 + q^{-4}(1 - q^2)v_1^2.$$

Remark 4.5 The polynomial  $p_{i_1,i_2,...,i_n}$  is related to the form  $f_{i_1,i_2,...,i_n}$  (cf. Example 4.5). By Lemma 3.1 we have the commutation relations

$$(0i)y_0^{-1} = \frac{1}{q}y_0^{-1}(0i), \qquad (0i)y_i^{-1} = qy_i^{-1}(0i), \qquad (0i)y_j^{-1} = q^3y_j^{-1}(0i)$$

for 0 < i < j. It follows that

$$p_{1,2,\dots,n} = [01][02] \cdots [0n] = y_0^{-1}(01)y_1^{-1}y_0^{-1}(02)y_2^{-1}y_0^{-1} \cdots y_0^{-1}(0n)y_n^{-1}$$
$$= q^{n^2}y_0^{-1}y_1^{-1}y_0^{-1}y_2^{-1} \cdots y_0^{-1}y_n^{-1}(01)(02) \cdots (0n)$$
$$= q^{\frac{n(3-n)}{2}}y_0^{-n}y_1^{-1}y_2^{-1} \cdots y_n^{-1}f_{1,2,\dots,n}.$$

In the following we restrict the consideration to polynomials  $p_{1,2,\dots,n}$ ,  $n \in \mathbb{N}$ .

**PROPOSITION 4.2** We have

$$p_{1,2,\dots,n} = q^{n(1-n)} \sum_{j=0}^{n} v_0^{n-j} A_j$$

with

$$A_j = (-1)^j \sum_{1 \le i_1 < i_2 < \dots < i_j \le n} q^{2j - 2(i_1 + i_2 + \dots + i_j)} v_{i_1} v_{i_2} \dots v_{i_j}$$

for j = 1, ..., n and  $A_0 = 1$ .

The statement can be shown by induction.

We say that  $A_j$  is the *j*-th symmetric function of  $v_{i_1}, v_{i_2}, ..., v_{i_j}$ . Furthermore we define the *j*-th power sum  $P_j, j \ge 1$  by

$$P_j = v_1^j + q^{-2}v_2^j + \dots + q^{2-2n}v_n^j.$$

PROPOSITION 4.3 Let  $k \ge 1$  and  $A_k := 0$  for k > n or k < 0. Then we have

$$P_k = -q^{2(k-1)(n-1)}(k)_{q^2}A_k - \sum_{i=1}^{k-1} q^{2(k-i)(n-1)} A_{k-i}P_i$$

(Newton relations).

*Proof.* Let n = 1. We have  $P_k = v_1^k$ ,  $A_0 = 1$ ,  $A_1 = -v_1$  and  $A_k = 0$  for  $k \neq 0, 1$ . It follows that  $P_1 = -A_1$  and  $P_k = -A_1P_{k-1}$  for  $k \geq 2$ . Therefore the assertion is true for n = 1.

We suppose that the assertion is true for n. Let  $A_i^{(n)} := A_i$  be the coefficients of  $p_{1,2,\dots,n}$ . By Proposition 4.2 and the definition of  $P_j$  we have

$$A_j^{(n+1)} = A_j^{(n)} - q^{-2n} A_{j-1}^{(n)} v_{n+1}$$

and

$$P_j^{(n+1)} = P_j^{(n)} + q^{-2n} v_{n+1}^j \quad \text{for } j \ge 1.$$

It follows that

$$P_{k}^{(n+1)} + q^{2(k-1)n}(k)A_{k}^{(n+1)} + \sum_{i=1}^{k-1} q^{2(k-i)n}A_{k-i}^{(n+1)}P_{i}^{(n+1)}$$

$$= P_k^{(n)} + q^{-2n} v_{n+1}^k + q^{2(k-1)n}(k) A_k^{(n)} - q^{2(k-1)n-2n}(k) A_{k-1}^{(n)} v_{n+1}$$
  
+  $\sum_{i=1}^{k-1} q^{2(k-i)n} A_{k-i}^{(n)} P_i^{(n)} - \sum_{i=1}^{k-1} q^{2(k-i)n-2n} A_{k-i-1}^{(n)} v_{n+1} P_i^{(n)}$   
+  $\sum_{i=1}^{k-1} q^{2(k-i)n-2n} A_{k-i}^{(n)} v_{n+1}^i - \sum_{i=1}^{k-1} q^{2(k-i)n-4n} A_{k-i-1}^{(n)} v_{n+1}^{i+1}.$ 

The last row gives  $q^{2(k-2)n} A_{k-1}^{(n)} v_{n+1} - q^{-2n} v_{n+1}^k$ . Because of the identity  $v_{n+1}v_j^i = q^{2i}v_j^i v_{n+1} + (1-q^{2i})v_j^{i+1}$  for j < n+1 we have

$$= P_k^{(n)} + q^{2(k-1)n}(k)A_k^{(n)} + q^{2(k-2)n}(1-(k))A_{k-1}^{(n)}v_{n+1} + \sum_{i=1}^{k-1} q^{2(k-i)n}A_{k-i}^{(n)}P_i^{(n)}$$

$$-\sum_{i=1}^{k-1} q^{2(k-i)n-2n+2i} A_{k-i-1}^{(n)} P_i^{(n)} v_{n+1} - \sum_{i=1}^{k-1} q^{2(k-i)n-2n} (1-q^{2i}) A_{k-i-1}^{(n)} P_{i+1}^{(n)}.$$

We unite the first and the third sum and rewrite the second sum.

$$= P_k^{(n)} + q^{2(k-1)n}(k)A_k^{(n)} + q^{2(k-2)n}(1-(k))A_{k-1}^{(n)}v_{n+1} + \sum_{i=1}^{k-1} q^{2(k-i)n+2i-2}A_{k-i}^{(n)}P_i^{(n)}$$
$$-(1-q^{2k-2})A_0^{(n)}P_k^{(n)} - q^{2k-2}\sum_{i=1}^{k-2} q^{2(k-1-i)(n-1)}A_{k-i-1}^{(n)}P_i^{(n)}v_{n+1} - q^{2k-2}P_{k-1}^{(n)}.$$

Because the assertion is true for n we can replace the first sum by  $-q^{2k-2}(P_k^{(n)}+q^{2(k-1)(n-1)}A_k^{(n)})$ and the last two terms by  $(k-1)_{q^2}q^{2(k-2)(n-1)+2k-2}A_{k-1}^{(n)}v_{n+1}$ . We obtain

$$= P_k^{(n)} + q^{2(k-1)n}(k)_{q^2} A_k^{(n)} + q^{2(k-2)n}(1-(k)_{q^2}) A_{k-1}^{(n)} v_{n+1} - q^{2k-2} (P_k^{(n)} + q^{2(k-1)(n-1)} A_k^{(n)}) -(1-q^{2k-2}) A_0^{(n)} P_k^{(n)} + (k-1)_{q^2} q^{2(k-2)(n-1)+2k-2} A_{k-1}^{(n)} v_{n+1} = (q^{2(k-2)n}(1-(k)_{q^2}) + (k-1)_{q^2} q^{2(k-2)(n-1)+2k-2}) A_{k-1} v_{n+1} = 0.$$

The assertion follows.  $\bullet$ 

# 5. INVARIANTS OF FORMS

#### 5.1 Definition

We consider a *n*-form  $f = \sum_{i=0}^{n} \begin{bmatrix} n \\ i \end{bmatrix} x^{n-i} y^{i} A_{i}.$ 

DEFINITION 5.1 We say that the polynomial expression of the coefficients of f

$$I_f = I_f(A_0, \dots, A_n)$$

is an *invariant*, if  $I_f \in H_{I\setminus\{0\}}^{Inv}$ . We say that an invariant  $I_f$  is an *invariant of degree* k, if  $I_f$  is a polynomial of degree k in the coefficients  $A_i$ . We say that the polynomial expression

$$C_f = C_f(A_0, ..., A_n, x_0, y_0)$$

is a *covariant* of f of degree m, if  $C_f$  is a m-form.

Example 5.1 Let f = q(01)(02) (cf. Example 4.3) and

$$I_f := \left(\frac{1}{q}AC + \frac{1}{q^3}CA\right) - \frac{1}{q}[2]B^2$$

A calculation yields

$$I_f = -\frac{1}{[2]}(12)^2.$$

Therefore  $I_f$  is an invariant of degree 2.

Sums and scalar multiples of invariants of degree k are again invariants of degree k. By (2) the product of two invariants of the degrees k and l is an invariant of degree k + l. Therefore the invariants of a *n*-form f form a graded algebra.

Furthermore we define invariants of a collection of forms.

DEFINITION 5.2 We say that a polynomial expression  $I_{f',f'',\dots}$  of the coefficients  $A'_i, A''_i, \dots$ of the binary forms  $f', f'', \dots \in H_I$  of the degrees  $n', n'', \dots$  is a simultaneous invariant, if  $I_{f',f'',\dots} \in H_{I\setminus\{0\}}^{Inv}$ . Analogous we define common covariants.

Example 5.2 Let f' = (01), f'' = (02) (cf. Example 4.1) and

$$I_{f',f''} = q^{-\frac{1}{2}}B'A'' - q^{\frac{1}{2}}A'B''.$$

We have

$$I_{f',f''} = -(12) \in H^{Inv}_{I \setminus \{0\}}.$$

### 5.2 Universal invariants

The Remark 4.2 demonstrates us that the structure of the coefficient algebra depends on the concrete type of the *n*-form. Therefore it is useful to define the notions of *universal* n-forms and *universal invariants*.

DEFINITION 5.3 We say that the expression  $\mathbf{D}$ 

$$\mathcal{F} := x^n \mathcal{A}_0 + \begin{bmatrix} n \\ 1 \end{bmatrix} x^{n-1} y \mathcal{A}_1 + \begin{bmatrix} n \\ 2 \end{bmatrix} x^{n-2} y^2 \mathcal{A}_2 + \dots + y^n \mathcal{A}_n$$

is an universal *n*-form, if the  $\mathcal{A}_i$  are abstract variables. We obtain a *n*-form as a realisation from an universal form, if we insert the elements  $A_i \in H_I$  for  $\mathcal{A}_i$ .

Consider k universal n-forms  $\mathcal{F}', ..., \mathcal{F}^{(k)}$ . We form the free algebra  $F_I$  generated by their coefficients  $\mathcal{A}_i^{(j)}$ . For the realisation  $f', ..., f^{(k)}$  we have the homomorphism  $h_{f',...,f^{(k)}}$ :  $F_I \to H_I$  determined by  $h_{f',...,f^{(k)}}(\mathcal{A}_i^{(j)}) := A_i^{(j)}$ .

The formulas (17) determine also an  $U_q(gl(2, \mathbb{R}))$ -action on  $F_I$  if we replace  $A_i$  by  $\mathcal{A}_i$ . We have

$$X(h_{f',\dots,f^{(k)}}(a)) = h_{f',\dots,f^{(k)}}(X(a)),$$
(18)

for  $X \in U_q(gl(2, \mathbb{R}) \text{ and } a \in F_I$ .

DEFINITION 5.4 We say that  $\mathcal{I} \in F_I$  is an *universal invariant* of  $\mathcal{F}$  if  $\mathcal{I}$  is an invariant element of the module algebra  $F_I$  (i.e.  $E(\mathcal{I}) = F(\mathcal{I}) = 0$  and  $K(\mathcal{I}) = \mathcal{I}$ ). Analogous we define *universal simultaneous invariants* and *universal covariants*.

Example 5.3 Consider the universal form

$$x^2\mathcal{A} + [2]_{q^2}xy\mathcal{B} + y^2\mathcal{C}$$

and

$$\mathcal{I} := \frac{1}{q}\mathcal{A}\mathcal{C} + \frac{1}{q^3}\mathcal{C}\mathcal{A} - \frac{1}{q}[2]\mathcal{B}^2.$$

By (17) and (18) we have  $E(\mathcal{A}) = 0, E(\mathcal{B}) = -q\mathcal{A}, E(\mathcal{C}) = -q^2\mathcal{B}$  and  $K(\mathcal{A}) = q\mathcal{A}, K(\mathcal{B}) = \mathcal{B}, K(\mathcal{C}) = q^{-1}\mathcal{C}$ . Therefore we obtain  $K(\mathcal{I}) = \mathcal{I}$  and

$$E(\mathcal{I}) = \frac{1}{q} K^{-1}(\mathcal{A}) E(\mathcal{C}) + \frac{1}{q} E(\mathcal{A}) K(\mathcal{C}) + \frac{1}{q^3} K^{-1}(\mathcal{C}) E(\mathcal{A})$$
$$+ \frac{1}{q^3} E(\mathcal{C}) K(\mathcal{A}) - \frac{1}{q} [2] K^{-1}(\mathcal{B}) E(\mathcal{B}) - \frac{1}{q} [2] E(\mathcal{B}) K(\mathcal{B})$$
$$= -[2] \mathcal{A} \mathcal{B} - [2] \mathcal{B} \mathcal{A} + [2] \mathcal{A} \mathcal{B} + [2] \mathcal{B} \mathcal{A} = 0.$$

Similarly we find  $F(\mathcal{I}) = 0$ .

LEMMA 5.1 Let  $\mathcal{I}$  be an universal invariant of  $\mathcal{F}$  and let f be a realisation of  $\mathcal{F}$ . Then  $I_f := h_f(\mathcal{I})$  is an invariant of f.

*Proof.* The assertion follows from (18).

Remark 5.1 Let  $\mathcal{I}$  be a simultaneous universal invariant of the universal n-forms  $\mathcal{F}', \mathcal{F}'', \dots$ of the same degree k. If we identify the coefficients of the different forms with the coefficients of  $\mathcal{F}$ , then  $\mathcal{I}$  becomes an invariant of the form  $\mathcal{F}$ . This follows, since the form of the  $U_q(sl_2)$ -action on the coefficients of the different forms  $\mathcal{F}^{(j)}$  is independent on j.

5.3 The symbolic method of Clebsch and Gordan

Consider the universal n-form

$$\mathcal{F} = x^n \mathcal{A}_0 + \begin{bmatrix} n \\ 1 \end{bmatrix} x^{n-1} y \mathcal{A}_1 + \begin{bmatrix} n \\ 2 \end{bmatrix} x^{n-2} y^2 \mathcal{A}_2 + \dots + y^n \mathcal{A}_n.$$

Our aim is to construct universal invariants  $\mathcal{I}$  of degree k.

We consider the collection of k realisations of  $\mathcal{F}$ 

$$f' := (01)^n, \quad f'' := (02)^n, \quad \dots, \quad f^{(k)} := (0k)^n.$$

By Lemma 3.3 (ii) the forms have the coefficients

$$A^{(j)}{}_{i} = (-1)^{i} q^{\frac{1}{2}n^{2} + (i-n)(1+i)} x_{j}^{i} y_{j}^{n-i}.$$
(19)

Consider an element d of  $H_I^{Inv}$  which is homogeneous of degree n with respect to 1, 2, ..., k. (For example (12)(13)(23) is homogeneous of degree 2 with respect to 1, 2, 3.)

Because of the PBW theorem for  $H_I$  and because the application of the reduction rules (5) does not change the degree of homogenity with respect to 1, 2, ..., k we can express d as a linear combination of monomials

$$x_1^{n-i_1}y_1^{i_1}x_2^{n-i_2}y_2^{i_2}\dots x_k^{n-i_k}y_k^{i_k}.$$
(20)

Using (19) we can represent d in terms of the coefficients  $A^{(j)}{}_i$ . We obtain the expression

$$d = \sum c_{i_1, i_2, \dots, i_k} A_{i_1}^{(1)} A_{i_2}^{(2)} \dots A_{i_k}^{(k)}.$$

Furthermore we consider the elements of  $F_I$ 

$$\mathcal{I}' = \sum c_{i_1, i_2, \dots, i_k} \mathcal{A}_{i_1}^{(1)} \mathcal{A}_{i_2}^{(2)} \dots \mathcal{A}_{i_k}^{(k)}$$

and

$$E(\mathcal{I}') = \sum d_{i_1, i_2, \dots, i_k} \mathcal{A}_{i_1}^{(1)} \mathcal{A}_{i_2}^{(2)} \dots \mathcal{A}_{i_k}^{(k)}.$$

By (18) it follows that

$$0 = E(d) = E(h_{f',\dots,f^{(k)}}(\mathcal{I}')) = h_{f',\dots,f^{(k)}}(E(\mathcal{I}')) = \sum d_{i_1,i_2,\dots,i_k} A_{i_1}^{(1)} A_{i_2}^{(2)} \dots A_{i_k}^{(k)}$$
$$= \sum d_{i_1,i_2,\dots,i_k} q^{\frac{1}{2}k_{i_1,i_2,\dots,i_k}} x_1^{n-i_1} y_1^{i_1} x_2^{n-i_2} y_2^{i_2} \dots x_k^{n-i_k} y_k^{i_k}$$

with certain numbers  $k_{i_1,i_2,\ldots,i_k} \in \mathbb{Z}$ . We have  $|q^{\frac{1}{2}k_{i_1,i_2,\ldots,i_k}}| = 1$ . By the PBW theorem we have  $d_{i_1,i_2,\ldots,i_k} = 0$ , i.e.  $E(\mathcal{I}') = 0$ . Similarly we encounter  $F(\mathcal{I}') = 0$  and  $K(\mathcal{I}') = \mathcal{I}'$ . Therefore  $\mathcal{I}'$  is an universal simultaneous invariant of k universal n-forms  $\mathcal{F}', \mathcal{F}'', \ldots, \mathcal{F}^k$ .

By Remark 5.1 we obtain again an invariant if we identify  $\mathcal{A}_i^{(j)}$  with  $\mathcal{A}_i$ . Therefore

$$\mathcal{I} = \sum c_{i_1, i_2, \dots, i_k} \mathcal{A}_{i_1} \mathcal{A}_{i_2} \dots \mathcal{A}_{i_k}$$

is an universal invariant of degree k of the universal *n*-form  $\mathcal{F}$ .

Therefore we have demonstrated the following Theorem.

THEOREM 5.2 Let  $\mathcal{F}$  be an universal n-form and let d be an element of  $H_I^{Inv}$  which is homogeneous of degree n with respect to 1, 2, ..., k. Then the above construction gives an universal invariant of  $\mathcal{F}$  of degree k.

Analogous we can use the Clebsch-Gordan method in order to construct universal covariants, universal common invariants and universal common covariants.

We say that the element of  $H_I^{Inv}$ 

$$d = (i_1 i_2)(i_3 i_4)\dots(i_{kn-1} i_{kn})$$

is a symbol of  $\mathcal{F}$ , if d is homogeneous of degree n with respect to 1, 2, ..., k.

The simplest symbols are  $(12)^2$ , (12)(13)(23) for the quadratic form,  $(12)^3$  for the cubic form and  $(12)^4$ ,  $(12)^2(13)^2(23)^2$  for the quartic form.

If  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ , then we can represent the generating element d by a symbol by Theorem 3.6.

Using Lemma 5.1, we obtain invariants of degree k for the realisations of the universal n-form.

Example 5.4 Consider the universal quadratic form

$$f = x^2 \mathcal{A} + [2]xy\mathcal{B} + y^2\mathcal{C}.$$

For the invariant  $(ij)^2$  we have (cf. Lemma 3.3 (ii))

$$(ij)^2 = x_i^2 y_j^2 - [2] x_i y_i x_j y_j + q^2 y_i^2 x_j^2.$$

Consider the realisations  $f = (0i)^2$ , i = 1, 2 with

$$A^{(i)} = y_i^2, \qquad B^{(i)} = -x_i y_i, \qquad C^{(i)} = q^2 x_i^2$$

We obtain for the symbol  $d := 1/q(12)^2$ 

$$\frac{1}{q}(12)^2 = \frac{1}{q}x_1^2y_2^2 - \frac{1}{q}[2]x_1y_1x_2y_2 + qy_1^2x_2^2$$
$$= \frac{1}{q^3}C'A'' - \frac{[2]}{q}B'B'' + \frac{1}{q}A'C''.$$

Therefore we have an universal common invariant of  $\mathcal{F}'$  and  $\mathcal{F}''$ .

$$\mathcal{I}' := \frac{1}{q^3} \mathcal{C}' \mathcal{A}'' - (1 + \frac{1}{q^2}) \mathcal{B}' \mathcal{B}'' + \frac{1}{q} \mathcal{A}' \mathcal{C}''.$$

If we identify the coefficients of  $\mathcal{F}'$  and  $\mathcal{F}''$  we obtain again the universal invariant of Example 5.3.

$$\mathcal{I} = \frac{1}{q^3} \mathcal{C} \mathcal{A} - \frac{|2|}{q} \mathcal{B} \mathcal{B} + \frac{1}{q} \mathcal{A} \mathcal{C}.$$

We can obtain all universal invariants by the symbolic method.

PROPOSITION 5.3 Every universal invariant of the n-form  $\mathcal{F}$  is a linear combination of invariants, which are derived from elements  $d \in H_I^{Inv}$  by the above method. If  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ , then we can derive all universal invariants from symbols of  $\mathcal{F}$ .

*Proof.* Let

$$\mathcal{I} = \sum c_{i_1, i_2, \dots, i_k} \mathcal{A}_{i_1} \mathcal{A}_{i_2} \dots \mathcal{A}_{i_k}$$

be an universal invariant of degree m of the *n*-form  $\mathcal{F}$  and let

$$f^{(j)} = (0j)^n = \sum_{i=1}^n \begin{bmatrix} n\\i \end{bmatrix} x^{n-i} y^i A_i^{(j)},$$

with j = 1, ..., k be realisations of  $\mathcal{F}$ . We set

$$d := \sum c_{i_1, i_2, \dots, i_k} A_{i_1}^{(1)} A_{i_2}^{(2)} \dots A_{i_k}^{(k)}.$$

 $d = h_{f',\dots,f^{(k)}}(\mathcal{I})$  is an invariant element of  $H_{I\setminus\{0\}}$ . From the construction of d it follows that we obtain again  $\mathcal{I}$  from d by the symbolic method. Using Theorem 3.6 we can express d as a linear combination of symbols of  $\mathcal{F}$  if  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ .

Remark 5.2 We can modify the symbolic method, if we use the elements

$$x_{\pi(1)}^{n-i_1}y_{\pi(1)}^{i_1}x_{\pi(2)}^{n-i_2}y_{\pi(2)}^{i_2}...x_{\pi(k)}^{n-i_k}y_{\pi(k)}^{i_k},$$

where  $\pi \in S_k$  is a permutation, as PBW basis elements instead of (20). It is a consequence of Proposition 5.3 that we obtain no further invariants by this modification.

### 5.4 The case of q being a root of unity

In this section we give some remarks for the case of q being a root of unity. We mention that the Clebsch-Gordan symbolic method gives all universal invariants even in this case. However, if  $q^{2d} = 1$  for  $d \in \mathbb{N}$ , we have invariant elements of  $H_I$ , which cannot be represented by symbols.

PROPOSITION 5.4 Let q be a primitive 2d-th root of unity  $(d \ge 2)$ . Then  $H_{\{1\}}^{Inv}$  is generated by the elements  $x_1^{kd}y_1^{k'd}$  with  $k, k' \in \mathbb{N}$  and 4|(k-k').

*Proof.* Let  $f = \sum_{i,j} a_{ij} x_1^i y_1^j$  be an invariant element. We have

$$Kf = \sum_{i,j} a_{ij} q^{\frac{-i+j}{2}} x_1^i y_1^j = f,$$
$$Ef = \sum_{i,j} a_{ij} [i] q^{\frac{2-i+j}{2}} x_1^{i-1} y_1^{j+1} = 0$$

and

$$Ff = \sum_{i,j} a_{ij}[j] q^{\frac{i-j}{2}} x_1^{i+1} y_1^{j-1} = 0$$

Note that [i] = 0, if and only if d|i. Therefore  $H_{\{1\}}^{Inv}$  is spanned by the monomials  $x^{kd}y^{k'd}$  with 4|(k-k')|.

Now we can use the Clebsch-Gordan symbolic method in order to determine the universal invariants of degree 1. Let again q be a primitive 2d-th root of unity  $(d \ge 2)$ , let  $\mathcal{F}$  be the universal *n*-form and let

$$f = (01)^n = \sum_{k=0}^n \begin{bmatrix} n\\k \end{bmatrix} (-1)^k q^{\frac{1}{2}n^2 + (k-n)(1+k)} x_0^{n-k} y_0^k x_1^k y_1^{n-k}$$

be a realisation. For the invariant element  $x_1^{kd}y_1^{k'd}$  with 4|k-k' we have (k+k')d = n.

Therefore  $\mathcal{F}$  has universal invariants only for even  $\frac{n}{d}$  and by the Clebsch-Gordan symbolic method certain coefficients are obtained as universal invariants: Case 1.:  $\frac{n}{d} \equiv 2 \mod 4$ :

$$\mathcal{A}_d, \quad \mathcal{A}_{3d}, \quad \mathcal{A}_{5d}, \quad \cdots, \quad \mathcal{A}_{n-d},$$
 (21)

Case 2.:  $\frac{n}{d} \equiv 0 \mod 4$ :

$$\mathcal{A}_0, \quad \mathcal{A}_{2d}, \quad \mathcal{A}_{4d}, \quad \cdots, \quad \mathcal{A}_n.$$
 (22)

By Proposition 5.4 these are all universal invariants of degree 1. We note that there are no invariants of degree 1 for  $q^{2d} \neq 1$  and for  $q^2 = 1$ .

### 5.5 Polarisation

Let f be the n-form

$$f = \sum_{i=0}^{n} \begin{bmatrix} n\\i \end{bmatrix} x^{n-i} y^{i} A_{i}$$
(23)

with  $A_i \in H_{I \setminus \{0,p\}}$  and  $k \in \mathbb{N}$ . We define the k-th polar  $f^{(k)}$  of f by

$$f^{(k)} := (P_{0p})^k f.$$

From this definition we infer the properties  $f^{(n)} \in H_{I \setminus \{0\}}, f^{(k)} = 0$  for k > n and

$$(f^{(k)})^{(l)} = (f^{(l)})^{(k)} = f^{(k+l)}$$

for  $k, l \in \mathbb{N}$ .

**PROPOSITION 5.5** Let f be the n-form (23). We can represent f in the form

$$f = \sum_{i=0}^{n} \begin{bmatrix} n\\i \end{bmatrix} A_i^- x^{n-i} y^i A_i^+$$

with  $A_i^- \in H_{\{i<0\}}$  and  $A_i^+ \in H_{\{i>0\}}$ . Furthermore let  $C_i^{n,k} \in H_{\{0,1\}}$  be the elements which are uniquely determined by the PBW expansion

$$\sum_{i=0}^{n} C_i^{n,k} A_i := (02)^{n-k} (12)^k.$$

with  $A_i \in H_{\{2\}}$ . Then we find for the k-th polar of f

$$f^{(k)} = \sum_{i=0}^{n} A_i^- C_i^{n,k} A_i^+.$$

Example 5.5 Let p = 1. For the quadratic form

$$f = x_0^2 A + [2]x_0 y_0 B + y_0^2 C$$

with  $A, B, C \in H_{\{i \in I | i > 1\}}$ , Lemma 3.3(i) implies that the first polar is

$$f^{(1)} = x_0(x_1A + qy_1B) + y_0(x_1B + y_1C)$$

For f = q(02)(03) we obtain the linear form

$$f^{(1)} = qx_0x_1y_2y_3 - \frac{1}{[2]}(q^2x_0y_1(qx_2y_3 + y_2x_3) + qy_0x_1(qx_2y_3 + y_2x_3)) + q^3y_0y_1x_2x_3$$
$$= \frac{1}{[2]}((02)(13) + (12)(03)).$$

Example 5.6 Let p = 1. For the cubic form

$$f = x_0^3 A + [3]x_0^2 y_0 B + [3]x_0 y_0^2 C + y_0^3 D$$

with  $A, B, C, D \in H_{\{i \in I | i > 1\}}$  we find

$$f^{(1)} = x_0^2 x_1 (A + (q^2 x_0^2 y_1 + (q^{-1} + q) x_0 y_0 x_1) B + ((1 + q^2) x_0 y_0 y_1 + y_0^2 x_1) C + y_0^2 y_1 D$$
  
=  $x_0^2 (x_1 A + q^2 y_1 B) + x_0 y_0 [2] (x_1 B + q y_1 C) + y_0^2 (x_1 C + y_1 D)$ 

and

$$f^{(2)} = x_0 x_1^2 A + ((1+q^2)x_0 x_1 y_1 + y_0 x_1^2) B + (q^2 x_0 y_1^2 + (q^{-1}+q)y_0 y_1 x_1) C + y_0 y_1^2 D$$
  
=  $x_0 (x_1^2 A + q[2]x_1 y_1 B + q^2 y_1^2 C) + y_0 (x_1^2 B + [2]y_1 x_1 C + y_1^2 D).$ 

For  $f = q^3(01)(02)(03)$  we obtain the quadratic form

$$f^{(1)} = \frac{q}{[3]}((02)(03)(14) + (02)(13)(04) + (12)(03)(04))$$

and the linear form

$$f^{(2)} = \frac{q}{[3]}((02)(13)(14) + (12)(03)(14) + (12)(13)(04)).$$

# 6. APPLICATION OF THE SYMBOLIC METHOD

### 6.1 Linear Forms

We consider the universal linear form

$$\mathcal{F} = x\mathcal{A} + y\mathcal{B}$$

In the classical case the linear form  $\mathcal{F}$  has no invariants. For the realisations f = xA + yB := (0i) (i = 1, 2) we have  $A = q^{-\frac{1}{2}}y_i$  and  $B = -q^{\frac{1}{2}}x_i$ . Therefore, for the symbol  $(12) = q^{-\frac{1}{2}}x_1y_2 - q^{\frac{1}{2}}y_1x_2$  we obtain the universal invariant

$$\mathcal{I}_1 = -q^{-\frac{1}{2}}\mathcal{B}\mathcal{A} + q^{\frac{1}{2}}\mathcal{A}\mathcal{B}.$$

Example 6.1 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ . We consider the simplest realisation f = (0i),  $i \neq 0$  (cf. Example 4.1). For an arbitrary universal invariant  $\mathcal{I}$  we obtain the realisation  $I_f = 0$ .

*Proof.* Since each universal invariant  $\mathcal{I}$  is generated by a symbol,  $\mathcal{I}$  is homogeneous with respect to  $\mathcal{A}$  and  $\mathcal{B}$  of the same degree. Therefore we find  $I_f = c x_i^k y_i^k$ ,  $k \ge 1$  with  $c \in \mathbb{C}$ . The assertion follows from the formulas (3). •

The next Example demonstrates, that there are nontrivial realisations of universal invariants for linear forms.

Example 6.2 For the realisation f = (01)(23) - c(03)(12) (cf. Example 4.2) we obtain

$$I_{1,f} = c(q^3 - q^5)(12)(13)(23) \neq 0,$$

(cf. formula (16)). One can show, that all invariants of f have the form  $\alpha I_{1,f}^k$  with  $\alpha \in \mathbb{C}$  and  $k \in \mathbb{N}$ . The proof is similar to the proof of Proposition 6.3 (cf. below).

A main task of the classical and the modern representation theory is to find finite generating sets of the algebras of invariants of forms. The following proposition shows, that there is no finite generating set for universal invariants even in the case of the universal linear form. In this article we give finitness theorems for realisations of universal forms only for simple Examples (cf. the Examples 6.1 and 6.2 above and the Propositions 6.2, 6.3 below).

PROPOSITION 6.1 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$ . The algebra of universal invariants of the universal linear form has not a finite set of generating elements.

*Proof.* Consider the sequence of symbols

$$(1, 2n)(2, 2n-1)(3, 2n-2)\cdots(n, n+1)$$

with  $n = 1, 2, \dots$ . By Lemma 3.3 (iv) we obtain by the symbolic method the sequence of universal invariants

$$\mathcal{I}_{n} = \sum_{i_{1}, i_{2}, \cdots, i_{n} = 0}^{1} -q^{\frac{3}{2}n(n-1)}(-q)^{i_{1}+i_{2}+\cdots+i_{n}}\mathcal{U}^{(i_{1})}\mathcal{U}^{(i_{2})}\cdots\mathcal{U}^{(i_{n})}\mathcal{U}^{(1-i_{n})}\mathcal{U}^{(1-i_{n-1})}\cdots\mathcal{U}^{(1-i_{1})}$$

of degree 2n with  $\mathcal{U}^{(0)} := \mathcal{B}$  and  $\mathcal{U}^{(1)} := \mathcal{A}$ . The  $\mathcal{I}_n$  are elements of the free algebra  $F_I$ . Since each universal invariant  $\mathcal{I}$  is generated by a symbol,  $\mathcal{I}$  is a linear combination of monomials, which are homogeneous with respect to  $\mathcal{A}$  and  $\mathcal{B}$  of degree  $k \in \mathbb{N}$ . Because  $\mathcal{I}_n$  contains the monomial  $-q^{\frac{3}{2}n(n-1)}\mathcal{B}^n\mathcal{A}^n$ , it is impossible to express  $\mathcal{I}_n$  in terms of universal invariants  $\mathcal{I}$  of degree 2k with k < n. The assertion follows  $\bullet$ .

#### 6.2 Quadratic forms

In Example 5.4 we obtained for the symbol  $d := 1/q(12)^2$  by the symbolic method the universal invariant

$$\mathcal{I}_1 = \frac{1}{q^3} \mathcal{C} \mathcal{A} - \frac{[2]}{q} \mathcal{B} \mathcal{B} + \frac{1}{q} \mathcal{A} \mathcal{C}.$$

Example 6.3 In Example 5.1 we found for the realisation f = q(01)(02)

$$I_{1,f} = -\frac{1}{[2]}(12)^2.$$
(24)

For  $f = (01)^2$  we find  $I_{1,f} = 0$ .

Now we consider an invariant without a classical analog. Let  $\mathcal{I}_2$  be the invariant, which arises from the symbol  $-\frac{1}{q^4}(12)(13)(23)$ . We obtain

$$\mathcal{I}_{2} = q^{\frac{-5}{2}} \mathcal{ABC} - q^{\frac{-9}{2}} \mathcal{ACB} - q^{\frac{-9}{2}} \mathcal{BAC} + q^{\frac{-9}{2}} \mathcal{BCA} + q^{\frac{-9}{2}} \mathcal{CAB} - q^{\frac{-13}{2}} \mathcal{CBA} - (q^{\frac{-7}{2}} - q^{\frac{-15}{2}}) \mathcal{BBB}.$$

Remark 6.1  $\mathcal{I}_2$  is associated to the combinant of three quadratic forms (cf. below).

For the realisation f = q(01)(02) we have

$$I_{2,f} = (q^2 - \frac{1}{q^2})(12)^3.$$
(25)

We note that  $I_{2,f} \neq 0$  if  $q^4 \neq 1$ .

Now we consider invariants of degree four. The fourth order symbols are generated by  $Z_1^2, Z_1Z_3, Z_3^2$  (c.f. Remark 3.1). We obtain by the symbolic method

$$\begin{split} \mathcal{I}_{3,1} &= q^8 \mathcal{A} \mathcal{C} \mathcal{A} \mathcal{C} - q^{10} [2] \mathcal{A} \mathcal{C} \mathcal{B} \mathcal{B} + q^6 \mathcal{A} \mathcal{C} \mathcal{C} \mathcal{A} - q^{10} [2] \mathcal{B} \mathcal{B} \mathcal{A} \mathcal{C} + q^{12} [2]^2 \mathcal{B} \mathcal{B} \mathcal{B} \mathcal{B} \\ &- q^8 [2] \mathcal{B} \mathcal{B} \mathcal{C} \mathcal{A} + q^6 \mathcal{C} \mathcal{A} \mathcal{A} \mathcal{C} - q^8 [2] \mathcal{C} \mathcal{A} \mathcal{B} \mathcal{B} + q^4 \mathcal{C} \mathcal{A} \mathcal{C} \mathcal{A}, \\ \\ \mathcal{I}_{3,2} &= q^{12} \mathcal{A} \mathcal{B} \mathcal{B} \mathcal{C} - q^{10} \mathcal{A} \mathcal{B} \mathcal{C} \mathcal{B} - q^7 \mathcal{A} \mathcal{C} \mathcal{A} \mathcal{C} + q^8 \mathcal{A} \mathcal{C} \mathcal{B} \mathcal{B} - q^{10} \mathcal{B} \mathcal{A} \mathcal{B} \mathcal{C} \\ &+ q^8 \mathcal{B} \mathcal{A} \mathcal{C} \mathcal{B} + q^8 \mathcal{B} \mathcal{B} \mathcal{A} \mathcal{C} - (q^9 + q^{15}) \mathcal{B} \mathcal{B} \mathcal{B} \mathcal{B} + q^{10} \mathcal{B} \mathcal{B} \mathcal{C} \mathcal{A} + q^{10} \mathcal{B} \mathcal{C} \mathcal{A} \mathcal{B} \\ &- q^8 \mathcal{B} \mathcal{C} \mathcal{B} \mathcal{A} + q^{10} \mathcal{C} \mathcal{A} \mathcal{B} \mathcal{B} - q^5 \mathcal{C} \mathcal{A} \mathcal{C} \mathcal{A} - q^8 \mathcal{C} \mathcal{B} \mathcal{A} \mathcal{B} + q^6 \mathcal{C} \mathcal{B} \mathcal{B} \mathcal{A}, \\ \\ \\ \mathcal{I}_{3,3} &= q^8 \mathcal{A} \mathcal{A} \mathcal{C} \mathcal{C} - q^{10} [2] \mathcal{A} \mathcal{B} \mathcal{B} \mathcal{C} + q^6 \mathcal{A} \mathcal{C} \mathcal{A} \mathcal{C} - q^{10} [2] \mathcal{B} \mathcal{A} \mathcal{C} \mathcal{B} + q^{12} [2]^2 \mathcal{B} \mathcal{B} \mathcal{B} \mathcal{B} \\ &- q^8 [2] \mathcal{B} \mathcal{C} \mathcal{A} \mathcal{B} + q^6 \mathcal{C} \mathcal{A} \mathcal{C} \mathcal{A} - q^8 [2] \mathcal{C} \mathcal{B} \mathcal{B} \mathcal{A} + q^4 \mathcal{C} \mathcal{C} \mathcal{A} \mathcal{A}. \end{split}$$

We have  $\mathcal{I}_{3,1} = q^{10}\mathcal{I}_1^2$ . Since  $F_I$  is a free algebra, the universal invariants  $\mathcal{I}_{3,1}$ ,  $\mathcal{I}_{3,2}$  and  $\mathcal{I}_{3,3}$  are independent.

We do not try to describe the algebraic structure of the algebra of universal invariants. Here we consider this subject for invariants of a certain realisation of a 2-form.

In the classical case the algebra of invariants is generated by the discriminant. However, if  $q^{2d} \neq 1$  we have two basic invariants for f = q(01)(02):

PROPOSITION 6.2 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$  and let f be the quadratic form q(01)(02). The algebra of invariants is generated by  $I_{1,f}$  and  $I_{2,f}$ . They are related by the equation

$$q^{2}[2]I_{2,f}^{2} + (q^{2} - 1)^{2}I_{1,f}^{3} = 0.$$
(26)

*Proof.* Let I be an invariant of f of degree k. Then we have  $I \in H^{Inv}_{\{1,2\}}$ . By Theorem 3.6 we have  $I = (12)^k$ . Since the symbol (12) is not a linear combination of A, B, C (cf. Example 4.3), we have  $k \ge 2$ . Let k = 2d + 3e,  $d, e \in \mathbb{N}$ . By (24) and (25) we have  $I = c I^d_{1,f} I^e_{2,f}$  with a certain constant  $c \in \mathbb{C}$ . The relation (26) follows from (24) and (25).

#### 6.3 Two and three quadratic forms

In Example 5.4 we obtained for the symbol  $\frac{1}{q}(12)^2$  the universal common invariant of two forms  $\mathcal{F}, \mathcal{F}'$ 

$$\mathcal{I}_{1,\mathcal{F},\mathcal{F}'} = \frac{1}{q^3} \mathcal{C} \mathcal{A}' - (1 + \frac{1}{q^2}) \mathcal{B} \mathcal{B}' + \frac{1}{q} \mathcal{A} \mathcal{C}'.$$

Example 6.4 The realisations  $I_{1,f,f'}$  and  $I_{1,f,f'}$  are in general not equal. For the realisation f = q(01)(03), f' = q(02)(04) we obtain

$$I_{1,f,f'} = \frac{1}{[2]} (q^5(12)(34) - \frac{1}{q}(14)(23))$$

and

$$I_{1,f',f} = \frac{1}{[2]}(q^3(12)(34) - q^3(14)(23)).$$

From the symbol (12)(01)(02) we obtain the common covariant of two forms  $\mathcal{F}, \mathcal{F}'$ 

$$\Delta_{\mathcal{F},\mathcal{F}'} = x^2 \mathcal{K} + xy \mathcal{L} + y^2 \mathcal{M}$$

with

$$\mathcal{K} = -q^7 \mathcal{B} \mathcal{A}' + q^9 \mathcal{A} \mathcal{B}',$$
  
 $\mathcal{L} = -q^7 \mathcal{C} \mathcal{A}' - q^6 \mathcal{B} \mathcal{B}' + q^{10} \mathcal{B} \mathcal{B}' + q^7 \mathcal{A} \mathcal{C}',$   
 $\mathcal{M} = -q^7 \mathcal{C} \mathcal{B}' + q^9 \mathcal{B} \mathcal{C}'.$ 

In the quantised case we have in general not the classical relation  $\Delta_{\mathcal{F},\mathcal{F}'} = -\Delta_{\mathcal{F}',\mathcal{F}}$ .

Example 6.5 Example 6.5: For f = f' = q(01)(02) we obtain

$$\Delta_{f,f} = (q^{\frac{23}{2}} - q^{\frac{15}{2}})(01)(02)(12).$$

 $\Delta_{f,f}$  has the discriminant

$$d_{f,f} = -q^{15}(q^2 - 1)^2 [2]^3 (12)^4.$$

This discriminant vanishes only for  $q^2 = 1$ .

Remark 6.2 In the classical situation the zeros of  $\Delta_{f,f'} = -\Delta_{f',f}$  are a point pair, which is harmonic to the zeros of f and to the zeros of f'. Furthermore the zeros of  $\Delta_{f,f'}$ are the fixpoints of the projective involution, which is determined by the zeros of f and f'. We have two real zeros if the intervalls  $[v_1, v_3]$ ,  $[v_2, v_4]$  are disjoint or if there is an inclusion between both intervals. Otherwise we have two complex conjugated zeros. This corresponds to a construction of complex elements due to von Staudt. In the classical situation the discriminant  $d_{f,f'}$  of  $\Delta_{f,f'}$  is the resultant of f and f', i.e.  $d_{f,f'}$  vanishes in the case of a common zero of f and f'.

Finally we consider an invariant of three quadratic forms

$$f^{(i)} = x^2 A^{(i)} + [2]xyB^{(i)} + y^2C^{(i)},$$

i = 1, 2, 3. From the symbol  $-\frac{1}{q^4}(12)(13)(23)$  we obtain

$$\begin{aligned} \mathcal{I}_{f',f'',f'''} &= q^{-\frac{5}{2}} \mathcal{A}' \mathcal{B}'' \mathcal{C}''' - q^{-\frac{9}{2}} \mathcal{A}' \mathcal{C}'' \mathcal{B}''' - q^{-\frac{9}{2}} \mathcal{B}' \mathcal{A}'' \mathcal{C}''' + q^{-\frac{9}{2}} \mathcal{B}' \mathcal{C}'' \mathcal{A}''' + q^{-\frac{9}{2}} \mathcal{C}' \mathcal{A}'' \mathcal{B}''' - q^{-\frac{13}{2}} \mathcal{C}' \mathcal{B}'' \mathcal{A}''' \\ &- (q^{-\frac{7}{2}} - q^{-\frac{15}{2}}) \mathcal{B}' \mathcal{B}'' \mathcal{B}'''. \end{aligned}$$

 $\mathcal{I}_{f',f'',f'''}$  is a quantum analog of the classical *combinant* 

$$\begin{vmatrix} A' & A'' & A''' \\ B' & B'' & B''' \\ C' & C'' & C''' \end{vmatrix}.$$

In the classical situation the three point pairs corresponding to f', f'', f''' are in involution if and only if their combinant vanishes.

6.4 A point figuration on the real line

In this section we apply the invariants and covariants of the preceding section to a point figuration on the real line. In the classical case we have  $I_{1,f,f'} = I_{1,f',f}$  and the zeros of the two quadratic forms are two point pairs, which are harmonic if and only if  $I_{1,f,f'} = 0$ . Furthermore we have  $I_{f,f',f''} = I_{f'',f,f'} = I_{f',f'',f} = -I_{f,f'',f'} = -I_{f',f,f''} = -I_{f'',f',f}$  and the three point pairs determined by the three quadratic forms are in involution if  $I_{f,f',f''} = 0$ . And the zeros of the quadratic form  $\Delta_{f,f'} = -\Delta_{f',f}$  are a point pair, which is harmonic to the zeros of f and f'.

In the following we discuss a quantum analog for points, which are harmonic or in involution. We consider three points  $(x_i, y_i)$ , i = 1, 2, 3. We form the six quadratic forms

$$f_{11} = (01)(01), \quad f_{22} = (02)(02), \quad f_{33} = (03)(03),$$
  
 $f_{12} = q(01)(02), \quad f_{13} = q(01)(03), \quad f_{23} = q(02)(03).$ 

We can form the three quadratic forms

$$f_{11|23} := \Delta_{f_{11}, f_{23}} - \Delta_{f_{23}, f_{11}},$$
  

$$f_{22|13} := \Delta_{f_{22}, f_{13}} - \Delta_{f_{13}, f_{22}},$$
  

$$f_{33|12} := \Delta_{f_{33}, f_{12}} - \Delta_{f_{12}, f_{33}}.$$

For  $f_{11|23}$  and the two quadratic forms  $f_{11}$ ,  $f_{23}$  we have the identities

$$I_{f_{11|23},f_{11}} = I_{f_{11},f_{11|23}} = 0,$$
  
$$q^4 I_{f_{11|23},f_{23}} + I_{f_{23},f_{11|23}} = 0.$$

They give the classical harmonicity of the corresponding forms in the classical limit. Similar results hold for  $f_{22|13}$  and  $f_{33|12}$ . We have

$$I_{f_{22|13},f_{22}} = I_{f_{22},f_{22|13}} = 0,$$
  

$$I_{f_{22|13},f_{13}} = I_{f_{13},f_{22|13}} = 0,$$
  

$$I_{f_{33|12},f_{33}} = I_{f_{33},f_{33|12}} = 0,$$
  

$$I_{f_{33|12},f_{12}} + q^4 I_{f_{12},f_{33|12}} = 0.$$

Next we consider the relation between  $f_{11|23}$ ,  $f_{22|13}$  and  $f_{33|12}$ . We find

$$I_{f_{11|23},f_{33|12},f_{22|13}} = -I_{f_{22|13},f_{11|23},f_{33|12}} = \frac{1}{q^4} I_{f_{22|13},f_{33|12},f_{11|23}} = -\frac{1}{q^4} I_{f_{33|12},f_{11|23},f_{22|13}}$$
$$= q^7 [2]^2 (q^4 + 1)(q^6 - 1)((12)(13)(23))^2$$

and furthermore

$$\begin{split} I_{f_{11|23},f_{22|13},f_{33|12}} &= 0, \qquad I_{f_{33|12},f_{22|13},f_{11|23}} = 0, \\ I_{f_{33|12},f_{11|23},f_{22|13}} + I_{f_{22|13},f_{33|12},f_{11|23}} &= 0, \\ I_{f_{11|23},f_{33|12},f_{22|13}} + I_{f_{22|13},f_{11|23},f_{33|12}} &= 0. \end{split}$$

The formulas were checked by a computer calculation. Each of these four equations gives the classical relation  $I_{f_{11|23},f_{22|13},f_{33|12}} = 0$  in the classical limit.

This is an analog of the following classical proposition. Let A, B, C be three points and let P, Q and R be the three fourth harmonic points of A, B and C with respect to the pair  $\{B, C\}, \{A, C\}$  and  $\{A, B\}$ , respectively. Then the three point pairs  $\{A, P\},$  $\{B, Q\}$  and  $\{C, R\}$  are in involution.

### 6.5 Cubic forms

We consider the universal cubic form

$$\mathcal{F} = x^3 \mathcal{A} + [3] x^2 y \mathcal{B} + [3] x y^2 \mathcal{C} + y^3 \mathcal{D}.$$

By Lemma 3.3(ii) we have

$$(ij)^3 = q^{\frac{3}{2}} x_i^3 y_j^3 - q^{\frac{1}{2}} [3] x_i^2 y_i x_j y_j^2 + q^{\frac{3}{2}} [3] x_i y_i^2 x_j^2 y_j - q^{\frac{9}{2}} y_i^3 x_j^3.$$

Applying this formula to  $f = (0i)^3$ , i > 0, we obtain

$$A = q^{\frac{3}{2}}y_i^3, \qquad B = -q^{\frac{1}{2}}x_iy_i^2, \qquad C = q^{\frac{3}{2}}x_i^2y_i, \qquad D = -q^{\frac{9}{2}}x_i^3.$$

The simplest invariant  $\mathcal{I}_1$  arises from the symbol  $\frac{1}{q^5}(ij)^3$ . The symbolic method gives

$$\mathcal{I}_1 = \left(-q^{\frac{-19}{2}}\mathcal{D}\mathcal{A} + q^{\frac{-13}{2}}\mathcal{A}\mathcal{D}\right) + [3](q^{\frac{-13}{2}}\mathcal{C}\mathcal{B} - q^{\frac{-11}{2}}\mathcal{B}\mathcal{C}).$$

Example 6.6 For the realisation  $f = q^3(01)(02)(03)$  we obtain

$$I_{1,f} = (q^{-3} - q^3)(12)(13)(23).$$
(27)

We have  $I_{1,f} \neq 0$  for  $q^6 \neq 1$ . For the realisations  $f = q^2(01)(01)(02)$ ,  $f = q^2(01)(02)(02)$ and f = (01)(01)(01) we obtain  $I_{1,f} = 0$ . Therefore  $\mathcal{I}_1$  is an invariant of degree two with the characterising property of the discriminant. It is well known that for the classical cubic form it is impossible to express the root of the discriminant (12)(13)(23) in terms of the coefficients.

There are no symbols, which generate invariants of degree three. By means of the Grassmann-Plücker relations (cf. Proposition 3.2(iii)) we can reduce the symbols for invariants of degree four to the following four symbols

$$Z_1^3, \quad Z_1^2 Z_3, \quad Z_1 Z_3^2, \quad Z_3^3$$
 (28)

(cf. Remark 3.1). The symbols (29) can be expressed by PBW basis elements using a computer calculation. Subsequently we can apply the symbolic method to obtain the following universal invariants  $\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3, \mathcal{D}_4$ :

$$\mathcal{D}_{1} := q^{24} \mathcal{A} \mathcal{D} \mathcal{A} \mathcal{D} - q^{21} \mathcal{A} \mathcal{D} \mathcal{D} \mathcal{A} - q^{21} \mathcal{D} \mathcal{A} \mathcal{A} \mathcal{D} + q^{18} \mathcal{D} \mathcal{A} \mathcal{D} \mathcal{A}$$
$$+ [3] (-q^{25} \mathcal{A} \mathcal{D} \mathcal{B} \mathcal{C} + q^{24} \mathcal{A} \mathcal{D} \mathcal{C} \mathcal{B} - q^{25} \mathcal{B} \mathcal{C} \mathcal{A} \mathcal{D} + q^{22} \mathcal{B} \mathcal{C} \mathcal{D} \mathcal{A}$$
$$+ q^{22} \mathcal{C} \mathcal{B} \mathcal{A} \mathcal{D} - q^{19} \mathcal{C} \mathcal{B} \mathcal{D} \mathcal{A} + q^{20} \mathcal{D} \mathcal{A} \mathcal{B} \mathcal{C} - q^{19} \mathcal{D} \mathcal{A} \mathcal{C} \mathcal{B})$$
$$+ (1 + 2q^{2} + 3q^{4} + 2q^{6} + q^{8}) (q^{22} \mathcal{B} \mathcal{C} \mathcal{B} \mathcal{C} - q^{21} \mathcal{B} \mathcal{C} \mathcal{C} \mathcal{B} - q^{21} \mathcal{C} \mathcal{B} \mathcal{B} \mathcal{C} + q^{20} \mathcal{C} \mathcal{B} \mathcal{C} \mathcal{B})$$

$$\begin{split} \mathcal{D}_{2} &:= -q^{19} \mathcal{D} \mathcal{A} \mathcal{D} \mathcal{A} - q^{23} \mathcal{A} \mathcal{D} \mathcal{A} \mathcal{D} \\ + [2](-q^{27} \mathcal{A} \mathcal{C} \mathcal{C} \mathcal{C} - q^{23} \mathcal{C} \mathcal{A} \mathcal{C} \mathcal{C} - q^{25} \mathcal{C} \mathcal{C} \mathcal{A} \mathcal{C} - q^{21} \mathcal{C} \mathcal{C} \mathcal{C} \mathcal{A}) \\ + q^{28} \mathcal{A} \mathcal{C} \mathcal{B} \mathcal{D} + q^{24} \mathcal{A} \mathcal{C} \mathcal{D} \mathcal{B} + q^{23} [2] \mathcal{A} \mathcal{D} \mathcal{B} \mathcal{C} - q^{21} \mathcal{A} \mathcal{D} \mathcal{C} \mathcal{B} \\ + q^{23} [2] \mathcal{B} \mathcal{C} \mathcal{A} \mathcal{D} - q^{25} \mathcal{B} \mathcal{C} \mathcal{D} \mathcal{A} + q^{26} \mathcal{B} \mathcal{D} \mathcal{A} \mathcal{C} + q^{22} \mathcal{B} \mathcal{D} \mathcal{C} \mathcal{A} \\ + q^{24} \mathcal{C} \mathcal{A} \mathcal{B} \mathcal{D} + q^{20} \mathcal{C} \mathcal{A} \mathcal{D} \mathcal{B} - q^{21} \mathcal{C} \mathcal{B} \mathcal{A} \mathcal{D} + q^{23} [2] \mathcal{C} \mathcal{B} \mathcal{D} \mathcal{A} \\ - q^{25} \mathcal{D} \mathcal{A} \mathcal{B} \mathcal{C} + q^{23} [2] \mathcal{D} \mathcal{A} \mathcal{C} \mathcal{B} + q^{22} \mathcal{D} \mathcal{B} \mathcal{A} \mathcal{C} + q^{18} \mathcal{D} \mathcal{B} \mathcal{C} \mathcal{A} \\ + [2](-q^{27} \mathcal{B} \mathcal{B} \mathcal{B} \mathcal{D} - q^{23} \mathcal{B} \mathcal{B} \mathcal{D} \mathcal{B} - q^{25} \mathcal{B} \mathcal{D} \mathcal{B} \mathcal{B} - q^{21} \mathcal{D} \mathcal{B} \mathcal{B} \mathcal{B}) \\ + q^{26} [2]^2 \mathcal{B} \mathcal{B} \mathcal{C} \mathcal{C} - q^{21} (1 + 2q^2 + q^4 + q^{10}) \mathcal{B} \mathcal{C} \mathcal{B} \mathcal{C} + q^{21} [2] (1 + q^8) \mathcal{B} \mathcal{C} \mathcal{C} \mathcal{B} \\ + q^{21} [2] (1 + q^8) \mathcal{C} \mathcal{B} \mathcal{B} \mathcal{C} - q^{19} (1 + q^6 + 2q^8 + q^{10}) \mathcal{C} \mathcal{B} \mathcal{C} \mathcal{B} + q^{24} [2]^2 \mathcal{C} \mathcal{C} \mathcal{B} \mathcal{B}, \end{split}$$

$$\begin{split} \mathcal{D}_{3} &:= q^{22}\mathcal{A}\mathcal{D}\mathcal{A}\mathcal{D} + q^{20}\mathcal{D}\mathcal{A}\mathcal{D}\mathcal{A} \\ &+ [2](q^{24}\mathcal{A}\mathcal{C}\mathcal{C}\mathcal{C} + q^{26}\mathcal{C}\mathcal{A}\mathcal{C}\mathcal{C} + q^{22}\mathcal{C}\mathcal{C}\mathcal{A}\mathcal{C} + q^{24}\mathcal{C}\mathcal{C}\mathcal{C}\mathcal{A}) \\ &+ q^{28}\mathcal{A}\mathcal{B}\mathcal{C}\mathcal{D} - q^{25}\mathcal{A}\mathcal{B}\mathcal{D}\mathcal{C} - q^{26}[2]\mathcal{A}\mathcal{C}\mathcal{B}\mathcal{D} - q^{21}\mathcal{A}\mathcal{D}\mathcal{B}\mathcal{C} \\ &- q^{25}\mathcal{B}\mathcal{A}\mathcal{C}\mathcal{D} + q^{22}\mathcal{B}\mathcal{A}\mathcal{D}\mathcal{C} - q^{21}\mathcal{B}\mathcal{C}\mathcal{A}\mathcal{D} - q^{24}[2]\mathcal{B}\mathcal{D}\mathcal{A}\mathcal{C} \\ &- q^{22}[2]\mathcal{C}\mathcal{A}\mathcal{D}\mathcal{B} - q^{25}\mathcal{C}\mathcal{B}\mathcal{D}\mathcal{A} + q^{24}\mathcal{C}\mathcal{D}\mathcal{A}\mathcal{B} - q^{21}\mathcal{C}\mathcal{D}\mathcal{B}\mathcal{A} \\ &- q^{25}\mathcal{D}\mathcal{A}\mathcal{C}\mathcal{B} - q^{20}[2]\mathcal{D}\mathcal{B}\mathcal{C}\mathcal{A} - q^{21}\mathcal{D}\mathcal{C}\mathcal{A}\mathcal{B} + q^{18}\mathcal{D}\mathcal{C}\mathcal{B}\mathcal{A} \\ &+ [2](+q^{24}\mathcal{B}\mathcal{B}\mathcal{B}\mathcal{D} + q^{26}\mathcal{B}\mathcal{B}\mathcal{D}\mathcal{B} + q^{22}\mathcal{B}\mathcal{D}\mathcal{B}\mathcal{B} + q^{24}\mathcal{D}\mathcal{B}\mathcal{B}\mathcal{B}) \\ &- q^{22}[2](1+q^8)\mathcal{B}\mathcal{B}\mathcal{C}\mathcal{C} + q^{20}(1+q^6+2q^8+q^{10})\mathcal{B}\mathcal{C}\mathcal{B}\mathcal{C} - q^{25}[2]^2\mathcal{B}\mathcal{C}\mathcal{C}\mathcal{B} \\ &- q^{25}[2]^2\mathcal{C}\mathcal{B}\mathcal{B}\mathcal{C} + q^{20}(1+2q^2+q^4+q^{10})\mathcal{C}\mathcal{B}\mathcal{C}\mathcal{B} - q^{20}[2](1+q^8)\mathcal{C}\mathcal{C}\mathcal{B}\mathcal{B}, \end{split}$$

$$\begin{split} \mathcal{D}_4 &:= q^{24} \mathcal{A} \mathcal{A} \mathcal{D} \mathcal{D} - q^{21} \mathcal{A} \mathcal{D} \mathcal{A} \mathcal{D} - q^{21} \mathcal{D} \mathcal{A} \mathcal{D} \mathcal{A} + q^{18} \mathcal{D} \mathcal{D} \mathcal{A} \mathcal{A} \\ & [3] (-q^{25} \mathcal{A} \mathcal{B} \mathcal{C} \mathcal{D} + q^{24} \mathcal{A} \mathcal{C} \mathcal{B} \mathcal{D} - q^{25} \mathcal{B} \mathcal{A} \mathcal{D} \mathcal{C} + q^{22} \mathcal{B} \mathcal{D} \mathcal{A} \mathcal{C} \\ & + q^{22} \mathcal{C} \mathcal{A} \mathcal{D} \mathcal{B} - q^{19} \mathcal{C} \mathcal{D} \mathcal{A} \mathcal{B} + q^{20} \mathcal{D} \mathcal{B} \mathcal{C} \mathcal{A} - q^{19} \mathcal{D} \mathcal{C} \mathcal{B} \mathcal{A}) \\ & + (1 + 2q^2 + 3q^4 + 2q^6 + q^8) (q^{22} \mathcal{B} \mathcal{B} \mathcal{C} \mathcal{C} - q^{21} \mathcal{B} \mathcal{C} \mathcal{B} \mathcal{C} - q^{21} \mathcal{C} \mathcal{B} \mathcal{C} \mathcal{B} + q^{20} \mathcal{C} \mathcal{C} \mathcal{B} \mathcal{B}). \end{split}$$

Remark 6.3 In the classical situation the discriminant

$$d = -a^2d^2 + 3b^2c^2 - 4b^3d - 4ac^3 + 6abcd$$

of the cubic form  $ax^3 + 3bx^2y + 3cxy^2 + dy^3$  generates the algebra of invariants (cf. [1]). If q is not a root of unity we get a four-dimensional space of invariants of degree four, generated by  $\mathcal{D}_1$ ,  $\mathcal{D}_2$ ,  $\mathcal{D}_3$  and  $\mathcal{D}_4$ . One can show that each element of this space vanishes for  $f = q^2(01)(01)(02)$ ,  $f = q^2(01)(02)(02)$  and f = (01)(01)(01). Therefore we have a four-dimensional space of invariants with the characterising property of the discriminant. We have  $\mathcal{D}_1 = q^{37}\mathcal{I}_1^2$ . In the classical case (i.e. for q = 1 and commuting  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ ) we have  $\mathcal{D}_2 = \mathcal{D}_3 = 2d$  and  $\mathcal{D}_1 = \mathcal{D}_4 = 0$ . Now we give a complete description of all invariants for the realisation  $f = q^3(01)(02)(03)$ of  $\mathcal{F}$ . The invariant  $\mathcal{I}_1$  plays in a certain sense the role of the root of the discriminant.

PROPOSITION 6.3 Let  $q^{2d} \neq 1$  for all  $d \in \mathbb{N}$  and let  $f = q^3(01)(02)(03)$ . Then the algebra of invariants is generated by  $I_{1,f}$ .

*Proof.* Let I be an invariant of degree k. Then we have  $I \in H_{\{1,2,3\}}^{Inv}$ . By Theorem 3.6, I is a linear combination of products  $(12)^{k_1}(13)^{k_2}(23)^{k_3}$  with  $k_1 + k_2 + k_3 = \frac{3}{2}k$ . Since I is homogeneous of degree k with respect to the indices 1, 2, 3, we have  $k_1 + k_2 = k_1 + k_3 = k_2 + k_3 = k$ . It follows that  $k_1 = k_2 = k_3 = \frac{k}{2}$ . By Proposition 3.2 (i) I is a linear combination of products

$$(12)^{\frac{k}{2}}(13)^{\frac{k}{2}}(23)^{\frac{k}{2}} = c((12)(13)(23))^{\frac{k}{2}} = c' \quad I_{1,f}^{\frac{k}{2}}$$

with certain nonvanishing constants  $c, c' \in \mathbb{C}$ , which proves the assertion.•

6.6 The quadratic covariant  $\Delta$ 

From the symbol  $(12)^2(01)(02)$  we derive the quadratic covariant

$$\Delta = x^2 \mathcal{K} + xy \mathcal{L} + y^2 \mathcal{M}$$

for the universal cubic form with

$$\mathcal{K} = q^{9} \mathcal{C} \mathcal{A} - q^{12} [2] \mathcal{B} \mathcal{B} + q^{13} \mathcal{A} \mathcal{C},$$
$$\mathcal{L} = q^{9} \mathcal{D} \mathcal{A} - (-q^{8} + q^{12} + q^{14}) \mathcal{C} \mathcal{B} - (q^{9} + q^{11} - q^{15}) \mathcal{B} \mathcal{C} + q^{10} \mathcal{A} \mathcal{D},$$
$$\mathcal{M} = q^{9} \mathcal{D} \mathcal{B} - q^{12} [2] \mathcal{C} \mathcal{C} + q^{13} \mathcal{B} \mathcal{D}.$$

For example for  $f = q^3(01)(02)(03)$  we have

$$\Delta_f = -q^{\frac{31}{2}}[2](01)(02)(13)(23) + q^{\frac{21}{2}}(1+q^8)(01)(03)(12)(23) - q^{\frac{23}{2}}[2](02)(03)(12)(13).$$

The quadratic form  $\Delta$  has the discriminant  $\mathcal{I}_{\Delta} := [2]\mathcal{M}\mathcal{K} - q^2\mathcal{L}\mathcal{L} + q^2[2]\mathcal{K}\mathcal{M}$  with

$$\begin{split} \mathcal{I}_{\Delta} &= -q^{20}(q^2\mathcal{A}\mathcal{D}\mathcal{A}\mathcal{D} + q\mathcal{A}\mathcal{D}\mathcal{D}\mathcal{A} + \mathcal{D}\mathcal{A}\mathcal{D}\mathcal{A} + q\mathcal{D}\mathcal{A}\mathcal{A}\mathcal{D}) \\ &-q^{21}[2]^2(q^6\mathcal{A}\mathcal{CCC} + q^2\mathcal{C}\mathcal{ACC} + q^4\mathcal{C}\mathcal{C}\mathcal{AC} + \mathcal{CCCA}) + q^{24}[2](q^4\mathcal{A}\mathcal{C}\mathcal{B}\mathcal{D} + \mathcal{A}\mathcal{C}\mathcal{D}\mathcal{B}) \\ &+ (q^{20} + q^{22} - q^{26})(q\mathcal{A}\mathcal{D}\mathcal{B}\mathcal{C} + \mathcal{D}\mathcal{A}\mathcal{B}\mathcal{C} + q\mathcal{B}\mathcal{C}\mathcal{A}\mathcal{D} + \mathcal{B}\mathcal{C}\mathcal{D}\mathcal{A}) \\ &+ q^{20}[2](q^6\mathcal{B}\mathcal{D}\mathcal{A}\mathcal{C} + q^2\mathcal{B}\mathcal{D}\mathcal{C}\mathcal{A} + q^4\mathcal{C}\mathcal{A}\mathcal{B}\mathcal{D} + \mathcal{C}\mathcal{A}\mathcal{D}\mathcal{B}) \\ &+ (-q^{19} + q^{23} + q^{25})(q\mathcal{C}\mathcal{B}\mathcal{A}\mathcal{D} + \mathcal{C}\mathcal{B}\mathcal{D}\mathcal{A} + q\mathcal{A}\mathcal{D}\mathcal{C}\mathcal{B} + \mathcal{D}\mathcal{A}\mathcal{C}\mathcal{B}) + q^{18}[2](q^4\mathcal{D}\mathcal{B}\mathcal{A}\mathcal{C} + \mathcal{D}\mathcal{B}\mathcal{C}\mathcal{A}) \\ &- q^{21}[2]^2(q^6\mathcal{B}\mathcal{B}\mathcal{B}\mathcal{D} + q^2\mathcal{B}\mathcal{B}\mathcal{D}\mathcal{B} + q^4\mathcal{B}\mathcal{D}\mathcal{B}\mathcal{B} + \mathcal{D}\mathcal{B}\mathcal{B}\mathcal{B}) + q^{24}[2]^3(q^2\mathcal{B}\mathcal{B}\mathcal{C}\mathcal{C} + \mathcal{C}\mathcal{C}\mathcal{B}\mathcal{B}) \\ &+ (-q^{20} - 2q^{22} - q^{24} + 2q^{26} + 2q^{28} - q^{32})\mathcal{B}\mathcal{C}\mathcal{B}\mathcal{C} \\ &+ (-q^{18} + 2q^{22} + 2q^{24} - q^{26} - 2q^{28} - q^{30})\mathcal{C}\mathcal{B}\mathcal{C}\mathcal{B} \\ &+ (q^{19} + q^{21} - q^{23} - 3q^{25} - q^{27} + q^{29} + q^{31})\mathcal{C}\mathcal{B}\mathcal{B}\mathcal{C}. \end{split}$$

Remark 6.4 We have  $\mathcal{I}_{\Delta} = \mathcal{D}_1 + [2]\mathcal{D}_2$ . Therefore we can distinguish the element  $\mathcal{I}_{\Delta}$  in our four-dimensional space of discriminants. In the classical case we have  $\mathcal{I}_{\Delta} = 2\mathcal{D}_2$ .

Remark 6.5 Let  $f = q^3(01)(02)(03)$  be a cubic form "with three real zeros". We form the realisation  $\Delta_f$  of  $\Delta$ . We can consider  $\Delta_f$  as an example of a quantised real quadratic form with two complex conjugated zeros. The form  $\Delta_f$  is related to a projective construction of complex elements due to F. Klein (cf. [20]). In the classical case the two complex conjugated zeros are those two unique complex points, from which one sees the intervals, determined by the three real zeros of f under an angle of  $\frac{\pi}{3}$ .

### 6.7 Quartic forms

We consider the universal quartic form

$$\mathcal{F} = x^4 \mathcal{A} + [4]x^3 y \mathcal{B} + \frac{[3][4]}{[2]} x^2 y^2 \mathcal{C} + [4]x y^3 \mathcal{D} + y^4 \mathcal{E}.$$

By Lemma 3.3 (ii) we have

$$(ij)^{4} = q^{4}x_{i}^{4}y_{j}^{4} - q^{2}[4]x_{i}^{3}y_{i}x_{j}y_{j}^{3} + q^{4}\begin{bmatrix} 4\\2 \end{bmatrix} x_{i}^{2}y_{i}^{2}x_{j}^{2}y_{j}^{2} - q^{4}[4]x_{i}y_{i}^{3}x_{j}^{3}y_{j} + q^{8}y_{i}^{4}x_{j}^{4}.$$
 (29)

Applying this formula to the realisations  $f^{(i)} = (0i)^4, i \neq 0$  we have

$$\begin{aligned} A^{(i)} &:= q^4 y_i^4, \qquad B^{(i)} := -q^2 x_i y_i^3, \qquad C^{(i)} := q^2 x_i^2 y_i^2, \\ D^{(i)} &:= -q^4 x_i^3 y_i, \qquad E^{(i)} := q^8 x_i^4. \end{aligned}$$

We consider the two symbols

$$I_1 := \frac{1}{q^6} (12)^4$$
 and  $I_2 := \frac{1}{q^{12}} (12)^2 (13)^2 (23)^2$ .

We can use formula (30) in order to construct the universal invariant  $\mathcal{I}_1$  for the symbol  $I_1$ , we obtain

$$I_{1} = q^{-6} (x_{1}^{4} A^{(2)} + x_{1}^{3} y_{1}[4] B^{(2)} + x_{1}^{2} y_{1}^{2} \frac{[3][4]}{[2]} C^{(2)} + x_{1}^{1} y_{1}^{3}[4] D^{(2)} + y_{1}^{4} E^{(2)})$$
  
$$= q^{-6} (q^{-4} A^{(1)} E^{(2)} - q^{-2}[4] B^{(1)} D^{(2)} + q^{-2} \frac{[3][4]}{[2]} C^{(1)} C^{(2)} - q^{-4}[4] D^{(1)} B^{(2)} + q^{-8} E^{(1)} A^{(2)}).$$

Therefore we have by the symbolic method

$$\mathcal{I}_{1} = q^{-10}\mathcal{A}\mathcal{E} - q^{-8}[4]\mathcal{B}\mathcal{D} + q^{-8}\frac{[3][4]}{[2]}\mathcal{C}\mathcal{C} - q^{-10}[4]\mathcal{D}\mathcal{B} + q^{-14}\mathcal{E}\mathcal{A}.$$

This is an analog of the invariant

$$2(ae - 4bd + 3c^2)$$

of the classical quartic form  $ax^4 + 4bx^3y + 6cx^2y^2 + 4dxy^3 + ey^4$ . In the classical case  $\mathcal{I}_1$  vanishes, if the cross ratio of the four zeros is one of the two third roots of unity  $exp(\pm \frac{2\pi i}{3})$ , i.e the four points are equiharmonic.

Now we construct the invariant for the symbol  $I_2 = q^{-12}(12)^2(13)^2(23)^2$ .

$$q^{12}I_2 = (12)^2 (13)^2 (23)^2$$

$$=q^{4}x_{1}x_{1}x_{1}x_{1}x_{2}x_{2}y_{2}y_{2}y_{3}y_{3}y_{3}y_{3}$$

$$-(q^{5}+q^{7})x_{1}x_{1}x_{1}x_{1}x_{2}y_{2}y_{2}y_{2}x_{3}y_{3}y_{3}y_{3}$$

$$+q^{10}x_{1}x_{1}x_{1}x_{1}y_{2}y_{2}y_{2}y_{2}x_{3}x_{3}y_{3}y_{3}$$

$$-(q^{5}+q^{7})x_{1}x_{1}x_{1}y_{1}x_{2}x_{2}x_{2}y_{2}y_{3}y_{3}y_{3}y_{3}$$

$$+(-1/q-q+q^{5}+2q^{7}+q^{9})x_{1}x_{1}x_{1}y_{1}x_{2}x_{2}y_{2}y_{2}x_{3}x_{3}y_{3}y_{3}y_{3}$$

$$+(q+2q^{3}+q^{5}-q^{9}-q^{11})x_{1}x_{1}x_{1}y_{1}x_{2}y_{2}y_{2}y_{2}x_{3}x_{3}y_{3}y_{3}$$

$$-(+q^{7}+q^{9})x_{1}x_{1}x_{1}y_{1}y_{2}y_{2}y_{2}y_{2}x_{3}x_{3}y_{3}y_{3}$$

$$+(q+2q^{3}+q^{5}-q^{9}-q^{11})x_{1}x_{1}y_{1}y_{1}x_{2}x_{2}y_{2}y_{2}x_{3}x_{3}y_{3}y_{3}$$

$$+(q^{2}-q^{2}-3q^{4}-3q^{6}-q^{8}+q^{12})x_{1}x_{1}y_{1}y_{1}x_{2}x_{2}y_{2}y_{2}x_{3}x_{3}y_{3}y_{3}$$

$$+(-q-q^{3}+q^{7}+2q^{9}+q^{11})x_{1}x_{1}y_{1}y_{1}x_{2}x_{2}y_{2}y_{2}x_{3}x_{3}x_{3}y_{3}$$

$$+(-q-q^{3}+q^{7}+2q^{9}+q^{11})x_{1}y_{1}y_{1}y_{1}x_{2}x_{2}y_{2}y_{2}x_{3}x_{3}x_{3}y_{3}$$

$$+(q^{3}+2q^{5}+q^{7}-q^{11}-q^{13})x_{1}y_{1}y_{1}y_{1}x_{2}x_{2}y_{2}y_{2}x_{3}x_{3}x_{3}y_{3}$$

$$-(q^{9}+q^{11})x_{1}y_{1}y_{1}y_{1}x_{2}x_{2}x_{2}y_{2}x_{3}x_{3}x_{3}y_{3}$$

$$-(q^{9}+q^{11})y_{1}y_{1}y_{1}y_{2}x_{2}x_{2}y_{2}x_{3}x_{3}x_{3}x_{3}$$

We obtain by the symbolic method the universal invariant

$$\begin{aligned}
\mathcal{I}_{2} &= q^{-12} \mathcal{ACE} + q^{-18} \mathcal{AEC} + q^{-18} \mathcal{CAE} \\
&+ q^{-16} \mathcal{CEA} + q^{-16} \mathcal{EAC} + q^{-22} \mathcal{ECA} \\
&+ q^{-19} (1 + 2q^{2} + q^{4} - q^{8} - q^{10}) (q^{2} \mathcal{BCD} + \mathcal{CDB} + \mathcal{DBC}) \\
&+ q^{-21} (-1 - q^{2} + q^{6} + 2q^{8} + q^{10}) (q^{2} \mathcal{BDC} + q^{2} \mathcal{CBD} + \mathcal{DCB}) \\
&- q^{-18} [2] (q^{4} \mathcal{BBE} + q^{2} \mathcal{BEB} + \mathcal{EBB}) \\
&- q^{-18} [2] (q^{4} \mathcal{ADD} + q^{2} \mathcal{DAD} + \mathcal{DDA}) \\
&+ q^{-20} (1 - q^{4} - 3q^{6} - 3q^{8} - q^{10} + q^{14}) \mathcal{CCC}.
\end{aligned}$$
(30)

 $\mathcal{I}_2$  is an analog of the catalecticant

of the classical quartic form (cf. [3]). In the classical case  $\mathcal{I}_2$  vanishes, if the cross ratio of the four zeros takes one of the values  $-1, 2, \frac{1}{2}$ , i.e. the four points are harmonic.

In the following we will consider forms "with real zeros" as realisations of the universal quartic form. First we consider the form with different zeros

$$f = q^{6}(0i)(0j)(0k)(0l), \qquad 0 < i < j < k < l.$$
(31)

**PROPOSITION 6.4** The algebra of invariants of the form (31) is commutative.

*Proof.* Let I be an invariant of f of degree k. By Theorem 3.6 we can represent I by the bracket symbols  $(\alpha\beta)$ ,  $\alpha, \beta = i, j, k, l, \alpha < \beta$ . By Proposition 3.2(i) we can reduce I to a linear combination of monomials

$$(ij)^{k_{ij}}(kl)^{k_{kl}}(ik)^{k_{ik}}(jl)^{k_{jl}}(il)^{k_{il}}(jk)^{k_{jk}}.$$

We have

$$k_{ij} + k_{ik} + k_{il} = 2k$$
  

$$k_{ij} + k_{jk} + k_{jl} = 2k$$
  

$$k_{ik} + k_{jk} + k_{kl} = 2k$$
  

$$k_{il} + k_{jl} + k_{kl} = 2k.$$

If we add two of the equations and subtract the remaining two equations in different ways, we obtain  $k_{ij} = k_{kl}$ ,  $k_{ik} = k_{jl}$  and  $k_{il} = k_{jk}$ . Therefore we can represent I by a linear combination of monomials

$$(ij)^{k_{ij}}(kl)^{k_{ij}}(ik)^{k_{ik}}(jl)^{k_{ik}}(il)^{k_{il}}(jk)^{k_{il}}.$$

If we use again Proposition 3.2(i) we can represent I as a linear combination of monomials

$$((ij)(kl))^{k_{ij}+i}((ik)(jl))^{k_{ik}-i}((il)(jk))^{k_{il}}, \qquad i \in \mathbb{N}_0.$$

The assertion follows by Proposition 3.2(ii).

Next we express the invariants  $I_{1,f}$ ,  $I_{2,f}$  of the quartic form (32) in terms of the "zeros".

We insert  $A_i$  for  $\mathcal{A}_i$  in  $\mathcal{I}_1$  and express the resulting invariant in terms of the bracket symbols. By the proof of Proposition 6.4 and Remark 3.1 we can represent the result as a linear combination of  $Z_1^2$ ,  $Z_1Z_3$  and  $Z_3^2$ . By a computer calculation we obtain

$$I_{1,f} = \frac{1}{[3][4]}([2](Z_1^2 + Z_3^2) + (-q^{-4} + q^{-2} + 2 + q^2 - q^4)Z_1Z_3).$$

Similarly we obtain

$$I_{2,f} = \frac{[2]^2}{[3]^3[4]^3} (aZ_1^3 + bZ_1^2Z_3 + cZ_1Z_3^2 + dZ_3^3)$$

with

$$a = q^{-6}[2][3](1 + q^2 - q^4 - q^8 + q^{10} + q^{12}),$$
  

$$b = q^{-12}(-2 + 7q^4 + 14q^6 + 7q^8 - 8q^{10} - 18q^{12} - 8q^{14} + 7q^{16} + 14q^{18} + 7q^{20} - 2q^{24}),$$
  

$$c = q^{-12}[2](-1 - q^2 + 3q^4 + 6q^6 + 2q^8 - 7q^{10} - 13q^{12} - 7q^{14} + 2q^{16} + 6q^{18} + 3q^{20} - q^{22} - q^{24}),$$
  

$$d = q^{-4}[2]^2[3](1 - 2q^2 + q^4 - 2q^6 + q^8).$$

Remark 6.6 Using these equations we can represent the absolut invariant  $I_{2,f}^2 I_{1,f}^{-3} \in Q_I$ in terms of the quantum cross ratio  $\lambda$  of the four zeros  $v_i := q^{\frac{1}{2}} x_i y_i^{-1}$  of the form f (cf. Remark 3.2).

At the end of this section we discuss polynomials with double zeros. We have seven cases.

(i)For  $f_{iijk} = q^5(0i)(0i)(0j)(0k)$  we have

$$I_1 = \frac{[2]}{[3][4]} q^4 (ij)^2 (ik)^2$$

and

$$I_2 = \frac{[2]^4(q^{-4} - 2q^{-2} + 1 - 2q^2 + q^4)}{[4]^3[3]^2}q^9(ij)^3(ik)^3.$$

(ii) For  $f_{ijjk}=q^5(0i)(0j)(0j)(0k)$  we have

$$I_1 = \frac{[2]}{[3][4]} q^8 (ij)^2 (jk)^2$$

and

$$I_2 = \frac{[2]^3(-q^{-6} - q^{-4} + q^{-2} + q^2 - q^4 - q^6)}{[4]^3[3]^2}q^{18}(ij)^3(jk)^3.$$

(iii) For  $f_{ijkk} = q^5(0i)(0j)(0k)(0k)$  we have

$$I_1 = \frac{[2]}{[3][4]} q^4 (ik)^2 (jk)^2$$

and

$$I_2 = \frac{[2]^4(q^{-4} - 2q^{-2} + 1 - 2q^2 + q^4)}{[4]^3[3]^2}q^9(ik)^3(jk)^3.$$

(iv) For  $f_{iijj} = q^4(0i)(0i)(0j)(0j)$  we have

$$I_1 = \frac{[2]}{[3][4]} (ij)^4$$

and

$$I_{2} = \frac{[2]^{4}(q^{-4} - 2q^{-2} + 1 - 2q^{2} + q^{4})}{[4]^{3}[3]^{2}}(ij)^{6}.$$

(v)-(vii) For  $f_{iiij} = q^3(0i)(0i)(0i)(0j)$ ,  $f_{ijjj} = q^3(0i)(0j)(0j)(0j)$  and  $f_{iiii} = (0i)(0i)(0i)(0i)$  we have  $I_1 = 0$  and  $I_2 = 0$ .

Remark 6.7 In the classical case  $6\mathcal{I}_2^2 - \mathcal{I}_1^3$  is the discriminant of the quartic form. In the quantised situation the invariants  $\mathcal{I}_1$  and  $\mathcal{I}_2$  commute in cases (i)-(vii). The question arises, if it is possible to combine  $\mathcal{I}_1$  and  $\mathcal{I}_2$  to an invariant of degree six with the discriminant property. Up to a nonvanishing factor,

$$\mathcal{D} := \frac{[3][4]^3}{[2]^5(-q^{-4} + 2q^{-2} - 1 + 2q^2 - q^4)^2} \mathcal{I}_2^2 - \mathcal{I}_1^3$$

is the only element of this kind which vanishes on the noncommutative polynomials with a double zero  $f_{iijj}$ ,  $f_{iijk}$ ,  $f_{ijkk}$ ,  $f_{iiij}$ ,  $f_{ijjj}$  and  $f_{iiii}$ , but  $\mathcal{D}$  does not vanish on  $f_{ijjk}$ . There is no invariant of degree one.  $\mathcal{I}_1$  and  $\mathcal{I}_2$  are the only invariants of degree two and three, respectively. Therefore in case there exists a sixth-degree element with the discriminant property, there need to be further basic invariants of degree four or six.

# References

- 1. A. Clebsch, "Theorie der binären algebraischen Formen," B.G. Teubner, Leipzig, 1872.
- P. Gordan, "Dr. Paul Gordan's Vorlesungen über Invariantentheorie," ed. by Dr. Georg Kerschensteiner, B.G. Teubner, Leipzig, 1885-87.
- 3. G. Salmon, "Vorlesungen über die Algebra der linearen Transformationen," B.G. Teubner, Leipzig, 1877.
- 4. J.A. Dieudonne and J.B. Carrell, "Invariant theory, old and new," Academic Press, New York, 1971.
- 5. H. Weyl, "The classical groups, their invariants and representations," Princeton University Press, Princeton, 1946.
- 6. A. Clebsch, "Vorlesungen über Geometrie," B.G. Teubner, Leipzig, 1871.
- N.Yu. Reshetikhin, L.A. Takhtajan and L.D. Faddeev, Quantization of Lie groups and Lie algebras, *Leningrad Math. J.* 1 (1990) 193-225.
- S. Klimek and A. Lesniewski, A Two-Parameter Quantum Deformation of the Unit Disc, J. Func. Anal. 115 (1993), 1-23.
- 9. S. Majid, "Foundations of Quantum Group Theory," Cambridge University Press, New York/Cambridge, 1995.
- J.C. McConnel and J.C. Robson, "Noncommutative Noetherian Rings," John Wiley & Sons, Chichester, 1987.
- 11. C. Kassel, "Quantum Groups," Springer Verlag, New York, 1995.

- 12. O. Ore, Linear equations in non-commutative fields, Ann. Math. 32 (1933), 480-508.
- 13. C.W. Curtis, A note on noncommutative polynomials, *Proc. Amer. Math. Soc.* **3** (1952), 965-969.
- F. Leitenberger, Quantum theory of the real and complexified projective line, J. Math. Phys. 39 (1998), 802-817.
- 15. A. Capelli, Uber die Zurückführung der Cayley'schen Operation  $\Omega$  auf gewöhnliche Polar-Operationen, *Math. Ann.* **29** (1887), 331-338.
- 16. W. Fulton and J. Harris, "Representation theory," Springer-Verlag, New York, 1991.
- R. Howe, Remarks on classical invariant theory, Trans. Amer. Math. Soc. 313 (1989), 539-570.
- T. Hibi and M. Wakayama, A q-Analogue of Capelli's identity for GL(2), Adv. in Math. 115 (1995) 49-53.
- 19. M. Noumi, T. Umeda and M. Wakayama, A quantum analogue of the Capelli identity and an elementary differential calculus on  $GL_q(n)$ . Duke Math. J. 76 (1994), 567-594.
- 20. F. Klein, Zur Interpretation der komplexen Elemente in der Geometrie, Göttinger Nachrichten vom 14. August 1872.
- 21. B. Sturmfels, "Algorithms in Invariant Theory," Springer-Verlag, Wien/New York, 1993.