

A CHARACTERIZATION OF REFLEXIVITY FOR DUAL BANACH SPACES

JAVIER H. GUACHALLA

Universidad Mayor de San Andrés¹
La Paz - Bolivia

August 2005

ABSTRACT. Given a Banach space. We show that its three times dual space can be written as a direct sum. Being one of the sumands null is a necessary and sufficient condition for the reflexivity of the dual space. We end with an application to the known problem of the relationship between the reflexivity of a Banach space and that of its dual space.

INTRODUCTION

Let X be a Banach Space. We denote X^* , X^{**} , \dots , the (topological) dual, bidual, \dots spaces of X , and

$$\begin{aligned}\pi : X &\rightarrow X^{**} \\ \pi' : X^* &\rightarrow X^{***}\end{aligned}$$

the canonical maps into their respective bidual spaces.

For X, Y Banach spaces, and $T : X \rightarrow Y$ a bounded linear operator given, we denote $Ran(T)$ the range of T and $Ker(T)$ the null space of T . Note that since T is bounded, $Ker(T)$ is a closed subspace of X .

The adjoint operator of the bounded linear operator T will be denoted T^* , which is defined by $T^*(\mu) = \mu \circ T$ for $\mu \in Y^*$ and is a bounded linear operator; thus for the map π , we have

$$\pi^*(\eta) = \eta \circ \pi \quad (\text{for all } \eta \in X^{***}).$$

1. Definition. Let X and Y be Banach spaces, $T : X \rightarrow Y$ be a bounded linear operator. We will say that T is an isomorphism if it has a bounded inverse linear operator.

2. Definition. Let X be a Banach space and M, N closed subspaces of X . We will say that X is the direct sum of M and N , denoted $X = M \oplus N$, if $X = M + N$ and $M \cap N = \{0\}$ where $M + N = \{x + y : x \in M \text{ and } y \in N\}$.

We begin with a theorem which extends a result from algebra.

3. Theorem. Let X, Y , and Z be Banach spaces, $T : X \rightarrow Y$ and $S : Y \rightarrow Z$ bounded linear operators. Let the composition ST be an isomorphism. Then the subspace $Ran(T)$ is closed and

$$Y = Ran(T) \oplus Ker(S).$$

Proof. Let us see that $Ran(T)$ is closed. Let $y \in \overline{Ran(T)}$. There exist a sequence $\{x_n\}$ in X such that

$$y = \lim_{n \rightarrow \infty} T(x_n)$$

1991 *Mathematics Subject Classification.* FA - Functional Analysis.

¹e-mail:jguachal@entelnet.bo

Since ST is an isomorphism, there exists a unique $x \in X$ such that $ST(x) = S(y)$. Let us prove that $x_n \rightarrow x$. We have that $(ST)^{-1}$ is bounded,

$$\begin{aligned} ST(x_n) &\rightarrow ST(x) \quad \text{and} \\ \|x_n - x\| &\leq \|(ST)^{-1}\| \|(ST)(x_n) - (ST)(x)\| \end{aligned}$$

Hence

$$x_n \rightarrow x$$

Therefore $\lim_{n \rightarrow \infty} T(x_n) = T(x)$. Which means $y = T(x)$, i.e. $\text{Ran}(T)$ is a closed subspace.

Since $\text{Ker}(S)$ is a closed subspace, let us verify the direct sum conditions. Let $y \in Y$. Then $\exists! x \in X$ such that $(ST)(x) = S(y)$. Let us call $y' = y - T(x)$. Then $y' \in \text{Ker}(S)$. Therefore, we can write $y = T(x) + y'$. Where $T(x) \in \text{Ran}(T)$ and $y' \in \text{Ker}(S)$. Thus $Y = \text{Ran}(T) + \text{Ker}(S)$.

Let us see that $\text{Ran}(T) \cap \text{Ker}(S) = \{0\}$. Suppose $y \in \text{Ran}(T) \cap \text{Ker}(S)$. Therefore $y = T(x)$, for some $x \in X$ and $S(y) = 0$. Thus $S(T(x)) = S(y) = 0$. Therefore $x = 0$, since ST is an isomorphism. So $y = T(0) = 0$. Hence, according to the definition

$$Y = \text{Ran}(T) \oplus \text{Ker}(S)$$

THE DUAL SPACE AND ITS CANONICAL MAP

In this section, we apply the above result to the canonical map of the dual space of a Banach space X and to the adjoint of the canonical map of X itself.

4. Theorem.. *Let X and X^* be a Banach space and its dual space. Consider the maps $\pi' : X^* \rightarrow X^{***}$ and the adjoint operator of the canonical map of X , $\pi^* : X^{***} \rightarrow X^*$. Then $\pi^* \circ \pi'$ is the identity map on X^* , is an isomorphism, and*

$$X^{***} = \text{Ran}(\pi') \oplus \text{Ker}(\pi^*).$$

Proof. Let $\lambda \in X^*$. Then, by definition we have

$$(\pi^* \circ \pi')(\lambda) = \pi^*(\pi'(\lambda)) = \pi'(\lambda) \circ \pi$$

Evaluating in an arbitrary $x \in X$.

$$\begin{aligned} [(\pi^* \circ \pi')(\lambda)](x) &= [\pi'(\lambda) \circ \pi](x) \\ &= \pi'(\lambda)(\pi(x)) \\ &= (\pi(x))(\lambda) = \lambda(x) \quad (\text{for all } x \in X) \end{aligned}$$

therefore

$$\pi^* \circ \pi'(\lambda) = \lambda \quad (\text{for all } \lambda \in X^*)$$

Then we conclude that $\pi^* \circ \pi' = 1_{X^*}$ the identity map in X^* . Since X^* has the same topology in both maps, the identity map, 1_{X^*} is an isomorphism. Hence, by theorem 3

$$X^{***} = \text{Ran}(\pi') \oplus \text{Ker}(\pi^*)$$

5. Corollary. *Let X be a Banach space. The dual space X^* of X is reflexive if and only if $\text{Ker}(\pi^*) = 0$. Where π^* is the adjoint map of the canonical map π of X .*

APPLICATION

Let X be a Banach space. We prove X is reflexive if and only if X^* is reflexive.

To do so, let us consider the following notations. Given a subset M of a Banach space. We denote M° the annihilator of M , that is, the closed subspace of functionals of the respective dual space, whose restriction to M are zero.

Similarly, we denote ${}^\circ N$, for a subset N of a dual space; the closed subspace of elements of the space itself, that annihilate N , i.e. if $N \subset X^*$ then ${}^\circ N$ is the set $\{x \in X : \forall f \in N f(x) = 0\}$, which in fact is a closed subspace.

Suppose first, that X^* is reflexive. By Corollary 5 $Ker(\pi^*) = 0$. Then by [T.4.6], and since $Ran(\pi)$ is a closed subspace, being π an isometry, we obtain

$$\begin{aligned} Ran(\pi) &= {}^\circ Ker(\pi^*) \\ &= {}^\circ \{0\} \\ &= X^{**} \end{aligned}$$

Which says that X is reflexive.

Let us now suppose that X is reflexive, by the later reference and $Ran(\pi)$ closed, we have

$$\begin{aligned} Ker(\pi^*) &= Ran(\pi)^\circ \\ &= (X^{**})^\circ \\ &= \{0\} \end{aligned}$$

by an application of the Hahn-Banach theorem, and therefore, Corollary 5 implies X^* is reflexive.

REFERENCES

[T] A. Taylor, *Analysis*, 6th. printing, John Wiley & Sons, Inc., London, 1967.