

# SOME EFFECTS OF VERONESE MAP ON SYZYGIES OF PROJECTIVE VARIETIES

EUISUNG PARK

ABSTRACT. Let  $X \subset \mathbb{P}^r$  be a nondegenerate projective variety and let  $\nu_\ell : \mathbb{P}^r \rightarrow \mathbb{P}^N$  be the  $\ell$ -th Veronese embedding. In this paper we study the higher normality, defining equations and syzygies among them for the projective embedding  $\nu_\ell(X) \subset \mathbb{P}^N$ . We obtain that for a very ample line bundle  $L \in \text{Pic}X$  such that  $X \subset \mathbb{P}H^0(X, L)$  is  $m$ -regular in the sense of Castelnuovo-Mumford,  $(X, L^\ell)$  satisfies property  $N_\ell$  for all  $\ell \geq m$  (Theorem 1.5). This is a generalization of M. Green's work that  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_\ell$ . Also our result refines works of Ein-Lazarsfeld in [EL], Gallego-Purnaprajna in [GP1] and E. Rubei in [Ru1], [Ru2] and [Ru3].

## CONTENTS

1. Introduction	1
2. Preliminaries	5
3. The main theorem	7
4. Applications and examples	11
References	15

## 1. INTRODUCTION

Let  $X \subset \mathbb{P}^r$  be a nondegenerate projective variety where the embedding is defined by an  $(r + 1)$ -dimensional subspace  $V \subset H^0(X, \mathcal{O}_X(1))$  and hence  $\mathbb{P}^r = \mathbb{P}(V)$ . Let  $V_\ell$  be the image of the natural homomorphism

$$S^\ell V \cong H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell)) \rightarrow H^0(X, \mathcal{O}_X(\ell)).$$

In this paper we are concerned with the problem to see which information on

$$X \subset \mathbb{P}(V_\ell) \text{ and } X \subset \mathbb{P}H^0(X, \mathcal{O}_X(\ell))$$

can be deduced from  $X \subset \mathbb{P}(V)$ . The first result to this problem is due to Mumford.

**Theorem 1.1** (D. Mumford, [Mum2]). *Let  $d$  be the degree of a nondegenerate projective variety  $X \subset \mathbb{P}^r$ . If  $\ell \geq d$ , then  $X \subset \mathbb{P}(V_\ell)$  and  $X \subset \mathbb{P}H^0(X, \mathcal{O}_X(\ell))$  are represented as an intersection of quadratic forms.*

Later the result about the embedding defined by  $H^0(X, \mathcal{O}_X(\ell))$  is generalized to more general statements about higher syzygies. For precise statements, we recall the definition of property  $N_p$ .

---

Mathematics Subject Classification (2000): 13D02, 14N15.

**Definition 1.1.** Let  $X$  be a projective variety and let  $L \in \text{Pic}X$  a very ample line bundle defining an embedding  $X \hookrightarrow \mathbb{P}H^0(X, L)$ . Denote by  $S = \text{Sym}^\bullet H^0(X, L)$  the homogeneous coordinate ring of  $\mathbb{P}H^0(X, L)$ , and consider the graded  $S$ -module  $R(L) = \bigoplus_{n \in \mathbb{Z}} H^0(X, L^n)$ . Let

$$\cdots \rightarrow \bigoplus_j S^{\beta_{i,j}}(-i-j) \rightarrow \cdots \rightarrow \bigoplus_j S^{\beta_{1,j}}(-1-j) \rightarrow \bigoplus_j S^{\beta_{0,j}}(-j) \rightarrow R(L) \rightarrow 0$$

be a minimal graded free resolution of  $R(L)$ . The line bundle  $L$  is said to satisfy property  $N_p$  if  $\beta_{i,j} = 0$  for  $0 \leq i \leq p$  and  $j \geq 2$ .

Therefore property  $N_0$  holds if and only if  $X \hookrightarrow \mathbb{P}H^0(X, L)$  is a projectively normal embedding, property  $N_1$  holds if and only if property  $N_0$  is satisfied and the homogeneous ideal is generated by quadrics, and property  $N_p$  holds for  $p \geq 2$  if and only if it has property  $N_1$  and the  $k^{\text{th}}$  syzygies among the quadrics are generated by linear syzygies for all  $1 \leq k \leq p-1$ . For  $(X, L) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$ , M. Green obtain

**Theorem 1.2** (M. Green, [Green2]).  $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(\ell))$  satisfies property  $N_\ell$ .

Also L. Ein and R. Lazarsfeld generalize Theorem 1.1 as follows:

**Theorem 1.3** (L. Ein and R. Lazarsfeld, [EL]). Let  $X$  be a smooth complex projective variety and let  $L \in \text{Pic}X$  be a very ample line bundle of degree  $d$ . Then  $(X, L^\ell)$  satisfies property  $N_{\ell+1-d}$  for all  $\ell \geq d-1$ .

Later F. J. Gallego and B. P. Purnaprajna obtain the following result:

**Theorem 1.4** (F. J. Gallego and B. P. Purnaprajna, [GP1]). Let  $X$  be a projective variety and let  $L \in \text{Pic}X$  be a base point free and ample line bundle. Assume that  $\mathcal{O}_X$  is  $m$ -regular with respect to  $L$ , i.e.,  $H^i(X, L^{m-i}) = 0$  for all  $i \geq 1$ . Then for  $\ell \geq \max\{m-1+p, m+1, p+1\}$ ,  $(X, L^\ell)$  satisfies property  $N_p$ . In particular, if  $m \geq 2$  and  $\ell \geq 2m-1$ , then  $(X, L^\ell)$  satisfies property  $N_{\ell-m+1}$ .

**Remark 1.1.** Theorem 1.4 extends Theorem 1.3. Indeed let  $L$  be a very ample line bundle of degree  $d$  on a smooth complex projective variety  $X$  of dimension  $n$ . Then the line bundle  $D = L^{d-n-2} \otimes K_X^{-1}$  is base point free. One can find the details at the proof of Proposition 3.3 in [EL]. Thus  $L^\ell = K_X \otimes L^{\ell+n+2-d} \otimes D$  and hence  $\mathcal{O}_X$  is  $(d-1)$ -regular with respect to  $L$  by Kodaira vanishing theorem. If  $d \geq 2$ , then Theorem 1.4 implies that

$$(X, L^\ell) \text{ satisfies } \begin{cases} \text{property } N_{\ell-1} & \text{for } d = 2 \text{ and } \ell \geq 2, \text{ and} \\ \text{property } N_{\ell+2-d} & \text{for } d \geq 3 \text{ and } \ell \geq d-1. \end{cases}$$

If  $d = 1$ , then  $(X, L) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$ . In this case, see Theorem 1.2.  $\square$

Roughly speaking, these works guarantee that for a very ample line bundle  $L$  on a projective variety  $X$ ,  $(X, L^\ell)$  satisfies property  $N_{\ell-c}$  for all sufficiently large  $\ell$  where  $c \geq 0$  is an invariant of  $(X, L)$ . And our first main theorem shows that  $(X, L^\ell)$  satisfies property  $N_\ell$  for all sufficiently large  $\ell$ .

**Theorem 1.5.** Let  $X$  be a projective variety and let  $L \in \text{Pic}X$  be a very ample line bundle such that  $X \subset \mathbb{P}H^0(X, L) = \mathbb{P}$  is  $m$ -regular in the sense of Castelnuovo-Mumford, that is,  $H^i(\mathbb{P}, \mathcal{J}_{X/\mathbb{P}}(m-i)) = 0$  for all  $i \geq 1$ . Then

- (1)  $L^\ell$  is normally generated for  $\ell \geq \frac{m-1}{2}$ .
- (2)  $L^{m-1}$  satisfies property  $N_{m-2}$ .
- (3)  $L^\ell$  satisfies property  $N_\ell$  for all  $\ell \geq m$ .

This generalizes Green's theorem for  $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$  to arbitrary  $(X, L)$ . Also this refines Theorem 1.3 and 1.4. In §4 we apply our theorems to various projective varieties. We refine or reprove some known facts about property  $N_p$  of K3 surfaces, Enriques surfaces and abelian varieties in [GP1], [Ru1] and [Ru3].

Theorem 1.5 is a consequence of a more general result on some effects of the Veronese map on higher normality, defining equations and syzygies among them. Let  $X \subset \mathbb{P}(V)$  be a nondegenerate projective variety defined by a subspace  $V \subset H^0(X, \mathcal{O}_X(1))$  and let  $V_\ell$  be the image of the homomorphism

$$S^\ell V \cong H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell)) \rightarrow H^0(X, \mathcal{O}_X(\ell)).$$

From the surjective map  $H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell)) \rightarrow V_\ell$ , we can consider  $\mathbb{P}(V_\ell)$  as a linear subspace of  $\mathbb{P}(S^\ell V)$ . Thus we get the following commutative diagram:

$$\begin{array}{ccc} X & \subset & \mathbb{P}(V) \\ \nu_\ell \downarrow & & \downarrow \nu_\ell \\ \mathbb{P}(V_\ell) & \subset & \mathbb{P}(S^\ell V) \end{array}$$

We call  $X \subset \mathbb{P}(V_\ell)$  the  $\ell$ -th Veronese embedding. Recall that as defined in [EGHP], a projective variety embedded in a projective space satisfies property  $N_{2,1}$  if the homogeneous ideal is generated by quadrics and it satisfies property  $N_{2,p}$  for  $p \geq 2$  if property  $N_{2,1}$  holds and the  $k$ -th syzygies among the quadrics are generated by linear syzygies for all  $1 \leq k \leq p-1$ . See §2.3. We prove the following

**Theorem 1.6.** *Let  $X \subset \mathbb{P}(V)$  be a nondegenerate projective variety defined by a subspace  $V \subset H^0(X, \mathcal{O}_X(1))$ . Let  $s, t$  and  $m$  be integers such that*

- ( $\alpha$ )  $k$ -normality holds for all  $\ell \geq s$ ,
- ( $\beta$ ) the homogeneous ideal is generated by forms of degree  $\leq t$ , and
- ( $\gamma$ )  $X \subset \mathbb{P}(V)$  is  $m$ -regular, i.e.,  $H^1(\mathbb{P}(V), \mathcal{J}_X(m-i)) = 0$  for all  $i \geq 1$ .

Then for the  $\ell$ -th Veronese embedding,

- (1)  $\nu_\ell(X)$  satisfies  $k$ -normality for all  $k \geq \frac{s}{\ell}$ .
- (2) The homogeneous ideal of  $\nu_\ell(X)$  is generated by forms of degree  $\leq \max\{2, \frac{t}{\ell}\}$ .  
In particular,  $\nu_\ell(X)$  is ideal-theoretically cut out by quadrics if  $\ell \geq \frac{t}{2}$ .

- (3)  $\nu_\ell(X)$  satisfies  $\begin{cases} \text{property } N_{2,2\ell-m} & \text{for } \frac{m+1}{2} \leq \ell < m, \text{ and} \\ \text{property } N_{2,\ell} & \text{for } \ell \geq m. \end{cases}$

For the proof we first investigate syzygies among the defining equations of the “degenerate” embedding  $X \subset \mathbb{P}(S^\ell V)$ . Let  $\mathcal{J}_{X/\mathbb{P}(V)}$ ,  $\mathcal{J}_{\mathbb{P}(V)/\mathbb{P}(S^\ell V)}$ ,  $\mathcal{J}_{X/\mathbb{P}(S^\ell V)}$  and  $\mathcal{J}_{X/\mathbb{P}(V_\ell)}$  be sheaves of ideals of  $X \subset \mathbb{P}(V)$ ,  $\nu_\ell(\mathbb{P}(V)) \subset \mathbb{P}(S^\ell V)$ ,  $\nu_\ell(X) \subset \mathbb{P}(S^\ell V)$  and  $\nu_\ell(X) \subset \mathbb{P}(V_\ell)$ , respectively. Then we have the short exact sequence

$$0 \rightarrow \mathcal{J}_{\mathbb{P}(V)/\mathbb{P}(S^\ell V)} \rightarrow \mathcal{J}_{X/\mathbb{P}(S^\ell V)} \rightarrow \mathcal{J}_{X/\mathbb{P}(V)} \rightarrow 0.$$

For details, see the proof of Theorem 1.6. For  $\mathcal{J}_{\mathbb{P}(V)/\mathbb{P}(S^\ell V)}$  we know that  $\nu_\ell(\mathbb{P}(V)) \subset \mathbb{P}(S^\ell V)$  satisfies property  $N_\ell$ . Also our assumptions ( $\alpha$ )  $\sim$  ( $\gamma$ ) give some information

about  $J_{X/\mathbb{P}(V)}$ . Thus we obtain some results about syzygy modules of the degenerate embedding  $X \subset \mathbb{P}(S^\ell V)$ . As discussed in §2.3, syzygies among defining equations of  $X \subset \mathbb{P}(S^\ell V)$  are closely related to those of  $X \subset \mathbb{P}(V_\ell)$ . These observations enables us to prove Theorem 1.6.

The proof of Theorem 1.6.(3) combines the theory of Castelnuovo-Mumford regularity with Green's theorem. Indeed we obtain the following more general statement:

**Theorem 1.7.** *Let  $L$  be a very ample line bundle on a projective variety  $X$  such that  $X \subset \mathbb{P}H^0(X, L) = \mathbb{P}^r$  is  $m$ -regular. For  $\ell \geq m$ , suppose that  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_p$ . Then  $(X, L^\ell)$  satisfies property  $N_p$ .*

This result guarantees that if Green's theorem is refined, then Theorem 1.5 is automatically refined. See the following two examples.

**Example 1.1.** In [Ru5], E. Rubei shows that  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(3))$  satisfies property  $N_4$ . Therefore if  $X \subset \mathbb{P}H^0(X, L)$  is 3-regular, then  $(X, L^3)$  satisfies property  $N_4$ . See §4.3 for applications to 3-regular varieties.  $\square$

**Example 1.2.** Let  $X \subset \mathbb{P}^r$  be a quadratic hypersurface or a rational normal scroll. Then  $(X, \mathcal{O}_X(2))$  satisfies property  $N_5$ . Indeed T. Jozefiak, P. Pragacz and J. Weyman prove that  $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(2))$ ,  $n \geq 3$ , satisfies property  $N_p$  if and only if  $p \leq 5$ [JPW]. Since  $X \subset \mathbb{P}^r$  is 2-regular, the assertion comes from Theorem 1.7.  $\square$

**Example 1.3.** Let  $C \subset \mathbb{P}^2$  be a plane cubic curve which is not necessarily smooth or irreducible. When  $C$  is smooth,  $(C, \mathcal{O}_C(\ell))$  satisfies property  $N_{3\ell-3}$  by Green's "2g + 1 + p" theorem in [Green1]. By Theorem 1.7, we can extend this fact for arbitrary cubic plane curves. Indeed  $(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(\ell))$ ,  $\ell \geq 3$ , satisfies property  $N_{3\ell-3}$ . For details, we refer the reader to see [OP]. Therefore  $(C, \mathcal{O}_C(\ell))$  satisfies property  $N_{3\ell-3}$ ,  $\ell \geq 3$ , satisfies property  $N_{3\ell-3}$ .  $\square$

Recently G. Ottaviani and R. Paoletti investigate syzygies of Veronese embedding of projective spaces. By Ottaviani-Paoletti's theorem in [OP], if  $r \geq 2$  and  $\ell \geq 3$  then  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  fails to satisfy property  $N_{3\ell-2}$ . Also they formulate the following

**Conjecture A.** Assume that  $r \geq 2$  and  $\ell \geq 3$ . Then  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_p$  if and only if  $p \leq 3\ell - 3$ .

Ottaviani-Paoletti's conjecture can be generalized as follows:

**Conjecture B.** Let  $L$  be a very ample line bundle on a projective variety  $X$  such that  $X \subset \mathbb{P}H^0(X, L)$  is  $m$ -regular. Then for  $\ell \geq m$ ,  $(X, L^\ell)$  satisfies property  $N_{3\ell-3}$ .

Clearly Theorem 1.7 guarantees that if "Conjecture A" is true, then "Conjecture B" is also true. This conjecture says that for all sufficiently large  $\ell$ , syzygies of  $(X, L^\ell)$  is closed related to those of  $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(\ell))$ . Also the following example shows that the assumption " $\ell \geq m$ " is necessary.

**Example 1.4.** Let  $X$  be an elliptic ruled surface with the numerical invariant  $e \geq 0$ . Then  $aC_0 + bf$  is very ample if and only if  $a \geq 1$  and  $b - ae \geq 3$ . Also  $aC_0 + bf$  satisfies property  $N_p$  if and only if  $a \geq 1$  and  $b - ae \geq 3 + p$  by Theorem 1.4 in [Park1]. Therefore Conjecture B is true for every very ample line bundle on  $X$ . Now let  $L = aC_0 + bf$  be such that  $a \geq 1$  and  $b - ae = 3$ . Then  $L$  is very ample and  $X \subset \mathbb{P}H^0(X, L)$  is 3-regular. Note that  $(X, L^2)$  satisfies property  $N_p$  if and only if  $p \leq 3$  while  $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(2))$ ,  $n \geq 3$ , satisfies property  $N_p$  if and only if  $p \leq 5$ .  $\square$

**Organization of the paper.** In §2, we review some basic facts to study syzygies. In §3 we deal with higher normality and defining equations of Veronese map of arbitrary projective varieties. Also we prove our main results on syzygies of powers of a very ample line bundle. Finally we present some applications of our results in §4.

**Notations.** Throughout this paper all varieties are defined over an algebraically closed field  $K$  with  $\text{char}K = 0$ . For a finite dimensional  $K$ -vector space  $V$ ,  $\mathbb{P}(V)$  denotes the projective space of one-dimensional quotients of  $V$ .

**Acknowledgement.** I thank Elena Rubei for useful comments about §4.5.

## 2. PRELIMINARIES

**2.1. Surjection of multiplication maps.** To prove our main result we need to show that a particular multiplication map of global sections on coherent sheaves is surjective. Here we present an elementary but useful lemma for this aim.

**Lemma 2.1.** *Let  $X$  be a projective variety and let  $\mathcal{E}$  be a locally free sheaf on  $X$ . For a short sequence  $0 \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0$  of coherent sheaves on  $X$ , assume that*

- ( $\alpha$ )  $\mathcal{F}$  and  $\mathcal{H}$  are globally generated (and hence  $\mathcal{G}$  is globally generated),
- ( $\beta$ ) the following two sequences are exact, and

$$\begin{cases} 0 \rightarrow H^0(X, \mathcal{F}) \rightarrow H^0(X, \mathcal{G}) \rightarrow H^0(X, \mathcal{H}) \rightarrow 0 \\ 0 \rightarrow H^0(X, \mathcal{F} \otimes \mathcal{E}) \rightarrow H^0(X, \mathcal{G} \otimes \mathcal{E}) \rightarrow H^0(X, \mathcal{H} \otimes \mathcal{E}) \rightarrow 0 \end{cases}$$

- ( $\gamma$ ) the following two multiplication maps are surjective.

$$\begin{cases} H^0(X, \mathcal{F}) \otimes H^0(X, \mathcal{E}) \rightarrow H^0(X, \mathcal{F} \otimes \mathcal{E}) \\ H^0(X, \mathcal{H}) \otimes H^0(X, \mathcal{E}) \rightarrow H^0(X, \mathcal{H} \otimes \mathcal{E}) \end{cases}$$

Then the multiplication map  $H^0(X, \mathcal{G}) \otimes H^0(X, \mathcal{E}) \rightarrow H^0(X, \mathcal{G} \otimes \mathcal{E})$  is surjective.

*Proof.* By our assumption we have the commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & H^0(X, \mathcal{F}) \otimes H^0(X, \mathcal{E}) & \rightarrow & H^0(X, \mathcal{G}) \otimes H^0(X, \mathcal{E}) & \rightarrow & H^0(X, \mathcal{H}) \otimes H^0(X, \mathcal{E}) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & H^0(X, \mathcal{F} \otimes \mathcal{E}) & \rightarrow & H^0(X, \mathcal{G} \otimes \mathcal{E}) & \rightarrow & H^0(X, \mathcal{H} \otimes \mathcal{E}) \rightarrow 0. \end{array}$$

where the two rows are exact. The assertion comes from snake lemma.  $\square$

**2.2. Cohomological criterion of property  $N_{2,p}$ .** Let  $X \subset \mathbb{P}^r$  be a nondegenerate projective variety and let  $I_X$  be the homogeneous ideal of  $X$  in the homogeneous coordinate ring  $S$  of  $\mathbb{P}^r$ . For  $p \geq 1$ ,  $X$  is said to satisfy property  $N_{2,p}$  if  $I_X$  admits the minimal free resolution of the form

$$\cdots \rightarrow S(-p-1)^{\beta_{p-1,2}} \rightarrow \cdots \rightarrow S(-3)^{\beta_{1,2}} \rightarrow S(-2)^{\beta_{0,2}} \rightarrow I_X \rightarrow 0.$$

Therefore property  $N_{2,1}$  holds if  $I_X$  is generated by quadrics. For  $p \geq 2$ , it is said to satisfy property  $N_{2,p}$  if property  $N_{2,1}$  holds and the  $k$ -th syzygies among the quadrics are generated by linear syzygies for all  $1 \leq k \leq p-1$ . Equivalently, property  $N_{2,p}$  holds if  $\text{Tor}_i^S(I_X, K)$  is a vector space concentrated in degrees  $\leq i+2$  for all  $i \leq p-1$ . It is well known that  $\text{Tor}_i^S(I_X, K)$  can be read off as follows:

**Theorem 2.2.** *Put  $V = H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(1))$  and  $\mathcal{M} = \Omega_{\mathbb{P}^r}(1)$ . Then for the sheaf of ideals  $\mathcal{J}_X$  of  $X$ , there is an exact sequence*

$$\begin{aligned} \bigwedge^{i+1} V \otimes H^0(\mathbb{P}^r, \mathcal{J}_X(j-1)) &\xrightarrow{\alpha_{i,j}} H^0(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{J}_X(j)) \rightarrow \\ H^1(\mathbb{P}, \bigwedge^{i+1} \mathcal{M} \otimes \mathcal{J}_X(j-1)) &\rightarrow \bigwedge^{i+1} V \otimes H^1(\mathbb{P}, \mathcal{J}_X(j-1)) \end{aligned}$$

with  $\text{Coker}(\alpha_{i,j}) = \text{Tor}_i^{i+j}(I_X, K)$ . Therefore we have the exact sequence

$$0 \rightarrow \text{Tor}_i^{i+j}(I_X, K) \rightarrow H^1(\mathbb{P}^r, \bigwedge^{i+1} \mathcal{M} \otimes \mathcal{J}_X(j-1)) \rightarrow \bigwedge^{i+1} V \otimes H^1(\mathbb{P}^r, \mathcal{J}_X(j-1)).$$

*Proof.* See Theorem (1.b.4) in [Green1] or Theorem 4.5 in [E].  $\square$

**Corollary 2.3.** (1) *If  $H^1(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{J}_X(j)) = 0$  for  $1 \leq i \leq p$  and  $j \geq 2$ , then  $X \subset \mathbb{P}^r$  satisfies property  $N_{2,p}$ .*

(2) *If  $X \subset \mathbb{P}^r$  is projectively normal, then it satisfies property  $N_{2,p}$  if and only if  $H^1(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{J}_X(j)) = 0$  for  $1 \leq i \leq p$  and  $j \geq 2$ .*

*Proof.* By definition property  $N_{2,p}$  holds if and only if

$$\text{Tor}_i^{i+j}(I_X, K) = 0 \quad \text{for } 0 \leq i \leq p-1 \text{ and } j \geq 3.$$

Thus the assertion comes immediately from the exact sequence in Theorem 2.2.  $\square$

**2.3. Syzygies of degenerate varieties.** Let  $X \subset \mathbb{P}^{r+e}$  ( $e \geq 1$ ) be a degenerate projective variety, i.e., there exists a linear subspace  $\Lambda \cong \mathbb{P}^r \subset \mathbb{P}^{r+e}$  such that  $X \subset \Lambda$ . To simplify notations, put  $\mathcal{M}_n = \Omega_{\mathbb{P}^n}(1)$ . It is easily checked that

$$\mathcal{M}_{r+e}|_{\Lambda} \cong \mathcal{M}_r \oplus \mathcal{O}_{\mathbb{P}^r}^e.$$

Let  $\mathcal{J}_{X/\mathbb{P}^{r+e}}$ ,  $\mathcal{J}_{\Lambda/\mathbb{P}^{r+e}}$  and  $\mathcal{J}_{X/\mathbb{P}^r}$  be sheaves of ideals of  $X \subset \mathbb{P}^{r+e}$ ,  $\Lambda \subset \mathbb{P}^{r+e}$  and  $X \subset \mathbb{P}^r$ , respectively.

**Lemma 2.4.** *Under the situation just stated, suppose that for some  $k \geq 0$  and  $j \geq 1$ ,*

$$H^1(\mathbb{P}^{r+e}, \bigwedge^k \mathcal{M}_{r+e} \otimes \mathcal{J}_{X/\mathbb{P}^{r+e}}(j)) = 0$$

*Then  $H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_r \otimes \mathcal{J}_{X/\mathbb{P}^r}(j)) = 0$ .*

*Proof.* We may assume that  $e = 1$  and hence  $\mathcal{J}_{\Lambda/\mathbb{P}^{r+1}} = \mathcal{O}_{\mathbb{P}^{r+1}}(-1)$ . By snake lemma we have the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{O}_{\mathbb{P}^{r+1}}(-1) & = & \mathcal{O}_{\mathbb{P}^{r+1}}(-1) & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \rightarrow & \mathcal{J}_{X/\mathbb{P}^{r+1}} & \rightarrow & \mathcal{O}_{\mathbb{P}^{r+1}} & \rightarrow & \mathcal{O}_X \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \rightarrow & \mathcal{J}_{X/\mathbb{P}^r} & \rightarrow & \mathcal{O}_{\mathbb{P}^r} & \rightarrow & \mathcal{O}_X \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 
 \end{array}$$

Recall that  $H^2(\mathbb{P}^{r+1}, \bigwedge^k \mathcal{M}_{r+1}(j-1)) = H^2(\mathbb{P}^{r+1}, \Omega_{\mathbb{P}^{r+1}}^k(k+j-1)) = 0$  for  $k+j \geq 2$ . The assertion comes from the cohomology long exact sequence of the left column.  $\square$

**Remark 2.3.1.** From the exact sequence  $0 \rightarrow \mathcal{O}_{\mathbb{P}^{r+1}}(-1) \rightarrow \mathcal{J}_{X/\mathbb{P}^{r+1}} \rightarrow \mathcal{J}_{X/\mathbb{P}^r} \rightarrow 0$ , it is easily proved that if the homogeneous ideal of  $X \subset \mathbb{P}^{r+e}$  is generated by forms of degree  $\leq d$ , then so does  $X \subset \mathbb{P}^r$ .  $\square$

**Corollary 2.5.** *Under the same situation, assume that  $X \subset \mathbb{P}^r$  is nondegenerate.*

(1) *If  $H^1(\mathbb{P}^{r+e}, \mathcal{J}_{X/\mathbb{P}^{r+e}}(j)) = 0$ , then  $H^1(\mathbb{P}^r, \mathcal{J}_{X/\mathbb{P}^r}(j)) = 0$  and hence  $X \subset \mathbb{P}^r$  satisfies  $j$ -normality.*

(2) *If  $H^1(\mathbb{P}^{r+e}, \bigwedge^k \mathcal{M}_{r+e} \otimes \mathcal{J}_{X/\mathbb{P}^{r+e}}(j)) = 0$  for  $1 \leq k \leq p$  and  $j \geq 2$ , then  $X \subset \mathbb{P}^r$  satisfies property  $N_{2,p}$ .*

*Proof.* The assertion comes immediately from Corollary 2.3 and Lemma 2.4.  $\square$

### 3. THE MAIN THEOREM

This section is devoted to prove Theorem 1.6. Let  $X \subset \mathbb{P}(V) = \mathbb{P}^r$  be a nondegenerate projective variety and let  $\ell \geq 1$  be an integer. Throughout this section we use the following notations:

- $N := \binom{r+\ell}{\ell}$  : the dimension of  $S^\ell V \cong H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$
- $Z := \nu_\ell(\mathbb{P}(V)) \subset \mathbb{P}^N$
- $V_\ell$  : the image of  $H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell)) \rightarrow H^0(X, \mathcal{O}_X(\ell))$

Thus we have the following commutative diagram:

$$\begin{array}{ccc}
 X & \subset & \mathbb{P}^r \\
 \nu_\ell \downarrow & & \downarrow \nu_\ell \\
 \mathbb{P}(V_\ell) & \subset & \mathbb{P}^N
 \end{array}$$

Let  $\mathcal{J}_{X/\mathbb{P}^r}$ ,  $\mathcal{J}_{Z/\mathbb{P}^N}$ ,  $\mathcal{J}_{X/\mathbb{P}^N}$  and  $\mathcal{J}_{X/\mathbb{P}(V_\ell)}$  be sheaves of ideals of  $X \subset \mathbb{P}^r$ ,  $Z \subset \mathbb{P}^N$ ,  $\nu_\ell(X) \subset \mathbb{P}^N$  and  $\nu_\ell(X) \subset \mathbb{P}(V_\ell)$ , respectively. By snake lemma we have the following:

$$\begin{array}{ccccccc}
& & & & 0 & & \\
& & & & \downarrow & & \\
& & & & 0 & & \mathcal{J}_{X/\mathbb{P}^r} \\
& & & & \downarrow & & \downarrow \\
0 & \rightarrow & \mathcal{J}_{Z/\mathbb{P}^N} & \rightarrow & \mathcal{O}_{\mathbb{P}^N} & \rightarrow & \mathcal{O}_{\mathbb{P}^r} \rightarrow 0 \\
& & \downarrow & & \parallel & & \downarrow \\
0 & \rightarrow & \mathcal{J}_{X/\mathbb{P}^N} & \rightarrow & \mathcal{O}_{\mathbb{P}^N} & \rightarrow & \mathcal{O}_X \rightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & \mathcal{J}_{X/\mathbb{P}^r} & & 0 & & \\
& & \downarrow & & & & \\
& & 0 & & & & 
\end{array}$$

In particular, we get the short exact sequence

$$(\diamond) \quad 0 \rightarrow \mathcal{J}_{Z/\mathbb{P}^N} \rightarrow \mathcal{J}_{X/\mathbb{P}^N} \rightarrow \mathcal{J}_{X/\mathbb{P}^r} \rightarrow 0.$$

Now we start to prove our main theorem.

**Proof of Theorem 1.6.** (1) Since  $Z \subset \mathbb{P}^N$  is projectively normal,

$$H^1(\mathbb{P}^N, \mathcal{J}_{Z/\mathbb{P}^N}(k)) = 0 \quad \text{for all } k \geq 1.$$

Also from the short exact sequence  $0 \rightarrow \mathcal{J}_{Z/\mathbb{P}^N} \rightarrow \mathcal{O}_{\mathbb{P}^N} \rightarrow \mathcal{O}_{\mathbb{P}^r} \rightarrow 0$ ,

$$H^2(\mathbb{P}^N, \mathcal{J}_{Z/\mathbb{P}^N}(k)) = 0 \quad \text{for all } k \geq 1.$$

From the cohomology long exact sequence of  $(\diamond)$ , this implies that

$$H^1(\mathbb{P}^N, \mathcal{J}_{X/\mathbb{P}^N}(k)) \cong H^1(\mathbb{P}^r, \mathcal{J}_{X/\mathbb{P}^r}(k\ell)) \quad \text{for all } k \geq 1.$$

Also our assumption  $(\alpha)$  guarantees that  $H^1(\mathbb{P}^r, \mathcal{J}_{X/\mathbb{P}^r}(k\ell)) = 0$  if  $k \geq \frac{s}{\ell}$ . Therefore we have  $H^1(\mathbb{P}^N, \mathcal{J}_{X/\mathbb{P}^N}(k)) = 0$  for all  $k \geq \frac{s}{\ell}$ . This implies that  $X \subset \mathbb{P}(V_\ell)$  satisfies  $k$ -normality for all  $k \geq \frac{s}{\ell}$  by Corollary 2.5.

(2) By Remark 2.2.1 we need to show that the homogeneous ideal of  $\nu_\ell(X) \subset \mathbb{P}^N$  is generated by forms of degree  $\leq \mu$  or, equivalently, the multiplication maps

$$H^0(\mathbb{P}^N, \mathcal{J}_{X/\mathbb{P}^N}(\mu)) \otimes H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(k)) \rightarrow H^0(\mathbb{P}^N, \mathcal{J}_{X/\mathbb{P}^N}(\mu + k))$$

are surjective for all  $k \geq 1$  where  $\mu = \max\{2, \frac{t}{\ell}\}$ . By applying Lemma 2.1 to the short exact sequence  $(\diamond)$  and the line bundle  $\mathcal{O}_{\mathbb{P}^N}(k)$  where  $k \geq 1$ , it suffices to check the followings:

- (a)  $\mathcal{J}_{Z/\mathbb{P}^N}(\mu)$  and  $\mathcal{J}_{X/\mathbb{P}^r}(\mu\ell)$  are globally generated.
- (b) The homomorphism

$$H^0(\mathbb{P}^N, \mathcal{J}_{X/\mathbb{P}^N}(k)) \rightarrow H^0(\mathbb{P}^r, \mathcal{J}_{X/\mathbb{P}^r}(k\ell))$$



is surjective for all  $k \geq \mu$ .

(c) The multiplication maps

$$\begin{cases} (\triangleleft) & H^0(\mathbb{P}^N, \mathcal{J}_{Z/\mathbb{P}^N}(\mu)) \otimes H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(k)) \rightarrow H^0(\mathbb{P}^N, \mathcal{J}_{Z/\mathbb{P}^N}(\mu+k)) \\ (\triangleright) & H^0(\mathbb{P}^r, \mathcal{J}_{X/\mathbb{P}^r}(\mu\ell)) \otimes H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(k)) \rightarrow H^0(\mathbb{P}^r, \mathcal{J}_{X/\mathbb{P}^r}((\mu+k)\ell)) \end{cases}$$

are surjective for all  $k \geq 1$ .

Since the homogeneous ideal of  $Z \subset \mathbb{P}^N$  is generated by quadrics,  $\mathcal{J}_{Z/\mathbb{P}^N}(\mu)$  is globally generated. Also the homomorphism in (b) and the multiplication map  $(\triangleleft)$  are surjective for all  $k \geq 1$  since  $Z \subset \mathbb{P}^N$  is projectively normal. The global generation of  $\mathcal{J}_{X/\mathbb{P}^r}(\mu\ell)$  and the surjectivity of the multiplication map  $(\triangleright)$  come from the assumption  $(\beta)$ .

(3) Let  $\mathcal{M}_\ell$  be the restriction of  $\mathcal{M}_{\mathbb{P}^N} := \Omega_{\mathbb{P}^N}(1)$  to  $Z$ . Clearly  $\mathcal{M}_\ell$  is the kernel of the evaluation homomorphism  $V_\ell \otimes \mathcal{O}_{\mathbb{P}^r} \rightarrow \mathcal{O}_{\mathbb{P}^r}(\ell) \rightarrow 0$  and the short exact sequence

$$0 \rightarrow \mathcal{M}_\ell \rightarrow V_\ell \otimes \mathcal{O}_{\mathbb{P}^r} \rightarrow \mathcal{O}_{\mathbb{P}^r}(\ell) \rightarrow 0$$

is the restriction of the Euler sequence on  $\mathbb{P}^N$  to  $Z$ . From the short exact sequence  $(\diamond)$ , we have the long exact sequence

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{Z/\mathbb{P}^N}(j)) \rightarrow H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{X/\mathbb{P}^N}(j)) \rightarrow H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}(j\ell))$$

of cohomology groups. Since  $Z \subset \mathbb{P}^N$  satisfies property  $N_\ell$  by Theorem 1.2,

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{\mathbb{P}^r/\mathbb{P}^N}(j)) = 0$$

for  $1 \leq k \leq \ell$  and all  $j \geq 2$  by Corollary 2.3. Also we claim that

$$H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}(j\ell)) = 0$$

for  $1 \leq k \leq \ell + 1$  and  $j\ell \geq 2\ell \geq m + k - 1$ . Indeed  $\bigwedge^k \mathcal{M}_\ell$  is  $(\ell + 1)$ -regular with respect to  $\mathcal{O}_{\mathbb{P}^r}(1)$  by the next Proposition 3.1 since  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_\ell$  by Green's theorem. Therefore  $\bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}$  is  $(m + \ell + 1)$ -regular with respect to  $\mathcal{O}_{\mathbb{P}^r}(1)$ . For details, we refer the reader to see Proposition 1.8.9 in [Laz2]. This completes the proof of our claim.

Case 1. Assume that  $\frac{m+1}{2} \leq \ell < m$ . Then  $2\ell \geq m + k - 1$  holds for  $1 \leq k \leq 2\ell - m + 1$ . Therefore

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{X/\mathbb{P}^N}(j)) = 0 \quad \text{for } 1 \leq k \leq 2\ell - m + 1 \text{ and all } j \geq 2.$$

Consequently Corollary 2.5 shows that  $X \subset \mathbb{P}(V_\ell)$  satisfies property  $N_{2,2\ell-m}$ .

Case 2. Assume that  $\ell \geq m$ . Then  $2\ell \geq m + k - 1$  holds for  $1 \leq k \leq \ell + 1$  and hence we have

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{X/\mathbb{P}^N}(j)) = 0 \quad \text{for } 1 \leq k \leq \ell \text{ and } j \geq 2.$$

Therefore  $X(X) \subset \mathbb{P}(V_\ell)$  satisfies property  $N_{2,\ell}$  by Corollary 2.5.  $\square$

**Proposition 3.1.** *If  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_p$ , then  $\bigwedge^k \mathcal{M}_\ell$  is  $(\ell + 1)$ -regular for every  $1 \leq k \leq p + 1$ .*

*Proof.* By definition of the Castelnuovo-Mumford regularity, we need to show that

$$H^i(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell(\ell + 1 - i)) = 0$$

for  $1 \leq k \leq p + 1$  and all  $i \geq 1$ . For each  $k \geq 0$ , consider the short exact sequence

$$(\star) \quad 0 \rightarrow \bigwedge^{k+1} \mathcal{M}_\ell \rightarrow \bigwedge^{k+1} V_\ell \otimes \mathcal{O}_{\mathbb{P}^r} \rightarrow \bigwedge^k \mathcal{M}_\ell(\ell) \rightarrow 0.$$

Step 1. Since  $H^r(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(j)) = 0$  for all  $j \geq -r$ , we have

$$H^r(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell(j)) = 0$$

for all  $k \geq 1$  and  $j \geq \ell - r$  by the cohomology long exact sequence induced from  $(\star)$ .

Step 2. If  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_p$ , then

$$(\star\star) \quad H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell(\ell)) = 0$$

for every  $1 \leq k \leq p + 1$  by the cohomological criterion in [EL].

Step 3. By  $(\star)$  and  $(\star\star)$ , we have

$$H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell(\ell)) \cong H^2(\mathbb{P}^r, \bigwedge^{k+1} \mathcal{M}_\ell) = 0$$

for every  $1 \leq k \leq p + 1$ . Then Proposition 1.7 in [OP] guarantees that

$$(\star\star\star) \quad H^2(\mathbb{P}^r, \bigwedge^{k+1} \mathcal{M}_\ell(t)) \cong H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell(\ell + t)) = 0$$

for every  $1 \leq k \leq p + 1$  and  $t \geq 0$ .

Step 4. Assume that  $1 \leq i \leq r - 1$ . If  $k \geq i$ , then

$$H^i(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell(\ell + 1 - i)) \cong \dots \cong H^1(\mathbb{P}^r, \bigwedge^{i-k+1} \mathcal{M}_\ell(\ell + (\ell - 1)(i - 1))) = 0$$

by  $(\star\star\star)$ . Also if  $k < i$ , then

$$H^i(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell(\ell + 1 - i)) \cong \dots \cong H^{i-k}(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell + 1 - i + k\ell)) = 0.$$

Therefore  $\bigwedge^k \mathcal{M}_d$  is  $(\ell + 1)$ -regular for every  $1 \leq k \leq p + 1$ .  $\square$

**Proof of Theorem 1.5.** It is well known that if  $X \subset \mathbb{P}H^0(X, L)$  is  $m$ -regular, then  $k$ -normality holds for all  $k \geq m - 1$  and the homogeneous ideal is generated by forms of degree  $\leq m$  (Lecture 14 in [Mum1]).

(1) Let  $V_\ell$  be the image of  $H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell)) \rightarrow H^0(X, L^\ell)$ . If  $\ell \geq \frac{m-1}{2}$ , then Theorem 1.6.(1) implies that  $X \subset \mathbb{P}(V_\ell)$  is  $k$ -normal for all  $k \geq 2$ . Therefore the linearly normal embedding  $X \subset \mathbb{P}H^0(X, L^\ell)$  is projectively normal.

(2) Since  $(m - 1)$ -normality holds for  $X \subset \mathbb{P}H^0(X, L)$ ,  $V_{m-1} = H^0(X, L^{m-1})$ . Also Theorem 1.6.(3) implies that  $X \subset \mathbb{P}H^0(X, L^{m-1})$  satisfies property  $N_{2, m-2}$ . Therefore  $(X, L^{m-1})$  satisfies property  $N_{m-2}$ .

(3) Since  $X \subset \mathbb{P}H^0(X, L)$  is  $\ell$ -normal for all  $\ell \geq m$ ,  $V_\ell = H^0(X, L^\ell)$ . Also Theorem 1.6.(3) implies that  $X \subset \mathbb{P}H^0(X, L^\ell)$  satisfies property  $N_{2, \ell}$  if  $\ell \geq m$ . Therefore  $(X, L^\ell)$  satisfies property  $N_\ell$ .  $\square$

**Remark 3.1.** In many cases, an  $m$ -regular projective variety  $X \subset \mathbb{P}H^0(X, L)$  satisfies  $s$ -normality for some  $s < m - 1$ . Suppose that  $s$ -normality holds for some  $\frac{m+1}{2} \leq s \leq m - 2$ . Then by the same reason as in the above proof, Theorem 1.6.(3) guarantees that  $(X, L^s)$  satisfies property  $N_{2s-m}$ .  $\square$

#### 4. APPLICATIONS AND EXAMPLES

In this section, we present some applications of our results.

**4.1. Generalized Ottaviani-Paoletti's conjecture.** By M. Green's Theorem 2.2 in [Green2],  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_\ell$ . We generalize this result to arbitrary projective varieties in Theorem 1.5. On the other hand, G. Ottaviani and R. Paoletti[OP] prove that if  $r \geq 2$  and  $\ell \geq 3$ , then  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  fails to satisfy property  $N_{3\ell-2}$ . Also they conjectured that for  $r \geq 2$  and  $\ell \geq 3$ ,  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_p$  if and only if  $p \leq 3\ell - 3$ . The cases  $r \geq 3$  are still open except  $r = \ell = 3$ . We generalize Ottaviani-Paoletti's conjecture to very ample line bundles on an arbitrary projective variety. See "Conjecture A" and "Conjecture B" in §1. And the aim of this subsection is to prove that if Conjecture A is true, then Conjecture B is also true.

**Proof of Theorem 1.7.** We use the short exact sequence  $(\diamond)$  in §3. Note that by Corollary 2.5 we need to show that

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{X/\mathbb{P}^N}(j)) = 0 \quad \text{for } 1 \leq k \leq p \text{ and } j \geq 2.$$

The short exact sequence  $(\diamond)$  gives the cohomology long exact sequence

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{Z/\mathbb{P}^N}(j)) \rightarrow H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{X/\mathbb{P}^N}(j)) \rightarrow H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}(j\ell)).$$

Since  $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell))$  satisfies property  $N_p$ ,

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{Z/\mathbb{P}^N}(j)) = 0 \quad \text{for } 1 \leq k \leq p \text{ and } j \geq 2.$$

Also  $\bigwedge^k \mathcal{M}_\ell$  is  $(\ell + 1)$ -regular for every  $1 \leq k \leq p + 1$  by Proposition 3.1. Since  $\mathcal{J}_{X/\mathbb{P}^r}$  is  $m$ -regular,  $\bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}$  is  $(m + \ell + 1)$ -regular by Proposition 1.8.9 in [Laz2].

Therefore  $H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}(j\ell)) = 0$  for  $1 \leq k \leq p$  and  $j \geq 2$  since we assume that  $\ell \geq m$ . As a consequence, we have

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{X/\mathbb{P}^N}(j)) = 0 \quad \text{for } 1 \leq k \leq p \text{ and } j \geq 2.$$

Therefore  $X(X) \subset \mathbb{P}(V_\ell)$  satisfies property  $N_{2,\ell}$  by Corollary 2.5.  $\square$

**4.2. Castelnuovo-Mumford regularity of projective varieties.** Our results are stated in terms of the Castelnuovo-Mumford regularity of projective varieties. Here we shortly review known results in this field. We refer to [Kwak1],[Kwak2] and [Kwak3] for details. There is a well known conjecture due to Eisenbud and Goto[EG] which gives a bound for regularity in terms of the degree and the codimension:

**Regularity conjecture.** Let  $X \subset \mathbb{P}^r$  be a nondegenerate integral projective variety of degree  $d$  and codimension  $e$ . Then it satisfies  $(d - e + 1)$ -regularity.

Let us recall the following works developed toward this conjecture.

- (L. Gruson, R. Lazarsfeld and C. Peskine, [GLP]) Let  $C \subset \mathbb{P}^r$  be a nondegenerate integral curve of degree  $d$ . Then  $C$  is  $(d + 2 - r)$ -regular.
- (R. Lazarsfeld, [Laz1]) Let  $S \subset \mathbb{P}^r$  be a nondegenerate smooth surface of degree  $d$ . Then  $S$  is  $(d + 3 - r)$ -regular.
- (Ziv Ran, [Ran]) Let  $X \subset \mathbb{P}^r$ ,  $r \geq 9$ , be a nondegenerate smooth threefold of degree  $d$ . Then  $X$  is  $(d + 4 - r)$ -regular.
- (Sijong Kwak, [Kwak1][Kwak2][Kwak3]) Let  $X \subset \mathbb{P}^r$  be a nondegenerate smooth projective variety of degree  $d$ .
  1. If  $X$  is a threefold and  $r = 5$ , then  $X$  is  $(d - 1)$ -regular.
  2. If  $X$  is a threefold and  $6 \leq r \leq 8$ , then  $X$  is  $(d + 5 - r)$ -regular.
  3. If  $X$  is a fourfold, then  $X$  is  $(d + 9 - r)$ -regular.
  4. If  $X$  is a fivefold, then  $X$  is  $(d + 16 - r)$ -regular.
  5. If  $X$  is a sixfold, then  $X$  is  $(d + 27 - r)$ -regular.
- (A. Bertram, L. Ein and R. Lazarsfeld, [BEL]) Let  $X \subset \mathbb{P}^r$  be a nondegenerate smooth projective variety. Let  $n$  be the dimension of  $X$  and let  $d$  be the degree of  $X \subset \mathbb{P}^r$ . Then  $X$  is  $(\min\{r - n, n + 1\})(d - 1) - n + 1$ -regular.

Note that Theorem 1.5 is stated in terms of the Castelnuovo-Mumford regularity of the linearly normal embedding. For a projective variety  $X$  of dimension  $n$  and a very ample line bundle  $L \in \text{Pic}X$  of degree  $d$ , the  $\Delta$ -genus  $\Delta(X, L)$  of  $(X, L)$  is defined to be  $d + n - h^0(X, L)$ . Therefore for linearly normal varieties, the Eisenbud-Goto's regularity conjecture is

**Regularity conjecture for linearly normal varieties.** Let  $X$  be an integral projective variety and let  $L$  be a very ample line bundle on  $X$ . Then the linearly normal embedding  $X \subset \mathbb{P}H^0(X, L)$  satisfies  $(\Delta(X, L) + 2)$ -regularity.

One can apply Theorem 1.5 to the above works about Castelnuovo-Mumford regularity. For an example, we obtain the following

**Corollary 4.1.** *Let  $S$  be a smooth surface and let  $L \in \text{Pic}S$  be a very ample line bundle. Put  $m = \Delta(S, L) + 2$ . Then*

- (1)  $L^\ell$  is normally generated for  $\ell \geq \frac{m-1}{2}$ .
- (2)  $L^{m-1}$  satisfies property  $N_{m-2}$ .
- (3)  $L^\ell$  satisfies property  $N_\ell$  for  $\ell \geq m$ .

*Proof.* By R. Lazarsfeld's result in [Laz1],  $S \subset \mathbb{P}H^0(S, L)$  is  $(\Delta(S, L) + 2)$ -regular. Thus the assertion follows immediately from Theorem 1.5.  $\square$

**4.3. 3-regular varieties.** Let  $L$  be a very ample line bundle on a smooth projective variety  $X$  where the linearly normal embedding  $X \subset \mathbb{P}H^0(X, L)$  is 3-regular. Then it is projectively normal and Theorem 1.5 implies that

$$(\dagger) \begin{cases} (i) & (X, L) \text{ satisfies property } N_0. \\ (ii) & (X, L^2) \text{ satisfies property } N_1. \\ (iii) & (X, L^3) \text{ satisfies property } N_4. \\ (iv) & (X, L^\ell), \ell \geq 4, \text{ satisfies property } N_\ell. \end{cases}$$

For (iii), see Example 1.1. Here we exhibits examples where 3-regularity holds.

(4.3.1) Let  $S$  be a K3 surface. Then any linearly normal embedding of  $S$  is 3-regular. Indeed for a very ample line bundle  $L$  on  $S$ , let  $C \in |L|$  be a smooth section. Since  $H^1(S, L^j) = 0$  for all  $j \geq 0$ ,  $S \subset \mathbb{P}H^0(S, L)$  and  $C \subset \mathbb{P}H^0(C, L|_C)$  has the same Betti table. Thus Noether's theorem implies that  $S \subset \mathbb{P}H^0(S, L)$  is projectively normal and hence it is 3-regular. Therefore  $(S, L)$  satisfies  $(\dagger)$ . We should mention Gallego-Purnaprajna's result in [GP2]. Recall that  $L^2 \geq 4$ ,  $L^2 = 4$  if and only if  $(S, L)$  defines a quartic hypersurface in  $\mathbb{P}^3$ , and  $L^2 = 6$  if and only if  $(S, L)$  defines the complete intersection of a quadric equation and a cubic equation in  $\mathbb{P}^4$ . Therefore  $L^2 \geq 8$  except these two cases.

**Theorem 4.2** (F. J. Gallego and B. P. Purnaprajna). *Let  $S$  be a K3 surface and let  $L \in \text{Pic}S$  be a very ample line bundle.*

- (1) *If  $L^2 = 4$  or 6, then  $(S, L^\ell)$  satisfies property  $N_{\ell-1}$ .*
- (2) *If  $L^2 \geq 8$  and the general member of  $|L|$  is non-trigonal, then  $(S, L^\ell)$  satisfies property  $N_\ell$ .*

Indeed their result is about the case when  $L$  is ample and base point free. Clearly  $(\dagger)$  refines or reproves their result for  $\ell \geq 3$ .

(4.3.2) Let  $S \subset \mathbb{P}^{g-1}$  be a smooth linearly normal Enriques surface. Then  $g \geq 6$ .  $\mathcal{O}_S(1)$  is called a Reye polarization if  $g = 6$  and  $S$  lies on a quadric. In [GLM], the projective normality of Enriques surfaces is studied.

**Theorem 4.3** (L. Giraldo, A. F. Lopez and R. Muñoz). (1) *If  $g = 6$  and  $\mathcal{O}_S(1)$  is a Reye polarization, then it is  $j$ -normal for every  $j \geq 3$ , 4-regular, and its homogeneous ideal is generated by forms of degree  $\leq 3$ .*

- (2) *If  $g \geq 7$  or  $g = 6$  and  $\mathcal{O}_S(1)$  is not a Reye polarization, then  $S \subset \mathbb{P}^{g-1}$  is 3-regular.*

Therefore we have

**Corollary 4.4.** *Let  $S \subset \mathbb{P}^{g-1}$  be a smooth linearly normal Enriques surface.*

(1) *If  $g = 6$  and  $\mathcal{O}_S(1)$  is a Reye polarization, then  $\mathcal{O}_S(2)$  satisfies property  $N_1$ ,  $\mathcal{O}_S(3)$  satisfies property  $N_2$ , and  $\mathcal{O}_S(\ell)$ ,  $\ell \geq 4$ , satisfies property  $N_\ell$ .*

(2) *If  $g \geq 7$  or  $g = 6$  and  $\mathcal{O}_S(1)$  is not a Reye polarization, then  $\mathcal{O}_S(1)$  satisfies property  $N_0$ ,  $\mathcal{O}_S(2)$  satisfies property  $N_1$ ,  $\mathcal{O}_S(3)$  satisfies property  $N_4$ , and  $\mathcal{O}_S(\ell)$ ,  $\ell \geq 4$ , satisfies property  $N_\ell$ .*

*Proof.* (1) By the Riemann-Roch theorem, the degree of  $S \subset \mathbb{P}^{g-1}$  is 10. Therefore Theorem 2.14 in [GP1] guarantees that  $\mathcal{O}_S(2)$  satisfies property  $N_1$  and  $\mathcal{O}_S(3)$  satisfies property  $N_2$ . Also  $\mathcal{O}_S(\ell)$ ,  $\ell \geq 4$ , satisfies property  $N_\ell$  by Theorem 1.5.

(2) This follows immediately from Theorem 4.3 and (†).  $\square$

**4.4. Complex torus.** In [Ru1] and [Ru3], E. Rubei studies syzygies of a power of normally generated very ample line bundles on a complex torus. She prove that (1) for a normally generated line bundle  $L \in \text{Pic}X$  on a complex torus  $X$ ,  $(X, L^\ell)$  satisfies property  $N_{\ell-1}$ , and (2) for a normally presented line bundle  $L \in \text{Pic}X$  on  $X$ ,  $(X, L^\ell)$  satisfies property  $N_\ell$ . This result is refined by Theorem 1.5 as follows:

**Corollary 4.5.** *Let  $X$  be a complex torus of dimension  $n$  and let  $L \in \text{Pic}X$  be a very ample line bundle such that  $X \subset \mathbb{P}H^0(X, L)$  satisfies  $(n+1)$ -normality. Then*

(1)  *$L^{n+1}$  satisfies property  $N_n$ .*

(2)  *$L^\ell$  satisfies property  $N_\ell$  for  $\ell \geq n+2$ .*

*Proof.* Since  $(n+1)$ -normality holds for  $X \subset \mathbb{P}H^0(X, L)$ , it is  $(n+2)$ -regular by Kodaira vanishing theorem. Thus Theorem 1.5 guarantees the assertion.  $\square$

**4.5. A remark on a Rubei's result.** In [Ru2], Rubei consider the problem to see how property  $N_p$  propagates through powers. More precisely she prove that if  $(X, L)$  satisfies property  $N_p$ , then  $(X, L^\ell)$  satisfies property  $N_{\min\{\ell, p\}}$ . In this subsection, we prove the following:

**Theorem 4.6.** *Let  $X$  be a projective variety and let  $L \in \text{Pic}X$  be a very ample line bundle such that  $(X, L)$  satisfies property  $N_p$ . Then for  $\ell \geq 2$ ,  $(X, L^\ell)$  satisfies property  $N_k$  where*

$$k = \begin{cases} \min\{5, p\} & \text{for } \ell = 2, \\ \min\{4, p\} & \text{for } \ell = 3, \text{ and} \\ \min\{\ell, p\} & \text{for } \ell \geq 4. \end{cases}$$

*Proof.* We use the short exact sequence ( $\diamond$ ) in §3. As in the proof of Theorem 1.6.(3), we have the long exact sequence

$$H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{Z/\mathbb{P}^N}(j)) \rightarrow H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{X/\mathbb{P}^N}(j)) \rightarrow H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}(j\ell))$$

of cohomology groups. Since  $(X, L)$  satisfies property  $N_p$ , we have

$$H^1(\mathbb{P}^r, \bigwedge^k \mathcal{M}_\ell \otimes \mathcal{J}_{X/\mathbb{P}^r}(j\ell)) = 0 \quad \text{for } 1 \leq k \leq p \text{ and } j\ell \geq 2.$$

For the desired vanishing of  $H^1(\mathbb{P}^N, \bigwedge^k \mathcal{M}_{\mathbb{P}^N} \otimes \mathcal{J}_{Z/\mathbb{P}^N}(j))$ , see Theorem 1.2, Example 1.1 and Example 1.2. Also we refer the reader to see [OP].  $\square$

**Remark 4.5.1.** As in the case of Theorem 1.5, if Ottaviani-Paoletti's conjecture is true then Theorem 4.6 is automatically refined. That is, for  $\ell \geq 3$ ,  $(X, L^\ell)$  satisfies property  $N_{\min\{3\ell-3, p\}}$ .  $\square$

**Remark 4.5.2.** As remarked in [Ru2] and [Ru4] it is not true that if  $(X, L)$  satisfies property  $N_p$  then any power of  $L$  satisfies property  $N_p$ . Indeed Rubei exhibits a concrete example. Also one can find more examples in [Park2]. More precisely, let  $(X, L)$  be a ruled scroll over a smooth curve  $C$  with the projection morphism  $\pi : X \rightarrow C$ . For an arbitrary  $p \geq 1$ , one can find  $(X, L)$  which satisfies property  $N_p$ . Now assume that  $\dim(X) \geq 3$ . Then for  $\ell \geq 3$ ,  $(X, L^\ell)$  fails to satisfy property  $N_{3\ell-2}$ . Therefore if  $p \geq 3\ell - 2$ , then  $(X, L)$  satisfies property  $N_p$  while  $(X, L^\ell)$  fails to satisfy property  $N_p$ . For details, see Corollary 3.7 in [Park2].  $\square$

**4.6. Complete intersection.** Let  $X \subset \mathbb{P}^r$  be a normal variety of dimension  $\geq 1$  which is a complete intersection. Then it is projectively normal. For the property  $N_1$  of  $\mathcal{O}_X(\ell)$ , we have the following

**Corollary 4.7.** *Under the situation just stated, assume that the homogeneous ideal of  $X \subset \mathbb{P}^r$  is generated by forms  $F_1, \dots, F_e$  of degrees  $d_1 \leq d_2 \leq \dots \leq d_e$  where  $e = \text{codim}(X, \mathbb{P}^r)$ . Then  $(X, \mathcal{O}_X(\ell))$  satisfies property  $N_1$  if and only if  $\ell \geq \frac{d_e}{2}$ .*

*Proof.* If  $\ell \geq \frac{d_e}{2}$ , then  $\mathcal{O}_X(\ell)$  satisfies property  $N_1$  by Theorem 1.6. Conversely, assume that  $\mathcal{O}_X(\ell)$  satisfies property  $N_1$ . We follow notations in §3. From the short exact sequence

$$(\diamond) \quad 0 \rightarrow \mathcal{J}_{Z/\mathbb{P}^N} \rightarrow \mathcal{J}_{X/\mathbb{P}^N} \rightarrow \mathcal{J}_{X/\mathbb{P}^r} \rightarrow 0.$$

if  $\mathcal{J}_{X/\mathbb{P}^N}(2)$  is globally generated, then  $\mathcal{J}_{X/\mathbb{P}^r}(2\ell)$  is also globally generated. Since  $\mathcal{J}_{X/\mathbb{P}^r}(2\ell)$  is globally generated if and only if  $2\ell \geq d_e$ , the converse is proved.  $\square$

## REFERENCES

- [BEL] A. Bertram, L. Ein and R. Lazarsfeld, *Vanishing theorems, a theorem of Severi, and the equations defining projective varieties*, J. Amer. Math. Soc. 4 (1991), no. 3, 587–602.
- [EL] L. Ein and R. Lazarsfeld, *Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension*, Invent. Math. 111 (1993), 51–67.
- [E] D. Eisenbud, *The Geometry of Syzygies*, no. 229, Springer-Verlag New York, (2005)
- [EG] D. Eisenbud and S. Goto, *Linear free resolutions and minimal multiplicity*, Journal of Algebra 88 (1984), 89–133.
- [EGHP] D. Eisenbud, M. Green, K. Hulek and S. Popescu, *Restriction linear syzygies: algebra and geometry*, math.AG/0404516, to appear in Compositio Math.
- [GP1] F. J. Gallego and B. P. Purnaprajna, *Projective normality and syzygies of algebraic surfaces*, J. Reine Angew. Math. 506, (1999), 145–180.
- [GP2] F. J. Gallego and B. P. Purnaprajna, *Vanishing theorems and syzygies for K3 surfaces and Fano varieties*, J. Pure Appl. Algebra 146 (2000), no. 3, 251–265.
- [GLM] L. Giraldo, A. F. Lopez and R. Muñoz, *On the projective normality of Enriques surfaces. With an appendix by Lopez and Alessandro Verra*, Math. Ann. 324 (2002), no. 1, 135–158.
- [Green1] M. Green, *Koszul cohomology and the geometry of projective varieties I*, J. Differ. Geom., 19 (1984), 125–171.
- [Green2] M. Green, *Koszul cohomology and the geometry of projective varieties II*, J. Differ. Geom., 20 (1984), 279–289.

- [GLP] L. Gruson, R. Lazarsfeld and C. Peskine, *On a theorem of Castelnuovo and the equations defining projective varieties*, Invent. Math. 72 (1983), 491-506.
- [JPW] T. Jozefiak, P. Pragacz and J. Weyman, *Resolutions of determinantal varieties and tensor complexes associated with symmetric and antisymmetric matrices*, Asterisque 87-88, (1981) 109-189.
- [Kwak1] Sijong Kwak, *Castelnuovo Regularity of Smooth Projective Varieties of Dimension 3 and 4*, J. Alg. Geom. 7 (1998), 195-206.
- [Kwak2] Sijong Kwak, *Castelnuovo-Mumford regularity bound for smooth threefolds in  $\mathbb{P}^5$  and extremal examples*, J. Reine Angew. Math. 509 (1999), 21-34.
- [Kwak3] Sijong Kwak, *Generic projections, the equations defining projective varieties and Castelnuovo regularity*, Math. Z. 234 (2000), no. 3, 413-434.
- [Laz1] R. Lazarsfeld, *A sharp Castelnuovo bound for smooth surfaces*, Duke Math. J. 55 (1987), no. 2, 423-429.
- [Laz2] R. Lazarsfeld, *Positivity in algebraic geometry. I. Classical setting: line bundles and linear series.*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics 48. Springer-Verlag Berlin, (2004)
- [Mum1] David Mumford, *Lectures on curves on an algebraic surface*, Ann. of Math. Studies. 59 (1966).
- [Mum2] D. Mumford, *Varieties defined by quadratic equations*, Corso CIME in questions on Algebraic Varieties, Rome, (1970), 30-100.
- [Ran] Ziv Ran, *Local differential geometry and generic projections of threefolds*, J. Differ. Geom. 32 (1990), no. 1, 131-137.
- [OP] G. Ottaviani and R. Paoletti, *Syzygies of Veronese embeddings*, Compositio Math., 125 (2001), 31-37.
- [Park1] Euisung Park, *On higher syzygies of ruled surfaces*, math.AG/0401100, to appear in Trans. Amer. Math. Soc.
- [Park2] Euisung Park, *Higher syzygies of ruled varieties over a curve*, to appear in Journal of Algebra.
- [Ru1] E. Rubei, *On syzygies of abelian varieties*, Trans. Amer. Math. Soc. 352 (2000), no. 6, 2569-2579.
- [Ru2] E. Rubei, *A note on Property  $N_p$* , Manuscripta Math. 101 (2000), 449-455
- [Ru3] E. Rubei, *On Koszul rings, syzygies, and abelian varieties*, Comm. Algebra 29 (2001), no. 12, 5631-5640.
- [Ru4] E. Rubei, *A strange example on Property  $N_p$* , Manuscripta Math. 108 (2002), 135-137
- [Ru5] E. Rubei, *A result on resolutions of Veronese embeddings*, Ann. Univ. Ferrara Sez. VII 50 (2004) 151-165

EUISUNG PARK : SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, 207-43 CHEONGRYANGRI-DONG, DONGDAEMUN-GU, SEOUL 130-722, REPUBLIC OF KOREA,  
*E-mail address:* puserdos@kias.re.kr