# ON THE DYNAMICAL SYMMETRIC ALGEBRA OF AGEING: LIE STRUCTURE, REPRESENTATIONS AND APPELL SYSTEMS

# MALTE HENKEL ,\* RENÉ SCHOTT, STOIMEN STOIMENOV <del>,</del>\* JÉRÉMIE UNTERBERGER<sup>§</sup>

The study of ageing phenomena leads to the investigation of a maximal parabolic subalgebra of  $\text{conf}_3$  which we call  $\text{alt.}$  We investigate its Lie structure, prove some results concerning its representations and characterize the related Appell systems.

### 1. Introduction

Ageing phenomena occur widely in physics: glasses, granular systems or phase-ordering kinetics are just a few examples. While it is well-accepted that they display some sort of dynamical scaling, the question has been raised whether their non-equilibrium dynamics might posses larger symmetries than merely scale-invariance. At first sight, the noisy terms in the Langevin equations usually employed to model these systems might appear to exclude any non-trivial answer, but it was understood recently that provided the deterministic part of a Langevin equation is Galilei-invariant, then all observables can be exactly expressed in terms of multipoint correlation functions calculable from the deterministic part only.<sup>7</sup> It is therefore of interest to study the dynamical symmetries of non-linear partial differential equations which extend dynamical scaling. In this context, the so-called Schrödinger algebra  $\mathfrak{sch}$  has been shown to play an important rôle in phase-ordering kinetics. In what follows we shall restrict to one space dimension and we recall in figure 1 through a root diagram the definition of  $\mathfrak{sch}$  as a parabolic subalgebra of the conformal algebra  $\mathfrak{conf}_3$ .<sup>6</sup>

<sup>\*</sup>lpm (cnrs umr 7556), université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france

<sup>&</sup>lt;sup>†</sup>iecn and loria, université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france <sup>‡</sup>lpm (cnrs umr 7556), université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france and institute of nuclear research and nuclear energy, bulgarian academy of sciences, 1784 sofia, bulgaria

 $\S$ iecn, université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france



Figure 1. (a) Root diagram of the complexified conformal Lie algebra  $(\text{conf}_3)$ <sub>C</sub> and the labelling of its generators. The double circle in the center denotes the Cartan subalgebra. The generators of the maximal parabolic subalgebras (b) sch and (c) alt are indicated by the filled dots.

## 2. A brief perspective on the algebra alt

There is a classification of semi-linear partial differential equations with a parabolic subalgebra of  $\text{conf}_3$  as a symmetry.<sup>9</sup> Here we shall study the abstract Lie algebra alt, which is the other maximal parabolic subalgebra of  $\mathfrak{conf}_3$  (see figure 1) and its representations; we shall also see that, like the algebra sch, it can be embedded naturally in an infinite-dimensional Lie algebra  $W$  which is an extension of the algebra  $Vect(S^1)$  of vector fields on the circle. Quite strikingly, we shall find on our way a 'no-go theorem' that proves the impossibility of a conventional extension of the embedding alt  $\subset$  conf $_3.$ 

#### 2.1. The abstract Lie algebra alt

Elementary computations make it clear that (see figure 1 and  $D = 2X_0 - N$ )

$$
\mathfrak{alt} = \langle V_+, D, Y_{-\frac{1}{2}} \rangle \ltimes \langle X_1, Y_{\frac{1}{2}}, M_0 \rangle := \mathfrak{g} \ltimes \mathfrak{h} \tag{1}
$$

is a semi-direct product of  $\mathfrak{g} \simeq \mathfrak{sl}(2,\mathbb{R})$  by a three-dimensional commutative Lie algebra  $\mathfrak h$ ; the vector space  $\mathfrak h$  is the irreducible spin-1 real representation of  $\mathfrak{sl}(2,\mathbb{R})$ , which can be identified with  $\mathfrak{sl}(2,\mathbb{R})$  itself with the adjoint action. So one has the following

## Proposition 2.1:

- (1) alt  $\simeq$   $\mathfrak{sl}(2,\mathbb{R})\otimes\mathbb{R}[\varepsilon]/\varepsilon^2$ , where  $\varepsilon$  is a 'Grassmann' variable;
- (2) alt  $\simeq \mathfrak{p}_3$  where  $\mathfrak{p}_3 \simeq \mathfrak{so}(2,1) \ltimes \mathbb{R}^3$  is the relativistic Poincaré algebra in  $(2+1)$ -dimensions.

$$
levicoproc 205
$$

**Proof :** The linear map  $\Phi$  :  $\mathfrak{alt} \to \mathfrak{sl}(2,\mathbb{R}) \otimes \mathbb{R}[\varepsilon]/\varepsilon^2$  defined by

$$
\Phi(V_+) = L_1, \ \Phi(D) = L_0, \ \Phi(Y_{-\frac{1}{2}}) = L_{-1}
$$
  

$$
\Phi(X_1) = \frac{1}{2}L_1^{\varepsilon}, \ \Phi(Y_{\frac{1}{2}}) = L_0^{\varepsilon}, \ \Phi(M_0) = L_{-1}^{\varepsilon}
$$

is easily checked to be a Lie isomorphism.

In particular, the representations of  $\mathfrak{alt} \simeq \mathfrak{p}_3$  are well-known since Wigner studied them in the 30'es.

#### 2.2. Central extensions: an introduction

Consider any Lie algebra  $\mathfrak g$  and an antisymmetric real two-form  $\alpha$  on g. Suppose that its Lie bracket [ , ] can be 'deformed' into a new Lie bracket  $\left[\right,$  on  $\tilde{\mathfrak{g}} := \mathfrak{g} \times \mathbb{R}K$ , where  $[K, \mathfrak{g}] = 0$ , by putting  $[(X, 0), (Y, 0)] =$  $([X, Y], \alpha(X, Y))$ . Then  $\tilde{\mathfrak{g}}$  is called a *central extension* of  $\mathfrak{g}$ . The Jacobi identity is equivalent with the nullity of the totally antisymmetric threeform  $d\alpha : \Lambda^3(\mathfrak{g}) \to \mathbb{R}$  defined by

$$
d\alpha(X, Y, Z) = \alpha([X, Y], Z) + \alpha([Y, Z], X) + \alpha([Z, X], Y).
$$

Now we say that two central extensions  $\mathfrak{g}_1, \mathfrak{g}_2$  of  $\mathfrak{g}$  defined by  $\alpha_1, \alpha_2$  are equivalent if  $\mathfrak{g}_2$  can be gotten from  $\mathfrak{g}_1$  by substituting  $(X, c) \mapsto (X, c +$  $\lambda(X)$ )  $(X \in \mathfrak{g})$  for a certain 1-form  $\lambda \in \mathfrak{g}^*$ , that is, by changing the nonintrinsic embedding of  $\mathfrak g$  into  $\tilde{\mathfrak g}_1$ . In other words,  $\alpha_1$  and  $\alpha_2$  are equivalent if  $\alpha_2 - \alpha_1 = d\lambda$ , where  $d\lambda(X, Y) = \langle \lambda, [X, Y] \rangle$ . The operator d can be made into the differential of a complex (called Chevalley-Eilenberg complex), and the preceding considerations make it clear that the classes of equivalence of central extensions of  $\mathfrak g$  make up a vector space  $H^2(\mathfrak g) = Z^2(\mathfrak g)/B^2(\mathfrak g)$ , where  $Z^2$  is the space of cocycles  $\alpha \in \Lambda^2(\mathfrak{g}^*)$  verifying  $d\alpha = 0$ , and  $B^2$  is the space of *coboundaries*  $d\lambda$ ,  $\lambda \in \mathfrak{g}^*$ . We have the well-known

Proposition 2.2: The Lie algebra alt has no non-trivial central extension:  $H^2(\mathfrak{alt})=0.$ 

All this becomes very different when one embeds alt into an infinitedimensional Lie algebra.

# 2.3. Infinite-dimensional extension of alt

The Lie algebra  $Vect(S^1)$  of vector fields on the circle has a long story in mathematical physics. It was discovered by Virasoro<sup>10</sup> in the  $70'$ es that



 $Vect(S<sup>1</sup>)$  has a one-parameter family of central extensions which yield the so-called Virasoro algebra

$$
\text{vir} := \text{Vect}(S^1) \oplus \mathbb{R}K = \langle (L_n)_{n \in \mathbb{Z}}, K \rangle \tag{2}
$$

with Lie brackets

$$
[K, L_n] = 0, \quad [L_n, L_m] = (n - m)L_{n+m} + \delta_{n+m,0} \ c \ n(n^2 - 1)K \quad (c \in \mathbb{R})
$$

When  $c = 0$ , one retrieves Vect(S<sup>1</sup>) by identifying the  $(L_n)$  with the usual Fourier basis  $(e^{in\theta} d\theta)_{n\in\mathbb{Z}}$  of periodic vector fields on  $[0, 2\pi]$ , or with  $-z^{n+1} \frac{d}{dz}$  with  $z := e^{i\theta}$ . Note in particular that  $\langle L_{-1}, L_0, L_1 \rangle$  is isomorphic to  $\mathfrak{sl}(2,\mathbb{R})$ , and that the Virasoro cocycle restricted to  $\mathfrak{sl}(2,\mathbb{R})$  is 0, as should be (since  $\mathfrak{sl}(2,\mathbb{R})$  has no non-trivial central extensions).

It is tempting to embed  $\mathfrak{alt} \simeq \mathfrak{sl}(2,\mathbb{R}) \otimes \mathbb{R}[\varepsilon]/\varepsilon^2$  into the Lie algebra

$$
\mathcal{W} := \text{Vect}(S^1) \otimes \mathbb{R}[\varepsilon]/\varepsilon^2 = \langle L_n \rangle_{n \in \mathbb{Z}} \ltimes \langle L_n^{\varepsilon} \rangle_{n \in \mathbb{Z}},\tag{3}
$$

with Lie brackets

$$
[L_n, L_m] = (n-m)L_{n+m}, [L_n, L_m^{\varepsilon}] = (n-m)L_{n+m}^{\varepsilon}, [L_n^{\varepsilon}, L_m^{\varepsilon}] = 0.
$$

These brackets come out naturally putting  $W$  in the  $2 \times 2$ -matrix form

$$
L_n \mapsto \begin{pmatrix} L_n & 0 \\ 0 & L_n \end{pmatrix}, \ L_n^{\varepsilon} \mapsto \begin{pmatrix} 0 & L_n \\ 0 & 0 \end{pmatrix}
$$
 (4)

leading to straightforward generalizations (there exists a deformation of sv that can be represented as upper-triangular  $3 \times 3$  Virasoro matrices<sup>8</sup>).

In terms of the standard representations of  $Vect(S^1)$  as modules of  $\alpha$ densities  $\mathcal{F}_{\alpha} = \{u(z)(dz)^{\alpha}\}\$  with the action

$$
f(z)\frac{\mathrm{d}}{\mathrm{d}z}(u(z)(dz)^{\alpha}) = (fu' + \alpha f'u)(z)(dz)^{\alpha},\tag{5}
$$

we have

Proposition 2.3:  $\mathcal{W} \simeq \text{Vect}(S^1) \ltimes \mathcal{F}_{-1}.$ 

There are two linearly independent central extensions of  $W$ :

1. the natural extension to W of the Virasoro cocycle on  $Vect(S^1)$ , namely  $[ , ] = \widetilde{[ , ]}$  except for  $\widetilde{[L_n, L_{-n}]} = n(n^2-1)K+2nL_0$ . In other words, Vect( $S<sup>1</sup>$ ) is centrally extended, but its action on  $\mathcal{F}_{-1}$  remains unchanged;

2. the cocycle  $\omega$  which is zero on  $\Lambda^2(\text{Vect}(S^1))$  and  $\Lambda^2(\mathcal{F}_{-1})$ , and defined by  $\omega(L_n, L_m^{\varepsilon}) = \delta_{n+m,0} n(n^2 - 1)K$  on  $\text{Vect}(S^1) \times \mathcal{F}_{-1}$ .

A natural related question is: can one deform the extension of  $Vect(S^1)$ by the Vect( $S^1$ )-module  $\mathcal{F}_{-1}$ ? The answer is: no, thanks to the triviality<sup>35</sup>

of the cohomology space  $H^2(\text{Vect}(S^1), \mathcal{F}_{-1})$ . Hence, any Lie algebra struc-

$$
[(X,\widetilde{\phi),(Y,\psi)}] = ([X,Y]_{\text{Vect}(S^1)}, \text{ad}_{\text{Vect}(S^1)}X.\psi - \text{ad}_{\text{Vect}(S^1)}Y.\phi + B(X,Y))
$$

is isomorphic to the Lie structure of  $W$  (where B is an antisymmetric twoform on  $Vect(S^1)$ ). So one may say that W and its central extensions are natural objects to look at.

### 2.4. Some results on representations of  $W$

ture [, ] on the vector space  $\text{Vect}(S^1) \oplus \mathcal{F}_{-1}$  such that

We now state two results which may deserve deeper thoughts and will be developed in the future.

Proposition 2.4 ('no-go theorem'): There is no way to extend the usual representation of alt as conformal vector fields into an embedding of W into the Lie algebra of vector fields on  $\mathbb{R}^3$ .

**Proposition 2.5:** The infinite-dimensional extension W of the algebra  $\mathfrak{alt}_1$ is a contraction of a pair of commuting Virasoro algebras  $\mathfrak{vir} \oplus \overline{\mathfrak{vir}} \to \mathcal{W}$ . In particular, we have the explicit differential operator representation

$$
L_n = -t^{n+1}\partial_t + (n+1)t^n r \partial_r - (n+1)xt^n - n(n+1)\gamma t^{n-1}r
$$
  

$$
L_n^{\varepsilon} = -t^{n+1}\partial_r - (n+1)\gamma t^n
$$
 (6)

where x and  $\gamma$  are parameters and  $n \in \mathbb{Z}$ .

## 3. Appell systems

**Definition 3.1.** Appell polynomials  $\{h_n(x); n \in \mathbb{N}\}\$  on  $\mathbb{R}$  are usually characterized by the two conditions

- $h_n(x)$  are polynomials of degree n,
- $Dh_n(x) = nh_{n-1}(x)$ , where D is the usual derivation opertor.

Interesting examples are furnished by the shifted moment sequences

$$
h_n(x) = \int_{-\infty}^{\infty} (x+y)^n p(\mathrm{d}y)
$$

where  $p$  is a probability measure on  $\mathbb R$  with all moments finite.

This definition generalizes to higher dimensions. On non-commutative algebraic structures, the shifting corresponds to left or right multiplication and in general,  $\{h_n\}$  is not a family of polynomials. We shall call it *Appell* systems. 2

Appell systems of the Schrödinger algebra  $\mathfrak{sch}$  have been investigated<sup>1</sup> but the algebra alt requires a specific study. alt has the following Cartan decomposition:

$$
\mathfrak{alt} = \mathfrak{P} \oplus \mathfrak{K} \oplus \mathfrak{L} = \{Y_1, X_1\} \oplus \{Y_0, X_0\} \oplus \{Y_{-1}, X_{-1}\} \tag{7}
$$

and there is a one to one correspondence between the subalgebras  $\mathfrak P$  and **L**. Write  $X \in \mathfrak{alt}$  in the form  $X = \sum_{i=1}^{6} \alpha_i b_i$ , where  $\{b_i, i = 1, \ldots, 6\}$  is a basis of alt. The  $\alpha_i$  are called coordinates of the first kind.

Here we use the basis  $b_1 = Y_1$ ,  $b_2 = X_1$ ,  $b_3 = Y_0$ ,  $b_4 = X_0$ ,  $b_5 = Y_{-1}$ ,  $b_6 = Y_1$ . Let ALT be the simply connected Lie group corresponding to alt. Group elements in a neighbourhood of the identity can be expressed as

$$
e^X = e^{A_1b_1} \dots e^{A_6b_6}
$$

The  $A_i$  are called *coordinates of the second kind.* 

Referring to the decomposition (7), we specialize variables, writing  $V_1, V_2, B_1, B_2$  for  $A_1, A_2, A_5, A_6$  respectively. Basic for our approach is to establish the partial group law:  $e^{B_1Y_{-1}+B_2X_{-1}}e^{V_1Y_1+V_2X_1} = ?$ . We get

$$
B_1Y_{-1} + B_2X_{-1} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ -B_2 & 0 & -B_1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -B_2 & 0 \end{pmatrix} , \quad V_1Y_1 + V_2X_1 = \begin{pmatrix} 0 & V_2 & 0 & V_1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & V_2 \\ 0 & 0 & 0 & 0 \end{pmatrix} (8)
$$

and finally:

$$
e^{B_1Y_{-1}+B_2X_{-1}}e^{V_1Y_1+V_2X_1} = \begin{pmatrix} 1 & V_2 & 0 & V_1 \ -B_2 1-B_2V_2-B_1-B_2V_1-B_1V_2 \ 0 & 0 & 1 & V_2 \ 0 & 0 & -B_2 & 1-B_2V_2 \end{pmatrix}
$$
(9)

Proposition 3.1. In coordinates of the second kind, we have the Leibnitz formula,

$$
g(0,0,0,0,B_1,B_2)g(V_1,V_2,0,0,0,0) = g(A_1, A_2, A_3, A_4, A_5, A_6) =
$$
  
\n
$$
g(\frac{B_1V_2^2 + V_1}{(1 - B_2V_2)}, \frac{V_2}{(1 - B_2V_2)}, -2\frac{B_1V_2 + B_2V_1}{(1 - B_2V_2)}, 2\ln(1 - B_2V_2),
$$
  
\n
$$
\frac{B_1 - 2B_1B_2V_2 - B_2^2V_1}{(1 - B_2V_2)^2}, \frac{B_2}{(1 - B_2V_2)})
$$
\n(10)

Now we are ready to construct the representation space and basis-the canonical Appell system. To start, define a vacuum state  $\Omega$ . The elements  $Y_1, X_1$ of  $\mathfrak P$  can be used to form basis elements

$$
|jk\rangle = Y_1^j X_1^k \Omega, j, k \geq 0
$$
\n(11)

of a Fock space  $\mathfrak{F} = \text{span}\{|jk\rangle\}$  on which  $Y_1, X_1$  act as raising operators,  $Y_{-1}, X_{-1}$  as lowering operator and  $Y_0, X_0$  as multiplication with the constants  $\gamma$ , x (up to the sign) correspondingly. That is,

$$
Y_1\Omega = |10\rangle, X_1\Omega = |01\rangle
$$
  
\n
$$
Y_{-1}\Omega = 0, X_{-1}\Omega = 0
$$
  
\n
$$
Y_0\Omega = -\gamma|00\rangle, X_0\Omega = -x|00\rangle
$$
\n(12)

The goal is to find an abelian subalgebra spanned by some selfadjoint operators acting on representation space, just constructed. Such a twodimensional subalgebra can be obtained by an appropriate "turn" of the plane  $\mathfrak P$  in the Lie algebra, namely via the adjoint action of the group element formed by exponentiating  $X_{-1}$ . The resulting plane,  $\mathfrak{P}_{\beta}$  say, is abelian and is spanned by

$$
\bar{Y}_1 = e^{\beta X_{-1}} Y_1 e^{-\beta X_{-1}} = Y_1 - 2\beta Y_0 + \beta^2 Y_{-1}
$$
\n
$$
\bar{X}_1 = e^{\beta X_{-1}} X_1 e^{-\beta X_{-1}} = X_1 - 2\beta X_0 + \beta^2 X_{-1}
$$
\n(13)

Next we determine our canonical Appell systems. We apply the Leibniz formula (10) with  $B_1 = 0, B_2 = \beta, V_1 = z_1, V_2 = z_2$  and (12). This yields

$$
e^{z_1\bar{Y}_1}e^{z_2\bar{X}_1}\Omega = e^{\beta X_{-1}}e^{z_1Y_1}e^{z_2X_1}e^{-\beta X_{-1}}\Omega = e^{\beta X_{-1}}e^{z_1Y_1}e^{z_2X_1}\Omega =
$$
  
= 
$$
e^{\frac{z_1Y_1}{(1-\beta z_2)^2}}e^{\frac{z_2X_1}{(1-\beta z_2)}}e^{\frac{2\gamma\beta z_1}{(1-\beta z_2)}}(1-\beta z_2)^{-2x}\Omega
$$
 (14)

To get the generating function for the basis  $|jk\rangle$  set in equation (14)

$$
v_1 = \frac{z_1}{(1 - \beta z_2)^2}, v_2 = \frac{z_2}{(1 - \beta z_2)}
$$
(15)

Substituting throughout, we have

Proposition 3.2. The generating function for the canonical Appell system,  $|jk\rangle = Y_1^j X_1^k \Omega$  is

$$
e^{v_1 Y_1 + v_2 X_1} \Omega = exp(y_1 \frac{v_1}{(1 + \beta v_2)^2}) exp(y_2 \frac{v_2}{(1 + \beta v_2)}) exp(-\frac{2\gamma \beta v_1}{(1 - \beta v_2)})
$$
  
(1 +  $\beta v_2$ )<sup>-2x</sup> \Omega (16)

where we identify  $\bar{Y}_1 \Omega = y_1 \cdot 1$  and  $\bar{X}_1 \Omega = y_2 \cdot 1$  in the realization as function of  $y_1, y_2$ .

With  $v_1 = 0$ , we recognize the generating function for the Laguerre polynomials, while  $v_2 = 0$  reduces to the generating function for Hermite polynomials.

7

### References

- 1. P. Feinsilver, Y. Kocik and R. Schott, Representations of the Schrödinger Algebra and Appell Systems, Progress of Physics, 52 (2004) 343-359.
- 2. P. Feinsilver and R. Schott, Algebraic Structures and Operator Calculus, Vol.3: Representations of Lie Groups. Kluwer Academic Publishers, Dordrecht, 1993.
- 3. D. B. Fuks, Cohohomology of infinite-dimensional Lie algebras, Contemporary Soviet Mathematics, Consultants Bureau, New York, 1986.
- 4. M. Henkel, R. Schott, S. Stoimenov and J. Unterberger, On the dynamical symmetric algebra of ageing: Lie structure, representations and Appell systems. Prépublication Institut Elie Cartan, 2005.
- 5. L. Guieu, C. Roger, preprint, available on http://www.math.univ-montp2.fr/~guieu/The Virasoro Project/Phase1/.
- 6. M. Henkel, J. Unterberger, Schrödinger-invariance and space-time symmetries, Nuclear Physics B660 (2003) 407-435.
- 7. A. Picone and M. Henkel, Local scale-invariance and ageing in noisy systems, Nuclear Physics B688 (2004) 217-265.
- 8. C. Roger and J. Unterberger, in progress (2005).
- 9. S. Stoimenov and M. Henkel, Dynamical symmetries of semi-linear Schrödinger and diffusion equations, Nuclear Physics B723 (2005) 205-233.
- 10. M. Virasoro, Phys. Rev. D1, 2933-2936 (1970).