ON THE POLAR DECOMPOSITION OF CIRCULAR VARIABLES

Teodor Banica

We find an elementary proof for Voiculescu's theorem on the polar decomposition of circular variables.

0. INTRODUCTION: In non commutative probability theory (see [1]) an important role is played by the Haar-unitaries, which appear naturally in the von Neumann algebras of free groups $W^*(\mathbb{Z}^{*n})$, and by the circular variables, which appear naturally in the algebras of creation operators on the full Fock spaces $W^*(\mathbb{N}^{*n})$.

In [2] Voiculescu finds the polar decomposition of the circular variables. In order to work at the same time with circular variables and with Haar-unitaries, he uses approximation by random Gaussian matrices.

Following an idea of G.Skandalis, we give in this note an elementary proof of this theorem: it turns out that the result is an immediate consequence of some combinatorial properties of the monoid $\mathbf{Z} * \mathbf{N}$.

In what follows, we introduce (see 4) a certain class of monoids, which contains \mathbf{N} , all the groups, and is stable under free products (see 8). Consequently, for such monoids M, many variables having interesting *-distributions (semicircular and circular variables, Haar-unitaries) appear naturally in the algebras $W^*(M)$.

For such monoids, some combinatorial properties (see 5 and 7) allow us to find (and easily manipulate) "many" circular systems in their algebras $W^*(M)$ (see 6.2). The polar decomposition of the circular variables follows easily.

1. NOTATION: In what follows, we will denote (by abuse of language) by "monoid" a countable unital monoid, which can be embedded in a group. For such a monoid M, the symbol M^{-1} will denote, when there are no confusions, the subset $\{m^{-1}, m \in M\}$ of some group containing M.

2. DEFINITIONS: Let M be a monoid and $l^2(M)$ the Hilbert space of square

summable functions from M to \mathbf{C} , with $(\delta_m)_{m \in M}$ the canonical orthonormal basis. Using the left simplifiability of M one can define, as for discrete groups, the embedding of monoids $(M, \cdot) \rightarrow (B(l^2(M)), \circ)$ by $\lambda_M(m)\delta_n = \delta_{mn}$. Let $W^*(M)$ be the Von Neumann algebra generated by $\lambda_M(M)$. Together with the canonical state $\tau_M(T) = \langle T\delta_e, \delta_e \rangle$ it is a non commutative W^* -probability space.

3.1. REMARK: The operators in $\lambda_M(M)$ are isometries, but not necessarely unitaries, as in the group case. Indeed, for every $m \in M$, $\lambda_M(m)^*$ is given by $\lambda_M(m)^*(\delta_n) = \sum_{x \in M} \langle \lambda_M(m)^* \delta_n, \delta_x \rangle \delta_x = \sum_{x \in M} \delta_{n,mx} \delta_x$, so that $\lambda_M(m)^* \lambda_M(m) = 1$.

3.2. REMARK: It is easy to see that $l^2(\mathbf{N}^{*I})$ is the full Fock space over \mathbf{C}^I . By this identification, $(W^*(\mathbf{N}^{*I}), \tau_{\mathbf{N}^{*I}})$ is the algebra of creation operators, with the canonical state associated to the vacuum vector.

3.3. REMARK: Let $M \subset N$ be monoids (so $l^2(M) \subset l^2(N)$). For $m, m' \in M$ one has $\lambda_M(m)\delta_{m'} = \lambda_N(m)\delta_{m'}$, so if we suppose M(N - M) = N - M then $\lambda_M(m)^*\delta_{m'} = \sum_{x \in M} \delta_{m',mx}\delta_x = \sum_{x \in N} \delta_{m',mx}\delta_x = \lambda_N(m)^*\delta_{m'}$. In particular, if $m_1...m_k \in M$ and $\alpha_1...\alpha_k$ are exponents $\in \{1, *\}$ then $\lambda_M(m_1)^{\alpha_1}...\lambda_M(m_k)^{\alpha_k}\delta_e = \lambda_N(m_1)^{\alpha_1}...\lambda_N(m_k)^{\alpha_k}\delta_e$.

It follows that if $M \subset N$ are monoids such that M(N - M) = N - M then for every family $\{a_i\}_{i \in I}$ of elements in M, the *-distribution joint to $\{\lambda_N(a_i)\}_{i \in I}$ is equal to the *-distribution joint to $\{\lambda_M(a_i)\}_{i \in I}$.

3.4. REMARK: Let $\prod_{i \in I} M_i$ be a direct product of monoids and $a \in M_k$, $k \in I$. Then, by (3.3) the *-distribution of $\lambda_{M_k}(a) \in W^*(M_k)$ is equal to the *-distribution of $\lambda_{\prod M_i}(a) \in W^*(\prod M_i)$. Moreover, let $b \in M_j$ with $j \neq k$. Then it is easy to see that $\lambda_{\prod M_i}(a)$ and $\lambda_{\prod M_i}(b)$ are independent.

3.5. REMARK: Let $*_{i \in I} M_i$ be a free product of monoids and $a \in M_k$, $k \in I$. Then, by (3.3) the *-distribution of $\lambda_{M_k}(a) \in W^*(M_k)$ is equal to the *-distribution of $\lambda_{*M_i}(a) \in W^*(*M_i)$.

We are now interested to find semicircular variables in the algebras of monoids. Let us introduce some definitions related to the combinatorics of free monoids:

4. **DEFINITION:** Let N be a monoid. Consider the following order relation on it: $a \leq_N b$ if and only if $b \in aN$. We say that N is in the class E if it satisfies one of the (obvious) equivalent conditions:

- (4.1.) for \preceq_N every bounded subset is totally ordered.
- (4.2.) $(a \leq c, b \leq c \Rightarrow a \leq b \text{ or } b \leq a).$
- (4.3.) $aN \cap bN \neq \emptyset \Rightarrow aN \subset bN$ or $bN \subset aN$.
- $(4.4.) NN^{-1} \cap N^{-1}N = N \cup N^{-1}.$

5. **DEFINITION:** Let $(a_i)_{i \in I}$ be a family of elements in a monoid N. We call it "code" if it satisfies the following conditions: (5.1.) the monoid M generated by the a_i 's is isomorphic to \mathbf{N}^{*I} by $a_i \mapsto e_i$.

(5.2.) M(N - M) = N - M.

We call it "prefix" if it satisfies the following condition:

(5.3.) $a_i \in a_j N \Rightarrow i = j$ (i.e. the a_i 's are not comparable by \leq_N).

6.1. REMARK: These are extensions of the classical notions of code and prefix, which already appeared in the combinatorial theory of free monoids. A well known result (see [3]) asserts that on free monoids the prefixes are the codes. We will extend this result to all monoids in the class E.

6.2. **REMARK:** Let $(a_i, b_i)_{i \in I}$ be a code. Then by (3.2) and (3.3), the family $(\lambda_N(a_i), \lambda_N(b_i))_{i \in I}$ has the same *-distribution as a family of creation operators associated to a family of 2*I* orthonormal vectors, acting on the Fock space. In particular $(1/2(\lambda_N(a_i) + \lambda_N(b_i)^*))_{i \in I}$ is a circular family. Thus the following proposition is a nice criterion for finding circular systems in the algebras $W^*(N)$ of monoids (in the class *E*).

7. PROPOSITION: For a monoid $N \in E$, a family $(a_i)_{i \in I} \subset N$ having at least two elements is a prefix if and only if it is a code.

PROOF: Let $(a_i)_{i \in I}$ be a code which is not a prefix. Suppose for instance that $a_i = a_j n$ with $i \neq j, n \in N$. By (5.2) n is in the monoid M generated by the a_k 's and $a_i = a_j n$ with $i \neq j$, so M cannot be free, contradiction.

Suppose now that $(a_i)_{i\in I}$ is a prefix and let $A = a_{i_1}^{\alpha_1} \dots a_{i_n}^{\alpha_n} m = a_{j_1}^{\beta_1} \dots a_{j_s}^{\beta_s}$ with $m \in N$. One has $a_{i_1} \leq A$, $a_{j_1} \leq A$, so by (4.2) and (5.3), $i_1 = j_1$. We simplify A to the left by a_{i_1} (recall that all the monoids we consider are bisimplifiable); a recurrence on $\Sigma \alpha_i$ shows that $n \leq s$, $a_{i_k} = a_{j_k} (\forall k \leq n)$, $\alpha_k = \beta_k (\forall k < n)$, $\alpha_n \leq \beta_n$, $m = a_{j_n}^{\beta_n - \alpha_n} a_{j_{n+1}}^{\beta_{n+1}} \dots a_{j_s}^{\beta_s}$. Finally, m is in the monoid generated by the a_i 's, so (5.2) is true. Moreover, for m = e we obtain n = s, $a_{j_k} = a_{i_k}$, $\alpha_k = \beta_k$, $(\forall k \leq n)$ so the a_i 's generate freely M and $(a_i)_{i \in I}$ is a code.

8. **PROPOSITION:** (8.1.) all the groups are in E.

(8.2.) the positive parts of totally ordered abelian groups are in E.

(8.3.) if G is a group and $M \in E$, then $M \times G \in E$.

(8.4.) if A_1 , A_2 are in E, then the free product $A_1 * A_2$ is in E.

PROOF: (8.1) et (8.2) are obvious (M is totally ordered by \preceq_M).

Remark: Reciprocally, if M is an abelian monoid in E, then one can easily construct a total order on its Grothendieck group K(M) such that $M = \{g \in K(M), g \succeq 0\}$.

Let G be a group and $M \in E$. Using (4.4) we have $(M \times G)(M \times G)^{-1} \cap (M \times G)^{-1}(M \times G) = (M \times G)(M^{-1} \times G) \cap (M^{-1} \times G)(M \times G) = (MM^{-1} \times G) \cap (M^{-1}M \times G) = (MM^{-1} \cap M^{-1}M) \times G = (M \cup M^{-1}) \times G = (M \times G) \cup (M^{-1} \times G) = (M \times G) \cup (M \times G)^{-1}$, so we proved (8.3).

We prove now (8.4). Let $a, b, c \in A_1 * A_2$ such that ab = c.

Write $a = x_1...x_n$, $b = y_1...y_m$, $c = z_1...z_p$ as reduced words. Let *s* be such that $x_ny_1 = 1,...,x_{n-s+1}y_s = 1$, but $x_{n-s}y_{s+1} \neq 1$. Let $u = x_{n-s+1}...x_n = (y_1...y_s)^{-1}$. Then $c = ab = x_1...x_{n-s}y_{s+1}...y_m$. Let $i \in \{1, 2\}$ be such that $z_{n-s} \in A_i$. There are two cases:

- if $x_{n-s} \in A_1$ and $y_{s+1} \in A_2$ or if $x_{n-s} \in A_2$ and $y_{s+1} \in A_1$, then $x_1 \dots x_{n-s} y_{s+1} \dots y_m$ is a reduced word. In particular, $x_1 = z_1, \dots, x_{n-s} = z_{n-s}$. Thus $a = z_1 \dots z_{n-s} u$ with u invertible. - if $x_{n-s}, y_{s+1} \in A_i$ then $x_1 = z_1, \dots, x_{n-s-1} = z_{n-s-1}$ and $x_{n-s} y_{s+1} = z_{n-s}$. In this

case $a = z_1...z_{n-s-1}x_{n-s}u$ with u invertible.

Remark that in both cases we obtained that a is of the form $z_1...z_f xu$ for some f, with u invertible and such that if $z_{f+1} \in A_i$, then there exists $y \in A_i$ with $xy = z_{f+1}$ (take f = n - s - 1 and $x = z_{n-s}, y = 1$ in the first case, $x = x_{n-s}, y = y_{s+1}$ in the second one).

Suppose now that $A_1, A_2 \in E$ and let $a, b, a', b' \in A_1 * A_2$ such that ab = a'b'. Let $z_1...z_p$ be the decomposition of ab = a'b' as a reduced word. Then we can decompose $a = z_1...z_f xu$ and $a' = z_1...z_{f'}x'u'$ as above. We have to show that a = a'm or that a' = amfor some $m \in A_1 * A_2$. There are three cases:

- if f < f', then $a' = au^{-1}yz_{f+2}...z_{f'}x'u'$.

- if f' < f, then $a = a'u'^{-1}z_{f'+2}...z_fxu$.

- if f = f', then $xy = x'y' = z_{f+1} \in A_i$ for some $i \in \{1, 2\}$. As $A_i \in E$, we have that x = x'm or x' = xm for some $m \in A_i$, so that $a = a'u'^{-1}mu$ or a' = aumu'.

The proof of (8.4) is now complete.

9. PROPOSITION: (9.1.) Let $M \subset N$ be two monoids in the class E such that M(N - M) = N - M. Let $\lambda = \lambda_N$. Then every element x of the *-algebra generated by $\lambda(M)$ could be written as $x = \sum a_i \lambda(p_i) \lambda(q_i)^*$, with $p_i, q_i \in M$.

(9.2.) Let $A, B \in E$, $\lambda = \lambda_{A*B}$, $\tau = \tau_{A*B}$, and let x be an element of the *-algebra generated by $\lambda(A)$ such that $\tau(x) = 0$. Denote by W_A the set of reduced words beginning by an element of A, and by W_B the set of reduced words beginning by an element of B. Then $x \text{ maps } l^2(W_B \cup \{e\}) \text{ into } l^2(W_A).$

(9.3.) Let $A, B \in E$. Then $\lambda_{A*B}(A)$ and $\lambda_{A*B}(B)$ are *-free.

PROOF: It is enough to prove (9.1) for $x = \lambda(m)^*\lambda(n)$ with $m, n \in M$; the general case will follow easily. Remark that $x = \lambda(m)^*\lambda(n)$ is different from 0 iff $\exists a, b \in N$ such that $\langle \lambda(m)^*\lambda(n)\delta_a, \delta_b \rangle \neq 0$, i.e. if na = mb. By (4.2), $\exists c \in N$ with n = mc or with m = nc. Moreover, as M(N - M) = N - M, it follows that $c \in M$. Thus $x = \lambda(m)^*\lambda(n) \neq 0 \Rightarrow x = \lambda(c)$ or $x = \lambda(c)^*$ with $c \in M$, and this finishes the proof.

For proving (9.2), we apply (9.1) with M = A and N = A * B for writing $x = \sum a_i \lambda(p_i) \lambda(q_i)^*$, with $p_i, q_i \in A$. Remark that $\tau(\lambda(p_i)\lambda(q_i)^*) = \sum_x \delta_{e,p_ix} \delta_{e,q_ix}$ is nonzero iff $p_i = q_i$ =invertible, and in this case $\lambda(p_i)\lambda(q_i)^* = 1$. As $\tau(x) = 0$, it follows that we may write $x = \sum a_i \lambda(p_i)\lambda(q_i)^*$, such that $\tau(\lambda(p_i)\lambda(q_i)^*) = 0$ for every *i*. By linearity, it is enough

to prove (9.2) for $x = \lambda(p_i)\lambda(q_i)^*$.

Let $m \in W_B \cup \{e\}$ and suppose that $x\delta_m \neq 0$. Then $\lambda(q_i)^*\delta_m \neq 0$ implies that $m = q_i c$ for some word $c \in A * B$. As $q_i \in A$ and $m \in W_B \cup \{e\}$, it follows that q_i is invertible. In this case, $x\delta_m = \delta_{p_i q_i^{-1}m} \in l^2(W_A)$ (recall that $p_i q_i^{-1} = 1 \Rightarrow \tau(x) = 1$).

Finally, (9.3) follows from (9.2). Indeed, let $P = x_n \dots x_1$ be a product of elements in $ker(\tau)$, such that x_{2k} is in the *-algebra generated by $\lambda(B)$ and x_{2k+1} is in the *-algebra generated by $\lambda(A)$. Then $x_1\delta_e \in l^2(W_A)$, so that $x_2x_1\delta_e \in l^2(W_B)$ etc. By a recourse, $P\delta_e$ is in $l^2(W_A)$ or in $l^2(W_B)$, and this implies that $\tau(P) = 0$.

10. PROPOSITION: Consider (in some non commutative probability space) a Haar-unitary u, *-free from a semicircular s. Then us is a circular variable.

PROOF: Denote by z the image of $1 \in \mathbb{Z}$ and by n the image of $1 \in \mathbb{N}$ by the canonical embeddings into the free product $\mathbb{Z} * \mathbb{N}$. Let $\lambda = \lambda_{\mathbb{Z}*\mathbb{N}}$. By (8), $\mathbb{Z} * \mathbb{N} \in E$. (zn, nz^{-1}) is obviously a prefix, so by the criterion (7), it is a code. By (6.2), $1/2(\lambda(zn) + \lambda(nz^{-1})^*)$ is circular. But $1/2(\lambda(zn) + \lambda(nz^{-1})^*) = us$ where:

- $u := \lambda(z)$ is a Haar-unitary (see (3.5)).

- $s := 1/2(\lambda(n) + \lambda(n)^*)$ is semicircular (see (3.5) and (3.2)).

- u and s are *-free (by (9.3)).

11. **PROPOSITION** : Let s = dq be the polar decomposition of a semicircular variable in some W^{*}-probability space with faithful normal state. Then (q, d) are independent, q is quarter-circular and the distribution of d is given by $\mu_d(X^{2k}) = 1$ and $\mu_d(X^{2k+1}) = 0$.

PROOF: Look for instance at the semicircular $(x \mapsto x) \in (L^{\infty}[-1, 1], \gamma_{0,1})$.

12. THEOREM [2]: Let x = vb be the polar decomposition of a circular variable in some W^{*}-probability space with faithful normal state. Then v is Haar-unitary, b is quartercircular and (v, b) is a *-free pair.

PROOF: The theorem is a fairly simple consequence of the proposition 10.

Consider the group $G = \mathbf{Z} * (\mathbf{Z} \times \mathbf{Z}/2\mathbf{Z})$ and denote by z, t, a the images of $1 \in \mathbf{Z}$, $(1, \hat{0}) \in \mathbf{Z} \times (\mathbf{Z}/2\mathbf{Z})$ and $(0, \hat{1}) \in \mathbf{Z} \times (\mathbf{Z}/2\mathbf{Z})$ by the canonical embeddings into G.

Let $u = \lambda_G(z)$, $d = \lambda_G(a)$ and choose a quarter-circular $q \in W^*(\lambda_G(t))$. Then (q, d) are independent, and the distribution of d is given by $\mu_d(X^{2k}) = 1$ and $\mu_d(X^{2k+1}) = 0$ (see (3.4)). By (11), dq is semicircular, so by (10), c := udq is circular, and:

- the module of c is q, which is a quarter-circular.

- the polar part of c is ud, which is obviously a Haar-unitary.

- consider the automorphism ψ of G which is the identity on $\mathbf{Z} \times \mathbf{Z}/2\mathbf{Z}$ and sends $z \mapsto za$. It extends to a trace-preserving automorphism $\tilde{\psi}$ of $W^*(G)$ which sends $u \mapsto ud$ and $q \mapsto q$. As u and q are *-free, it follows that ud and q are *-free.

13. We give in the end another kind of result which seems to be non-trivial, but

which follows easily by using our formalism:

Let $P \in \mathbb{C}[X]$ be a polynomial. We can write $P = \sum_{j=1}^{j=k} m_j X^{p_j}$ with $p_i \neq p_j, m_j \neq 0$.

Let z and $n_1...n_k$ be the images of $1 \in \mathbb{Z}$ and of the $1 \in \mathbb{N}$'s by the canonical embeddings into the free product $\mathbb{Z} * \mathbb{N}^{*k}$; let $\lambda = \lambda_{\mathbb{Z}*\mathbb{N}^{*k}}$. By (7), $\{zn_1^{p_1}, ..., zn_k^{p_k}, n_1z^{-1}, ..., n_kz^{-1}\}$ is a prefix, so it is a code, and $\{\lambda(z)(\lambda(n_j^{p_j}) + \lambda(n_j)^*)/2\}_{1 \leq j \leq k}$ is a circular system.

Let a_j be complex numbers such that $m_j = a_j^{p_j+1}$. By [1], proposition 2.2., the sum $\sum a_j \lambda(z)(\lambda(n_j)^* + \lambda(n_j)^{p_j})$ is a (non centered) circular variable. By [1], example 3.4.3., the *R*-transform of $a_j(\lambda(n_j)^* + \lambda(n_j)^{p_j})$ is $m_j X^{p_j}$. The additivity of the *R*-transform ([1], theorem 3.2.3.), implies that the *R*-transform of $x = \sum a_j(\lambda(n_j)^* + \lambda(n_j)^{p_j})$ is *P*.

Thus, for any polynomial P, we can find a random variable x such that:

- the R-transform of x is P.

- if u is a Haar-unitary *-free from x, then ux is a (non centered) circular variable.

ACKNOWLEDGEMENTS: I would like to thank G. Skandalis, for directing this work and for many suggestions and advices; A. Boutet de Monvel, for invinting me in her laboratory during the acomplishment of this research; E. Germain, for many helpful suggestions; P. Biane, for pointing out an error in a preliminary version of this paper.

REFERENCES:

[1] Voiculescu, Dykema, Nica - Free random variables, CRM Monograph Series $n^{\circ}1$, AMS (1993)

[2] Voiculescu - Circular and semicircular systems and free product factors, Progress in Math. 92, Birkhäuser (1990)

[3] Lothaire - Combinatorics on Words, Addison-Wesley (1983)

Université Paris 7, Aile 45-55, 5^{eme} étage, 2 place Jussieu, 75251 Paris Cedex 05. banica@mathp6.jussieu.fr

AMS Classification: 46L50