# A note on the smoothness of energy-minimizing incompressible deformations

Nirmalendu Chaudhuri

Mathematical Sciences Research Institute 17 Gauss Way Berkeley, CA 94720-5070, USA chaudhur@msri.org

&

Centre for Mathematics and its Applications Australian National University Canberra, ACT 0200 Australia chaudhur@maths.anu.edu.au

#### Abstract

In this note we prove that any  $W^{1,2}$  mapping u in the plane that minimizes an appropriate quasiconvex energy functional subject to the Jacobian constraint det  $\nabla u = 1$  a.e., are necessarily Lipschitz. Furthermore we show that the minimizers corresponding to uniformly convex energy are affine and give an example of non-affine minimizers subject to affine boundary data corresponding to a convex energy. We also discuss the regularity issues in dimension greater than or equal to 3.

#### 1 Introduction

Let  $\Omega \subset \mathbb{R}^2$  be a bounded incompressible material body, that is, every  $W^{1,2}$  deformation of  $\Omega$  locally preserves its volume, in particular, the Jacobian of every such deformation is 1 almost everywhere. For an incompressible neo-Hookean material [Og 84], [BOP 92] such as vulcanized rubber, in the equilibrium, one is interested minimizing the potential energy

$$I[u] := \int_{\Omega} F(\nabla u), \qquad (1.1)$$

for incompressible  $W^{1,2}$  deformations  $u : \Omega \to \mathbb{R}^2$  with prescribed boundary conditions corresponding to a given bulk energy  $F : \mathbb{R}^{2 \times 2} \to \mathbb{R}$ . The simplest F is the Dirichlet energy  $F(X) := \frac{1}{2} \operatorname{tr} (X^T X)$ . Let us denote the incompressible or so-called the area-preserving mappings

$$\mathcal{A} := \{ u \in W^{1,2}(\Omega, \mathbb{R}^2) : \det \nabla u(x) = 1, \text{a.e. in } \Omega \},$$
(1.2)

 $u = (u^1, u^2), \ \nabla u = (u^i_{x_j})_{1 \le i,j \le 2}$ , the gradient and det  $\nabla u := u^1_{x_1} u^2_{x_2} - u^1_{x_2} u^2_{x_1}$ , is the Jacobian of u. A function  $f : \mathbb{R}^{m \times n} \to \mathbb{R}$  is said to be *quasiconvex* if

$$\int_{(0,1)^n} f(X + \nabla \phi(x)) \, dx \ge f(X)$$

for each  $m \times n$  matrices X and each smooth compactly supported  $\phi : (0,1)^n \to \mathbb{R}^m$ . This definition was introduced by Morrey [Mo 52], as a necessary and sufficient condition for weak lower semicontinuity of the energy functional associated to f with respect to uniform convergence of Lipschitz functions. Under the basic assumption that  $F : \mathbb{R}^{2\times 2} \to \mathbb{R}$  is smooth, quasiconvex with quadratic growth, together with the weak continuity [Mu 89] of the Jacobian, the functional I admits *local* (see for example [EG 99]) minimizers in the class  $\mathcal{A}$ . It remains a difficult problem (due to the hard Jacobian constraint) to understand the regularity properties of the local minimizes of I. Under the additional assumption (the so-called uniform quasiconvexity, see, [Ev 86])

$$\int_{(0,1)^2} F(X + \nabla \phi(x)) - F(X) \, dx \ge C \int_{(0,1)^2} |\nabla \phi|^2 dx \tag{1.3}$$

for some C > 0, for each  $2 \times 2$  matrices X and each smooth compactly supported  $\phi$ :  $(0,1)^2 \to \mathbb{R}^2$ , Evans and Gariepy [EG 99] proved that any *non-degenerate*, *Lipschitz area-preserving* local minimizers of I are  $C^{1,\alpha}$  on a dense open subset. It remains to understand whether area-preserving local minimizers are Lipschitz. Here we consider only the global minimizers. A map  $u \in \mathcal{A}$  is said to be global minimizer of I subject to its own boundary if

$$I[u] \leq I[v], \text{ for all } v \in \mathcal{A}.$$

**Theorem 1.1.** Let  $F : \mathbb{R}^{2 \times 2} \to \mathbb{R}$  be uniformly quasiconvex,  $C^2$ , and  $D^2F$  is bounded. Then global minimizers of I in the area-preserving class  $\mathcal{A}$  are Lipschitz. Furthermore, if F is uniformly convex and frame indifference then the minimizers are affine

The proof follows by reducing the minimization problem to a partial differential relation of the form

$$\nabla u(x) \in K$$
 a.e. in  $\Omega$ ,

for suitable subset K of  $\mathbb{R}^{2\times 2}$ . As a consequence of this observation and a theorem of Müller and Šverák [MS 96] on convex integration (also see, [DM 97]), we give an example of non-affine minimizers subject to affine boundary data corresponding to a convex (non-uniform) energy.

#### 2 Proof of Theorem 1.1.

We recall the set of area-preserving mappings

$$\mathcal{A} := \{ u \in W^{1,2}(\Omega, \mathbb{R}^2) : \det \nabla u(x) = 1, \text{a.e. in } \Omega \}.$$

Let  $SL(2) := \{ P \in \mathbb{R}^{2 \times 2} : \det P = 1 \}$ , the special linear group and

$$Z_{\min}(F) := \{ Q \in SL(2) : F(Q) = \min_{P \in SL(2)} F(P) \},$$
(2.4)

be the minimizing set. Since F is uniformly quasiconvex, the minimizing set  $Z_{\min}(F)$  is non-empty and compact. Without loss of generality assume  $|\Omega| = 1$ . Observe that for any  $v \in \mathcal{A}$ , we have

$$I[v] = \int_{\Omega} F(\nabla v) \ge \min\{F(P) : \det P = 1\}.$$
(2.5)

Let  $Q \in Z_{\min}(F)$  and u(x) = Qx, be a linear deformation. Then  $u \in \mathcal{A}$  and

$$I[u] = F(Q) = \min_{P \in SL(2)} F(P).$$

Therefore

$$\min_{v \in \mathcal{A}} I[v] = \min_{P \in SL(2)} F(P).$$
(2.6)

Hence  $u \in \mathcal{A}$  is a minimizer of I if and only if it satisfies the partial differential inclusion

$$\nabla u(x) \in Z_{\min}(F)$$
 a.e. in  $\Omega$ , (2.7)

Since  $Z_{\min}(F)$  is compact, u is Lipschitz. This proves first part of the theorem.

**Lemma 2.1.** Let  $F : \mathbb{R}^{2\times 2} \to \mathbb{R}$  be uniformly convex, that is,  $D^2F(X)Y : Y \ge 2\lambda|Y|^2$ for some  $\lambda > 0$ , for all  $X, Y \in \mathbb{R}^{2\times 2}$ . Suppose further, F(RX) = F(X) for each  $X \in \mathbb{R}^{2\times 2}$ , and each rotations R. Then the minimizing set  $Z_{\min}(F)$  is simply the coset SO(2)P, for some det P = 1, where  $SO(2) := \{R \in \mathbb{R}^{2\times 2} : R^TR = Id, \det R = 1\}$ , the special orthogonal group.

*Proof.* Here we follow the standard uniqueness arguments. Suppose there exists  $Q_1, Q_2 \in SL(2)$  that  $F(Q_1) = F(Q_2) = \min_{SL(2)} F(P)$ . Since F is frame indifferent, it follows that any

$$Q \in K := SO(2)Q_1 \cup SO(2)Q_2$$

also minimizes F over SL(2). We claim that  $Q_1$  and  $Q_2$  are conformally equivalent, i.e.,  $Q_1 = RQ_2$ , for some  $R \in SO(2)$ . Suppose,  $Q_1$  and  $Q_2$  are not conformally equivalent. Since det  $Q_1 = \det Q_2 = 1$ , a simple calculation shows that the cosets  $SO(2)Q_1$  and  $SO(2)Q_2$  are rank-one connected. Therefore for each  $P_1 \in SO(2)$  there exists  $P_2 \in SO(2)$  such that

$$\det \left( P_1 Q_1 - P_2 Q_2 \right) = 0.$$

Since  $X \mapsto \det X$  is linear along rank-one directions, it follows that

$$\det \left(\lambda P_1 Q_1 + (1-\lambda) P_2 Q_2\right) = 1, \text{ for all } 0 \le \lambda \le 1.$$

From the uniform convexity of F, we have

$$F(X) \ge F(Y) + DF(Y) : (X - Y) + \lambda |X - Y|^2$$
, for all  $X, Y \in \mathbb{R}^{2 \times 2}$ , (2.8)

for some  $\lambda > 0$ , where  $A : B := tr(A^T B)$  is the scalar product. By taking  $X = P_1 Q_1$ and  $Y = (P_1 Q_1 + P_2 Q_2)/2$ , and vice versa and adding these two inequalities, we obtain

$$F(P_1Q_1) + F(P_2Q_2) \ge 2F\left(\frac{P_1Q_1 + P_2Q_2}{2}\right) + 2\lambda|P_1Q_1 - P_2Q_2|^2.$$

Since  $F(P_1Q_1) = F(P_2Q_2) = \min_{SL(2)} F$  and  $(P_1Q_1 + P_2Q_2)/2 \in SL(2)$ , it follows that  $P_1Q_1 = P_2Q_2$ , a contradiction. Hence the minimizing set  $Z_{\min}(F) := \{P \in SL(2) : F(P) = \min_{SL(2)} F(Q)\}$  is just one copy of the special orthogonal group SO(2). This proves the lemma.

Suppose F is uniformly convex and  $u \in \mathcal{A}$  is a minimizer of I. Then by (2.7) and lemma 2.1, it follows that

$$\nabla u(x) \in SO(2)P$$
 a.e. in  $\Omega$ , (2.9)

for some det P = 1. From the Liouville Theorem of Reshetnyak [Re 68], it follows that u is affine. However for the convenience of the readers we give a proof, which is due to Kinderlehrer [Ki 88] (the same proof works in all dimensions). Let us make the change of variables,  $v : P^{-1}(\Omega) \to \mathbb{R}^2$ , by  $v(P^{-1}x) = u(x), x \in \Omega$ . Then  $\nabla v(P^{-1}x) = \nabla u(x) P$ . Hence

$$\nabla v(y) \in SO(2)$$
 a.e. in  $P^{-1}(\Omega)$ .

Since  $\operatorname{cof} Q = Q$  on SO(2), and

div cof 
$$\nabla v = 0$$
,

(div is taken in each rows) it follows that v is harmonic, i.e.,  $\Delta v = (\Delta v^1, \Delta v^2) = (0, 0)$ and hence smooth. Since  $|\nabla v|^2 = 2$ , the identity

$$\frac{1}{2}\Delta|\nabla v|^2 = |\nabla^2 v|^2 + \nabla v : \nabla\Delta v$$

yields  $\nabla^2 v = 0$  in  $P^{-1}(\Omega)$  and hence u is affine in  $\Omega$ . This proves the theorem.  $\Box$ 

**Remark 1.** The proof shows that the energy minimizing volume-preserving  $W^{1,n}$  deformations on bounded open subsets of  $\mathbb{R}^n$ , for  $n \geq 2$  are Lipschitz.

**Remark 2.** However, for  $n \geq 3$ , we are unable to conclude whether minimizers corresponding to frame indifferent uniformly convex functions on  $\mathbb{R}^{n \times n}$  are necessarily affine. Let us briefly discuss the case n = 3. For any given  $3 \times 3$  matrices  $Q_1$  and  $Q_2$ with determinant 1, the cosets  $SO(3) Q_1$  and  $SO(3) Q_2$  are not necessarily rank-one connected (this is the main difference with the two dimension), so the above proof fails to conclude  $SO(3) Q_1 = SO(3) Q_2$ . If the cosets are not rank-one connected (for example, the SO(3) and  $SO(3) Q_2$ ,  $Q_2 = \text{diag}(\lambda_1, \lambda_2, \lambda_3)$ ,  $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3$ ,  $\lambda_2 \neq 1$ ,  $\lambda_1 \lambda_2 \lambda_3 = 1$  are not rank-one connected) it is natural to determine whether any Lipschitz map  $u : \Omega \subset \mathbb{R}^3 \to \mathbb{R}^3$  satisfying

$$\nabla u(x) \in Z := SO(3) Q_1 \cup SO(3) Q_2 \quad \text{a.e.} \quad \text{in } \Omega \subset \mathbb{R}^3 \tag{2.10}$$

are necessarily affine. From the separation lemma [CM 04, Lemma 2.4] it is enough to show that such solutions u are  $W_{loc}^{2,2}$ . In order to obtain such regularity one usually tries to find a suitable system of partial differential equations for u satisfying (2.10). However, it follows that there are no uniformly elliptic system of PDEs (as the set Zis not strongly incompatible, see, [CM 04]) for u in (2.10). This suggests that there are no obvious way of getting  $W^{2,2}$  regularity.

### 3 Non-affine Minimizers

In this section we show that there are convex functions F for which the functional I in (1.1) admits area-preserving non-affine minimizers even with prescribed affine boundary data. To obtain such minimizers, idea is to look for a smooth convex (not uniform) function F such that the minimizing set  $Z_{\min}(F)$  strictly contains two copies of SO(2), which are rank-one connected. Then trivially one obtains non-affine minimizers, for example, simple laminates. But interestingly, by a theorem of Müller and Šverák [MS 96], on convex integration, we can find non-affine minimizers with prescribed affine boundary.

Let  $H := \text{diag}(\lambda, \mu)$ , be a diagonal matrix such that  $0 < \lambda < 1 < \mu$  and  $\lambda \mu = 1$ . Set

$$K := SO(2) \cup SO(2) H,$$

the two wells. Then a simple calculation shows that each matrix in K is rank-one connected with exactly two other matrices in K. Define  $F : \mathbb{R}^{2 \times 2} \to \mathbb{R}$  by

$$F := \sup\{g : g \text{ convex on } \mathbb{R}^{2 \times 2}, \ g \le \operatorname{dist}^2(\cdot, K)\},\$$

the convex envelope of the square of the distance function  $dist(\cdot, K)$ . Therefore F is smooth, convex and the second derivative of F uniformly bounded. Notice that F(X) = 0 if and only if

$$X \in K^{c} = \left\{ \left( \begin{array}{cc} x_{1} & -x_{2} \\ x_{2} & x_{1} \end{array} \right) + \left( \begin{array}{cc} y_{1} & -y_{2} \\ y_{2} & y_{1} \end{array} \right) \left( \begin{array}{cc} \lambda & 0 \\ 0 & \mu \end{array} \right) : x, y \in \mathbb{R}^{2}, \ |x| + |y| \le 1 \right\},$$

the convex hull of the set K. Therefore the minimizing set  $Z_{\min}(F) = K^c \cap SL(2)$ given by

$$Z_{\min}(F) = \left\{ \begin{pmatrix} x_1 + \lambda y_1 & -x_2 - \mu y_2 \\ x_2 + \lambda y_2 & x_1 + \mu y_1 \end{pmatrix} : |x| + |y| \le 1, \ |x|^2 + |y|^2 + (\lambda + \mu)\langle x, y \rangle = 1 \right\},$$

is the so-called rank-one convex hull (see [Sv 93] for more details about rank-one convex or quasiconvex hulls of general two wells energy) of K. It is clear that the set K is strictly contained in  $Z_{\min}(F)$ . For  $R \in Z_{\min}(F) \setminus K$ . Müller and Šverák [MS 96] on convex integration, the following boundary value partial differential inclusion

$$\begin{cases} \nabla u(x) \in SO(2) \cup SO(2) H \text{ a.e. in } \Omega \\ u(x) = Rx + b \text{ on } \partial\Omega \end{cases}$$
(3.11)

admit solutions. Therefore solutions to the problem (3.11) are clearly non-affine and minimizes of the energy functional I in 1.1) over the class of functions

 $\mathcal{A}_R := \{ u \in W^{1,2}(\Omega, \mathbb{R}^2) : \det \nabla u(x) = 1, \text{a. e. in } \Omega, \ u(x) = Rx + b \text{ on } \partial \Omega \}.$ 

This shows that the uniform convexity assumption in the Theorem 1.1 is sharp.

## References

- [BOP 92] P. Bauman, N. C. Owen and D. Phillips, Maximum principles and an a priori estimates for an incompressible material in nonlinear elasticity, *Comm. PDE* 17 (1992), 1185-1212.
- [CM 04] N. Chaudhuri and S. Müller, Rigidity estimate for two incompatible wells, Calc. Var. 19 (2004), 379–390.
- [DM 97] B. Dacorogna and P. Marcellini, General existence theorems for Hamilton-Jacobi equations in the scalar and vectorial cases, *Acta Math.* **178** (1997), 1-37.
- [Ev 86] L. C. Evans, Quasiconvexity and partial regularity in the calculus of variations, Arch. Rational. Mech. Anal., **95** (1986), 227-252.
- [EG 99] L. C. Evans and R. F. Gariepy, On the partial regularity of energyminimizing, area-preserving maps, *Calc. Var. Partial Differential Equa*tions, 9 (1999), 357-372.
- [Ki 88] D. Kinderlehrer, Remarks about equilibrium configurations of crystals, in: Material instabilities in continuum mechanics and related mathematical problems (J.M. Ball, ed.) Oxford Univ. Press, 1988, 217-242.
- [Mo 52] C. B. Morrey, Quasiconvexity and the semicontinuity of multiple integrals. *Pacific. J. Math.*, **2**, (1952), 25-52.
- [Mu 89] S. Müller, A surprising higher integrability property of mappings with positive determinant, *Bull. Amer. Math. Soc.* (N.S.) **21** (1989), 245–248.
- [MS 96] S. Müller and V. Šverák, Attainment results for the two-well problem by convex integration, in *Geometric Analysis and the Calculus of Variations* (J. Jost, ed.), Internat. Press, Cambridge, MA, 1996, 239-251.
- [Og 84] R. W. Ogden, *Non-linear elastic deformations*, Ellis Horwood Ltd. Chichester, 1984.
- [Re 68] Yu. G. Reshetnyak, Liouville's theorem under minimal regularity assumptions, Sib. Math. J. 9 (1968), 1039-1045.
- [Sv 93] V. Šverák, On the problem of two wells. Microstructure and phase transition, 183–189, IMA Vol. Math. Appl., 54, Springer, New York, 1993.