# Empirical Bayes estimators for the reproduction parameter of Borel-Tanner distribution

George P.  $Yanev^1$ 

University of South Florida, Tampa, Florida

**ABSTRACT** We construct empirical Bayes estimators for the reproduction parameter of Borel-Tanner distribution assuming LINEX loss and prove their asymptotic optimality. Some properties of the estimators regret risk are illustrated through simulations. **AMS Subject Classification** 62C10, 62F15, 60J80.

#### 1. INTRODUCTION

The probability mass function of the Borel-Tanner distribution is

$$p(x|\theta, r) = a_r(x)\theta^{x-r}e^{-\theta x}, \qquad (x = r, r+1, \ldots)$$

$$(1)$$

where  $0 < \theta < 1$ , r is a positive integer and  $a_r(x) = rx^{x-r-1}/(x-r)!$ 

Initially (1) was derived as the probability distribution of the number of customers served in a queuing system. It also appears in random trees and branching processes. More specifically, it is the distribution of the total progeny in a Galton-Watson process assuming Poisson reproduction, see Aldous [1] for recent applications. Our interest in estimating  $\theta$  stems from its role as reproduction number of an epidemic infection modeled by a branching process, see Farrington et al. [2]. We study nonparametric (with respect to the prior) empirical Bayes (NPEB) estimators for  $\theta$ . The NPEB estimation procedures rely on the assumption for existence of a prior distribution G which, however, is unknown. Consider independent copies  $(X_1, \theta_1), \ldots, (X_{n+1}, \theta_{n+1})$  of  $(X, \theta)$ , where  $\theta$  has a distribution G, and conditional on  $\theta$ , X has a Borel–Tanner distribution given by (1). The "past" data consist of independent observations  $x_1, x_2, \ldots, x_n$  obtained with independent realizations  $\theta_1, \theta_2, \ldots, \theta_n$  of  $\theta$ , where the  $X_i$ s are observable and the  $\theta_i$ s are not observable. Denote by  $\theta_n(x)$  an empirical Bayes estimator for  $\theta$  based on the "past" data and the "present" observation  $x_{n+1} = x$ . As Maritz and Lwin [5] point out, an advantage of using NPEB estimators is the minimum assumptions on the class of prior distributions. It turns out that in the case of Borel-Tanner distribution the Bayes rule assuming LINEX loss depends on the prior through the marginals only. This remarkable

<sup>&</sup>lt;sup>1</sup>Address: Mathematics Dept., 4202 E. Fowler Ave, PHY114, University of South Florida, Tampa, FL 33620, USA. E-mail: gyanev@cas.usf.edu.

fact allows us to construct simple NPEB estimators estimating the Bayes rule directly without estimating the prior itself.

Notice that NPEB estimators for  $\theta$  under weighted squared-error loss are studied in Yanev [7]. In the next section we use the asymmetric LINEX loss function, instead. In Section 3 we prove the estimators asymptotic optimality. The last section contains numerical results concerning the estimators performance measured by their regret risk.

### 2. EMPIRICAL BAYES ESTIMATION USING LINEX LOSS

In some applications (e.g. surveillance of infectious diseases) the squared-error loss function seems inappropriate in that it assigns the same loss to overestimates as to equal underestimates. A well-known alternative (see Huang et al. [3] and the references therein) is the LINEX loss function defined, for  $\gamma_l \leq \gamma \leq \gamma_u$  and  $\gamma \neq 0$  by

$$L_{\gamma}(\hat{\theta},\theta) = e^{\gamma(\theta-\theta)} - \gamma(\hat{\theta}-\theta) - 1 , \qquad (2)$$

where  $\hat{\theta}$  is an estimator for  $\theta$ . It is clear that the LINEX loss function is convex, asymmetric and for  $\gamma > 0$  it increases almost linearly for negative errors and almost exponentially for positive errors. Thus, it penalizes an overestimation more seriously than an underestimation. This is reversed when  $\gamma < 0$ . For small values of  $|\gamma|$  the LINEX loss is close to the squared– error loss. From now on we assume that  $\gamma$  is a positive integer; the case  $\gamma < 0$  can be treated similarly.

Based on a single observation, the maximum likelihood estimator  $\theta_{MLE}(x)$  for  $\theta$  is (e.g., Kumar & Consul [4])

$$\theta_{MLE}(x) = \frac{x-r}{x}.$$
(3)

Denote by  $\mathcal{I}_A$  the indicator of the event A.

**Theorem 1** Assume LINEX loss with  $\gamma > 0$ , integer. A NPEB estimator for  $\theta$  in (1) is

$$\theta_n(x) = \gamma^{-1} \ln \tau_n(x) \mathcal{I}_{\{\tau_n(x) \in (1, e^{\gamma})\}} + (x - r) / x \mathcal{I}_{\{\tau_n(x) \notin (1, e^{\gamma})\}}, \tag{4}$$

where

$$\tau_n(x) = \frac{r+\gamma}{r} \left(\frac{x+\gamma}{x}\right)^{x-r-1} \frac{m_n(x|r)}{m_n(x+\gamma|r+\gamma)}$$

and  $m_n(z|y)$  is an estimate for the marginal distribution  $m_G(z|y) = \int_0^1 p(z|\theta, y) dG(\theta)$ . **Proof** The Bayesian estimator  $\theta_G(x)$  under LINEX loss is (e.g. Huang et al. [3])

$$\theta_G(x) = -\gamma^{-1} \ln E_{G|x} e^{-\gamma\theta},\tag{5}$$

provided that  $E_{G|x}e^{-\gamma\theta} < \infty$ , where  $E_{G|x}(\cdot)$  is the expectation w.r.t. the posterior. Since

$$E_{G|x}e^{-\gamma\theta} = \frac{1}{m_G(x|r)} \int_0^1 e^{-\gamma\theta} a_r(x)\theta^{x-r}e^{-x\theta} dG(\theta)$$
  
=  $\frac{a_r(x)}{a_{r+\gamma}(x+\gamma)} \frac{m_G(x+\gamma|r+\gamma)}{m_G(x|r)},$ 

we can write the Bayesian estimator  $\theta_G(x)$  from (5) as

$$\theta_G(x) = \gamma^{-1} \ln \left\{ \frac{r+\gamma}{r} \left( \frac{x+\gamma}{x} \right)^{x-r-1} \frac{m_G(x|r)}{m_G(x+\gamma|r+\gamma)} \right\}$$
$$= \gamma^{-1} \ln \tau_G(x), \quad \text{say.}$$

Note that,  $\theta_G(x)$  depends upon the prior through the marginal distribution only. Therefore, estimating the marginals, we can construct a NPEB estimator  $\theta_n(x)$  for  $\theta$  as given in (4).  $\Box$ 

One possible form of the estimators  $m_n(z|y)$  in Theorem 1 can be obtained as follows. In addition to the current  $X_{n+1}(r) = x$ , let us have observed n independent pairs

$$(X_1(r), X_1(\gamma)), (X_2(r), X_2(\gamma)), \dots, (X_n(r), X_n(\gamma)),$$
 (6)

where  $X_i(r)$  and  $X_i(\gamma)$  are independent and Borel-Tanner distributed with  $p(x|\theta_i, r)$  and  $p(x|\theta_i, \gamma)$ , respectively. It is known (e.g., Kumar & Consul [4]) that  $X_i(r) + X_i(\gamma)$  has pmf  $p(x|\theta_i, r+\gamma)$ . Let  $f_n(y|r+\gamma)$  be the number of pairs, such that  $X_i(r)+X_i(\gamma)=y$ ,  $(i=1,\ldots,n)$ . Consistent estimators for the marginals  $m_G(x+\gamma|r+\gamma)$  and  $m_G(x|r)$  are the relative frequencies

$$m_n(x+\gamma|r+\gamma) = \frac{f_n(x+\gamma|r+\gamma)}{n+1}$$
 and  $m_n(x|r) = \frac{1+f_n(x|r)}{n+1}$ . (7)

Let us notice here that a NPEB estimator  $\hat{\theta}_n(x)$  for  $\theta$  under the squared-error loss  $L(\hat{\theta}, \theta) = (\hat{\theta} - \theta)^2$  is constructed in Yanev [7] as follows

$$\tilde{\theta}_n(x) = \kappa_n(x)\mathcal{I}_{\{\kappa_n(x) \in (0,1)\}} + (x-r)/x\mathcal{I}_{\{\kappa_n(x) \notin (0,1)\}},$$

where

$$\kappa_n(x) = \frac{a_r(x)}{m_n(x)} \sum_{j=0}^{\infty} \frac{(j+1)^{j-1}}{j!} \frac{m_n(x+j+1)}{a_r(x+j+1)},$$

where, as before,  $a_r(y) = ry^{y-r-1}/(y-r)!$ 

#### 3. ASYMPTOTIC OPTIMALITY

The Bayes risk of an estimator  $\hat{\theta}$  can be written as

$$R(G,\hat{\theta}) = \int_X \int_{\Theta} L(\hat{\theta},\theta) p(x|\theta,r) dG(\theta) dx = \sum_{x=r}^{\infty} \int_{\Theta} L(\hat{\theta},\theta) p(\theta|x) dG(\theta) m_G(x|r),$$

where  $p(\theta|x)$  is the posterior distribution. If  $R(G, \theta_n | \underline{X}_n)$  is the conditional Bayes risk of the estimator  $\theta_n(x)$  given  $\underline{X}_n = (X_1, \ldots, X_n)$ , then  $R(G, \theta_n) = E_n\{R(G, \theta_n | \underline{X}_n)\}$  is the (unconditional) Bayes risk of  $\theta_n$ , where the expectation  $E_n(\cdot)$  is taken with respect to  $\underline{X}_n$ . The estimator  $\theta_n(x)$  is asymptotically optimal for given G if  $\lim_{n\to\infty} R(G, \theta_n) = R(G, \theta_G)$ . We shall prove the asymptotic optimality of  $\theta_n(x)$ .

First, let us find the minimum Bayes risk  $R(G, \theta_G)$  attained by the Bayesian estimator  $\theta_G(x)$ . Since (5) implies  $\exp(\gamma \theta_G(x)) \int_0^1 \exp(-\gamma \theta) p(\theta|x) d\theta = 1$ , we have

$$\begin{aligned} R(G,\theta_G) &= \sum_{x=r}^{\infty} \left\{ \int_0^1 \left\{ e^{\gamma(\theta_G(x)-\theta)} - \gamma(\theta_G(x)-\theta) - 1 \right\} p(\theta|x) d\theta \right\} m_G(x|r) \\ &= \sum_{x=r}^{\infty} \left\{ e^{\gamma\theta_G(x)} \int_0^1 e^{-\gamma\theta} p(\theta|x) d\theta - \gamma\theta_G(x) + \int_0^1 \gamma\theta p(\theta|x) d\theta - 1 \right\} m_G(x|r) \\ &= \sum_{x=r}^{\infty} \left\{ \int_0^1 \gamma\theta p(\theta|x) d\theta - \gamma\theta_G(x) \right\} m_G(x|r). \end{aligned}$$

Next, using  $\exp(\gamma \theta_G(x)) \int_0^1 \exp(-\gamma \theta) p(\theta|x) d\theta = 1$  again, we obtain

$$R(G,\theta_n) = \sum_{x=r}^{\infty} E_n \left\{ e^{\gamma \theta_n(x)} \int_0^1 e^{-\gamma \theta} p(\theta|x) d\theta - \gamma \theta_n(x) + \int_0^1 \gamma \theta p(\theta|x) d\theta - 1 \right\} m_G(x|r)$$
  
$$= \sum_{x=r}^{\infty} E_n \left\{ e^{\gamma (\theta_n(x) - \theta_G(x))} - \gamma \theta_n(x) + \int_0^1 \gamma \theta p(\theta|x) d\theta - 1 \right\} m_G(x|r).$$

Therefore,

$$R(G,\theta_n) - R(G,\theta_G) = \sum_{x=r}^{\infty} E_n \left\{ e^{\gamma(\theta_n(x) - \theta_G(x))} - \gamma(\theta_n(x) - \theta_G(x)) - 1 \right\} m_G(x|r)$$
(8)

Let us truncate the Borel–Tanner distribution (1) starting with r = k as follows

$$p^*(x|\theta,k) = \begin{cases} p(x|\theta,k), & \text{if } k \le x \le k+N-1;\\ \sum_{x=k+N}^{\infty} p(x|\theta,k), & \text{if } x=k+N. \end{cases}$$
(9)

where N is a positive integer. Denote the truncated marginal by  $m_G^*(x) = \int_0^1 p^*(x|\theta, y) dG(\theta)$ . Similar to the non-truncated case, if  $r \leq x \leq r + N - 1$  then

$$E_{G|x}(e^{-\gamma\theta}) = \frac{a_r(x)}{a_{r+\gamma}(x+\gamma)} \frac{m_G^*(x+\gamma|r+\gamma)}{m_G^*(x|r)} = \frac{a_r(x)}{a_{r+\gamma}(x+\gamma)} \frac{m_G(x+\gamma|r+\gamma)}{m_G(x|r)} = \frac{1}{\tau_G(x)}.$$

If x = r + N then

$$E_{G|x}(e^{-\gamma\theta}) = \frac{1}{m_G^*(r+N|r)} \int_0^1 e^{-\gamma\theta} \sum_{k=r+N}^\infty a_r(k)\theta^{k-r}e^{-\theta k} dG(\theta)$$
  
$$= \frac{1}{m_G^*(r+N|r)} \sum_{k=r+N}^\infty \frac{a_r(k)}{a_{r+\gamma}(k+\gamma)} m_G^*(k+\gamma|r+\gamma)$$
  
$$= \left(\sum_{k=r+N}^\infty m_G(k|r)/\tau_G(k)\right) / \sum_{k=r+N}^\infty m_G(k|r).$$

Let  $\tau_G^*(x) = \tau_G(x)$  if  $r \leq x \leq r + N - 1$ ;  $= \sum_{k=r+N}^{\infty} m_G(k|r) / \sum_{k=r+N}^{\infty} (m_G(k|r) / \tau_G(k))$  if x = r + N. The Bayesian estimator in the truncated case is given by  $\theta_G^*(x) = \gamma^{-1} \ln \tau_G^*(x)$ . Let us estimate  $m_G(x|y)$  by  $m_n(x|y)$  as in Theorem 1 and set  $\tau_n^*(x) = \tau_n(x)$  if  $r \leq x \leq r + N - 1$ ;

 $=\sum_{k=r+N}^{\infty} m_n(k|r) / \sum_{k=r+N}^{\infty} (m_n(k|r) / \tau_n(k))$  if x = r + N. We construct a NPEB estimator in the truncated case as follows

$$\theta_n^*(x) = \gamma^{-1} \ln \tau_n^*(x) \mathcal{I}_{\{\tau_n^*(x) \in (1, e^{\gamma})\}} + (x - r) / x \mathcal{I}_{\{\tau_n^*(x) \notin (1, e^{\gamma})\}}.$$

Now, we are in a position to prove the asymptotic optimality of  $\theta_n(x)$ .

**Theorem 2** Assume prior G with finite first moment. If  $m_n(z|y)$  is a consistent estimator for  $m_G(z|y)$ , then the NPEB estimator  $\theta_n(x)$  given by (4) is asymptotically optimal, i.e.,

$$\lim_{n \to \infty} R(G, \theta_n) = R(G, \theta_G).$$

**Proof** Since  $\theta_G(x)$  is the Bayesian estimator, we have  $R(G, \theta_n) > R(G, \theta_G)$  and thus

$$R(G,\theta_n) - R(G,\theta_G) \le |R(G,\theta_n) - R(G,\theta_n^*)| + |R(G,\theta_n^*) - R(G,\theta_G^*)| + |R(G,\theta_G^*) - R(G,\theta_G)|$$
(10)

To prove the theorem it is sufficient to show that the right hand side of (10) has  $\lim_{N\to\infty} \limsup_{n \to \infty} \lim_{N\to\infty} \sup_{n \to \infty} \lim_{N\to\infty} \sup_{n \to \infty} \lim_{N\to\infty} \sup_{n \to \infty} \lim_{N\to\infty} \lim_{N\to\infty} \lim_{N\to\infty} \sup_{n \to \infty} \lim_{N\to\infty} \lim_{N\to\infty}$ 

$$|R(G,\theta_n^*) - R(G,\theta_G^*)| = \sum_{x=r}^{r+N} E_n \left\{ e^{\gamma(\theta_n^*(x) - \theta_G^*(x))} - \gamma(\theta_n^*(x) - \theta_G^*(x)) - 1 \right\} m_G(x|r)$$

Since  $m_n(z|y)$  is a consistent estimator for  $m_G(z|y)$ , we have  $\lim_{n\to\infty} \theta_n^*(x) = \theta_G^*(x)$ ,  $F^{\infty}$ - a.s., where  $F^{\infty}$  is the product measure induced by  $X_1, X_2, \ldots, X_n, \ldots$ . Notice that, both  $\theta_n^*$  and  $\theta_G^*$  are bounded. Indeed,  $\theta_n^*$  is bounded by definition and  $0 < \theta_G^*(x) = -(1/\gamma) \ln E_{G|x}(e^{-\gamma\theta}) < (1/\gamma) \ln e^{\gamma} = 1$ . Therefore, by the Lebesgue dominated convergence theorem we can pass to the limit inside the expectation in the right hand side above and obtain

$$\lim_{n \to \infty} |R(G, \theta_n^*) - R(G, \theta_G^*)| = 0.$$
(11)

Also, since  $p^*(\theta|x, r) = p(\theta|x, r)$ ,  $m^*_G(x) = m_G(x)$  for  $r \le x \le r + N - 1$ , and  $m^*_G(r + N) = \sum_{x=r+N}^{\infty} m_G(x)$  it is not difficult to obtain

$$|R(G,\theta_G^*) - R(G,\theta_G)| = \sum_{x=r+N}^{\infty} \left\{ \int_0^1 \gamma \theta \left( p^*(\theta|r+N) - p(\theta|x) \right) d\theta - \gamma \left( \theta_G^*(r+N) - \theta_G(x) \right) \right\} m_G(x).$$

Since  $|p^*(\theta|r+N) - p(\theta|x)| < 1$ ,  $|\theta^*_G(r+N) - \theta_G(x)| < 1$ , and  $E\theta < \infty$  we have

$$\lim_{N \to \infty} |R(G, \theta_G^*) - R(G, \theta_G)| = 0.$$
(12)

Similar to (12) one can prove that  $\lim_{N\to\infty} |R(G,\theta_n) - R(G,\theta_n^*)| = 0$ . This along with (10)-(12) completes the proof.

#### 4. NUMERICAL EXAMPLES

Using the notation introduced before (7) we set

$$\tau_n^f(x) = \frac{r+\gamma}{r} \left(\frac{x+\gamma}{x}\right)^{x-r-1} \frac{f_n(x|r)}{f_n(x+\gamma|r+\gamma)}$$

Let  $A = \left\{ \tau_n^f(x) \in (1, e^{\gamma}) \cap f_n(x + \gamma | r + \gamma) \neq 0 \right\}$  and  $A^c$  be its complement. Making use of the relative frequency estimators (7) consider  $\theta_n^f(x)$  to be defined by

$$\theta_n^f(x) = \gamma^{-1} \ln \tau_n^f(x) \mathcal{I}_A + (x - r) / x \mathcal{I}_A^c.$$

That is, if A occurs, then we estimate  $\theta$  by  $\gamma^{-1} \ln \tau_n^f(x)$ ; whereas if  $A^c$  occurs then we use the MLE (3) for  $\theta$  instead.

A popular measure of the performance of one estimator  $\hat{\theta}(x)$  is its regret risk  $S(\hat{\theta}) = R(G, \hat{\theta}) - R(G, \theta_G) > 0$ . For our simulation study we take r = 5, Uniform (0.5, 1) prior and LINEX loss with  $\gamma = 3$ . Then the minimum Bayes risk attained by the Bayesian estimator

$$\theta_U(x) = \frac{1}{3} \ln \frac{\int_{0.5}^1 \theta^{x-5} e^{-x\theta} d\theta}{\int_{0.5}^1 \theta^{x-5} e^{-(x+3)\theta} d\theta}$$

is  $R(U_{(0.5,1)}, \theta_U) = 0.0622.$ 

In the empirical Bayes scheme (6), let us set n = 50. Selecting 50 random values for  $\theta_i \sim U_{(0.5,1)}$ , i = 1, 2, ..., 50, we generate two sets of 50 branching processes starting with r = 5 and  $\gamma = 3$  ancestors, respectively and both having  $Poisson(\theta_i), i = 1, 2, ..., 50$  offspring distributions. Notice that the total progeny of each process is a realization of a Borel-Tanner  $(\theta_i, \cdot)$  random variable. Repeating the above procedure 100 times, we obtain 100 samples of 50 pairs Borel-Tanner observations,  $(X_i(5), X_i(3)), i = 1, 2, ..., 50$ . Each sample gives us a NPEB estimate  $\theta_{50}^f(x)$  with regret risk  $S_i(\theta_{50}^f), i = 1, 2, ..., 100$ . We estimate the regret risk  $S(\theta_{50}^f)$  with the average  $\bar{S}(\theta_{50}^f) = \sum_{i=1}^{100} S_i(\theta_{50}^f)/100$ .

The above scheme is repeated with n = 75 and n = 100. As an illustration, we present in Table 1 results for one sample with n = 100. For this particular sample,  $S_i(\theta_{100}^f) = 0.0980$ , which is less than  $S(\theta_{MLE}) = 0.1327$ .

x		5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$ heta_{100}^f($	(x)	.46	.69	.92	.65	.58	.51	.55	.53	.62	.96	.61	.69	.16	.72	.79	.75
$ heta_U(z)$	x)	.63	.64	.65	.65	.66	.67	.67	.68	.69	.69	.70	.71	.71	.72	.73	.73
$\theta_{MLE}$	(x)	0	.16	.28	.38	.44	.50	.55	.58	.62	.64	.67	.69	.71	.72	.74	.75

Table 1: Estimates  $\theta_n^f(x)$ ,  $\theta_U(x)$ , and  $\theta_{MLE}(x)$  for  $\theta$  from a sample with n = 100.

		$0 \le x \le 15$		$0 \le x \le 200$				
n	$\bar{S}(\theta_n^f)$	$STD(\bar{S}(\theta_n^f))$	$S(\theta_{MLE})$	$\bar{S}(\theta_n^f)$	$STD(\bar{S}(\theta_n^f))$	$S(\theta_{MLE})$		
50	0.1211	0.0037	0.1292	0.1397	0.0037	0.1327		
75	0.1076	0.0036	0.1292	0.1300	0.0037	0.1327		
100	0.1038	0.0033	0.1292	0.1299	0.0036	0.1327		

Table 2: Numerical results on regret risks of  $\theta_n^f(x)$  and  $\theta_{MLE}(x)$ .

The numerical results for the regret risks are given in Table 2. Several comments are in place. For small x, (columns 2-4) and n = 75 or 100, the improvement of  $\theta_n^f$  over  $\theta_{MLE}$  is substantial. Overall, (columns 5-7), the regret risk of  $\theta_n^f$  is not higher than that of  $\theta_{MLE}$ .

Finally, note that the Borel-Tanner distribution (1) has monotone likelihood ratio in x, i.e.,  $p(x|\theta', r)/p(x|\theta, r)$  is an increasing function of x whenever  $0 < \theta < \theta' < 1$ . This suggests that the NPEB  $\theta_n(x)$  can be improved on by the monotonizing procedure of Van Houwelingen and Stijnen [6].

### ACKNOWLEDGEMENTS

I thank R. Gueorguieva for helping me with the simulations done with an Ox version 3.30. I also thank the referee for the valuable comments. This research is partially supported by NFSI-Bulgaria, Grant No. MM-1101/2001.

## References

- [1] Aldous, D.J. Deterministic and stochastic models for coalescence (aggregation and coagulation): a review of the mean-field theory for probabilists. Bernoulli **1999**, *5*, 3-48.
- [2] Farrington, C.P.; Kanaan, C.P.; Gay, N.J. Branching process models for surveillance of infectious diseases controlled by mass vaccination. Biostatistics 2003, 4(2), 279-295.
- [3] Huang, S.Y.; Liang, T.C. Empirical Bayes estimation of the truncation parameter with Linex loss. Statist. Sinica 1997, 7, 755–769.
- [4] A. Kumar, A.; Consul, P.C. Minimum variance unbiased estimation for modified power series distribution. Comm. Statist. A - Theory Methods 1980, 9, 1261-1275.
- [5] Maritz, J.S.; Lwin, T. Empirical Bayes Methods, 2nd Ed.; Chapman and Hall: London, 1989.
- [6] Van Houwelingen, J.C.; Stijnen, T. Monotone empirical Bayes estimators based on more informative samples. J. Amer. Statist. Assoc. 1993, 88, 1438-1443.
- [7] Yanev, G.P. Statistical modeling of epidemic disease propagation via branching processes and Bayesian inference, Dissertation, 2001, University of South Florida.