

Groupoid Approach to Quantum Projective Spaces

Albert Jeu-Liang Sheu*
 Department of Mathematics
 University of Kansas
 Lawrence, KS 66045
 U. S. A.

1 Introduction

The theory of groupoid C*-algebras [Re] has proved to be a very useful tool and provide a nearly universal context for the study of operator algebras, especially in connection with geometric structures. There have been many successful examples of such uses of groupoid C*-algebras [Co, CuM, MRe, SaShU, Sh1].

Recently interesting and important examples of C*-algebras of quantum groups G_q and quantum spaces M_q arise from the quantization [Dr, RTF, So, Wo1, Wo2, Po, Ri3, Ri1, Ri2, VaSo, Sh2, Na2] of Poisson Lie groups G and their Poisson homogeneous spaces $M = H \backslash G$ with H a Poisson Lie subgroup of G . Known results [So, Sh2] show that they are closely related to the underlying singular symplectic foliation [We, LuWe1]. It is shown in [Sh4] that the C*-algebra of compact quantum groups can be effectively described and studied in the context of groupoid C*-algebras with the groupoid summarizing the underlying singular symplectic foliation. In fact, the structures of the algebras of quantum groups $SU(n)_q$ and quantum spheres \mathbb{S}_q^{2n-1} are analyzed in detail in [Sh4, Sh5].

More recently, nonstandard quantum complex projective spaces $\mathbb{C}P_{q,c}^n$ have been studied by Dijkhuizen and Noumi [DiNo]. In this paper, we show how the C*-algebras of quantum complex projective spaces (standard or nonstandard) are related to groupoids.

2 Quantum groups $SU(n)_q$

In this section, we briefly recall the notations and results needed from [Sh4]. For more complete information, we refer to the paper [Sh4].

*Partially supported by NSF Grant DMS-9623008.

First we recall that the transformation group groupoid $\mathbb{Z}^m \times \overline{\mathbb{Z}}^m$ (with \mathbb{Z}^m acting on $\overline{\mathbb{Z}}^m$ by translation) when restricted to the positive cone $\overline{\mathbb{Z}}_{\geq}^m$ gives an important (Toeplitz) groupoid

$$\mathcal{T}_m := \mathbb{Z}^m \times \overline{\mathbb{Z}}_{\geq}^m |_{\overline{\mathbb{Z}}_{\geq}^m} = \{(j, k) \in \mathbb{Z}^m \times \overline{\mathbb{Z}}_{\geq}^m \mid j + k \in \overline{\mathbb{Z}}_{\geq}^m\}$$

where $\overline{\mathbb{Z}} = \mathbb{Z} \cup \{+\infty\}$ and $\overline{\mathbb{Z}}_{\geq} := \{0, 1, 2, 3, \dots\} \cup \{+\infty\}$.

Recall that the C*-algebra $C(SU(n+1)_q)$ is generated by u_{ij} , $1 \leq i, j \leq n+1$, satisfying $u^*u = uu^* = I$ and some other relations [Wo3, So]. In particular, the C*-algebra $C(SU(2)_q)$ is generated by u_{ij} , with $1 \leq i, j \leq 2$, satisfying $u_{22} = u_{11}^*$, $u_{12} = -q^{-1}u_{21}^*$, and $u^*u = uu^* = 1$ [Wo1, VaSo]. An important irreducible (non-faithful) *-representation π_0 of $C(SU(2)_q)$, $q > 1$, on $\ell^2(\mathbb{Z}_{\geq})$ is given by

$$\pi_0(u) = \begin{pmatrix} \alpha & -q^{-1}\gamma \\ \gamma & \alpha^* \end{pmatrix}$$

where $\alpha(e_j) = (1 - q^{-2j})^{1/2}e_{j-1}$ and $\gamma(e_j) = q^{-j}e_j$ for $j \geq 0$. Here π_0 is applied to $u = (u_{ij})$ entrywise.

Irreducible one-dimensional *-representations of $C(SU(n+1)_q)$ are defined by $\tau_t(u_{ij}) = \delta_{ij}t_j$ for $t \in \mathbb{T}^n$ (with $t_{n+1} = t_1^{-1}t_2^{-1}\dots t_n^{-1}$). The \mathbb{T}^n -family $\{\tau_t\}_{t \in \mathbb{T}^n}$ of one-dimensional irreducible *-representations of $C(SU(n+1)_q)$ can be viewed as a C*-algebra homomorphism $\tau_{n+1} = \tau : C(SU(n+1)_q) \rightarrow C(\mathbb{T}^n) \cong C^*(\mathbb{Z}^n)$. There are n fundamental *-representations $\pi_i = \pi_0\phi_i$ with $\phi_i : C(SU(n+1)_q) \rightarrow C(SU(2)_q)$ given by $\phi_i(u_{jk}) = u_{j-i+1, k-i+1}$ if $\{j, k\} \subseteq \{i, i+1\}$ and $\phi_i(u_{jk}) = \delta_{jk}$ if otherwise.

The unique maximal element in the Weyl group of $SU(n+1)$ can be expressed in the reduced form

$$s_1 s_2 s_1 s_3 s_2 s_1 \dots s_n s_{n-1} \dots s_2 s_1.$$

So by Soibelman's classification of irreducible *-representations [So], $C(SU(n+1)_q)$ can be embedded into

$$C^*(\mathcal{G}^n) \subseteq C^*(\mathbb{Z}^n) \otimes \mathcal{B}(\ell^2(\mathbb{Z}_{\geq}^N))$$

by

$$(\tau_{n+1} \otimes \pi_{121321\dots n(n-1)..21})\Delta^N$$

where $N = n(n+1)/2$,

$$\pi_{i_1 i_2 \dots i_m} := \pi_{i_1} \otimes \pi_{i_2} \otimes \dots \otimes \pi_{i_m},$$

Δ is the comultiplication on $C(SU(n+1)_q)$ with Δ^m defined recursively as $\Delta^k = (\Delta \otimes \text{id})\Delta^{k-1}$, and \mathcal{G}^n is the groupoid $\mathbb{Z}^n \times \mathbb{Z}^N \times \overline{\mathbb{Z}}_{\geq}^N |_{\overline{\mathbb{Z}}_{\geq}^N}$ with \mathbb{Z}^n acting trivially on $\overline{\mathbb{Z}}_{\geq}^N$, and \mathbb{Z}^N acts by translation on $\overline{\mathbb{Z}}_{\geq}^N$.

3 Quantum spheres \mathbb{S}_q^{2n+1}

Recall that the C*-algebra of the quantum sphere $S_q^{2n+1} = SU(n)_q \backslash SU(n+1)_q$ (as a homogeneous quantum space) is defined by

$$C(S_q^{2n+1}) = \{f \in C(SU(n+1)_q) : (\Phi \otimes id)(\Delta f) = 1 \otimes f\}$$

where $\Phi : C(SU(n+1)_q) \rightarrow C(SU(n)_q)$ is the quotient map defined by $\Phi(u_{ij}) = u_{ij}$, $\Phi(u_{n+1,k}) = \Phi(u_{k,n+1}) = 0$, and $\Phi(u_{n+1,n+1}) = 1$, for $1 \leq i, j, k \leq n$. From

$$u_{n+1,m} = (h_n \otimes 1)(1 \otimes u_{n+1,m}) = (h_n \otimes 1)(\Phi \otimes 1)(\Delta(u_{n+1,m})),$$

we have

$$u_{n+1,m} \in (h_n \otimes 1)(\Phi \otimes 1)\Delta(C(SU(n+1)_q)) = C(S_q^{2n+1})$$

by Nagy's result [Na1], where h_n is the Haar functional on $C(SU(n)_q)$ [Wo2, Sh4] such that $h_n(1) = 1$.

As discussed in [Sh4], we can prove that $C(S_q^{2n+1})$ is generated by $u_{n+1,m}$'s, by verifying that the monomials

$$p^{i,j,k} = (u_{n+1,n}^*)^{i_n} \dots (u_{n+1,1}^*)^{i_1} y_1^{j_1} \dots y_n^{j_n} y_{n+1}^{j_{n+1}} (u_{n+1,1})^{k_1} \dots (u_{n+1,n})^{k_n}$$

are linearly independent, where $y_m = u_{n+1,m} u_{n+1,m}^*$ and $i_m, j_m, k_m \geq 0$ with $i_m k_m = 0$ (this last condition was accidentally missing in [Sh4] for $1 \leq m \leq n$, $y_{n+1}^{j_{n+1}} = (u_{n+1,n+1})^{j_{n+1}}$ if $j_{n+1} \geq 0$, and $y_{n+1}^{j_{n+1}} = (u_{n+1,n+1}^*)^{-j_{n+1}}$ if $j_{n+1} \leq 0$).

So we have

$$C(S_q^{2n+1}) = C^*(\{u_{n+1,m} \mid 1 \leq m \leq n+1\}) \subset C(SU(n+1)_q)$$

(used as the definition of S_q^{2n+1} in [VaSo]) which can be embedded into the groupoid C*-algebra $C^*(\mathcal{F}^n)$, where \mathcal{F}^n is the augmented n -dimensional Toeplitz groupoid $\mathbb{Z} \times \mathcal{T}_n = \mathbb{Z} \times (\mathbb{Z}^n \times \overline{\mathbb{Z}}^n |_{\overline{\mathbb{Z}}^n})$, through the faithful representation

$$(\tau'_{n+1} \otimes \pi_{n(n-1)..21}) \Delta^n$$

where $\tau'_k(u_{ij}) = \delta_{ki} \delta_{ij} \text{id}_{\mathbb{T}} \in C(\mathbb{T}) \cong C^*(\mathbb{Z})$ for $1 \leq k \leq n+1$.

It turns out that $C(S_q^{2n+1})$ is actually isomorphic to the C*-algebra of a subquotient groupoid of \mathcal{F}^n [Sh5]. In fact, let

$$\widetilde{\mathfrak{F}}_n := \{(z, x, w) \in \mathcal{F}^n \mid w_i = \infty \implies$$

$$x_i = -z - x_1 - x_2 - \dots - x_{i-1} \text{ and } x_{i+1} = \dots = x_n = 0\}$$

be a subgroupoid of \mathcal{F}^n . Define $\mathfrak{F}_n := \widetilde{\mathfrak{F}}_n / \sim$ where \sim is the equivalence relation generated by

$$(z, x, w) \sim (z, x, w_1, \dots, w_i = \infty, \infty, \dots, \infty)$$

for all (z, x, w) with $w_i = \infty$ for an $1 \leq i \leq n$. Then we have $C(S_q^{2n+1}) \simeq C^*(\mathfrak{F}_n)$.

Since $SU(n)$ is a Poisson Lie subgroup of $SU(n+1)$, the multiplicative Poisson structure on $SU(n+1)$ induces a natural covariant Poisson structure on the homogeneous space $\mathbb{S}^{2n+1} = SU(n) \backslash SU(n+1)$, i.e. the $SU(n+1)$ -action

$$\mathbb{S}^{2n+1} \times SU(n+1) \rightarrow \mathbb{S}^{2n+1}$$

on \mathbb{S}^{2n+1} is a Poisson map [LuWe1]. Endowed with this covariant Poisson structure, the sphere \mathbb{S}^{2n+1} decomposes into symplectic leaves [We] which are the symplectic leaves in the canonically embedded Poisson \mathbb{S}^{2n-1} plus a circle family of leaves symplectically isomorphic to the complex space \mathbb{C}^n . Analyzing the above groupoid structure, we can easily see that this Poisson foliation structure is reflected on the quantum algebras, namely, there is a short exact sequence of C*-algebras

$$0 \rightarrow C(\mathbb{T}) \otimes \mathcal{K}(\ell^2(\mathbb{Z}_{\geq}^n)) \rightarrow C(S_q^{2n+1}) \rightarrow C(S_q^{2n-1}) \rightarrow 0.$$

In fact, \mathfrak{F}_{n-1} can be identified with

$$\left\{ [(z, x, w)] : (z, x, w) \in \widetilde{\mathfrak{F}}_n \text{ and } w_n = \infty \right\},$$

i.e. \mathfrak{F}_n restricted to the invariant closed subset X consisting of $[w] \in Z$ with $w_n = \infty$ in the unit space Z of \mathfrak{F}_n , through the groupoid monomorphism sending $(z, x', w') \in \widetilde{\mathfrak{F}}_{n-1}$ to

$$(z, x', -z - x'_1 - x'_2 - \dots - x'_{n-1}, w', \infty) \in \widetilde{\mathfrak{F}}_n,$$

and $\mathfrak{F}_n \backslash \mathfrak{F}_{n-1} = \mathfrak{F}_n|_{Z \setminus X}$ (the restriction of \mathfrak{F}_n to the invariant open set $Z \setminus X$) is isomorphic to the groupoid $\mathbb{Z}^1 \times (\mathbb{Z}^n \times \mathbb{Z}^n|_{\mathbb{Z}_{\geq}^n})$. It is well known [Re] that there is a short exact sequence of groupoid C*-algebras

$$0 \rightarrow C^*(\mathfrak{F}_n \backslash \mathfrak{F}_{n-1}) \rightarrow C^*(\mathfrak{F}_n) \rightarrow C^*(\mathfrak{F}_{n-1}) \rightarrow 0$$

which leads to the above short exact sequence for the quantum spheres since

$$C^*(\mathfrak{F}_n \backslash \mathfrak{F}_{n-1}) = C^*\left(\mathbb{Z}^1 \times (\mathbb{Z}^n \times \mathbb{Z}^n|_{\mathbb{Z}_{\geq}^n})\right) \cong C(\mathbb{T}) \otimes \mathcal{K}(\ell^2(\mathbb{Z}_{\geq}^n)).$$

4 Standard quantum projective spaces $\mathbb{C}P_q^n$

The C*-algebra of the standard quantum complex projective space

$$\mathbb{C}P_q^n := U(n)_q \backslash SU(n+1)_q,$$

corresponding to the Poisson Lie subgroup $U(n)$ of $SU(n+1)$, can be identified with

$$C(\mathbb{C}P_q^n) = C^*(\{u_{n+1,i}^* u_{n+1,j} \mid 1 \leq i, j \leq n+1\}) \subset C(\mathbb{S}_q^{2n+1}).$$

In fact, as a quantum homogeneous space, $\mathbb{C}P_q^n$ is defined by

$$C(\mathbb{C}P_q^n) = \{f \in C(SU(n+1)_q) : (\Phi' \otimes id)(\Delta f) = 1 \otimes f\}$$

where $\Phi' : C(SU(n+1)_q) \rightarrow C(U(n)_q)$ is the quotient map defined by $\Phi(u_{ij}) = u_{ij}$, $\Phi(u_{n+1,k}) = \Phi(u_{k,n+1}) = 0$, and

$$\Phi(u_{n+1,n+1}) = t_{n+1} = (t_1 \dots t_n)^{-1} \in C(U(n)_q) \subset C^*(\mathbb{Z}^n \times \mathcal{T}_{n(n-1)/2})$$

for $1 \leq i, j, k \leq n$. By Nagy's result [Na1], we have

$$(h'_n \otimes 1)(\Phi' \otimes 1)\Delta(C(SU(n+1)_q)) = C(\mathbb{C}P_q^n)$$

where h'_n is the Haar functional on $C(U(n)_q)$ [Wo2] such that $h'_n(1) = 1$. From

$$\begin{aligned} (h'_n \otimes 1)(\Phi' \otimes 1)(\Delta(u_{n+1,i}^* u_{n+1,j})) &= (h'_n \otimes 1)(\overline{t_{n+1}} t_{n+1} \otimes u_{n+1,i}^* u_{n+1,j}) \\ &= (h'_n \otimes 1)(1 \otimes u_{n+1,i}^* u_{n+1,j}) = u_{n+1,i}^* u_{n+1,j}, \end{aligned}$$

we get $u_{n+1,i}^* u_{n+1,j} \in C(\mathbb{C}P_q^n)$ for $1 \leq i, j \leq n+1$. On the other hand, similar to the discussion in [Sh4], we can prove that $C(\mathbb{C}P_q^n)$ is generated by $u_{n+1,i}^* u_{n+1,j}$, by verifying that the monomials

$$P^{r,i,j,m} = z_{11}^{r_1} \dots z_{nn}^{r_n} z_{i_1, j_1} \dots z_{i_m, j_m}$$

(i) are linearly independent in the quantum case (i.e. when $q > 1$) and (ii) linearly span the *-algebra generated by $\overline{u_{n+1,i}} u_{n+1,j}$, or the *-subalgebra of \mathbb{T} -invariant polynomials in $\overline{u_{n+1,i}}$'s and $u_{n+1,j}$'s, in the classical case (i.e. when $q = 1$), where $m, r_1, \dots, r_n \geq 0$, $z_{i,j} = u_{n+1,i}^* u_{n+1,j}$, and the two sets of indices $i_1 \geq i_2 \geq \dots \geq i_m$ and $j_1 \leq j_2 \leq \dots \leq j_m$ are disjoint subsets of $\{1, 2, \dots, n+1\}$. Note that if we allow all $u_{n+1,i}^*$'s and $u_{n+1,j}$'s in $P^{i,j,m}$ to commute, then we can rewrite it, by simply permuting such factors, in the form

$$p^{i,j,k} = (u_{n+1,n}^*)^{i_n} \dots (u_{n+1,1}^*)^{i_1} y_1^{j_1} \dots y_n^{j_n} y_{n+1}^{j_{n+1}} (u_{n+1,1})^{k_1} \dots (u_{n+1,n})^{k_n}$$

where $y_m = u_{n+1,m} u_{n+1,m}^*$ and $i_m, j_m, k_m \geq 0$ with $i_m k_m = 0$ for $1 \leq m \leq n$, $y_{n+1}^{j_{n+1}} = (u_{n+1,n+1})^{j_{n+1}}$ if $j_{n+1} \geq 0$, and $y_{n+1}^{j_{n+1}} = (u_{n+1,n+1}^*)^{-j_{n+1}}$ if $j_{n+1} \leq 0$. So condition (ii) can be verified easily by integrating \mathbb{T} -action on polynomials $p^{i,j,k}$ which form a linear basis of polynomials in $\overline{u_{n+1,i}}$'s and $u_{n+1,j}$'s on \mathbb{S}^{2n+1} . On the other hand, condition (i) can be verified by considering the weight functions and the orders of the weighted shifts of

$$(\tau'_{n+1} \otimes \pi_{n(n-1) \dots 21}) \Delta^n (P^{r,i,j,m})$$

which can be determined by

$$(\tau'_{n+1} \otimes \pi_{n(n-1) \dots 21}) \Delta^n (u_{n+1,i}) = \text{id}_{\mathbb{T}} \otimes \underbrace{\gamma \otimes \dots \otimes \gamma}_{n+1-i} \otimes \underbrace{\alpha^* \otimes 1 \otimes \dots \otimes 1}_{i-1}.$$

Under the faithful *-representation $(\tau'_{n+1} \otimes \pi_{n(n-1)\dots 21}) \Delta^n$, we have

$$\begin{aligned}
& (\tau'_{n+1} \otimes \pi_{n(n-1)\dots 21}) \Delta^n (u_{n+1,i}^* u_{n+1,j}) \\
= & [(\tau'_{n+1} \otimes \pi_{n(n-1)\dots 21} \Delta^{n-1}) (\Delta u_{n+1,i})]^* (\tau'_{n+1} \otimes \pi_{n(n-1)\dots 21} \Delta^{n-1}) (\Delta u_{n+1,j}) \\
& = (\text{id}_{\mathbb{T}} \otimes \pi_{n(n-1)\dots 21} \Delta^{n-1} u_{n+1,i})^* (\text{id}_{\mathbb{T}} \otimes \pi_{n(n-1)\dots 21} \Delta^{n-1} u_{n+1,j}) \\
& = 1 \otimes (\pi_{n(n-1)\dots 21} \Delta^{n-1} u_{n+1,i})^* (\pi_{n(n-1)\dots 21} \Delta^{n-1} u_{n+1,j}) \\
& = 1 \otimes \pi_{n(n-1)\dots 21} \Delta^{n-1} (u_{n+1,i}^* u_{n+1,j}).
\end{aligned}$$

So it is clear that the *-representation

$$\pi_{n(n-1)\dots 21} \Delta^{n-1} = (\pi_n \otimes \pi_{n-1} \otimes \dots \otimes \pi_1) \Delta^{n-1}$$

gives an embedding of $C(\mathbb{C}P_q^n)$ into the groupoid C*-algebra $C^*(\mathcal{T}_n)$, where

$$\mathcal{T}_n = \mathbb{Z}^n \times \overline{\mathbb{Z}}^n |_{\overline{\mathbb{Z}}_{\geq}^n}.$$

Applying a similar analysis as used in [Sh5], we can get the following results. Let

$$\widetilde{\mathfrak{T}}_n := \{(j, k) \in \mathcal{T}_n \mid k_i = \infty \implies$$

$$j_i = -j_1 - j_2 - \dots - j_{i-1} \text{ and } j_{i+1} = \dots = j_n = 0\}$$

be a subgroupoid of \mathcal{T}_n . Define a subquotient groupoid $\mathfrak{T}_n := \widetilde{\mathfrak{T}}_n / \sim$ where \sim is the equivalence relation generated by

$$(j, k) \sim (j, k_1, \dots, k_i = \infty, \infty, \dots, \infty)$$

for all (j, k) with $k_i = \infty$ for an $1 \leq i \leq n$.

Theorem 1 $C(\mathbb{C}P_q^n) \simeq C^*(\mathfrak{T}_n)$ and hence is independent of q .

Corollary 2 There is a short exact sequence

$$0 \rightarrow \mathcal{K} \rightarrow C(\mathbb{C}P_q^k) \rightarrow C(\mathbb{C}P_q^{k-1}) \rightarrow 0$$

for $k \geq 1$ with $C(\mathbb{C}P_q^0) \simeq \mathbb{C}$.

Corollary 3 The C*-algebra $C(\mathbb{C}P_q^n)$ has the following composition sequence,

$$C(\mathbb{C}P_q^n) = \mathcal{I}_0 \supseteq \mathcal{I}_1 \supseteq \dots \supseteq \mathcal{I}_n \supseteq \mathcal{I}_{n+1} := 0,$$

with

$$\mathcal{I}_k / \mathcal{I}_{k+1} \simeq \mathcal{K}(\ell^2(\mathbb{Z}^k))$$

for $k > 0$ and $\mathcal{I}_0 / \mathcal{I}_1 \simeq \mathbb{C}$.

The above short exact sequences reflect faithfully the underlying singular foliation defined by the canonical $SU(n+1)$ -covariant Poisson structure on $\mathbb{C}P^n = U(n) \backslash SU(n+1)$.

5 Nonstandard quantum projective spaces $\mathbb{C}P_{q,c}^n$

Recently Korogodsky and Vaksman studied nonstandard quantum projective spaces $\mathbb{C}P_{q,c}^n$ [KoVa]. Here we use the approach and result of Dijkhuizen and Noumi [DiNo]. They define $\mathbb{C}P_{q,c}^n$ by

$$C(\mathbb{C}P_{q,c}^n) := \{v \in C(SU(n+1)_q) : v \cdot \mathfrak{k}^c = 0\}$$

for some coideal \mathfrak{k}^c of the quantized universal enveloping algebra $\mathcal{U}_q(\mathfrak{su}(n+1))$, $c \in (0, \infty)$, and show that

$$C(\mathbb{C}P_{q,c}^n) \cong C^*(\{x_i^* x_j \mid 1 \leq i, j \leq n+1\}) \subset C(SU(n+1)_q)$$

where

$$x_i = \sqrt{c}u_{1,i} + u_{n+1,i}.$$

We remark that when $c = 0$, we have $C(\mathbb{C}P_{q,c}^n) \cong C(\mathbb{C}P_q^n)$ the standard case, but for $c \in (0, \infty)$, $\mathbb{C}P_{qc}^n$ does not come from $SU(n+1)_q$ modulo a quantum subgroup.

To embed $C(\mathbb{C}P_{q,c}^n)$ into a groupoid C^* -algebra, we first note that the unique maximal element in the Weyl group of $SU(n+1)$ can also be expressed in the reduced form

$$s_2 s_3 s_2 s_4 s_3 s_2 \dots s_n s_{n-1} \dots s_2 s_1 s_2 \dots s_{n-1} s_n.$$

So by Soibelman's classification of irreducible $*$ -representations [So],

$$(\tau_{n+1} \otimes \pi_{232432\dots n(n-1)\dots 2123\dots(n-1)n}) \Delta^N$$

is a faithful $*$ -representation of $C(SU(n+1)_q)$.

Note that

$$\begin{aligned} & (\tau_{n+1} \otimes \pi_{232432\dots n(n-1)\dots 2123\dots(n-1)n}) \Delta^N (u_{k,i}) \\ &= t_k \otimes 1 \otimes \dots \otimes 1 \otimes (\pi_{n(n-1)\dots 2123\dots(n-1)n} \Delta^{2n-2} (u_{k,i})) \end{aligned}$$

when $k = 1$ or $k = n+1$. Thus $(\tau_{n+1} \otimes \pi_{n(n-1)\dots 2123\dots(n-1)n}) \Delta^{2n-1}$ is a faithful $*$ -representation of $C^*(\{u_{1,i}, u_{n+1,i}\}_{i=1}^{n+1}) \supset C(\mathbb{C}P_{qc}^n)$. Furthermore, since

$$(\tau_{n+1} \otimes \pi_{n(n-1)\dots 2123\dots(n-1)n}) \Delta^{2n-1} (x_i) = t_1 \otimes \sqrt{c} \pi_{n(n-1)\dots 2123\dots(n-1)n} \Delta^{2n-2} (u_{1,i})$$

$$+ (t_1 t_2 \dots t_n)^{-1} \otimes \pi_{n(n-1)\dots 2123\dots(n-1)n} \Delta^{2n-2} (u_{n+1,i})$$

$$\in \{t_1^l (t_2 \dots t_n)^m : l - 2m = 1\} \otimes C^*(\mathbb{Z}^{2n-1} \times \overline{\mathbb{Z}}_{\geq}^{2n-1}),$$

we have

$$\begin{aligned}
& (\tau_{n+1} \otimes \pi_{n(n-1)..2123...(n-1)n}) \Delta^{2n-1} (x_i^* x_j) \\
& \in \{t_1^l (t_2 \dots t_n)^m : l - 2m = 0\} \otimes C^* \left(\mathbb{Z}^{2n-1} \times \overline{\mathbb{Z}}^{2n-1} \Big|_{\overline{\mathbb{Z}}_{\geq}^{2n-1}} \right) \\
& = \left\{ (t_1^2 t_2 \dots t_n)^m : m \in \mathbb{Z} \right\} \otimes C^* \left(\mathbb{Z}^{2n-1} \times \overline{\mathbb{Z}}^{2n-1} \Big|_{\overline{\mathbb{Z}}_{\geq}^{2n-1}} \right).
\end{aligned}$$

Thus $(\tau_{n+1} \otimes \pi_{n(n-1)..2123...(n-1)n}) \Delta^{2n-1}$ embeds $C(\mathbb{C}P_{qc}^n)$ into

$$\begin{aligned}
& C^*(\mathbb{Z}) \otimes C^* \left(\mathbb{Z}^{2n-1} \times \overline{\mathbb{Z}}^{2n-1} \Big|_{\overline{\mathbb{Z}}_{\geq}^{2n-1}} \right) \\
& \cong C^* \left(\{t_1^2 t_2 \dots t_n\} \right) \otimes C^* \left(\mathbb{Z}^{2n-1} \times \overline{\mathbb{Z}}^{2n-1} \Big|_{\overline{\mathbb{Z}}_{\geq}^{2n-1}} \right) \\
& \subset C(\mathbb{T}^n) \otimes C^* \left(\mathbb{Z}^{2n-1} \times \overline{\mathbb{Z}}^{2n-1} \Big|_{\overline{\mathbb{Z}}_{\geq}^{2n-1}} \right).
\end{aligned}$$

Theorem 4 For $0 < c < \infty$, $C(\mathbb{C}P_{qc}^n)$ can be embedded into the groupoid C^* -algebra $C^*(\mathfrak{G}_n)$ where

$$\mathfrak{G}_n := \mathbb{Z} \times \left(\mathbb{Z}^{2n-1} \times \overline{\mathbb{Z}}^{2n-1} \Big|_{\overline{\mathbb{Z}}_{\geq}^{2n-1}} \right).$$

When $n = 1$, we have $\mathbb{C}P^1 = \mathbb{S}^2$ in the classical case, while in the quantum case, $\mathbb{C}P_{qc}^1$ is indeed a Podleś' quantum sphere $\mathbb{S}_{\mu c}^2$ [Po] (with $\mu = q^{-1}$) for some (different) c . In [LuWe2] Lu and Weinstein classified all $SU(2)$ -covariant Poisson structures on \mathbb{S}^2 by a real parameter and showed that each 'nonstandard' $SU(2)$ -covariant Poisson sphere (corresponding to a suitable parameter $c \in (0, \infty)$) contains the trivial Poisson 1-sphere \mathbb{S}^1 (consisting of a circle family of 0-dimensional symplectic leaves) and exactly two 2-dimensional symplectic leaves. This geometric structure is again reflected faithfully in the algebraic structure of the algebra $C(\mathbb{S}_{\mu c}^2)$ of the nonstandard quantum spheres $\mathbb{S}_{\mu c}^2$ as follows.

Theorem 5 For $0 < c < \infty$, $C(\mathbb{S}_{\mu c}^2) \cong C^*(\mathfrak{G}')$, where

$$\mathfrak{G}' = \{(j, j, k_1, k_2) \mid k_1 = \infty \text{ or } k_2 = \infty\}$$

is a subgroupoid of the groupoid $\mathbb{Z}^2 \times \overline{\mathbb{Z}}^2 \Big|_{\overline{\mathbb{Z}}_{\geq}^2}$, and there is a short exact sequence

$$0 \rightarrow \mathcal{K} \oplus \mathcal{K} \rightarrow C(\mathbb{S}_{\mu c}^2) \rightarrow C(\mathbb{S}^1) \rightarrow 0.$$

On higher dimensional projective spaces $\mathbb{C}P^n$, we also have a one-parameter family of nonstandard $SU(n+1)$ -covariant Poisson structures suitably parametrized by $c \in (0, \infty)$, and similar to the case of $n = 1$, each such nonstandard Poisson $\mathbb{C}P^n$ contains an embedded copy of a standard Poisson \mathbb{S}^{2n-1} [Sh6]. On the quantum level, one would then expect that $C(\mathbb{C}P_{qc}^n)$ should have $C(\mathbb{S}_q^{2n-1})$ as a quotient to reflect this geometric fact. Using the above groupoid description of $C(\mathbb{C}P_{qc}^n)$, we can indeed show that this is the case.

Theorem 6 *For $0 < c < \infty$, there is a short exact sequence*

$$0 \rightarrow \mathcal{I} \rightarrow C(\mathbb{C}P_{qc}^n) \rightarrow C(\mathbb{S}_q^{2n-1}) \rightarrow 0$$

for some ideal \mathcal{I} .

Proof. First we note that since $(\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1}$ is faithful on $C(\mathbb{S}_q^{2n+1})$,

$$\begin{aligned} & (\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1}(u_{n+1,i}) = t_{n+1} \otimes \pi_{n(n-1)..2123...(n-1)n} \Delta^{2n-2} u_{n+1,i} \\ &= (1 - \delta_{n+1,i}) t_{n+1} \otimes \gamma^{\otimes n-i} \otimes \underbrace{\alpha^* \otimes 1 \otimes \dots \otimes 1}_{i-1} \otimes 1^{\otimes i-1} \otimes \underbrace{\gamma \otimes 1 \otimes \dots \otimes 1}_{(n+1)-i} + (1 - \delta_{1,i})(1 - \delta_{2,i}) \\ & \left[\sum_{k=n+1-i}^{n-2} t_{n+1} \otimes \gamma^{\otimes k} \otimes \alpha^* \otimes 1^{\otimes n-k-1} \otimes 1^{\otimes n-k-2} \otimes \alpha^* \otimes (-q^{-1}\gamma)^{\otimes k-(n+1)+i} \otimes \underbrace{\alpha \otimes 1 \otimes \dots \otimes 1}_{(n+1)-i} \right] \\ & + (1 - \delta_{1,i}) t_{n+1} \otimes \gamma^{\otimes n-1} \otimes \alpha^* \otimes (-q^{-1}\gamma)^{\otimes i-2} \otimes \underbrace{\alpha \otimes 1 \otimes \dots \otimes 1}_{(n+1)-i} \end{aligned}$$

with $1 \leq i \leq n+1$ generate a C^* -algebra isomorphic to $C(\mathbb{S}_q^{2n+1})$.

Now let $\rho : C^*(\mathbb{Z} \times \overline{\mathbb{Z}}|_{\overline{\mathbb{Z}}_{\geq}}) \rightarrow \mathbb{C}$ be the composition of the C^* -homomorphism

$$C^*(\mathbb{Z} \times \overline{\mathbb{Z}}|_{\overline{\mathbb{Z}}_{\geq}}) \rightarrow C^*(\mathbb{Z} \times \overline{\mathbb{Z}}|_{\{\infty\}}) = C^*(\mathbb{Z}) \cong C(\mathbb{T})$$

induced by the restriction to the invariant closed subset $\{\infty\}$ of the unit space $\overline{\mathbb{Z}}_{\geq}$ and the evaluation map $C(\mathbb{T}) \rightarrow \mathbb{C}$ at $1 \in \mathbb{T}$, and let

$$\tilde{\rho} := \underbrace{1 \otimes \dots \otimes 1}_{n-1} \otimes \rho \otimes \rho \otimes \underbrace{1 \otimes \dots \otimes 1}_{n-2} : C^*(\mathcal{T}_{2n-1}) \rightarrow C^*(\mathcal{T}_{2n-3})$$

be the homomorphism that ‘removes’ the middle two tensor factors. Then by direct calculation, we get

$$(1 \otimes \tilde{\rho})(\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1}(\sqrt{c}u_{1,1} + u_{n+1,1})$$

$$= \sqrt{c} t_1 \otimes 1 \otimes \dots \otimes 1$$

while

$$\begin{aligned} & (1 \otimes \tilde{\rho}) (\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1} (\sqrt{c} u_{1,i} + u_{n+1,i}) \\ &= t_{n+1} \otimes \pi_{n(n-1)..323...(n-1)n} \Delta^{2n-4} (u_{n+1,i}) \end{aligned}$$

for $i > 1$. So

$$(1 \otimes \tilde{\rho}) (\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1} (x_1^* x_1) = c,$$

and for $i > 1$,

$$\begin{aligned} & [(1 \otimes \tilde{\rho}) (\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1} (x_i^* x_1)]^* \\ &= (1 \otimes \tilde{\rho}) (\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1} (x_1^* x_i) \\ &= \sqrt{c} (t_1^2 t_2 \dots t_n)^{-1} \otimes \pi_{n(n-1)..323...(n-1)n} \Delta^{2n-4} (u_{n+1,i}) \\ &= \sqrt{c} (\tau'' \otimes \pi_{n(n-1)..323...(n-1)n}) \Delta^{2n-3} (u_{n+1,i}) \end{aligned}$$

while for $i, j > 1$,

$$\begin{aligned} & (1 \otimes \tilde{\rho}) (\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1} (x_i^* x_j) \\ &= 1 \otimes \pi_{n(n-1)..323...(n-1)n} \Delta^{2n-4} (u_{n+1,i}^* u_{n+1,j}) \\ &= (\tau'' \otimes \pi_{n(n-1)..323...(n-1)n}) \Delta^{2n-3} (u_{n+1,i}^* u_{n+1,j}), \end{aligned}$$

where $\tau''(u_{ij}) = \delta_{i,j} \delta_{n+1,i} (t_1^2 t_2 \dots t_n)^{-1}$. It is straight forward to verify that

$$(\tau'' \otimes \pi_{n(n-1)..323...(n-1)n}) \Delta^{2n-3} (u_{n+1,i}),$$

with $1 < i \leq n+1$, coincides with the operators

$$(\tau_n \otimes \pi_{(n-1)..212...(n-1)}) \Delta^{2n-3} (u_{n,i-1})$$

whose formula is explicitly written above for the case of dimension $n+1$. So

$$\begin{aligned} & (1 \otimes \tilde{\rho}) (\tau_{n+1} \otimes \pi_{n(n-1)..212...(n-1)n}) \Delta^{2n-1} (C(\mathbb{C}P_{qc}^n)) \\ & \cong (\tau_n \otimes \pi_{(n-1)..212...(n-1)}) \Delta^{2n-3} (C(\mathbb{S}_q^{2n-1})) \cong C(\mathbb{S}_q^{2n-1}) \end{aligned}$$

and hence $C(\mathbb{S}_q^{2n-1})$ is a quotient of $C(\mathbb{C}P_{qc}^n)$. ■

References

- [Co] A. Connes, *A survey of foliation and operator algebras*, Proc. Symp. Pure Math. Vol. 38, Part I, AMS, Providence, 1982, 521-628.
- [CuM] R. E. Curto and P. S. Muhly, *C*-algebras of multiplication operators on Bergman spaces*, J. Func. Anal. 64 (1985), 315-329.
- [DiNo] M. S. Dijkhuizen and M. Noumi, *A family of quantum projective spaces and related q-hypergeometric orthogonal polynomials*, preprint.
- [Dr] V. G. Drinfeld, *Quantum groups*, Proc. I.C.M. Berkeley 1986, Vol. 1, 789-820, Amer. Math. Soc., Providence, 1987.
- [KoVa] L. I. Korogodsky and L. L. Vaksman, *Quantum G-spaces and Heisenberg algebra*, in "Quantum Groups", P. P. Kulish (Ed.), Lecture Notes in Mathematics 1510, Springer-Verlag, Berlin, 1992, pp. 56-66.
- [LuWe1] J. H. Lu and A. Weinstein, *Poisson Lie groups, dressing transformations and Bruhat decompositions*, J. Diff. Geom. 31 (1990), 501-526.
- [LuWe2] ———, *Classification of SU(2)-covariant Poisson structures on \mathbb{S}^2* , Comm. Math. Phys. 135 (1991), 229-231.
- [MRe] P. S. Muhly and J. N. Renault, *C*-algebras of multivariable Wiener-Hopf operators*, Trans. Amer. Math. Soc. 274 (1982), 1-44.
- [Na1] G. Nagy, *On the Haar measure of the quantum SU(N) group*, Comm. Math. Phys. 153 (1993), 217-228.
- [Na2] ———, *A deformation quantization procedure for C*-algebras*, preprint.
- [Po] P. Podleś, *Quantum spheres*, Letters Math. Phys. 14 (1987), 193-202.
- [Re] J. Renault, *A Groupoid Approach to C*-algebras*, Lecture Notes in Mathematics, Vol. 793, Springer-Verlag, New York, 1980.
- [RTF] N. Yu. Reshetikhin, L. A. Takhtadzhyan, and L. D. Faddeev, *Quantization of Lie groups and Lie algebras*, Leningrad Math. J. 1 (1990), 193-225.
- [Ri1] M. A. Rieffel, *Deformation quantization and operator algebras*, in "Proc. Symp. Pure Math., Vol. 51", AMS, Providence, 1990, pp. 411-423.
- [Ri2] ———, *Deformation quantization for actions of \mathbb{R}^d* , Memoirs of AMS, Vol. 106, No. 506, 1993.
- [Ri3] ———, *Compact quantum groups associated with toral subgroups*, in "Contemporary Mathematics", Vol. 145, AMS, Providence, 1993, pp. 465-491.
- [SaShU] N. Salinas, A. J. L. Sheu, and H. Upmeyer, *Toeplitz operators on pseudoconvex domains and foliation C*-algebras*, Annals of Mathematics, 130 (1989), 531-565.

- [Sh1] A. J. L. Sheu, *Reinhardt domains, boundary geometry and Toeplitz C^* -algebras*, Journal of Functional Analysis, 92 (1990), 264-311.
- [Sh2] ———, *Quantization of the Poisson $SU(2)$ and its Poisson homogeneous space – the 2-sphere*, Comm. Math. Phys. 135 (1991), 217-232.
- [Sh3] ———, *Symplectic leaves and deformation quantization*, Proc. Amer. Math. Soc., 124 (1996), 95-100.
- [Sh4] ———, *Compact quantum groups and groupoid C^* -algebras*, J. Func. Anal. 144 (1997), 371-393.
- [Sh5] ———, *Quantum spheres as groupoid C^* -algebras*, to appear in Quarterly J. Math.
- [Sh6] ———, *Covariant Poisson structures on complex projective spaces*, preprint.
- [So] Ya. S. Soibelman, *The algebra of functions on a compact quantum group, and its representations*, Algebra Analiz. 2 (1990), 190-221. (Leningrad Math. J., 2 (1991), 161-178.)
- [VaSo] L. L. Vaksman and Ya. S. Soibelman, *The algebra of functions on the quantum group $SU(n + 1)$, and odd-dimensional quantum spheres*, Leningrad Math. J. 2 (1991), 1023-1042.
- [We] A. Weinstein, *The local structure of Poisson manifolds*, J. Diff. Geom. 18 (1983), 523-557.
- [Wo1] S. L. Woronowicz, *Twisted $SU(2)$ group: an example of a non-commutative differential calculus*, Publ. RIMS. 23 (1987), 117-181.
- [Wo2] ———, *Compact matrix pseudogroups*, Comm. Math. Phys. 111 (1987), 613-665.
- [Wo3] ———, *Tannaka-Krein duality for compact matrix pseudogroups, twisted $SU(N)$ groups*, Invent. Math. 93 (1988), 35-76.