

Spherical 2-categories and 4-manifold invariants

Marco Mackaay

Area Departamental de Matematica
Universidade do Algarve, UCEH
Campus de Gambelas
8000 Faro
Portugal
e-mail: mmackaay@ualg.pt

Introduction

In this paper I give a formal construction of a 4-manifold invariant out of what I call a *non-degenerate finitely semi-simple semi-strict spherical 2-category*.

For some time now people have had the feeling that 4-manifold invariants and certain kinds of monoidal 2-categories have a relation with each other similar to that of 3-manifold invariants and certain kinds of monoidal categories.

The first evidence for this feeling can be found in the work of Crane and collaborators. In [15] Crane and Frenkel give a formal construction of 4-manifold invariants out of Hopf categories and indicate where one should look for such algebraic objects, namely in the work on crystal bases by Saito and the work on canonical bases and perverse sheaves by Lusztig. One could argue that it should be possible to use the 2-category of representations of a Hopf category instead of the Hopf category itself; the reason for this thought being that for the construction of 3-manifolds one can use Hopf algebras, as Kuperberg [24] and Chung, Fukuma and Shapere did [14], or the category of representations of Hopf algebras as people like Turaev and Viro [29], Yetter [30], and Barret and Westbury [9] did. Recently Neuchl [26] showed that the representations of a Hopf category do form a monoidal 2-category indeed.

In [17] Crane and Yetter proposed a construction of 4-manifold invariants out of the semi-simple sub-quotient of the category of finite dimensional representations of the quantum group $U_q(sl(2))$ for q a principal $4r^{th}$ root of unity. In [16] Crane, Kauffman and Yetter generalized this construction for any tortile semi-simple category and gave detailed proofs. These Crane-Yetter invariants can be seen as a special case in which the authors use a 2-category of a certain type with only one object.

Another piece of evidence for the aforementioned “feeling” is the work of Baez and collaborators. In a series of papers [1,2,3,4] they have tried to persuade people that the study of monoidal n -categories is a promising road towards the construction of Topological Quantum Field Theories, i.e. n -functors from the n -category of n -cobordisms to the n -category of n -vector spaces. Here of course it is reasonable to start with the case $n = 2$.

Based on the work of Carter, Rieger and Saito [12,13], several people [5,6,20,23,25] have shown that braided 2-categories with duals form the right algebraic context in which to study invariants of 2-tangles. This should be closely related to the construction of 4-manifold invariants, at least formally.

So altogether one could say that there is more than enough reason to believe that it is possible to construct 4-manifold invariants out of certain kinds of 2-categories. But nowhere in the literature can one find a paper with an explicit construction. This paper is meant to fill this gap.

In this paper I use triangulations of 4-manifolds for the construction of a state sum. I will show that this state sum is independent of the chosen triangulation by using Pachner's theorem [27], that relates triangulations of piece-wise linear homeomorphic manifolds. The whole construction should be considered as a lift to the fourth dimension of Barrett's and Westbury's [9] construction of 3-manifold invariants out of non-degenerate finitely semi-simple spherical categories. For that reason I have given the kind of 2-categories that I use the name above.

I have to stress that it is a purely formal construction because I do not have any interesting examples of such categories yet. Examples of spherical 2-categories are not hard to find. The 2-category of 2-Hilbert spaces, 2Hilb (see [3]), is spherical, for example. It is likely that 2Tang , the 2-category of 2-tangles (see [25]) is spherical, but I have not checked it. But it is hard to find spherical 2-categories that are finitely semi-simple. It is likely that the 2-category of representations of the right kind of Hopf category will be such a 2-category. But we have only one Hopf category that has been worked out in detail, namely the categorification $C(D(G))$ of the quantum double of a finite group [18]. Probably $C(D(G))\text{-mod}$, the 2-category of finite dimensional representations of $C(D(G))$, is an example of a non-degenerate finitely semi-simple semi-strict spherical 2-category. But, again, I have not worked out the details yet. Carter, Kauffman and Saito are working out the Crane-Frenkel invariant for this particular example [11]. In [15] Crane and Frenkel indicate that it is possible to construct more examples of Hopf categories $C(U_0(g))$ using the crystal bases of quantum groups at $q = 0$. However, they do not explain how to obtain the right categorification of the antipode in these cases. In the aforementioned case of $C(D(G))$ this antipode looks to be straight forward, but Carter, Kauffman and Saito found that it is necessary to impose extra conditions on the cocycles defining the structure isomorphisms of the Hopf category $C(D(G))$ in order to obtain invariance of the state sum under permutation of the vertices of a chosen triangulation of the 4-manifold. This was not foreseen in [15] and, as far as I know, it is not known how to obtain invariance under permutation of vertices, or equivalently how to define the right categorification of the antipode, in the case of $C(U_0(g))$.

The strongest argument for the possible existence of non-degenerate finitely semi-simple semi-strict spherical 2-categories lies in a future paper in which Crane and Yetter show how to build a monoidal 2-category out of the modules of a quantum group at $q = 0$ using their crystal bases. Here they avoid the Hopf categories and build the 2-categories directly, which makes a construction of 4-manifold invariants out of a certain kind of 2-categories, as presented in this paper, even more desirable. It is definitely a good place to look for interesting examples. It seems that the "right" definition of duality, i.e. the one that ensures invariance under permutation of vertices, here is easier to find than the "right" definition of the antipode for Hopf categories. But it is not likely that these 2-categories are already the ones we are looking for. It is like having the non-finitely semi-simple "trivially" spherical category $U(g)\text{-mod}$, where g is a finite dimensional semi-simple Lie algebra, and

having an abstract Turaev-Viro-like construction of 3-manifold invariants which requires finitely semi-simple spherical categories. In that case the missing link comes from the deformation theory of $U(g)$ which shows that there are deformations $U_q(g)$, where q is a certain root of unity, such that there is a certain non-degenerate quotient of the category of *tilting modules* of $U_q(g)$ that is a finitely semi-simple spherical category. For details about this see [9] and some references therein. Likewise, in the case of 2-categories, one should first define the deformation theory of monoidal 2-categories analogously to what Crane and Yetter have done for monoidal categories [19]. Then one has to find actual deformations of the aforementioned 2-categories and finally one has to “melt” them, i.e. get back to generic q . These last two problems are of a very deep nature and certainly far beyond the scope of this paper. In the meanwhile it is worthwhile, I think, to study the kind of 2-category we are looking for from an abstract point of view.

1. The basic idea

Throughout this paper a manifold means an oriented piece-wise linear compact 4-manifold without boundary. A triangulated manifold (M, T) is a manifold M together with a given simplicial 4-complex T , the triangulation, such that its polytope is pl-homeomorphic to M . Throughout this paper we will always assume that there is a total ordering on the vertices of the triangulation of a manifold. A combinatorial isomorphism between two triangulated manifolds (M, T) and (M', T') will always mean an isomorphism between the simplicial complexes T and T' . A simplicial isomorphism between two triangulated manifolds (M, T) and (M', T') will always mean a combinatorial isomorphism that preserves the ordering on the vertices. I also want to fix some notation. The letter \mathbf{F} will always denote a fixed field of characteristic 0 and any vector space in this paper will be a finite dimensional vector space over \mathbf{F} . My notation for the simplices follows Barrett’s and Westbury’s convention in [9]. The standard n -simplex $(012\dots n)$ with vertices $\{0, 1, 2, \dots, n\}$ has the standard orientation $(+)$. The opposite orientation is denoted by $(-)$. The standard 4-simplex $+(01234)$ has boundary

$$(1234) - (0234) + (0134) - (0124) + (0123).$$

The 4-simplex $-(01234)$ has the same boundary but with the opposite signs. The sign with which a tetrahedron appears in the boundary of a 4-simplex I will call the induced orientation of the tetrahedron. The total ordering on the vertices of the simplicial complex that defines the triangulation of a manifold induces an ordering on all 4-simplices of course and the ordering on the vertices of a 4-simplex induces an orientation on its underlying polytope. If the orientation of the underlying polytope of a 4-simplex induced by the orientation of the manifold is equal to the orientation of this polytope induced by the total ordering on the vertices of the 4-simplex, then our convention will be that the 4-simplex has the positive orientation, as a simplex, and if the two induced orientations of the polytope are opposite, we take the 4-simplex to be negatively oriented, as a simplex.

Remark 1.1. Notice that in a triangulated manifold each tetrahedron lies in the boundary of exactly two 4-simplices, occuring once with induced orientation $(+)$ and once with $(-)$.

Let (M, T) be a triangulated manifold. For the definition of my state sum I need two sets of labels, \mathcal{E} and \mathcal{F} respectively. The edge (ij) with vertices i, j is labelled with

$e_{ij} \in \mathcal{E}$, the face (ijk) with vertices i, j, k is labelled with $f_{ijk} \in \mathcal{F}$. Let $T((0123), e, f)$ be the labelled standard oriented tetrahedron $+(0123)$. We do not take the different orientations of the edges and the faces into account for the labelling yet.

Notation 1.2. *The state space of this labelled tetrahedron is a certain vector space*

$$H((0123), e, f).$$

The state space of $-(0123)$ is defined to be the dual vector space

$$H((0123), e, f)^*.$$

Notation 1.3. *Let $+(01234)$ be the standard oriented 4-simplex with the triangle (ijk) labelled by $f_{ijk} \in \mathcal{F}$ and the edge (ij) by $e_{ij} \in \mathcal{E}$. Denote the state space of the labelled tetrahedron $(ijkl)$ by $H(ijkl)$ just as a shorthand. The partition function of this labelled 4-simplex is a certain linear map*

$$Z(+01234): H(0234) \otimes H(0124) \rightarrow H(1234) \otimes H(0134) \otimes H(0123).$$

The partition function of $-(01234)$ is also a certain linear map

$$Z(-01234): H(1234) \otimes H(0134) \otimes H(0123) \rightarrow H(0234) \otimes H(0124).$$

Notice that $Z(\pm(01234))$ is defined for a fixed labelling of T , although this dependence does not show up in the notation. This is a deliberate choice, or a deliberate flaw in the notation as you wish, that I want to allow myself in order to write down formulas that are not too polluted and incomprehensibly complex by a high number of sub- and superscripts. I think the context will leave no doubt of what depends on what in my formulas.

Assume that $M = (M, T)$ is labelled with a fixed labelling ℓ . Notice that the ordering on the vertices of T induces a natural total ordering on the tetrahedra, by means of the boundary operator, within each 4-simplex. For example in the ordered 4-simplex $(abcde)$ the ordering is given by 1. $(bcde)$, 2. $(acde)$, 3. $(abde)$, 4. $(abce)$, 5. $(abcd)$, which is independent of the orientation. In the same way we get a fixed ordering on the triangles within each tetrahedron etc. Fix also a total ordering on the 4-simplices, for example the one induced by the total ordering on the vertices of the whole triangulation. Take out of each 4-simplex the tetrahedra that appear with a negative sign in its boundary in the induced order described above. Together with the choosen ordering on the 4-simplices this fixes an ordering of all the tetrahedra of M . Notice that each tetrahedron appears exactly once in this way by remark 1.1.

Definition 1.4. *Let $V(M, T, \ell)$ be the tensor product of the state spaces of all the tetrahedra of T in which the ordering of the factors is as described above. The tensor product of the respective partition functions applied to $V(M, T, \ell)$ gives a permuted tensor product $V(M, T, \ell)'$. Again remark 1.1 is vital. Now compose this linear map with the linear map $P(M, T, \ell): V(M, T, \ell)' \rightarrow V(M, T, \ell)$ induced by the standard flip $P: x \otimes y \rightarrow y \otimes x$. The*

result is a linear map $L(M, T, \ell): V(M, T, \ell) \rightarrow V(M, T, \ell)$. The element $Z(M, T, \ell) \in \mathbf{F}$ is defined to be the trace of $L(M, T, \ell)$.

Notice that $Z(M, T, \ell)$ does not depend on the ordering on the 4-simplices, because it is defined by a conjugation invariant trace. In the next section we will prove that $Z(M, \ell)$ is a combinatorial invariant of M .

The state sum $I(M, T)$ is a certain weighted sum over all labellings of the numbers $Z(M, T, \ell)$.

So far we have only sketched the basic idea of our approach without telling anyone where to get these state spaces and these partition functions. In section 3 we will show how they appear naturally out of a certain kind of 2-categories. Therefore we have to study this kind of 2-categories in the next section first.

2. Spherical 2-categories

In this paragraph I define what I call a spherical 2-category. The underlying 2-category will always be assumed to be strict. This means that the composition is strictly associative and the composition of a 1-morphism with an identity 1-morphism is equal to the 1-morphism itself. I will denote the composition of two 1-morphisms f, g by fg , the vertical composition of two 2-morphisms α, β by $\alpha \cdot \beta$ and their horizontal composition by $\alpha \circ \beta$. If $f, f': A \rightarrow B$ and $g, g': B \rightarrow C$ are 1-morphisms and $\alpha: f \rightarrow f'$ and $\beta: g \rightarrow g'$ are 2-morphisms, then $f \circ \beta: fg \rightarrow fg'$ denotes $1_f \circ \beta$ and $\alpha \circ g: fg \rightarrow f'g$ denotes $\alpha \circ 1_g$.

So let C be a strict 2-category. This is not a too restrictive assumption because a weak 2-category can always be strictified, see [KV]. We also assume that C has a semi-strict monoidal structure. Loosely speaking this means that for every pair of objects A, B in C there is a unique object $A \otimes B$. For every object A and every 1-morphism $f: X \rightarrow Y$ there is a unique 1-morphism $A \otimes f: A \otimes X \rightarrow A \otimes Y$ and a unique 1-morphism $f \otimes A: X \otimes A \rightarrow Y \otimes A$. In the notation of the 1-morphism we identify A with 1_A . For every object A and every 2-morphism $\alpha: f \Rightarrow g$ there is a unique 2-morphism $A \otimes \alpha: A \otimes f \Rightarrow A \otimes g$ and a unique 2-morphism $\alpha \otimes A: f \otimes A \Rightarrow g \otimes A$. Here we identify A with 1_{1_A} in the notation of the 2-morphisms. Also there is an identity object I such that $I \otimes X = X \otimes I = X$ for all objects, 1- and 2-morphisms X . All the usual structural 2-isomorphism are identities except one: given a pair of 1-morphisms $f: A \rightarrow C$, $g: B \rightarrow D$ in C there is a 2-isomorphism

$$\otimes_{f,g}: (f \otimes B)(C \otimes g) \Rightarrow (A \otimes g)(f \otimes D).$$

This 2-isomorphism is required to satisfy a some conditions. These conditions guarantee that $\otimes_{f,g}$ behaves well under the tensor product and composition. The obvious condition that tells us how to obtain $\otimes_{fg,h}$ from $\otimes_{f,h}$ and $\otimes_{g,h}$, and analogously how to obtain $\otimes_{f,gh}$ from $\otimes_{f,g}$ and $\otimes_{f,h}$, resembles the condition defining a braiding in a monoidal category. For the exact definition of a semi-strict monoidal structure see [22].

Definition 2.1. *A semi-strict monoidal 2-category is a strict 2-category with a semi-strict monoidal structure.*

Let C always be a semi-strict monoidal 2-category. Again this is a legitimate assumption, since every weak monoidal 2-category is equivalent in a well defined sense to a semi-strict one [22]. The following lemma is well known. For a proof see [22].

Lemma 2.2. *Let I be the identity object in C . Then $\text{End}(I)$ is a category with objects all 1-morphism $I \rightarrow I$ and morphisms all 2-morphisms between such 1-morphisms. The composition in $\text{End}(I)$ is given by the vertical composition of the 2-morphisms. Furthermore $\text{End}(I)$ is a braided monoidal category with the tensor product being the horizontal composition of 1- and 2-morphisms and the braiding being defined by the 2-isomorphisms $\otimes_{\bullet, \bullet}$.*

In order to get to the spherical condition I first have to define duality in C . For this I will copy the definition Langford gives in her dissertation [25].

Definition 2.3. *C is called a semi-strict monoidal 2-category with duals if it is equipped with the following structures:*

1. *For every 2-morphism $\alpha: f \Rightarrow g$ there is a 2-morphism $\alpha^*: g \Rightarrow f$ called the dual of α .*
2. *For every 1-morphism $f: A \rightarrow B$ there is a 1-morphism $f^*: B \rightarrow A$ called the dual of f , and 2-morphisms $i_f: 1_A \Rightarrow f f^*$ and $e_f: f^* f \Rightarrow 1_B$, called the unit and counit of f , respectively.*
3. *For any object A , there is an object A^* called the dual of A , 1-morphisms $i_A: I \rightarrow A \otimes A^*$ and $e_A: A^* \otimes A \rightarrow I$ called the unit and counit of A , respectively, and a 2-morphism $T_A: (i_A \otimes A)(A \otimes e_A) \Rightarrow 1_A$ called the triangulator of A .*

We say that a 2-morphism α is *unitary* if it is invertible and $\alpha^{-1} = \alpha^*$. Given a 2-morphism $\alpha: f \Rightarrow g$, we define the *adjoint* $\alpha^\dagger: g^* \Rightarrow f^*$ by

$$\alpha^\dagger = (g^* \circ i_f) \cdot (g^* \circ \alpha \circ f^*) \cdot (e_g \circ f^*).$$

In addition, the structures above are also required to satisfy the following conditions:

1. $X^{**} = X$ for any object, 1-morphism or 2-morphism.
2. $1_X^* = 1_X$ for any object or 1-morphism X .
3. For all objects A, B , 1-morphisms f, g , and 2-morphisms α, β for which both sides of the following equations are well-defined, we have

$$(\alpha \cdot \beta)^* = \beta^* \cdot \alpha^*,$$

$$(\alpha \circ \beta)^* = \alpha^* \circ \beta^*,$$

$$(fg)^* = g^* f^*,$$

$$(A \otimes \alpha)^* = A \otimes \alpha^*,$$

$$(A \otimes f)^* = A \otimes f^*,$$

$$(\alpha \otimes A)^* = \alpha^* \otimes A,$$

$$(f \otimes A)^* = f^* \otimes A,$$

$$(A \otimes B)^* = B^* \otimes A^*.$$

4. For all 1-morphisms f, g the 2-isomorphism $\otimes_{f, g}$ is unitary.
5. For any object or 1-morphism X we have $i_{X^*} = e_X^*$ and $e_{X^*} = i_X^*$.

6. For any object A , the 2-morphism T_A is unitary.
7. For any objects A and B we have

$$i_{A \otimes B} = i_A \cdot (A \otimes i_B \otimes A^*),$$

$$e_{A \otimes B} = (B^* \otimes e_A \otimes B) \cdot e_B,$$

$$T_{A \otimes B} = [(i_A \otimes A \otimes B)(A \otimes \bigotimes_{i_B, e_A}^{-1} \otimes B)(A \otimes B \otimes e_B)] \cdot [(T_A \otimes B) \circ (A \otimes T_B)].$$

8. $T_I = 1_I$.
9. For any object A and 1-morphism f , we have

$$i_{A \otimes f} = A \otimes i_f,$$

$$i_{f \otimes A} = i_f \otimes A,$$

$$e_{A \otimes f} = A \otimes e_f,$$

$$e_{f \otimes A} = e_f \otimes A.$$

10. For any 1-morphisms f, g , $i_{fg} = i_f \cdot (f \circ i_g \circ f^*)$ and $e_{fg} = (g^* \circ e_f \circ g) \cdot e_g$.
11. For any 1-morphism f , $(i_f \circ f) \cdot (f \circ e_f) = 1_f$ and $(f^* \circ i_f) \cdot (e_f \circ f^*) = 1_{f^*}$.
12. For any 2-morphism α , $\alpha^{\dagger*} = \alpha^* \dagger$.
13. For any object A we have

$$[i_A \circ (A \otimes T_{A^*}^{\dagger})] \cdot [\bigotimes_{i_A, i_A}^{-1} (A \otimes e_A \otimes A^*)] \cdot [i_A \circ (T_A \otimes A^*)] = 1_{i_A}.$$

It would be worthwhile to study weaker notions of duality and prove a coherence theorem that allows one to strictify 2-categories with such duality up to a semi-strict monoidal 2-category with duals as defined above. In [7] Barrett and Westbury prove such a coherence theorem for monoidal categories with duals. In this paper duality is always assumed to be semi-strict.

For our purpose we need a little bit of extra structure. Let $f: A \rightarrow B$ a 1-morphism in C . We define the 1-morphism $\#f: B^* \rightarrow A^*$ by

$$B^* \xrightarrow{i_{A^*} \otimes B^*} A^* \otimes A \otimes B^* \xrightarrow{A^* \otimes f \otimes B^*} A^* \otimes B \otimes B^* \xrightarrow{A^* \otimes e_{B^*}} A^*.$$

Analogously we define $f^\#: B^* \rightarrow A^*$ by

$$B^* \xrightarrow{B^* \otimes i_A} B^* \otimes A \otimes A^* \xrightarrow{B^* \otimes f \otimes A^*} B^* \otimes B \otimes A^* \xrightarrow{e_{B^*} \otimes A^*} A^*.$$

Notice that given a 2-morphism $\alpha: f \Rightarrow g$ in C we also have the 2-morphisms $\alpha^\#: f^\# \Rightarrow g^\#$ and $\#\alpha: \#f \Rightarrow \#g$ in a natural way. The following definition reminds us of the definition of a pivotal category.

Definition 2.4. A semi-strict pivotal 2-category is a semi-strict monoidal 2-category with duals C such that for any 1-morphism $f: A \rightarrow B$ there exists a unitary 2-morphism

$$\phi_f: f^\# \Rightarrow \#f$$

satisfying the obvious condition

$$\alpha^\# \cdot \phi_g = \phi_f \cdot \# \alpha.$$

Additionally we require the following conditions to be satisfied for any 2-morphism α

$$\#(\alpha^\dagger) = \dagger(\alpha^\#) \quad \text{and} \quad (\# \alpha)^\dagger = (\dagger \alpha)^\#.$$

Note that, by definition, we have

$$\#(f^*) = (f^\#)^* \quad \text{and} \quad (\#f)^* = (f^*)^\#$$

for any 1-morphism f , so these conditions make sense. Note also that, by definition, any 2-morphism α satisfies

$$(\alpha^\dagger)^* = \dagger(\alpha^*) \quad \text{and} \quad (\dagger \alpha)^* = (\alpha^*)^\dagger,$$

where

$$\dagger \alpha = (i_{f^*} \circ g^*) \cdot (f^* \circ \alpha \circ g^*) \cdot (f^* \circ e_{g^*}).$$

This, together with conditions 1 and 12 in the definition of duality (def. 2.3), implies the equalities

$$\dagger \alpha = \dagger(\alpha^{**}) = ((\alpha^*)^\dagger)^* = (\alpha^\dagger)^{**} = \alpha^\dagger.$$

Of course this implies $\alpha^{\dagger\dagger} = \alpha$ in its turn.

The next thing to define is the notion of a *trace-functor* in a semi-strict pivotal 2-category. Let C be such a 2-category for the rest of this section. For any object A in C its endomorphisms and the 2-morphisms between them form the category $\text{End}(A)$ with the composition defined by the vertical composition of the 2-morphisms. The difference with $\text{End}(I)$ is that $\text{End}(A)$ does not have a monoidal structure in general.

Definition 2.5. For any object A in C there is a left trace functor $\text{Tr}_L: \text{End}(A) \rightarrow \text{End}(I)$. For any object $f \in \text{End}(A)$ define $\text{Tr}_L(f)$ by

$$I \xrightarrow{i_{A^*}} A^* \otimes A \xrightarrow{A \otimes f} A^* \otimes A \xrightarrow{e_A} I.$$

For a morphism $\alpha: f \Rightarrow g$ in $\text{End}(A)$ define $\text{Tr}_L(\alpha)$ by

$$1_{i_{A^*}} \circ (A^* \otimes \alpha) \circ 1_{e_A}.$$

Analogously there is a right trace functor Tr_R defined by

$$I \xrightarrow{i_A} A \otimes A^* \xrightarrow{f \otimes A} A \otimes A^* \xrightarrow{e_{A^*}} I,$$

and

$$1_{i_A} \circ (\alpha \otimes A^*) \circ 1_{e_{A^*}}.$$

The following lemma is the analogue of the well known lemma that says that traces are conjugation invariant in pivotal categories.

Lemma 2.6. *For any two objects A, B in C and any two 1-morphisms $f: A \rightarrow B$ and $g: B \rightarrow A$ we have a unitary 2-morphism $\theta(f, g): \text{Tr}_R(fg) \Rightarrow \text{Tr}_R(gf)$. For any 1-morphisms $f': A \rightarrow B$ and $g': B \rightarrow A$ and any 2-morphisms $\alpha: f \rightarrow f'$ and $\beta: g \rightarrow g'$ we have*

$$\theta(f, g) \cdot \text{Tr}_R(\beta \circ \alpha) = \text{Tr}_R(\alpha \circ \beta) \cdot \theta(f', g').$$

The proof is identical to the proof in the case of pivotal categories except that the essential identities are now 2-isomorphisms. I leave the details to the reader. Of course there is an analogous lemma for the left trace functor. Now if we take $g = f^*$ then we can write

$$\text{Tr}_R(ff^*) = i_A(f \otimes A^*)(f^* \otimes A^*)i_A^* = i_A(f \otimes A^*)(i_A(f \otimes A^*))^*,$$

so we have the 2-morphism

$$i_{i_A(f \otimes A^*)}: 1_I \Rightarrow \text{Tr}_R(ff^*).$$

Analogously we have the 2-morphisms

$$i_{i_A(f \otimes A^*)}^*: \text{Tr}_R(ff^*) \Rightarrow 1_I.$$

We will call these 2-morphisms *cap* and *cup* respectively. These names are of course inspired by what these 2-morphisms stand for in the 2Tang. For our purposes we want these cups and caps to be compatible with the pivotal condition.

Definition 2.7. *A semi-strict pivotal 2-category is called consistent if the following conditions are satisfied*

- 1) $i_{i_A(f \otimes A^*)} \cdot \theta(f, f^*) = i_{i_B(f^* \otimes B^*)}$
- 2) $\theta(f, f^*) \cdot i_{i_B(f^* \otimes B^*)}^* = i_{i_A(f \otimes A^*)}^*$

and the two analogous conditions with respect to $\text{Tr}_L(ff^*)$ and $\text{Tr}_L(f^*f)$.

In the sequel pivotal 2-categories will always be assumed to be consistent.

Definition 2.8. *A semi-strict spherical 2-category is a semi-strict consistent pivotal 2-category C such that for any object A in C and any 1-morphism $f \in \text{End}(A)$ we have a unitary 2-morphism $\sigma_f: \text{Tr}_L(f) \Rightarrow \text{Tr}_R(f)$. For any 1-morphism $g \in \text{End}(A)$ and any 2-morphism $\alpha: f \Rightarrow g$ these 2-isomorphisms are required to satisfy*

$$\sigma_f \cdot \text{Tr}_R(\alpha) = \text{Tr}_L(\alpha) \cdot \sigma_g.$$

Furthermore do we require the following identities to be satisfied:

$$i_{i_A^*} \cdot \sigma_{1_A} = i_{i_A} \quad \text{and} \quad \sigma_{1_A} \cdot i_{i_A^*}^* = i_{i_A^*}^*.$$

The last two conditions just mean that “cupping” and “capping” is compatible with the spherical condition.

In a semi-strict spherical 2-category C we can define a symmetric pairing

$$\langle \cdot, \cdot \rangle: 2\text{Hom}(f, g) \otimes 2\text{Hom}(g, f) \rightarrow 2\text{End}(1_I).$$

Definition 2.9. For any 1-morphisms $f, g: A \rightarrow B$ and any objects A, B in C . For any $\alpha \in 2\text{Hom}(f, g)$ and any $\beta \in 2\text{Hom}(g, f)$ we define a pairing by

$$\langle \alpha, \beta \rangle = i_{i_A(f \otimes A^*)} \cdot \text{Tr}_R((\alpha \cdot \beta) \circ 1_{f^*}) \cdot i_{i_A(f \otimes A^*)}^*.$$

Notice that the analogous definition of a pairing, using the left trace functor, gives exactly the same pairing because C is assumed to be spherical. Another way to understand this pairing is by noting that for any 2-morphism $\alpha: f \Rightarrow f$, with $f: A \rightarrow B$ an arbitrary 1-morphism, there is “a kind of trace” defined by

$$1_A \xrightarrow{i_f} f f^* \xrightarrow{\alpha \circ f^*} f f^* \xrightarrow{i_f^*} 1_A.$$

In order to get a real trace we have to map this to $2\text{End}(1_I)$. The right way to do this is by first applying the trace-functor to the “traced” 2-morphism above and then close this up by adding cups and caps. This leads precisely to our definition of the pairing. The following lemmas show that the pivotal condition implies that one gets the same trace if one uses

$$1_B \xrightarrow{i_{f^*}} f^* f \xrightarrow{f^* \circ \alpha} f^* f \xrightarrow{i_{f^*}^*} 1_B.$$

Lemma 2.10. $\langle \alpha, \beta \rangle = \langle \beta, \alpha \rangle$ for any $\alpha \in 2\text{Hom}(f, g)$ and any $\beta \in 2\text{Hom}(g, f)$.

Proof. The proof is identical to the proof in the case of pivotal categories. Just as in that case it is easy to show that

$$\begin{aligned} \langle \alpha, \beta \rangle &= i_{i_A(f \otimes A^*)} \cdot \text{Tr}_R(\alpha \circ \dagger \beta) \cdot i_{i_A(g \otimes A^*)}^* \\ &= i_{i_A(f \otimes A^*)} \cdot \text{Tr}_R(\alpha \circ \beta^\dagger) \cdot i_{i_A(g \otimes A^*)}^* = \langle \beta, \alpha \rangle. \end{aligned}$$

□

Lemma 2.11. $\langle \alpha, \beta \rangle^* = \langle \alpha^*, \beta^* \rangle$ for any $\alpha \in 2\text{Hom}(f, g)$ and any $\beta \in 2\text{Hom}(g, f)$.

Proof. From the definition of the pairing we see

$$\langle \alpha, \beta \rangle^* = \langle \beta^*, \alpha^* \rangle.$$

Now use the previous lemma. □

Lemma 2.12. $\langle \alpha, \beta \rangle = \langle \alpha^\dagger, \beta^\dagger \rangle$ for any $\alpha \in 2\text{Hom}(f, g)$ and any $\beta \in 2\text{Hom}(g, f)$.

Proof. From the pivotal condition we get

$$\langle \alpha, \beta \rangle = i_{i_A(f \otimes A^*)} \cdot \text{Tr}_R(1_f \circ (\beta^\dagger \cdot \alpha^\dagger)) \cdot i_{i_A(f \otimes A^*)}^*.$$

Now use $\text{Tr}_R(f f^*) \cong \text{Tr}_R(f^* f)$ and compatibility with cupping and capping. We get

$$i_{i_B(f^* \otimes B^*)} \cdot \text{Tr}_R((\beta^\dagger \cdot \alpha^\dagger) \circ 1_f) \cdot i_{i_B(f^* \otimes B^*)}^* = \langle \beta^\dagger, \alpha^\dagger \rangle.$$

Finally use lemma 2.10 again. \square

Lemma 2.13. $\langle \alpha^\#, \beta^\# \rangle = \langle \# \alpha, \# \beta \rangle = \langle \alpha, \beta \rangle$ for any $\alpha \in 2\text{Hom}(f, g)$ and any $\beta \in 2\text{Hom}(g, f)$.

Proof. If $\alpha = \beta = 1_{1_A}$, then this follows immediately from condition 13 in definition 2.3 and the spherical condition. The general case now follows from this particular case by using the pivotal condition. \square

The following lemma shows that our setup is really a generalization of the setup of Crane, Kauffman and Yetter in [16].

Lemma 2.14. *Let C be a semi-strict spherical 2-category, then $\text{End}(I)$ is a braided spherical category.*

Proof. We already know that $\text{End}(I)$ is a braided monoidal category (see lemma 2.2). Duality of course follows from the duality on C . The dual object of $f \in \text{End}(I)$ is $f^* \in \text{End}(I)$. The evaluation on f is defined by i_f and the coevaluation by e_f and the identities they should satisfy are exactly those of condition 11 in Definition 2.3. The dual morphism of $\alpha \in 2\text{Hom}(f, g)$ is $\alpha^\dagger \in 2\text{Hom}(g^*, f^*)$.

$\text{End}(I)$ is pivotal since we have imposed the condition $\alpha^\dagger = {}^\dagger \alpha$, which is equivalent to condition 12 in definition 2.3 as we explained. $\text{End}(I)$ is spherical because the left trace of $\alpha \in 2\text{End}(f)$ in $\text{End}(I)$ is equal to $\langle \alpha, 1_{f^*} \rangle$ and the right trace is equal to $\langle 1_{f^*}, \alpha \rangle$ and these two are equal as shown in lemma 2.10. \square

Before we go on let us have a look at an example of a spherical 2-category. In [3] Baez defines the 2-category of 2-Hilbert spaces, 2Hilb , and shows that it is a monoidal 2-category, although not semi-strict, with duals. Let us recall his definitions and some of his results and then indicate why the pivotal and the spherical conditions are satisfied in 2Hilb .

Example 2.15. The objects in 2Hilb , which are called 2-Hilbert spaces, are all finite dimensional abelian H^* -categories. An abelian H^* -category H is an abelian category such that the Hom-spaces are Hilbert spaces and composition is bilinear and additionally H is equipped with antilinear maps $*$: $\text{hom}(x, y) \rightarrow \text{hom}(y, x)$ for all objects x, y in H such that

1. $f^{**} = f$,
2. $(fg)^* = g^* f^*$,
3. $\langle fg, h \rangle = \langle g, f^* h \rangle$,
4. $\langle fg, h \rangle = \langle f, hg^* \rangle$

for all $f: x \rightarrow y$, $g: y \rightarrow z$ and $h: x \rightarrow z$. It is shown in [3] that any abelian H^* -category H is semi-simple as an abelian category and if it is finitely semi-simple then its dimension is defined as the number of objects in a basis of H . It is also shown that any basis of a 2-Hilbert space has the same number of objects, so its dimension is well defined. Furthermore does Baez show that two 2-Hilbert spaces are equivalent if and only if they have the same dimension.

A 1-morphism $F: H \rightarrow H'$ in 2Hilb is an exact functor such that $F: \text{hom}(x, y) \rightarrow \text{hom}(F(x), F(y))$ is linear and $F(f^*) = F(f)^*$ for all $f \in \text{hom}(x, y)$.

A 2-morphism $\alpha: F \Rightarrow F'$ in 2Hilb is a natural transformation.

This all looks a little abstract, but, since we can always choose bases in 2-Hilbert spaces, 1-morphisms correspond to matrices with integer coefficients. This correspondence is reliable because any 2-Hilbert space is unitarily equivalent to a skeletal 2-Hilbert space, which is one where isomorphic objects are equal. So given a basis $\{a_i\}$ in a 2-Hilbert space A and a basis $\{b_i\}$ in a 2-Hilbert space B any 1-morphism $F: A \rightarrow B$ can be presented by the matrix with coefficients $F_{ij} \in \mathbf{N}$, where

$$F(a_i) = F_{ij}b_j.$$

2-morphisms now correspond to matrices of matrices. If $(F_{ij})_{ij}$ presents the 1-morphism $F: A \rightarrow B$ and $(G_{ij})_{ij}$ the 1-morphism $G: A \rightarrow B$, then we can write a 2-morphism $\alpha: F \rightarrow G$ as the matrix $(\alpha_{ij})_{ij}$, where α_{ij} is a $G_{ij} \times F_{ij}$ matrix with complex coefficients.

The tensor product is a little bit complicated in 2Hilb and I will not define it here in a basis invariant way. As a matter of fact Baez does not work out what kind of a monoidal 2-category 2Hilb becomes, but he assures the reader that it should be possible “to strictify 2Hilb, obtaining a semistrict braided monoidal 2-category”. Roughly speaking the tensor product of two 2-Hilbert spaces A and B can be obtained in the obvious way: choose a basis $\{a_i\}$ in A and a basis $\{b_i\}$ in B and “define” $A \otimes B$ as the 2-Hilbert space with basis $\{a_i \otimes b_j\}$ and define $\text{hom}(a_i \otimes b_j, a_i \otimes b_j) = \text{hom}(a_i, a_i) \otimes \text{hom}(b_j, b_j)$. It is obvious that Hilb, the category of finite dimensional Hilbert spaces, is the identity object. The braiding comes from the ordinary transposition of factors in the tensor product and I will not say more about it because it is not important for our purpose.

Let us now have a look at the duality in 2Hilb. The dual of a 2-Hilbert space H is the 2-Hilbert space $H^* = \text{Hom}(H, \text{Hilb})$. There is always a unique dual basis in H^* for each basis in H up to isomorphism of course. The coevaluation i_H and evaluation e_H are now defined as usual. Let us assume that H is skeletal. Given a basis $\{h_i\}$ in H and its dual basis $\{h^i\}$ in H^* we define $i_H: \text{Hilb} \rightarrow H \otimes H^*$ by

$$\mathbf{C} \mapsto \bigoplus_i h_i \otimes h^i$$

and $e_H: H^* \otimes H \rightarrow \text{Hilb}$ by

$$h^i \otimes h_j \mapsto \delta_j^i \mathbf{C},$$

where δ_j^i is the Kronecker-delta. If H is skeletal, then T_H , the triangulator, is trivial. Since, as already mentioned, any 2-Hilbert space is unitarily equivalent to a skeletal one, this defines the triangulator in general.

The dual of a 1-morphism $F: A \rightarrow B$ presented by the matrix $(F_{ij})_{ij}$ is defined by the transpose of this matrix, i.e.

$$F^*(b_j) = \bigoplus_i F_{ij}a_i.$$

Baez shows that this really defines a left and right adjoint to F , so the 2-coevaluation i_F and the 2-evaluation e_F are easy to define. The Hilbert space $\text{hom}(F(a_i), F(a_i))$ is isomorphic to $\text{hom}(a_i, FF^*(a_i))$ for every i , so $i_F: 1_A \Rightarrow FF^*$ is simply defined as the natural transformation corresponding to the identity on $F(a_i)$ for each basis element a_i

under this isomorphism. In the same way we obtain the 2-evaluation $e_F: F^*F \Rightarrow 1_B$ by the isomorphism $\text{hom}(F^*(b_i), F^*(b_i)) \cong \text{hom}(F^*F(b_i), b_i)$. Note that here we continue to use the convention under which FF^* means first F and then F^* and not the other way around as is the more usual convention for functors and natural transformation but not for 2-categories. In order to understand what i_H is in more concrete terms one should note that the matrix corresponding to FF^* has diagonal coefficients that are sums of squares of coefficients of F and that 1_A is just a diagonal matrix with all diagonal coefficients equal to 1. So i_H is defined by the sums of the coevaluation maps on the terms in the diagonal coefficients and by the zero map for all non-diagonal coefficients of FF^* . In the same way we see that e_H is defined by the “ordinary” evaluation maps on the terms of the diagonal coefficients of F^*F , which are squares also of course, and by the zero map on all the non-diagonal coefficients.

The dual of a 2-morphism $\alpha: F \Rightarrow G$ presented by the matrix $(\alpha_{ij})_{ij}$ is the 2-morphism $\alpha^*: G \Rightarrow F$ presented by the matrix $(\alpha_{ij}^*)_{ij}$, where α_{ij}^* is the adjoint of α_{ij} , obtained by taking the transpose and then the complex conjugate of each coefficient. A little thinking shows that α^\dagger is presented by the adjoint of the whole matrix $(\alpha_{ij})_{ij}$. It is now obvious that $\alpha^{\dagger*} = \alpha^{*\dagger}$ corresponds to the matrix $(\alpha_{ij}^{\dagger*})_{ij}$ where $\alpha_{ij}^{\dagger*} = \alpha_{ji}$.

It is easy to check that with these definitions 2Hilb becomes a monoidal 2-category with duals, and hopefully Baez is right and we will be able to strictify the monoidal structure so that it really becomes a semi-strict monoidal 2-category with duals. It is also easy to check that 2Hilb is a spherical 2-category. As a matter of fact it turns out that $F^\#$ also corresponds to the transpose of $(F_{ij})_{ij}$, i.e.

$$F^\#(b^j) = \bigoplus_i F_{ij} a^i.$$

The same is true for ${}^\#F$, so for a 1-morphism F between skeletal 2-Hilbert spaces we see that $F^\# = {}^\#F$, which suffices to conclude that 2Hilb is pivotal. Of course $\alpha^\#$ corresponds to the matrix $(\alpha_{ij}^\#)_{ij}$ where $\alpha_{ij}^\# = \alpha_{ji}$, like for $\alpha^{\dagger*}$. Note that this does not mean that $\alpha^\#$ is equal to $\alpha^{\dagger*}$, because $\alpha^\#$ is a 2-morphism from $F^\#$ to $G^\#$ and $\alpha^{\dagger*}$ is a 2-morphism from F^* to G^* .

The left trace-functor applied to FF^* gives the same result as when it is applied to F^*F , since this is the ordinary trace of the matrices corresponding to FF^* and F^*F respectively. So compatibility with cupping and capping is guaranteed. As a matter of fact it is easy to see directly what the cup and cap are for $\text{Tr}_L(FF^*) = \text{Tr}_L(F^*F)$; The 2-morphism 1_I is just equal to 1, corresponding to the identity on \mathbf{C} , and $\text{Tr}_L(FF^*) = \text{Tr}(F^*F)$ is just the sum of the squares of the coefficients of F , so the cup $1_I \Rightarrow \text{Tr}_L(FF^*)$ is nothing but the sum of the respective coevaluations and the cap $\text{Tr}_L(FF^*) \Rightarrow 1_I$ is nothing but the sum of the respective evaluations.

The left and the right trace functor are equal when applied to 1-morphisms between skeletal 2-Hilbert spaces, again because these are just ordinary matrix traces, so the spherical condition is also satisfied.

Having defined a semi-strict spherical 2-category we now have to go a little further and add some linear structure. In order to define this linear structure we can work with a semi-strict 2-category C first.

Definition 2.16. C is called additive if $\text{Hom}(A, B)$ for any two objects A, B in C is an additive 2-vector space of finite rank, i.e. additive as a category, and the composition and the tensor product are Vect-bilinear, with Vect the ring category of finite dimensional vector spaces over the fixed field \mathbf{F} .

For the definitions of a 2-vector space and its rank and the definition of a ring category see [22]. The following lemma is straight forward.

Lemma 2.17. The category $\text{End}(I)$ is a braided monoidal ring category and $2\text{End}(1_I)$ is a commutative ring.

Each $\text{Hom}(A, B)$ becomes an $\text{End}(I)$ -module category with the action

$$\text{End}(I) \times \text{Hom}(A, B) \rightarrow \text{Hom}(A, B)$$

on objects defined by

$$f \times g \mapsto (f \otimes A)(I \otimes g).$$

The isomorphism $(f_1 f_2) \triangleright g \Rightarrow f_1 \triangleright (f_2 \triangleright g)$ is defined by $\otimes_{f_1 f_2, g}$ in the definition of a semi-strict monoidal 2-category, see condition $(\rightarrow \rightarrow \otimes)$ in [22]. Notice that

$$f \times g \mapsto (I \otimes g)(f \otimes B)$$

defines another action. This action is isomorphic to the one we have chosen by means of the isomorphisms $\otimes_{f, g}$. The action on morphisms is defined in an obvious way now. For the rest of this paper I will assume that $\text{End}(I)$ is isomorphic to Vect and that $2\text{End}(1_I)$ is a field isomorphic to \mathbf{F} . It is now easy to prove that the composition and the tensor product in C are $\text{End}(I)$ -bilinear. It is obvious that the action of $F = 2\text{End}(1_I)$ on $2\text{Hom}(f, g)$ for any $f, g: A \rightarrow B$ and for any A and B is the one induced by the action above. Notice that it makes sense to write $\text{Hom}(A, B) \oplus \text{Hom}(C, D)$ as the direct sum of two $\text{End}(I)$ -module categories and $\text{Hom}(A, B) \otimes \text{Hom}(C, D)$ as the tensor product of two $\text{End}(I)$ -module categories, see [22].

The following condition that we should impose on our 2-categories concerns the non-degeneracy of the pairing defined in 2.9. Assume that C is an additive semi-strict spherical 2-category. Notice that the pairing is bi-linear.

Definition 2.18. We call C non-degenerate if the pairing defined in 2.9 is non-degenerate.

As in the case of additive spherical categories [7,9] one can always take a non-degenerate quotient of an additive semi-strict spherical 2-category.

Lemma 2.19. Let C be as above. Let J be the additive subcategory with the same objects and 1-morphisms, but with $2\text{Hom}_J(f, g)$ being the sub-vector space of $2\text{Hom}_C(f, g)$ defined by

$$2\text{Hom}_J(f, g) = \{\alpha \in 2\text{Hom}_C(f, g) \mid \langle \alpha, \beta \rangle = 0 \quad \forall \beta \in 2\text{Hom}_C(g, f)\}.$$

Then C/J is an additive semi-strict spherical 2-category.

Proof. It is clear that the vertical composition of 2-morphisms is well defined in C/J .

Let us now prove that the horizontal composition is well defined also. Let $f, g: A \rightarrow B$ and $f', g': B \rightarrow C$ be 1-morphisms in C . For any $\alpha \in 2\text{Hom}_J(f, g)$, any $\beta \in 2\text{Hom}_C(f', g')$ and any $\gamma \in 2\text{Hom}_C(gg', ff')$ we have

$$\langle \alpha \circ \beta, \gamma \rangle = \langle \alpha, (g \circ i_{f'}) \cdot (g \circ \beta \circ (f')^*) \cdot (\gamma \circ (f')^*) \cdot (f \circ i_{f'}^*) \rangle = 0.$$

Next we show that the tensor product is well defined in C/J . If $\alpha \in 2\text{Hom}_J(f, g)$, then $\alpha \otimes X \in 2\text{Hom}_J(f \otimes X, g \otimes X)$ and $X \otimes \alpha \in 2\text{Hom}_J(X \otimes f, X \otimes g)$. The proof of these facts is not difficult, but is a bit cumbersome because we have to keep track of both the vertical and the horizontal composition. Writing out everything carefully gives

$$\begin{aligned} & \langle \alpha \otimes X, \beta \rangle \\ &= \langle \alpha, (g \circ i_{(B \otimes i_X)}) \cdot (\otimes_{g, i_X} \circ (B \otimes 1_{i_X}^*)) \\ & \cdot ((A \otimes 1_{i_X}) \circ (\beta \otimes X^*) \circ (B \otimes 1_{i_X}^*)) \cdot (\otimes_{f, i_X}^{-1} \circ (B \otimes 1_{i_X}^*)) \cdot (f \circ i_{(B \otimes i_X)}^*) \rangle = 0. \end{aligned}$$

Notice that the long formula defining the second 2-morphism in the second pairing is really a 2-morphism from g to f , so that its pairing with α makes sense. This shows our first assertion. The proof of the second is analogous and we leave the details to the reader.

Lemma 2.11 shows that $\alpha^* \in 2\text{Hom}_J(g, f)$ if and only if $\alpha \in 2\text{Hom}_J(f, g)$, lemma 2.12 shows that $\alpha^\dagger \in 2\text{Hom}_J(g^*, f^*)$ if and only if $\alpha \in 2\text{Hom}_J(f, g)$ and lemma 2.13 shows that $\alpha^\# \in 2\text{Hom}_J(f^\#, g^\#)$ if and only if $\alpha \in 2\text{Hom}_J(f, g)$.

Now take the quotients in C/J of all the structural 2-morphisms involved in the definition of the tensor product, the duality and the pivotal and the spherical condition. Then C/J becomes an additive non-degenerate semi-strict spherical 2-category. \square

Now let us define semi-simplicity for additive 2-categories. A non-zero object A in C is an object for which we have $\text{End}(A) \neq 0$.

Definition 2.20. Let C be an additive semi-strict monoidal 2-category. We say that C is finitely semi-simple if the following two conditions are satisfied:

1. For any pair of objects A, B in C the category $\text{Hom}(A, B)$ is finitely semi-simple. Here we choose a finite basis of non-isomorphic 1-morphisms $\mathcal{F}_{A, B}$.
2. There is a finite set of non-equivalent non-zero objects \mathcal{E} such that for any pair of objects A, B in C we have

$$\bigotimes_{X \in \mathcal{E}} \text{Hom}(A, X) \otimes \text{Hom}(X, B) \cong \text{Hom}(A, B).$$

The isomorphism is given by the obvious composition of 1- and 2-morphisms.

The first condition in the definition of semi-simplicity we will usually refer to as “vertical semi-simplicity” and the second condition we will refer to as “horizontal semi-simplicity”.

Definition 2.21. Let C be an additive semi-strict monoidal 2-category. An object A in C is called simple if $\text{rk}(\text{End}(A)) = 1$.

Here $\text{rk}(\text{End}(A))$ is the rank of $\text{End}(A)$ as a Vect-module, see [22].

Lemma 2.22. *Assume that C is an additive semi-strict monoidal finitely semi-simple 2-category. An object A in C is simple if and only if $A \in \mathcal{E}$.*

Proof. The proof is identical to the one that proves the analogous statement about finitely semi-simple categories. It follows from the following facts:

$$\mathrm{rk}(X \oplus Y) = \mathrm{rk}(X) + \mathrm{rk}(Y), \quad \mathrm{rk}(X \otimes Y) = \mathrm{rk}(X)\mathrm{rk}(Y),$$

for any Vect-modules X and Y . These identities can be found in [22].

Let us define the quantum dimension of objects and 1-morphisms in a finitely semi-simple non-degenerate semi-strict spherical 2-category.

Definition 2.23. *Let A be an object in C . Then we define its quantum dimension to be*

$$\dim_q(A) = \langle 1_{1_A}, 1_{1_A} \rangle.$$

Definition 2.24. *Let f be a 1-morphism in C . Then we define its quantum dimension to be*

$$\dim_q(f) = \langle 1_f, 1_f \rangle.$$

Lemma 2.25. *For any simple objects A, B, C, D, E and any 1-morphisms $f: A \rightarrow B \otimes C$, $g: B \rightarrow D \otimes E$ and $h: C \rightarrow D \otimes E$ we have*

$$\dim_q(f(g \otimes C)) = \dim_q(f)\dim_q(B)^{-1}\dim_q(g),$$

and

$$\dim_q(f(B \otimes h)) = \dim_q(f)\dim_q(C)^{-1}\dim_q(h).$$

Proof. Let us first look closely at $\dim_q(f)$. The 1-morphism $ff^*: A \rightarrow A$ is just a “multiple” of the identity, because A is simple. Multiple here means that there is a vector space $V(f)$ such that $ff^* = V(f)1_A$. So we get

$$\dim_q(f) = \dim(V(f))\dim_q(A).$$

Of course the same holds for g and h .

Thus we get

$$\dim_q(f(g \otimes C)) = \dim_q(f)\dim(V(g)) = \dim_q(f)\dim_q(B)^{-1}\dim_q(g)$$

and

$$\dim_q(f(B \otimes h)) = \dim_q(f)\dim(V(h)) = \dim_q(f)\dim_q(C)^{-1}\dim_q(h).$$

□

Note also that $\dim_q(A^*) = \dim_q(A)$ for any object A by the spherical condition, that $\dim_q(f^*) = \dim_q(f)$ for any 1-morphism f by lemma 2.12 and the fact that $1_f^\dagger = 1_{f^*}$, and that $\dim_q(f^\#) = \dim_q(\#f) = \dim_q(f)$ for any 1-morphism f by lemma 2.13. From this last observation it easily follows that $1_{A^*}^\# = \#1_{A^*} = 1_A$ and that the triangulator T_A is trivial for all simple objects. This is because $1_{A^*}^\# = \#1_{A^*} = 1_A = \mathbf{F}$ and $T_A = T_A^{\dagger*}$ is an invertible element of \mathbf{F} for any simple object A , which we will call λ . Now $\dim_q(1_{A^*}^\#) = \lambda\dim_q(A)$ but on the other hand $\dim_q(1_{A^*}^\#) = \dim_q(1_{A^*}) = \dim_q(A^*) = \dim_q(A)$, so $\lambda = 1$.

Finally we have to define the dimension of C .

Definition 2.26. Let C be a finitely semi-simple non-degenerate semi-strict spherical 2-category. Its dimension K is defined by

$$K = \sum_{\substack{A, B, C, \\ f_{A, B \otimes C}}} \dim_q(A)^{-1} \dim_q(B)^{-1} \dim_q(C)^{-1} \dim_q^2(f_{A, B \otimes C}),$$

where we sum over all the objects $A, B, C \in \mathcal{E}$ and all 1-morphisms $f_{A, B \otimes C} \in \mathcal{I}_{A, B \otimes C}$, the finite basis of non-isomorphic 1-morphisms in $\text{Hom}(A, B \otimes C)$.

3. The definitions of $Z_C(\pm(ijklm))$ and $I_C(M, T)$

In this section $M = (M, T)$ still denotes a triangulated manifold. We also assume that there is a total ordering on the vertices of the simplicial complex that defines the triangulation and a total ordering on the 4-simplices of the same complex. Let C be a finitely semi-simple semi-strict non-degenerate spherical 2-category. The linear maps $Z_C(\pm(ijklm))$, the number $Z_C(M, T, \ell)$ and the state sum $I_C(M, T)$ obviously depend on the given 2-category C , but we will suppress the subscript C at all places where this does not lead to any confusion.

We now label the edges (ij) of the triangulation with simple objects e_{ij} of C and the triangles (ijk) with simple 1-morphisms $f_{ijk} \in \text{Hom}(e_{ik}, e_{jk} \otimes e_{ij}) = H(ijk)$. We will use the following notation for the different compositions of the 1-morphisms:

$$f_{(ijk)l} = f_{ikl}(e_{kl} \otimes f_{ijk}),$$

$$f_{i(jkl)} = f_{ijl}(f_{jkl} \otimes e_{ij}),$$

$$f_{((ijk)l)m} = f_{(ikl)m}(e_{lm} \otimes e_{kl} \otimes f_{ijk}) = f_{ilm}(e_{lm} \otimes f_{ikl})(e_{lm} \otimes e_{kl} \otimes f_{ijk}),$$

$$f_{i(j(klm))} = f_{i(jkm)}(f_{klm} \otimes e_{jk} \otimes e_{ij}) = f_{ijm}(f_{jkm} \otimes e_{ij})(f_{klm} \otimes e_{jk} \otimes e_{ij}).$$

Let $+(ijklm)$ be the positively oriented standard 4-simplex labelled as described above. We are going to define the linear map

$$Z(+(ijklm)): 2\text{Hom}(f_{(ikl)m}, f_{i(klm)}) \otimes 2\text{Hom}(f_{(ijk)m}, f_{i(jkm)}) \rightarrow$$

$$2\text{Hom}(f_{(jkl)m}, f_{j(klm)}) \otimes 2\text{Hom}(f_{i(jl)m}, f_{i(jlm)}) \otimes 2\text{Hom}(f_{(ijk)l}, f_{i(jkl)})$$

using the pairing $\langle \cdot, \cdot \rangle$ described in the previous section. Consider the linear map

$$2\text{Hom}(f_{(ikl)m}, f_{i(klm)}) \otimes 2\text{Hom}(f_{(ijk)m}, f_{i(jkm)}) \rightarrow 2\text{Hom}(f_{((ijk)l)m}, f_{i(j(klm))})$$

by

$$\beta \otimes \delta \mapsto$$

$$(\beta \circ (e_{lm} \otimes e_{kl} \otimes f_{ijk})) \cdot (f_{ikm} \circ \otimes_{f_{klm}, f_{ijk}}) \cdot (\delta \circ (f_{klm} \otimes e_{jk} \otimes e_{ij})).$$

Let us write this as $\beta\delta$ as a shorthand. Consider also the linear map

$$2\text{Hom}(f_{j(klm)}, f_{(jkl)m}) \otimes 2\text{Hom}(f_{i(jlm)}, f_{i(jl)m}) \otimes 2\text{Hom}(f_{i(jkl)}, f_{(ijk)l}) \rightarrow$$

$$2\mathrm{Hom}(f_{i(jklm)}, f_{((ijk)l)m})$$

defined by

$$\alpha \otimes \gamma \otimes \epsilon \mapsto (f_{ijm} \circ (\alpha \otimes e_{ij})) \cdot (\gamma \circ (e_{lm} \otimes f_{ijk} \otimes e_{ij})) \cdot (f_{ilm} \circ (e_{lm} \otimes \epsilon)).$$

Let us write this as $\alpha\gamma\epsilon$ as a shorthand. Now define the linear map

$$2\mathrm{Hom}(f_{j(klm)}, f_{(jkl)m}) \otimes 2\mathrm{Hom}(f_{i(kl)m}, f_{i(klm)}) \otimes 2\mathrm{Hom}(f_{i(jlm)}, f_{(ijl)m}) \\ \otimes 2\mathrm{Hom}(f_{(ijk)m}, f_{i(jkm)}) \otimes 2\mathrm{Hom}(f_{i(jkl)}, f_{(ijk)l}) \rightarrow \mathbf{F}$$

defined by


$$\alpha \otimes \beta \otimes \gamma \otimes \delta \otimes \epsilon \mapsto \langle \beta\delta, \alpha\gamma\epsilon \rangle.$$

Using $2\mathrm{Hom}(g, f) = 2\mathrm{Hom}(f, g)^*$, following from the non-degeneracy of $\langle \cdot, \cdot \rangle$, this gives us $Z(+ (ijklm))$.

Before we go on, let us have a look at a kind of diagrammatic picture of $Z(+ (01234))$. In picture 1 we have depicted a diagram that resembles a $15j$ symbol as defined in [16]. The dumbbells are the 2-morphisms $\alpha, \beta, \gamma, \delta, \epsilon$. The triples (ijk) stand for the 1-morphisms f_{ijk} . Finally the crossing is just $\otimes_{f_{234}, f_{012}}$. In this diagram we do not see the objects involved and so we do not see the effect of the trace functor and the final caps and cups either. But we do see the “kind of trace” mentioned after the definition of the pairing. Remember, if $f, g: A \rightarrow B$ are 1-morphisms and $\zeta: f \Rightarrow g$ and $\xi: g \Rightarrow f$ are 2-morphisms, then one obtains $\langle \zeta, \xi \rangle$ by first taking a “kind of trace”, namely

$$1_A \xRightarrow{i_f} f f^* \xRightarrow{\zeta \circ f^*} g f^* \xRightarrow{\xi \circ f^*} f f^* \xRightarrow{i_f^*} 1_A.$$

This is exactly the diagram we see when we take $\zeta = \beta\delta$ and $\xi = \alpha\gamma\epsilon$, the respective 2-morphisms involved in the definition of $Z(+ (01234))$. The properties satisfied by the duality of the 1-morphisms, condition 11 in definition 2.3, and the properties of the composition of \otimes, \cdot , analogous to the properties of the braiding in a braided category, allow us to apply isotopic transformations to this diagram corresponding to Reidemeister moves 0, 2 and 3. Since our 2-category C is also assumed to be spherical, we can also exchange closing strands on the left hand side for closing strands to the right hand side of the diagram. The pivotal condition alone justifies these manipulations of the diagrams, but the spherical condition is needed for the different ways of nesting the cups and the caps that can not be seen in the diagrams. These are the ones we put on our pairing after applying the trace functor. This nice behaviour under different ways of nesting turns out to be necessary for our purposes, as can be seen in the proof of lemma 4.4. There this kind of diagrams allow us to prove invariance of our invariant under any permutation of the vertices of the chosen triangulation. The formulas, which I had written out completely by hand at first, are far too extensive to fit on ordinary sheets of paper. So, although these diagrams do not define a complete diagrammatic calculus for my invariants, since they do not show the objects,



Picture 1

they are certainly very helpful to see what is going on. Better diagrams would be very welcome, but so far I have not been able to invent them.

Let us now define

$$Z(-(ijklm)): 2\text{Hom}(f_{(jkl)m}, f_{j(klm)}) \otimes 2\text{Hom}(f_{(ijl)m}, f_{i(jlm)}) \otimes 2\text{Hom}(f_{(ijk)l}, f_{i(jkl)}) \rightarrow \\ 2\text{Hom}(f_{(ikl)m}, f_{i(klm)}) \otimes 2\text{Hom}(f_{(ijk)m}, f_{i(jkm)}).$$

Consider the linear map

$$2\text{Hom}(f_{(jkl)m}, f_{j(klm)}) \otimes 2\text{Hom}(f_{(ijl)m}, f_{i(jlm)}) \otimes 2\text{Hom}(f_{(ijk)l}, f_{i(jkl)}) \rightarrow \\ 2\text{Hom}(f_{((ijk)l)m}, f_{i(j(klm))})$$

defined by

$$\alpha \otimes \gamma \otimes \epsilon \mapsto \\ (f_{ilm} \circ (e_{lm} \otimes \epsilon)) \cdot (\gamma \circ (e_{lm} \otimes f_{jkl} \otimes e_{ij})) \cdot (f_{ijm} \circ (\alpha \otimes e_{ij})).$$

Call this $\epsilon\gamma\alpha$. Consider also the linear map

$$2\text{Hom}(f_{i(klm)}, f_{(ikl)m}) \otimes 2\text{Hom}(f_{i(jkm)}, f_{(ijk)m}) \rightarrow 2\text{Hom}(f_{i(j(klm))}, f_{((ijk)l)m})$$

by

$$\beta \otimes \delta \mapsto \\ (\delta \circ (f_{klm} \otimes e_{jk} \otimes e_{ij})) \cdot (f_{024} \circ \otimes_{f_{klm}, f_{ijk}}^{-1}) \cdot (\beta \circ (e_{lm} \otimes e_{kl} \otimes f_{ijk})).$$

Call this $\delta\beta$. Now define the linear map

$$2\text{Hom}(f_{(jkl)m}, f_{j(klm)}) \otimes 2\text{Hom}(f_{i(klm)}, f_{(ikl)m}) \otimes 2\text{Hom}(f_{(ijl)m}, f_{i(jlm)}) \\ \otimes 2\text{Hom}(f_{i(jkm)}, f_{(ijk)m}) \otimes 2\text{Hom}(f_{(ijk)l}, f_{i(jkl)}) \rightarrow \mathbf{F}$$

defined by

$$\alpha \otimes \beta \otimes \gamma \otimes \delta \otimes \epsilon \mapsto \\ < \delta\beta, \epsilon\gamma\alpha > .$$

Using $2\text{Hom}(g, f) = 2\text{Hom}(f, g)^*$ again this gives us $Z(-(ijklm))$.

In the sequel let us write $2H(ijkl)$ for $2\text{Hom}(f_{(ijk)l}, f_{i(jkl)})$. Likewise let us write $2H(ijklm)$ for $2\text{Hom}(f_{((ijk)l)m}, f_{i(j(klm))})$. The next lemma is the analogue of the Crossing Lemma 5.4 in [9].

Lemma 3.1(crossing). *The following diagram is commutative:*

$$\begin{array}{ccc}
 \bigoplus_{f_{024}} 2H(0234) \otimes 2H(0124) & \xrightarrow{\Phi_{01234}} & \bigoplus_{\substack{e_{13}, f_{013}, \\ f_{123}, f_{134}}} 2H(1234) \otimes 2H(0134) \otimes 2H(0123) \\
 \downarrow & & \downarrow \\
 2H(01234) & = & 2H(01234)
 \end{array}$$

Here the vertical linear maps are the isomorphisms defined by the composition of the respective 2-morphisms and Φ_{01234} is defined by

$$\Phi_{01234} = \bigoplus_{\substack{e_{13}, f_{024}, f_{013}, \\ f_{123}, f_{134}}} Z(+ (01234)) \dim_q(e_{13})^{-1} \dim_q(f_{013}) \dim_q(f_{123}) \dim_q(f_{134}).$$

Furthermore the inverse of Φ_{01234} is given by

$$\Phi_{01234}^{-1} = \bigoplus_{\substack{e_{13}, f_{024}, f_{013}, \\ f_{123}, f_{134}}} Z(- (01234)) \dim_q(f_{024}).$$

Proof. First of all let us explain why the vertical maps are isomorphisms. Using the same notation as above we must show that for any 2-morphism $\alpha \in 2\text{Hom}(f_{((012)3)4}, f_{0(1(234))})$ there exist 2-morphisms $\beta_{024} \in 2\text{Hom}(f_{(023)4}, f_{0(234)})$ and $\gamma_{024} \in 2\text{Hom}(f_{(012)4}, f_{0(124)})$ such that

$$\sum_{f_{024}} (\beta_{024} \circ (e_{34} \otimes e_{23} \otimes f_{012})) \cdot (f_{024} \circ \otimes_{f_{234}, f_{012}}) \cdot (\gamma_{024} \circ (f_{234} \otimes e_{12} \otimes e_{01})) = \alpha.$$

By (vertical) semi-simplicity we get the following (vertical) decomposition of α :

$$\begin{array}{ccc}
 f_{034}(e_{34} \otimes f_{023})(e_{34} \otimes e_{23} \otimes f_{012}) & & f_{034}(e_{34} \otimes f_{023})(e_{34} \otimes e_{23} \otimes f_{012}) \\
 & & \Downarrow \\
 & & \sum_{f_{024}} f_{024}(f_{234} \otimes e_{02})(e_{34} \otimes e_{23} \otimes f_{012}) \\
 \Downarrow & = & \Downarrow \otimes_{f_{234}, f_{012}}^{-1} \\
 & & \sum_{f_{024}} f_{024}(e_{24} \otimes f_{012})(f_{234} \otimes e_{12} \otimes e_{01}) \\
 & & \Downarrow \\
 f_{014}(f_{124} \otimes e_{01})(f_{234} \otimes e_{12} \otimes e_{01}) & & f_{014}(f_{124} \otimes e_{01})(f_{234} \otimes e_{12} \otimes e_{01})
 \end{array}$$

Notice that the 1-morphisms $f_{024}(f_{234} \otimes e_{02})(e_{34} \otimes e_{23} \otimes f_{012})$ do not form a basis of simple 1-morphisms of $2\text{Hom}(e_{04}, e_{34} \otimes e_{23} \otimes e_{12} \otimes e_{01})$ in general, but they are generators by

(horizontal) semi-simplicity and the fact that the f_{024} , the f_{234} and the f_{012} form bases of their respective Hom-spaces. So by summing the projections on the simple components of each $f_{024}(f_{234} \otimes e_{02})(e_{34} \otimes e_{23} \otimes f_{012})$ we get the decomposition in the diagram. By (horizontal) semi-simplicity we can now decompose the 2-morphisms on the right side in the diagram above and we get the β_{024} and γ_{024} we were looking for. This proves that the left vertical map in the lemma is an isomorphism. In an analogous way one proves that the other vertical map in the lemma is an isomorphism.

The commutativity of the diagram in the lemma now follows by taking arbitrary elements in $H(0234)$ and in $H(0124)$ respectively and pairing it with an arbitrary element in $H(01234)^*$. In order to get the multiplicative factors in the definition of Φ we have to use the identity

$$\dim_q(f_{ikl}(e_{kl} \otimes f_{ijk})) = \dim_q(f_{ikl})\dim_q(e_{ik})^{-1}\dim_q(f_{ikl}).$$

This identity follows from lemma 2.25.

The inverse of Φ can be computed by reading the diagram the other way around and using similar arguments. \square

We now define our state sum. Let C be a finitely semi-simple non-degenerate semi-strict spherical 2-category with non-zero dimension.

Definition 3.2. *With the data and notations as above we define for every closed compact piece-wise linear 4-manifold M with triangulation T the state sum*

$$I_C(M, T) = K^{-v} \sum_{\ell} Z_C(M, T, \ell) \prod_e \dim_q(\ell(e))^{-1} \prod_f \dim_q(\ell(f)).$$

Here v is the number of vertices in T . We sum over all the labellings ℓ and take the products over all the edges e and all the faces f in the triangulation.

4. $Z_C(M, T, \ell)$ is a combinatorial invariant

In this section we will show that $Z(M, T, \ell) = Z_C(M, T, \ell)$ is equal for all “isomorphic” labellings and $Z(M, T, \ell) = Z(M', T', \ell')$ for any pair of triangulated manifolds M and M' with triangulations T and T' that are isomorphic under a combinatorial isomorphism and any pair of “compatible” labellings ℓ and ℓ' .

First of all we have to show that $Z(M, T, \ell)$ does not depend on the choice of representatives $f_{ijk} \in \mathcal{F}_{ijk}$ nor on the choice of representatives $e_{ij} \in \mathcal{E}_{ij}$.

Lemma 4.1. *Let ℓ and ℓ' be labellings with the same choice of objects \mathcal{E}_{ij} but different choice of 1-morphisms in \mathcal{F}_{ijk} . Then we have $Z(M, T, \ell) = Z(M, T, \ell')$.*

Proof. Let us denote the labels in ℓ by f_{ijk} and the labels in ℓ' by f'_{ijk} . They are representatives of the same isomorphism classes so for any triple ijk there is a 2-isomorphism $\phi_{ijk}: f_{ijk} \Rightarrow f'_{ijk}$. These 2-isomorphisms induce an isomorphism of vector spaces

$$2\text{Hom}(f_{(ijk)l}, f_{i(jkl)}) \cong 2\text{Hom}(f'_{(ijk)l}, f'_{i(jkl)})$$

given by

$$\alpha \mapsto \phi_{(ijk)l}^{-1} \alpha \phi_{i(jkl)}.$$

Here $\phi_{(ijk)l}$ stands for $\phi_{ikl}(e_{kl} \otimes f_{ijk})$ and $\phi_{i(jkl)}$ for $\phi_{ijl}(\phi_{jkl} \otimes e_{ij})$. Likewise there is an isomorphism between $2\text{Hom}(f_{((ijk)l)m}, f_{i(jklm)})$ and $2\text{Hom}(f'_{((ijk)l)m}, f'_{i(jklm)})$.

Now without writing out the explicit formulas, which is not very difficult but extremely tedious, one can see immediately the result of the lemma. The crossing lemma implies that the following diagram is commutative.

$$\begin{array}{ccc} \bigoplus_{f_{ikm}} 2H(iklm) \otimes 2H(ijkm) & \xrightarrow{Z(ijklm)} & \bigoplus_{\substack{e_{jl}, f_{ijl}, \\ f'_{jkl}, f'_{jlm}}} 2H(jklm) \otimes 2H(ijlm) \otimes 2H(ijkl) \\ \downarrow & & \downarrow \\ \bigoplus_{f'_{ikm}} 2H'(iklm) \otimes 2H'(ijkm) & \xrightarrow{Z'(ijklm)} & \bigoplus_{\substack{e'_{jl}, f'_{ijl}, \\ f'_{jkl}, f'_{jlm}}} 2H'(jklm) \otimes 2H'(ijlm) \otimes 2H'(ijkl) \end{array}$$

So $Z'(ijklm)$ is a conjugate of $Z(ijklm)$. Taking the respective traces over the tensor product of all the partition functions now shows that $Z(M, T, \ell) = Z(M, T, \ell')$. \square

Now suppose we take a different choice of simple objects in C . In other words, suppose we have two labellings ℓ and ℓ' such that $\ell((ij)) = e_{ij}$ is equivalent to $\ell'((ij)) = e'_{ij}$ for every edge. Let us assume that there is an isomorphism $\phi_{ij}: e_{ij} \rightarrow e'_{ij}$ for every i, j actually. Equivalent simple objects are always isomorphic, so this assumption is justified. These isomorphisms induce a linear isomorphism

$$\phi_{ijk}: \text{Hom}(e_{ik}, e_{jk} \otimes e_{ij}) \rightarrow \text{Hom}(e'_{ik}, e'_{jk} \otimes e'_{ij})$$

defined by

$$f_{ijk} \mapsto \phi_{ik}^{-1} f_{ijk} (\phi_{jk} \otimes \phi_{ij})$$

for any ijk . Denote these 1-morphisms by f'_{ijk} . It is clear that the simplicity of f_{ijk} implies the simplicity of f'_{ijk} for any ijk . Since we have proved that $Z(M, T, \ell)$ is independent of the choice of simple 1-morphisms in lemma 4.1 we may assume that $\ell'((ijk)) = f'_{ijk}$.

Lemma 4.2. *Let ℓ and ℓ' be two labellings as described above. Then $Z(M, T, \ell) = Z(M, T, \ell')$.*

Proof. The identities

$$\begin{aligned} f'_{ikl}(e'_{kl} \otimes f'_{ijk}) &= \phi_{il}^{-1} f_{ikl}(\phi_{kl} \otimes \phi_{ik})(\phi_{kl}^{-1} \otimes \phi_{ik}^{-1}(e_{kl} \otimes f_{ijk}))(\phi_{kl} \otimes \phi_{jk} \otimes \phi_{ij}) \\ &= \phi_{il}^{-1} f_{ikl}(e_{kl} \otimes f_{ijk})(\phi_{kl} \otimes \phi_{jk} \otimes \phi_{ij}) \end{aligned}$$

and

$$\begin{aligned} f'_{ijl}(f'_{jkl} \otimes e'_{ij}) &= \phi_{il}^{-1} f_{ijl}(\phi_{jl} \otimes \phi_{ij})(\phi_{jl}^{-1} \otimes \phi_{ij}^{-1}(f_{jkl} \otimes e_{ij}))(\phi_{kl} \otimes \phi_{jk} \otimes \phi_{ij}) \\ &= \phi_{ij}^{-1} f_{ijl}(f_{jkl} \otimes e_{ij})(\phi_{kl} \otimes \phi_{jk} \otimes \phi_{ij}) \end{aligned}$$

show the existence of a linear isomorphism

$$2H(ijkl) = 2\text{Hom}(f_{(ijk)l}, f_{i(jkl)}) \cong 2\text{Hom}(f'_{(ijk)l}, f'_{i(jkl)}) = 2H'(ijkl).$$

Again by applying the crossing lemma we get the commutative diagram of the previous lemma, although the vertical arrows now represent different linear maps. So again $Z(ijklm)$ and $Z'(ijklm)$ are conjugates, which implies that $Z(M, T, \ell)$ and $Z(M, T, \ell')$ are equal. \square



Finally let us prove that $Z(M, T, \ell)$ does not depend on the ordering of the vertices in the triangulation T of M .

Definition 4.3. *Let $\phi: T \rightarrow T'$ be a combinatorial isomorphism of two triangulations of M . Let the edge (ij) of T be labelled with a simple object e_{ij} for all pairs of different vertices i and j of T and the edge (ij) of T' with a simple object e'_{ij} for all pairs of different vertices i and j of T' . Let a triangle (ijk) of T be labelled with a simple 1-morphism f_{ijk} for all triples of different vertices i, j, k of T and a triangle (ijk) of T' with a simple 1-morphism f'_{ijk} for all triples of different vertices i, j, k of T' . We say that ϕ is compatible with the labellings if the following conditions are satisfied:*

- 1) $e'_{\phi(i)\phi(j)} = e_{ij}$ if ϕ preserves the orientation of the edge and $e'_{\phi(i)\phi(j)} = e_{ij}^*$ if it reverses the orientation.
- 2) If ϕ decomposes into the transposition $(ijk) \mapsto (ikj)$ and a simplicial isomorphism, then $f'_{\phi(i)\phi(j)\phi(k)}$ is unitarily isomorphic to $(i_{e_{kj}} \otimes e_{ij})(e_{kj} \otimes f_{ijk}^*)$. If the transposition in the decomposition of ϕ is $(ijk) \mapsto (kji)$, then $f'_{\phi(i)\phi(j)\phi(k)}$ is unitarily isomorphic to $(f_{ijk}^\#)^* = \#(f_{ijk}^*)$.


If we represent f_{ijk} and f_{ijk}^* by the diagrams in picture 2, then we can represent $(i_{e_{kj}} \otimes e_{ij})(e_{kj} \otimes f_{ijk}^*)$ as in picture 3 and $(f_{ijk}^\#)^*$ as in picture 4. Note that any combinatorial

isomorphism ϕ , when restricted to a triangle (ijk) in T always decomposes into a permutation of (ijk) and a simplicial isomorphism. Since S_3 is generated by the two transpositions in condition 2, we have really defined compatibility for any combinatorial isomorphism. Note also that in condition 2 we have only required the isomorphy of the corresponding 1-morphisms. The reason for this is that we want the composition of a combinatorial isomorphism compatible with the labellings ϕ with its inverse ϕ^{-1} to be compatible with the labellings. This would not be the case if we required the corresponding 1-morphisms to be equal because $\#(f_{ijk}^\#)$ is only isomorphic to f_{ijk} . For our purpose these isomorphisms do not matter, because we have already shown that $Z(M, T, \ell)$ is independent of the choice of representative simple 1-morphisms in each isomorphism class. So, given a combinatorial isomorphism ϕ and a labelling of T , there is a unique compatible labelling only up to isomorphism.


 Picture 2. a) f_{ijk} b) f_{ijk}^*

 Picture 3. $(i_{e_{kj}} \otimes e_{ij})(e_{kj} \otimes f_{ijk}^*)$

Since $Z(M, T, \ell)$ does not depend on the chosen ordering on the 4-simplices themselves, as we have explained already at the end of section 1, we can restrict our attention to what happens when we permute the vertices of a 4-simplex. The symmetric group on 5 elements S_5 is generated by the transpositions interchanging i and 5 for $1 \leq i \leq 4$, so we will restrict ourselves to showing that $Z(M, T, \ell)$ is invariant under the transposition $\sigma: (01234) \mapsto (01432)$. The cases of the other transpositions are similar. Remember the definition of the boundary of $-(01432)$

$$-\partial(01432) = -(1432) + (0432) - (0132) + (0142) - (0143).$$


 Picture 4. $(f_{ijk}^\#)^*$

We have to show that σ induces linear isomorphisms $\sigma_{0432}: 2H(0234) \rightarrow 2H(0432)^*$ and $\sigma_{0142}: 2H(0124) \rightarrow 2H(0142)$ and linear isomorphisms $\sigma'_{1432}: 2H(1234)^* \rightarrow 2H(1432)$, $\sigma'_{0143}: 2H(0134)^* \rightarrow 2H(0143)$ and $\sigma'_{0132}: 2H(0123)^* \rightarrow 2H(0132)$ such that the following diagram

$$\begin{array}{ccc}
 2H(+ (01234)) & \xrightarrow{\sigma_{01432}} & 2H(- (01432)) \\
 \downarrow & & \downarrow \\
 \mathbf{F} & = & \mathbf{F}
 \end{array} \tag{4.4}$$

commutes. In this diagram $2H(+ (01234))$ stands for

$$2H(1234)^* \otimes 2H(0234) \otimes 2H(0134)^* \otimes 2H(0124) \otimes 2H(0123)^*$$

and $2H(- (01234))$ for

$$2H(1432) \otimes 2H(0432)^* \otimes 2H(0132) \otimes 2H(0142)^* \otimes 2H(0143).$$

The linear map σ_{01432} is the tensor product of the $\sigma_{\sigma(i)\sigma(j)\sigma(k)\sigma(l)}$ and the $\sigma'_{\sigma(i)\sigma(j)\sigma(k)\sigma(l)}$ composed with some flips $P: x \otimes y \mapsto y \otimes x$ so that the respective factors in the tensor product appear in the right order. The vertical linear maps are the ones defining $Z(+ (01234))$ and $Z(- (01432))$ respectively by means of the pairing (see section 3).


First of all we will show what the $\sigma_{\sigma(i)\sigma(j)\sigma(k)\sigma(l)}$ and the $\sigma'_{\sigma(i)\sigma(j)\sigma(k)\sigma(l)}$ are. Using the diagrammatic conventions as established by the diagrams in the pictures 2, 3 and 4, we are now going to give the definition of σ'_{1432} . Remember that $2H(1234)^*$ is the vector space of 2-morphisms

$$\alpha: f_{124}(f_{234} \otimes e_{12}) \mapsto f_{134}(e_{34} \otimes f_{123}).$$


Diagrammatically we denote such a 2-morphism by the diagram in picture 5.

The vector space $2H(1432)$ was defined by

$$2\text{Hom}(f_{132}(e_{32} \otimes f_{143}), f_{142}(f_{432} \otimes e_{14})).$$



Picture 5.


 Picture 6. $f_{142}(f_{234} \otimes e_{14})$

In picture 6 we show how to get a 1-morphism

$$e_{12} \mapsto e_{42} \otimes e_{14} \mapsto e_{32} \otimes e_{43} \otimes e_{14}$$

from $f_{124}(f_{234} \otimes e_{12})$.

Of course the cups and caps represent coevaluations i . and evaluations e . respectively, defined by the duality on the objects in C . The way we have drawn our diagrams already shows that we assume that the 1-morphism we obtain in this way is equal to $f_{142}(f_{432} \otimes e_{14})$. This is justified by two arguments. In the first place we can decompose the 1-morphism into simple 1-morphisms in $\text{Hom}(e_{12}, e_{42} \otimes e_{14})$ and $\text{Hom}(e_{42}, e_{32} \otimes e_{43})$ by semi-simplicity. Since we have shown in lemma 4.1 that $Z(M, T, \ell)$ does not depend on the particular choice of simple 1-morphisms in the various isomorphism classes, we may assume that the decomposition gives f_{142} and f_{432} actually.


In the same way we obtain a 1-morphism

$$e_{12} \mapsto e_{32} \otimes e_{13} \mapsto e_{32} \otimes e_{43} \otimes e_{14}$$

from $f_{134}(e_{34} \otimes f_{123})$, which we take to be equal to $f_{132}(e_{32} \otimes f_{143})$ (see picture 7).

Now by horizontal composition of the identity 2-morphisms on the respective coevaluations and evaluations and α^\dagger we get the image of α under σ'_{1432} . Explicitly we get the 2-morphism

$$(i_{e_{32}} \otimes e_{12})(e_{32} \otimes i_{e_{43}} \otimes e_{23} \otimes e_{12})(e_{32} \otimes e_{43} \otimes f_{234}^* \otimes e_{12})$$


 Picture 7. $f_{132}(e_{32} \otimes f_{143})$

$$\begin{aligned}
 & (e_{32} \otimes e_{43} \otimes e_{24} \otimes i_{e_{42}} \otimes e_{12})(e_{32} \otimes e_{43} \otimes i_{e_{24}}^* \otimes e_{24} \otimes e_{12})(e_{32} \otimes e_{43} \otimes f_{124}^*) \\
 & \xRightarrow{T_{e_{42}}} (i_{e_{32}} \otimes e_{12})(e_{32} \otimes i_{e_{43}} \otimes e_{23} \otimes e_{12})(e_{32} \otimes e_{43} \otimes f_{234}^* \otimes e_{12})(e_{32} \otimes e_{43} \otimes f_{124}^*) \\
 & \xRightarrow{\alpha^\dagger} (i_{e_{32}} \otimes e_{12})(e_{32} \otimes i_{e_{43}} \otimes e_{23} \otimes e_{12})(e_{32} \otimes e_{43} \otimes e_{34} \otimes f_{123}^*)(e_{32} \otimes e_{43} \otimes f_{134}^*).
 \end{aligned}$$

Note that by taking the dual of the 1-morphisms in the diagrams we have to use α^\dagger , which corresponds to the fact that σ is an odd permutation. Actually the first 2-isomorphism is an equality, because the triangulator of a simple object is trivial (see remarks after lemma 2.25). We have left the 2-isomorphism just to indicate what happens precisely. It is clear that σ'_{1432} is a linear isomorphism, although it is only an involution up to a 2-isomorphism. But, in the end when we take the pairing that defines our partition function $Z(ijklm)$, these isomorphisms do not harm us because the pairing is defined by means of a trace that is invariant under conjugation, so we always get the same result anyhow. The linear isomorphism σ_{0432} is defined in an analogous way. As a matter of fact one only has to reverse the morphisms in the definition of σ'_{1432} and substitute 1 by 0. Let us now define the linear isomorphism $\sigma'_{0143}: 2H(0134)^* \rightarrow 2H(0143)$. Remember that the vector space $2H(0134)^*$ was defined by

$$2\text{Hom}(f_{014}(f_{134} \otimes e_{01}), f_{034}(e_{34} \otimes f_{013}))$$

and $2H(0143)$ by

$$2\text{Hom}(f_{043}(e_{43} \otimes f_{014}), f_{013}(f_{143} \otimes e_{01})).$$

Picture 8 shows how we map $f_{014}(f_{134} \otimes e_{01})$ to $(f_{143} \otimes e_{01})(e_{43} \otimes f_{014}^*)$.



Picture 9 shows how we map $f_{034}(e_{34} \otimes f_{013})$ to $f_{013}^* f_{043}$.

So given $\epsilon \in 2H(0134)^*$ we get a 2-morphism

$$f_{013}^* f_{043} \Rightarrow (f_{143} \otimes e_{01})(e_{43} \otimes f_{014}^*).$$

Again we have composed the identities on the coevaluations and evaluations with ϵ^\dagger . Now the image of ϵ under σ'_{0143} is defined by the obvious composition

$$f_{043}(e_{43} \otimes f_{014}) \Rightarrow f_{013} f_{013}^* f_{043}(e_{43} \otimes f_{014})$$


 Picture 8. $(f_{143} \otimes e_{01})(e_{43} \otimes f_{014}^*)$

 Picture 9. $f_{013}^* f_{043}$

$$\Rightarrow f_{013}(f_{143} \otimes e_{01})(e_{43} \otimes f_{014}^*)(e_{43} \otimes f_{014}) \Rightarrow f_{013}(f_{143} \otimes e_{01}).$$

Explicitly we get

$$\begin{aligned} f_{043}(e_{43} \otimes f_{014}) &= (i_{e_{43}} \otimes e_{03})(e_{43} \otimes f_{034}^*)(e_{43} \otimes f_{014}) \\ &\xRightarrow{i_{f_{013}}} f_{013} f_{013}^* (i_{e_{43}} \otimes e_{03})(e_{43} \otimes f_{034}^*)(e_{43} \otimes f_{014}) \\ &\xRightarrow{\otimes_{i_{43}, f_{013}}^{-1}} f_{013}(i_{43} \otimes e_{13} \otimes e_{01})(e_{43} \otimes e_{34} \otimes f_{013}^*) \\ &\quad (e_{43} \otimes f_{034}^*)(e_{43} \otimes f_{014}) \\ &\xRightarrow{\epsilon^\dagger} f_{013}(i_{43} \otimes e_{13} \otimes e_{01})(e_{43} \otimes f_{134}^* \otimes e_{01}) \\ &\quad (e_{43} \otimes f_{014}^*)(e_{43} \otimes f_{014}) \\ &\xRightarrow{e_{f_{014}}} f_{013}(i_{43} \otimes e_{13} \otimes e_{01})(e_{43} \otimes f_{134}^* \otimes e_{01}) \\ &= f_{013}(f_{143} \otimes e_{01}). \end{aligned}$$

Again this is an isomorphism, because C is pivotal. Again σ'_{0143} is only involutive up to a 2-isomorphism. The other linear isomorphisms σ_{0142} and σ'_{0132} are defined analogously. So now all linear maps in diagram 4.4 are defined. Let us now prove the commutativity of the diagram.

Lemma 4.4. *With the definitions as above diagram 4.4 is commutative.*


Proof. As I already announced, the proof is essentially diagrammatic. I actually worked out all the formulas by hand, but these are far too extensive to fit on ordinary A4 sheets. Since I have given all the explicit isomorphisms and identifications used in the horizontal linear isomorphism of our diagram, the reader can work out the explicit formulas from them and check that my diagrammatics are correct in that way.

When one works out the image of an element $\alpha \otimes \beta \otimes \gamma \otimes \delta \otimes \epsilon \in 2H(01234)$ under σ_{01432} , as described above, and the right vertical linear map in our diagram explicitly, one can read off the diagram in picture 10.


We have explained the diagrammatics after our definition of $Z(+ (ijklm))$ in section 3 and we have explained what kind of moves we can apply to them. It is now immediately clear that the diagram in picture 10 can be transformed into the diagram in picture 11. We have already explained how these transformations work, but let us explain it again in this particular case. For the transformation of the diagram in picture 10 to the diagram in picture 11 one only has to use the rules for duality on 1-morphisms (condition 11 in def. 2.3), which resemble the same properties for duality on objects in a 1-category, and the evident properties of $\otimes, .$ which precisely resemble the properties of a braiding in a 1-category. So far we do not need the condition that C is spherical. But in order to transform the diagram in picture 11 into the one in picture 1 one definitely needs it. “Swinging” around the closing strands of our diagram requires both the pivotal condition and the spherical condition. In order to get the strands labelled by f_{023}^* and f_{034}^* to the other side of the diagram one needs the pivotal condition. The spherical condition is needed because the way in which the caps and cups are nested differs in the last two diagrams. This change of nesting can only be obtained by changing the left trace functor into the right trace functor on some levels of the diagram. All these operations of course need to be compatible with the cupping and capping, but we have included these compatibilities in the axioms of the pivotal and the spherical conditions. The proof now finishes with the observation that the last diagram is exactly the one describing the $Z(01234)$, i.e. the left arrow in the diagram of this lemma. \square

The following theorem is now an immediate result of the previous lemmas in this section.

Theorem 4.5. *Let $\Phi: (M, T, \ell) \rightarrow (M', T', \ell')$ be a combinatorial isomorphism of labelled triangulated manifolds that is compatible with the labellings. Then $Z(M, T, \ell) = Z(M', T', \ell')$.*



Picture 10.



Picture 11.

5. Invariance under the Pachner moves


For the proof of invariance under the Pachner moves we should really look at the 4D Pachner moves as equalities between series of 3D Pachner moves. This idea has been worked out in [10] and can be seen in the figures 12, 13 and 14. In these figures an arrow should be interpreted as the boundary of the 4-simplex representing a 3D Pachner move. The source diagram contains the simplicial 3-complex defining one side of the Pachner move and the target diagram contains the simplicial 3-complex defining the other side. For example the arrow labelled by (01235) in the first picture represents the $2 \rightleftharpoons 3$ 3D Pachner move that inserts the edge (13) in the 3-complex given by the two tetrahedra (0125) and (0235) glued over the triangle (025).

It shows that the algebraic categorification of going from a certain kind of categories up to a certain kind of 2-categories, as first predicted and sketched in [15], goes hand in hand with a geometrical kind of categorification. From a very abstract point of view we have substituted the identities in the categories which are equivalent to the 3D Pachner moves by isomorphisms in the 2-categories which we will prove to satisfy identities equivalent to the 4D Pachner moves. The Φ in the crossing lemma 3.1 is the isomorphism that substitutes the identity which is equivalent to the $2 \rightleftharpoons 3$ move of the 3D Pachner moves. Its inverse is of course the substitute of the inverse move. The isomorphism substituting the $1 \rightleftharpoons 4$ move is not so easy to describe but will come out of our calculations below. However vague these remarks may seem, they describe the deeper reason of why everything works as nicely as it does. The notion of categorification is really central in this whole setup and causes the proofs of invariance under the 4D Pachner moves to become almost tautological.


Let T_1 and T_2 be two triangulations of M that can be obtained from one another by one 4D Pachner move. Let $D_1 \subseteq T_1$ be the simplicial 4-complex on one side of the Pachner move and $D_2 \subseteq T_2$ the simplicial 4-complex on the other. We denote the complement of the interior of D_1 in T_1 by X , which by definition is equal to the complement of the interior of D_2 in T_2 . Notice that $\partial X = \partial D_1 = \partial D_2$. Also $D_1 \cup D_2$ is the boundary of a 5-simplex (012345). Now any labelling of $X \cup \partial(012345)$ defines a labelling ℓ_X on X , a labelling ℓ_1 on D_1 and a labelling ℓ_2 on D_2 which are equal on intersections. We define $Z(X)$ as the linear map obtained by taking the partial trace over all the state spaces of all the tetrahedra in the interior of X of the tensor product of the partition functions for each labelled 4-simplex in X . Define $Z(D_1)$ and $Z(D_2)$ analogously. Then we can decompose our state sum $I(M, T_i) = I_C(M, T_i)$ for $i = 1, 2$ in the following way:

$$I(M, T_i) = K^{-v_X} \text{tr} \left(\sum_{l_X} Z(X) \left(K^{-v'_i} \sum_{\ell'_i} Z(D'_i) \prod_{e_{D'_i}} \dim_q(\ell(e_{D'_i}))^{-1} \prod_{f_{D'_i}} \dim_q(\ell(f_{D'_i})) \right) \right) \\ \prod_{e_X} \dim_q(\ell(e_X))^{-1} \prod_{f_X} \dim_q(\ell(f_X)).$$


The number v_X is the number of vertices of X , v'_i is the number of vertices internal to D_i for $i = 1, 2$. The first summation is over all labellings on X , the second is over all labellings on D_i fixed on ∂D_i but ranging over all the simple objects in \mathcal{E} and all the simple 1-morphisms in all the \mathcal{F}_{ijk} for all the edges and faces internal to D_i . The trace is



Picture 12. Pachner move $3 \rightleftharpoons 3$



Picture 13. Pachner move $2 \rightleftharpoons 4$



Picture 14. Pachner move $1 \rightleftharpoons 5$

of course the trace over all the state spaces of all the tetrahedra in ∂X . Therefore proving invariance of our state sum under a 4D Pachner move means showing that the following identity holds

$$\begin{aligned} & K^{-v'_1} \sum_{\ell'_1} Z(D'_1) \prod_{e_{D'_1}} \dim_q(\ell(e_{D'_1}))^{-1} \prod_{f_{D'_1}} \dim_q(\ell(f_{D'_1})) \\ &= K^{-v'_2} \sum_{\ell'_2} Z(D'_2) \prod_{e_{D'_2}} \dim_q(\ell(e_{D'_2}))^{-1} \prod_{f_{D'_2}} \dim_q(\ell(f_{D'_2})). \end{aligned}$$

The lemmas in the rest of this section prove this identity for all the 4D Pachner moves.

The equation proving invariance under the $3 \rightleftharpoons 3$ move is the analogue of the Biedernharn-Elliot equation.

Lemma 5.1($3 \rightleftharpoons 3$).

$$\begin{aligned} & \sum_{f_{135}} \dim_q(f_{135})(1 \otimes Z(+ (01235)))(P \otimes 1 \otimes 1)(1 \otimes Z(+ (01345)) \otimes 1) \\ & (P \otimes 1 \otimes 1 \otimes 1)(Z(+ (12345)) \otimes 1 \otimes 1 \otimes 1)(1 \otimes 1 \otimes P \otimes 1 \otimes 1) \\ &= \sum_{f_{024}} \dim_q(f_{024})(Z(+ (02345))(1 \otimes 1 \otimes P)(1 \otimes Z(+ (01245)) \otimes 1) \\ & (1 \otimes 1 \otimes 1 \otimes P)(1 \otimes 1 \otimes 1 \otimes Z(+ (01234)))). \end{aligned}$$

Proof. Just write down the hexagon of which the left hand side is

$$\begin{aligned} & \bigoplus_{f_{035}, f_{025}} 2H(0345) \otimes 2H(0235) \otimes 2H(0125) \\ & \quad \downarrow 1 \otimes \Phi_{01235} \\ & \bigoplus_{\substack{e_{13}, f_{035}, f_{135}, \\ f_{123}, f_{013}}} 2H(0345) \otimes 2H(1235) \otimes 2H(0135) \otimes 2H(0123) \\ & \quad \downarrow (P \otimes 1 \otimes 1)(1 \otimes \Phi_{01345} \otimes 1) \\ & \bigoplus_{\substack{e_{13}, e_{14}, f_{135}, f_{123}, \\ f_{013}, f_{145}, f_{134}, f_{014}}} 2H(1235) \otimes 2H(1345) \otimes 2H(0145) \otimes 2H(0134) \otimes 2H(0123) \\ & \quad \downarrow (P \otimes 1 \otimes 1 \otimes 1)(\Phi_{12345} \otimes 1 \otimes 1 \otimes 1)(1 \otimes 1 \otimes P \otimes 1 \otimes 1) \\ & \bigoplus_{\substack{e_{13}, e_{14}, e_{24}, \\ f_{245}, f_{234}, f_{145}, f_{124}, \\ f_{014}, f_{134}, f_{123}, f_{013}}} 2H(2345) \otimes 2H(1245) \otimes 2H(0145) \otimes 2H(1234) \otimes 2H(0134) \otimes 2H(0123), \end{aligned}$$

and the right hand side

$$\begin{aligned}
 & \bigoplus_{f_{035}, f_{025}} 2H(0345) \otimes 2H(0235) \otimes 2H(0125) \\
 & \quad \downarrow \Phi_{02345} \otimes 1 \\
 & \bigoplus_{\substack{e_{24}, f_{025}, f_{245} \\ f_{024}, f_{234}}} 2H(2345) \otimes 2H(0245) \otimes 2H(0234) \otimes 2H(0125) \\
 & \quad \downarrow (1 \otimes 1 \otimes P)(1 \otimes \Phi_{01245} \otimes 1) \\
 & \bigoplus_{\substack{e_{24}, e_{14}, f_{245}, f_{024}, \\ f_{234}, f_{145}, f_{124}, f_{014}}} 2H(2345) \otimes 2H(1245) \otimes 2H(0145) \otimes 2H(0124) \otimes 2H(0234) \\
 & \quad \downarrow (1 \otimes 1 \otimes 1 \otimes P)(1 \otimes 1 \otimes 1 \otimes \Phi_{01234}) \\
 & \bigoplus_{\substack{e_{13}, e_{14}, e_{24}, \\ f_{245}, f_{234}, f_{145}, f_{124}, \\ f_{014}, f_{134}, f_{123}, f_{013}}} 2H(2345) \otimes 2H(1245) \otimes 2H(0145) \otimes 2H(1234) \otimes 2H(0134) \otimes 2H(0123).
 \end{aligned}$$

Applying the crossing lemma six times, i.e. for each Φ_{ijklm} separately, shows that the hexagon is commutative. The result now follows from restriction to the components which appear in the lemma. \square

In the next lemma we prove the analogue of the orthogonality equation.

Lemma 5.2(Orthogonality).

$$\begin{aligned}
 \dim_q(f_{024}) \sum_{\substack{e_{13}, f_{013}, \\ f_{123}, f_{134}}} Z(+ (01234)) Z(- (01234)) \dim_q(e_{13})^{-1} \dim_q(f_{013}) \dim_q(f_{123}) \dim_q(f_{134}) \\
 = id_{2H(0234) \otimes 2H(0124)}.
 \end{aligned}$$

$$\begin{aligned}
 \dim_q(e_{13})^{-1} \dim_q(f_{013}) \dim_q(f_{123}) \dim_q(f_{134}) \sum_{f_{024}} Z(- (01234)) Z(+ (01234)) \dim_q(f_{024}) \\
 = id_{2H(1234) \otimes 2H(0134) \otimes 2H(0123)}.
 \end{aligned}$$

Proof. This follows from the formulas in the crossing lemma for Φ and Φ^{-1} . \square

Now the other two Pachner moves follow from the $3 \rightleftharpoons 3$ lemma and the orthogonality lemma.

Lemma 5.3($2 \Rightarrow 4$).

$$\begin{aligned}
 & \sum_{f_{135}} \dim_q(f_{135})(1 \otimes Z(-(02345)) \otimes 1)(1 \otimes 1 \otimes Z(+ (01235))(\text{ev}_{2H(0345)} \otimes 1 \otimes 1 \otimes 1)) \\
 &= \sum_{e_{14}, f_{014}, f_{124}, f_{145}, f_{134}} \dim_q(e_{14})^{-1} \dim_q(f_{014}) \dim_q(f_{124}) \dim_q(f_{145}) \dim_q(f_{134}) \\
 & \quad (1 \otimes 1 \otimes 1 \otimes P)(1 \otimes 1 \otimes Z(+ (01245)) \otimes 1)(1 \otimes 1 \otimes 1 \otimes 1 \otimes P) \\
 & \quad (1 \otimes 1 \otimes 1 \otimes 1 \otimes Z(+ (01234)))(1 \otimes 1 \otimes 1 \otimes P \otimes 1 \otimes 1)(1 \otimes Z(-(12345)) \otimes 1 \otimes 1 \otimes 1) \\
 & \quad (1 \otimes P \otimes 1 \otimes 1 \otimes 1)(1 \otimes 1 \otimes Z(-(01345)) \otimes 1)(1 \otimes P \otimes 1 \otimes 1)(\text{ev}_{2H(0345)} \otimes 1 \otimes 1 \otimes 1).
 \end{aligned}$$

Proof. Multiply each side of the $3 \Rightarrow 3$ equation with

$$\begin{aligned}
 & \dim_q(e_{24})^{-1} \dim_q(e_{14})^{-1} \dim_q(f_{234}) \dim_q(f_{245}) \dim_q(f_{014}) \dim_q(f_{124}) \\
 & \quad \dim_q(f_{145}) \dim_q(f_{134})(Z(-(02345)) \otimes 1)
 \end{aligned}$$

on the left and multiply with

$$\begin{aligned}
 & (1 \otimes 1 \otimes 1 \otimes P \otimes 1 \otimes 1)(1 \otimes Z(-(12345)) \otimes 1 \otimes 1 \otimes 1)(1 \otimes P \otimes 1 \otimes 1 \otimes 1) \\
 & (1 \otimes 1 \otimes Z(-(01345)) \otimes 1)(1 \otimes P \otimes 1 \otimes 1)(\text{ev}_{2H(0345)} \otimes 1 \otimes 1 \otimes 1)
 \end{aligned}$$

on the right and sum over all the edges, i.e. simple objects, and all the faces, i.e. simple 1-morphisms, involved. Using the orthogonality lemma once on the left hand side and twice on the right hand side you get the $2 \Rightarrow 4$ equation. \square

Lemma 5.4($1 \Rightarrow 5$).

$$\begin{aligned}
 & Z(+ (01235)) \\
 &= K^{-1} \sum_{\substack{e_{04}, e_{14}, e_{24}, e_{34}, e_{45}, \\ f_{014}, f_{024}, f_{034}, f_{045}, \\ f_{124}, f_{134}, f_{145}, f_{245}, f_{345}}} \text{tr}_1 [(Z(+ (02345)) \otimes 1)(1 \otimes 1 \otimes P)(1 \otimes Z(+ (01245)) \otimes 1) \\
 & (1 \otimes 1 \otimes 1 \otimes P)(1 \otimes 1 \otimes 1 \otimes Z(+ (01234)))(1 \otimes 1 \otimes P \otimes 1 \otimes 1)(Z(-(12345)) \otimes 1 \otimes 1 \otimes 1) \\
 & (P \otimes 1 \otimes 1 \otimes 1)(1 \otimes Z(-(01345)) \otimes 1)(P \otimes 1 \otimes 1)] \prod_{\substack{i < j \\ i=4 \vee j=4}} \dim_q(e_{ij})^{-1} \prod_{\substack{i < j < k \\ j=4 \vee k=4 \\ (ijk) \neq (024)}} \dim_q(f_{ijk}).
 \end{aligned}$$

Proof. Multiply each side of the the $3 \Rightarrow 3$ equation on the right with

$$K^{-1} \prod_{\substack{i < j \\ i=4 \vee j=4}} \dim_q(e_{ij})^{-1} \prod_{\substack{i < j < k \\ j=4 \vee k=4 \\ (ijk) \neq (024)}} \dim_q(f_{ijk})(1 \otimes 1 \otimes P \otimes 1 \otimes 1)$$

$$(Z(-(12345)) \otimes 1 \otimes 1 \otimes 1)(P \otimes 1 \otimes 1 \otimes 1)(1 \otimes Z(-(01345)) \otimes 1)(P \otimes 1 \otimes 1),$$

take the trace on the first factor and sum over all the edges and faces involved.

The right hand side is now equal to the right hand side of the $1 \Rightarrow 5$ equation. Using the orthogonality lemma the left hand side becomes

$$\frac{K^{-1}}{\dim_q(f_{035})} \sum_{\substack{e_{04}, e_{34}, e_{45}, \\ f_{034}, f_{045}, f_{345}}} \dim_q(e_{04})^{-1} \dim_q(e_{34})^{-1} \dim_q(e_{45})^{-1} \dim_q(f_{034}) \\ \dim_q(f_{045}) \dim_q(f_{345}) \text{tr}_1(1 \otimes Z(+ (01235))).$$

Now use the identity

$$\text{tr}_1(1 \otimes Z(+ (01235))) = Z(+ (01235)) \dim(2\text{Hom}(f_{045}(e_{45} \otimes f_{034}), f_{035}(f_{345} \otimes e_{03})))$$

and the identity

$$\sum_{e_{04}, f_{034}, f_{045}} \dim_q(e_{04})^{-1} \dim_q(f_{034}) \dim_q(f_{045}) \dim(2\text{Hom}(f_{045}(e_{45} \otimes f_{034}), f_{035}(f_{345} \otimes e_{03}))) \\ = \dim_q(e_{35})^{-1} \dim_q(f_{035}) \dim_q(f_{345}),$$

which will follow from lemma 5.5. Finally we get

$$\frac{K^{-1}}{\dim_q(f_{035})} \sum_{\substack{e_{34}, e_{35}, e_{45}, \\ f_{345}}} \dim_q(e_{34})^{-1} \dim_q(e_{35})^{-1} \dim_q(e_{45})^{-1} \dim_q^2(f_{345}) \\ \dim_q(f_{035}) Z(+ (01235)) = Z(+ (01235)).$$

□

Lemma 5.5. *With the same notation as everywhere in this section we have*

$$\sum_{e_{04}, f_{034}, f_{045}} \dim_q(e_{04})^{-1} \dim_q(f_{034}) \dim_q(f_{045}) \dim(2\text{Hom}(f_{045}(e_{45} \otimes f_{034}), f_{035}(f_{345} \otimes e_{03}))) \\ = \dim_q(e_{35})^{-1} \dim_q(f_{035}) \dim_q(f_{345}).$$

Proof. Consider $f_{035}(f_{345} \otimes e_{03})$. Its quantum dimension is equal to

$$\dim_q(f_{035}) \dim_q(e_{35})^{-1} \dim_q(f_{345})$$

by lemma 2.25. Now use semi-simplicity to write

$$1_{f_{035}(f_{345} \otimes e_{03})} = \sum_{\substack{e_{04} \\ f_{034}, f_{045}}} \alpha_{0345} \cdot \bar{\alpha}_{0345},$$

where

$$\alpha_{0345} \in 2\text{Hom}(f_{035}(f_{345} \otimes e_{03}), f_{045}(e_{45} \otimes f_{034}))$$

and

$$\bar{\alpha}_{0345} \in 2\text{Hom}(f_{045}(e_{45} \otimes f_{034}), f_{035}(f_{345} \otimes e_{03})).$$

We can always take the α 's to be the projections and $\bar{\alpha}$ the inclusions, so

$$\bar{\alpha}_{0345} \cdot \alpha_{0345} = 1_{f_{045}(e_{45} \otimes f_{034})}.$$

Therefore we get

$$\begin{aligned} & \dim_q(f_{035}(f_{345} \otimes e_{03})) \\ &= \sum_{\substack{e_{04} \\ f_{034}, f_{045}}} \langle \alpha_{0345}, \bar{\alpha}_{0345} \rangle \\ &= \sum_{\substack{e_{04} \\ f_{034}, f_{045}}} \langle \bar{\alpha}_{0345}, \alpha_{0345} \rangle \\ &= \sum_{\substack{e_{04}, \\ f_{034}, f_{045}}} \dim_q(f_{045}(e_{45} \otimes f_{034})) \dim(2\text{Hom}(f_{045}(e_{45} \otimes f_{034}), f_{035}(f_{345} \otimes e_{03}))) \\ &= \sum_{\substack{e_{04}, \\ f_{034}, f_{045}}} \dim_q(e_{04})^{-1} \dim_q(f_{034}) \dim_q(f_{045}) \dim(2\text{Hom}(f_{045}(e_{45} \otimes f_{034}), f_{035}(f_{345} \otimes e_{03}))). \end{aligned}$$

□

Thus we can conclude this section by the following theorem, summarizing all the results obtained in this paper.

Theorem 5.6. *Let C be a non-degenerate finitely semi-simple semi-strict spherical 2-category of non-zero dimension. For any compact closed piece-wise linear oriented 4-manifold M there exists a state sum $I_C(M)$. For any two such manifolds M and N that are pl-homeomorphic we have $I_C(M) = I_C(N)$.*

6. Concluding remarks

In this last section I want to sketch my plans for further research based on some (obvious) remarks concerning the results in this paper.

First of all let me address the question of examples. Unfortunately this paper has not broken the tradition of writing about (quantum) 4-manifold invariants without giving any specific example. My reason is the same as that of the other followers of this tradition; it is extremely hard to find interesting examples of Hopf categories or 2-categories that are likely to give any new kind of invariant. Any “simple” attempt seems to be doomed to lead to the Euler characteristic or, if you are a bit luckier, the signature of the 4-manifolds (see [16]). Crane, Yetter and I are working on the definition of monoidal 2-categories built from representations of quantum groups at $q = 0$ using their crystal bases and their deformation theory. These results will be published in a separate paper. It might well be that, in order to get the desired examples of 2-categories, we have to find actual deformations of these 2-categories and then find our way back to a generic q . Although this is the hard way of trying to find examples it certainly is worth a try.

Next there is the question of the relation of the construction presented in this paper and the one using Hopf categories [15]. First of all let me recall that in the construction of 4-manifold invariants out of Hopf categories there is this difficult question, not addressed in [15], of the right categorified notion of an antipode. As I already mentioned in my introduction, the only example of a Hopf category of which we are sure that it gives an invariant is the categorification of the quantum double of a finite group [18]. And even in that case we do not know what kind of an invariant it gives us. In order to resolve that problem and to establish a concrete relation between the construction in this paper and the one in [15] I think one should first try to reconstruct the Hopf category out of a spherical 2-category. By Neuchl’s result [27] we know that it is going to be a bitensor category. The duality used in the construction in this paper, which is the right one for the purpose of 4-manifold invariants, will hopefully lead to the reconstruction of the right notion of the antipode. When we really know what the antipode looks like, we can try to see if the invariant coming from the Hopf category is the same as the one coming from the 2-category of its representations. For involutory Hopf algebras and their category of representations it has been shown in [8] that the respective 3-manifold invariants that they induce are the same. This result was my original motivation for the “categorification” of Barrett’s and Westbury’s construction of 3-manifold invariants.

Finally I want to say something about diagrams. If we ever want to compute a real quantum invariant of 4-manifolds we probably better start looking for some diagrammatic way of doing this. Every reader will remember the major advantage that the “skein approach” brought to the computation of 3-manifold invariants. It even enabled Justin Roberts to show that the Crane-Yetter 4-manifold invariant for $U_q(sl(2))$ was “just” the signature of the 4-manifolds [28]. In this paper I have used some diagrams, but, as I already admitted, they are rather poor diagrams and certainly not good enough for computational aims. For both the construction and the computation of 3-manifold invariants it is very convenient to know that tri-valent graphs in R^2 form the free pivotal category on one object [21]. As a matter of fact Barrett and Westbury [9] do assume, and more or less use, that tri-valent graphs in S^2 form the free spherical category on one object without

proving it. The question arises what the free spherical 2-category on one object would be geometrically. If one dualizes a tetrahedron and looks at the 2-skeleton of the dual complex, one gets a four-valent vertex of a graph with 2-cells. So maybe the answer to my question is that “four-valent graphs with 2-cells” in S^3 form the free spherical 2-category on one object. Here I am deliberately being sloppy in my definition of the sort of graph I am speaking about. Probably one has to define it by something like a set of elements, the vertices, and a family of two-element subsets, the edges, and a family of three-element subsets, the 2-cells and everything in such a way that the vertices are 4-valent with respect to the edges and 6-valent with respect to the 2-cells, and the edges are 3-valent with respect to the 2-cells. But I am not trying to make a precise conjecture here. I just want to point out a possible topic for further research that can lead to a better insight in the relation between 4-dimensional topology and combinatorics and algebra. In order to get such a result we could try to study this kind of graphs, or hyper-graphs, in a way similar to that in which Carter, Rieger and Saito [12,13] have studied 2-tangles and see if they provide the diagrammatical tools for the computation of 4-manifold invariants.

Acknowledgements

First of all I want to thank Louis Crane and David Yetter for the helpful discussions with them. I also want to thank the Mathematics Department of Kansas State University and Louis Crane in particular for their hospitality during my stay in Manhattan, where I wrote this paper. Finally I want to thank the Centro de Matematica Aplicada (CMA) and the Fundação Luso-Americana para o Desenvolvimento (FLAD) in Portugal for their generous financial support that enabled me to visit KSU.

References

- [1] J. Baez, J. Dolan, *Higher-Dimensional Algebra and Topological Quantum Field Theory*, Jour. Math. Phys. **36**, 1995, 6073-6105.
- [2] J. Baez, M. Neuchl, *Higher-Dimensional Algebra I: Braided Monoidal 2-Categories*, Adv. Math. **121**, 1996, 196-244.
- [3] J. Baez, *Higher-Dimensional Algebra II: 2-Hilbert Spaces*, Adv. Math. **127**, 1997, 125-189.
- [4] J. Baez, J. Dolan, *Higher-Dimensional Algebra III: n-Categories and the Algebra of Opetopes*, to appear in Adv. Math., preprint available as q-alg/9702014 and at <http://math.ucr.edu/home/baez/>.
- [5] J. Baez, L. Langford, *Higher-Dimensional Algebra IV: 2-Tangles*, to appear in Adv. Math.
- [6] J. Baez, L. Langford, *2-Tangles*, Lett. Math. Phys., **43**, 1998, 187-197.
- [7] J.W. Barrett, B.W. Westbury, *Spherical Categories*, preprint, University of Nottingham, 1993. Available as hep-th/9310164.
- [8] J.W. Barrett, B.W. Westbury, *The Equality of 3-Manifold Invariants*, Math. Proc. Camb. Phil. Soc. **118**, 1995, 503-510.
- [9] J.W. Barrett, B.W. Westbury, *Invariants of Piecewise-Linear 3-Manifolds*, Trans. Amer. Math. Soc., Vol. 348, No. 10, 1996, 3997-4022.

- [10] J.S. Carter, L.H. Kauffman, M. Saito, *Diagrammatics, Singularities, and Their Algebraic Interpretations*, to appear in *Matemática Contemporânea*, Sociedade Brasileira de Matemática. Available at <http://www.math.usf.edu/~saito/home/html>.
- [11] J.S. Carter, L.H. Kauffman, M. Saito, Talks given by Carter and Saito at the AMS-conference at Kansas State University, Manhattan Kansas, 1998.
- [12] J.S. Carter, J.H. Rieger, M. Saito, *A Combinatorial Description of Knotted Surfaces and Their Isotopies*, *Adv. Math.*, Vol. 127, No. 1, 1997, 1-51.
- [13] J. Carter, M. Saito, *Knotted Surfaces, Braid Movies, and Beyond*, in *Knots and Quantum Gravity*, ed. J. Baez, Oxford U. Press, 1994, 191-229.
- [14] S. Chung, M. Fukuma, A. Shapere, *The Structure of Topological Field Theories in Three Dimensions*, *Int. J. Mod. Phys. A*, 1994, 1305-1360.
- [15] L. Crane, I. Frenkel, *Four Dimensional Topological Quantum Field Theory, Hopf Categories, and the Canonical Bases*, *Jour. Math. Phys.* **35**, 1994, 5136-5154.
- [16] L. Crane, L.H. Kauffman, D.N. Yetter, *State-Sum Invariants of 4-Manifolds*, *JKTR*, Vol. 6, No. 2, 1997, 177-234.
- [17] L. Crane, D.N. Yetter, *A Categorical Construction of 4D Topological Quantum Field Theories*, in *Quantum Topology*, L.H. Kauffman and R.A. Baadhio eds., World Scientific Press, 1993, 131-138.
- [18] L. Crane, D.N. Yetter, *Examples of Categorification*, to appear in *Cahiers de Top. Geom. Diff. Cat.*
- [19] L. Crane, D.N. Yetter, *Deformations of (Bi)tensor Categories*, to appear in *Cahiers de Top. Geom. Diff. Cat.*
- [20] J. Fisher, *2-Categories and 2-Knots*, *Duke Math. Jour.* **75**, 1994, 493-526.
- [21] P.J. Freyd, D.N. Yetter, *Braided Compact Closed Categories with Applications to Low Dimensional Topology*, *Adv. in Math.* **77**, 1989, 156-182.
- [22] M. Kapranov, V. Voevodsky, *2-Categories and Zamolodchikov Tetrahedra Equations*, in *Proc. Symp. Pure Math.* **56** Part 2, 1994, AMS, Providence, 177-260.
- [23] V. Karlanov, V. Turaev, *On the Definition of 2-category of 2-Knots*, LOMI preprint.
- [24] G. Kuperberg, *Involutory Hopf Algebras and 3-Manifold Invariants*, *Int. J. of Math.* **2** (1), 1991, 41-66.
- [25] L. Langford, *2-Tangles as a Free Braided Monoidal 2-category with Duals*, Ph.D. Dissertation, UCR, 1997. Available at <http://math.ucr.edu/home/baez/>.
- [26] M. Neuchl, *Representation Theory of Hopf Categories*, Ph.D. Dissertation, University of Munich, 1997.
- [27] U. Pachner, *P.L. Homeomorphic Manifolds are Equivalent by Elementary Shellings*, *European Journal of Combinatorics* (**12**), 1991, 129-145.
- [28] J. Roberts, *Skein Theory and Turaev-Viro Invariants*, *Topology* **34**, 1995, 771-787.
- [29] V.G. Turaev, O.Y. Viro, *State Sum Invariants of 3-Manifolds and Quantum 6j-Symbols*, *Topology* **31** (4), 1992, 865-902.
- [30] D.N. Yetter, *State-Sum Invariants of 3-Manifolds Associated to Artinian Semisimple Tortile Categories*, *Topology and Its Applications* **58** (1), 1994, 47-80.