### SOME CRYSTAL ROGERS-RAMANUJAN TYPE IDENTITIES

### Mirko Primc

ABSTRACT. By using the KMN<sup>2</sup> crystal base character formula for the basic  $A_2^{(1)}$ -module, and the principally specialized Weyl-Kac character formula, we obtain a Rogers-Ramanujan type combinatorial identity for colored partitions. The difference conditions between parts are given by the energy function of certain perfect  $A_2^{(1)}$ -crystal. We also recall some other identities for this type of colored partitions, but coming from the vertex operator constructions and with no apparent connection to the crystal base theory.

#### 1. Introduction

J. Lepowsky and R. Wilson gave in [LW] a Lie-theoretic interpretation of Rogers-Ramanujan identities in terms of representations of affine Lie algebra  $\tilde{\mathfrak{g}} = \mathfrak{sl}(2,\mathbb{C})^{\sim}$ . The product sides of Rogers-Ramanujan identities follow from the principally specialized Weyl-Kac character formula for level 3 standard  $\tilde{\mathfrak{g}}$ -modules, the sum sides follow from the vertex operator construction of bases of level 3 standard  $\tilde{\mathfrak{g}}$ -modules, parameterized by partitions satisfying difference 2 conditions.

The Lepowsky-Wilson approach is also possible for other affine Lie algebras and for other constructions of vertex operators, various combinatorial consequences are illustrated by constructions given, for example, in [C3], [LP], [Ma] and [Mi].

In [P1] there is a construction of the basic  $A_{\ell}^{(1)}$ -module based on the Frenkel-Kac vertex operator formula. It was noted there that the combinatorial difference conditions arising from the vertex operator formula coincide with the energy function in the construction by paths of the basic representation of  $A_{\ell-1}^{(1)}$  given in [DJKMO]. In [P2] this combinatorial connection for basic modules is extended to other classical affine Lie algebras, this time by using more general crystal base character formula for standard modules due to S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki [KMN<sup>2</sup>].

This combinatorial connection of the crystal base theory and the vertex operator constructions suggested that it might be interesting to study Rogers-Ramanujan type combinatorial identities for colored partitions, where differences are given by energy functions of perfect crystals. So we start, roughly speaking, with colored partitions  $(n_1)_{\beta_1} \geq (n_2)_{\beta_2} \geq \cdots \geq (n_s)_{\beta_s} > 0$ , where each number  $n_r$  is "colored"

<sup>1991</sup> Mathematics Subject Classification. Primary 05A19; Secondary 17B37, 17B67.

Key words and phrases. Rogers-Ramanujan identities, colored partitions, partition ideals, perfect crystals, affine Lie algebras, basic modules.

Partially supported by the Ministry of Science and Technology of the Republic of Croatia, grant 037002.

with a "color"  $\beta_r$  from the set of nine "colors"  $\{1, \ldots, 9\}$ . Analogous to the Rogers-Ramanujan case, we consider colored partitions satisfying difference conditions

$$(n_r)_{\beta_r} \geqslant (n_{r+1})_{\beta_{r+1}} + E_{\beta_r \beta_{r+1}}$$
,

where differences  $E_{\beta_r\beta_{r+1}} \in \{0, 1, 2\}$  are the values of an energy function of certain perfect  $\mathfrak{sl}(3, \mathbb{C})^{\sim}$ -crystal. We obtain an identity for such colored partitions (Theorem 2.1) by using the principally specialized Weyl-Kac character formula and the crystal base character formula [KMN<sup>2</sup>].

In the last section we recall some other identities for this type of colored partitions, but coming from the vertex operator construction [MP2] and with no apparent connection to the crystal base theory.

I am grateful to Jim Lepowsky and Arne Meurman for many ideas and results implicit in this work, and to Ivica Siladić for testing the identities numerically. I thank the ICTP in Trieste, where this work started, for their kind hospitality.

#### 2. A COMBINATORIAL IDENTITY

Let A be a nonempty set and denote by  $\mathcal{P}(A)$  the set of all maps  $\pi: A \to \mathbb{N}$ , where  $\pi(a)$  equals zero for all but finitely many  $a \in A$ . Clearly  $\pi$  is determined by its values  $(\pi(a) \mid a \in A)$  and we shall also write  $\pi$  as a monomial

$$\pi = \prod_{a \in A} a^{\pi(a)}.$$

We shall say that  $\pi$  is a partition and for  $\pi(a) > 0$  we shall say that a is a part of  $\pi$ . We define the length  $\ell(\pi)$  of  $\pi$  by  $\ell(\pi) = \sum_{a \in A} \pi(a)$ . We consider elements of A as partitions of length 1, i.e.  $A \subset \mathcal{P}(A)$ . For  $\rho, \pi \in \mathcal{P}(A)$  we write  $\pi \supset \rho$  if  $\pi(a) \geqslant \rho(a)$  for all  $a \in A$  and we say that  $\pi$  contains  $\rho$ .

For  $\rho, \pi \in \mathcal{P}(A)$  we define  $\pi \rho$  in  $\mathcal{P}(A)$  by  $(\pi \rho)(a) = \pi(a) + \rho(a)$ ,  $a \in A$ . We shall say that  $1 = \prod_{a \in A} a^0$  is the partition with no parts and length 0. Clearly  $\mathcal{P}(A)$  is a monoid. For lack of a better terminology, we shall say that  $\mathcal{I} \subset \mathcal{P}(A)$  is an ideal in the monoid  $\mathcal{P}(A)$  if  $\rho \in \mathcal{I}$  and  $\pi \in \mathcal{P}(A)$  implies  $\rho \pi \in \mathcal{I}$ . For such an  $\mathcal{I}$  we call the difference of sets  $\mathcal{P}(A) \setminus \mathcal{I}$  a partition ideal in  $\mathcal{P}(A)$  (cf. [A1, Chapter 8]). Later on we shall consider an ideal  $\mathcal{I} = \mathcal{P}(A)\mathcal{D}$  generated by a set  $\mathcal{D}$ , and the corresponding partition ideal we shall denote as

$$\mathcal{P}_{\mathcal{D}} = \mathcal{P}(A) \setminus (\mathcal{P}(A)\mathcal{D}).$$

Let  $a \mapsto |a|$  be a map from A to N. Then we define the degree  $|\pi|$  of  $\pi$  by

$$|\pi| = \sum_{a \in A} |a| \, \pi(a),$$

and we say that a part a of  $\pi$  has the degree |a|.

Now let

$$\Gamma = \{1, 2, 3, 4, 5, 6, 7, 8, 9\},\$$

we shall think of  $\Gamma$  as a set of colors, and let  $A = \Gamma \times \mathbb{Z}_{<0}$ . We shall write  $(\alpha, i) = i_{\alpha}$ ,

$$A = \{i_{\alpha} \mid \alpha \in \Gamma, i \in \mathbb{Z}_{<0}\},\$$

and we shall think that  $i_{\alpha}$  has a color  $\alpha$ . We define a map  $(-i)_{\alpha} \mapsto |(-i)_{\alpha}|$  for i > 0 by

$$(-i)_1 \mapsto 3i - 2, \qquad (-i)_4 \mapsto 3i, \qquad (-i)_9 \mapsto 3i + 2,$$

$$(2.1) \qquad (-i)_2 \mapsto 3i - 1, \qquad (-i)_5 \mapsto 3i, \qquad (-i)_8 \mapsto 3i + 1,$$

$$(-i)_3 \mapsto 3i - 1, \qquad (-i)_6 \mapsto 3i, \qquad (-i)_7 \mapsto 3i + 1.$$

So, for example,  $(-5)_1$  has the color 1 and the degree  $|(-5)_1| = 13$ . In general, we can think of  $\pi \in \mathcal{P}(A)$  as a colored partition of the nonnegative integer  $n = |\pi|$ .

Let  $E = (E_{\alpha\beta})_{\alpha,\beta\in\Gamma}$  be a matrix

Define a subset  $\mathcal{D}$  in  $\mathcal{P}(A)$  by

$$\mathcal{D} = \{i_{\alpha}i_{\beta} \mid E_{\alpha\beta}E_{\beta\alpha} \geqslant 1\} \bigcup \{(i-1)_{\alpha}i_{\beta} \mid E_{\alpha\beta} = 2\}.$$

We shall say that a colored partition  $\pi$  satisfies the difference  $\mathcal{D}$  condition if  $\pi$  does not contain any colored partition from  $\mathcal{D}$ , or, equivalently, if  $\pi$  is an element of the partition ideal  $\mathcal{P}_{\mathcal{D}} = \mathcal{P}(A) \setminus (\mathcal{P}(A)\mathcal{D})$ . So, for example,  $\pi = (-5)_1(-3)_8(-2)_9$  does not satisfy the difference  $\mathcal{D}$  condition since  $E_{89} = 2$  and  $(-3)_8(-2)_9 \in \mathcal{D}$ .

Now we can state the following Rogers-Ramanujan type identity:

# Theorem 2.1.

$$\sum_{n=0}^{\infty} \sharp \{ |\pi| = n \mid \pi \in \mathcal{P}_{\mathcal{D}} \} q^n = \prod_{r=1}^{\infty} (1 - q^r)^{-1}.$$

The product side follows from the principally specialized Weyl-Kac character formula for the basic  $\mathfrak{sl}(3,\mathbb{C})^{\sim}$ -module [L], the sum side follows from the crystal base character formula [KMN<sup>2</sup>]. The proof is given in the next section.

3. The principally specialized character for the basic  $A_2^{(1)}$ -module Let us consider a multiple of the character of the basic  $\mathfrak{sl}(3,\mathbb{C})^{\sim}$ -module  $L(\Lambda_0)$ 

(3.1) 
$$\prod_{r=1}^{\infty} (1 - e^{-r\delta})^{-1} \cdot e^{-\Lambda_0} \operatorname{ch} L(\Lambda_0).$$

Then the principal specialization  $e^{-\alpha_i} \mapsto q$ , i = 0, 1, 2, of this product gives

$$\prod_{r=1}^{\infty} (1 - q^{3r})^{-1} \prod_{r \not\equiv 0 \mod 3} (1 - q^r)^{-1} = \prod_{r=1}^{\infty} (1 - q^r)^{-1}.$$

Here we use notions, notation and results as in [K] or [L].

On the other side, we may use the results in [KMN<sup>2</sup>] to express (3.1) in terms of colored partitions. We consider a perfect crystal  $\Gamma$  for  $\mathfrak{sl}(3,\mathbb{C})^{\sim}$  coming from the tensor product of the vector representation and its dual:

For  $\beta \in \Gamma$  let us denote by  $\operatorname{wt}(\beta)$  the  $\mathfrak{h}$ -weight of  $\beta$ , that is, the restriction of the classical weight of  $\beta$  on the fixed Cartan subalgebra  $\mathfrak{h} \subset \mathfrak{g}$ . Note that

wt 
$$1 = - \text{ wt } 9 = \alpha_1 + \alpha_2$$
,  
wt  $2 = - \text{ wt } 8 = \alpha_2$ ,  
wt  $3 = - \text{ wt } 7 = \alpha_1$ ,  
wt  $4 = \text{ wt } 5 = \text{ wt } 6 = 0$ .

The crystal  $\Gamma$  has the unique energy function H with values in  $\{0,1,2\}$ , and we have chosen

$$E_{\alpha\beta}=H(\beta\otimes\alpha).$$

In particular, we have

(3.2) 
$$E_{44} = H(4 \otimes 4) = 0, \quad \text{wt } 4 = 0.$$

The ground state path for the basic module  $L(\Lambda_0)$  is

$$p_{\Lambda_0} = 4, 4, 4, 4, 4, 4, \dots$$

By [KMN<sup>2</sup>, Proposition 4.6.4], the set  $\mathcal{P}(\Lambda_0, \Gamma)$  of sequences  $p = (p(j); j \ge 1)$  in  $\Gamma$  such that  $p(j) = p_{\Lambda_0}(j) = 4$  for  $j \gg 0$  parameterizes a basis of  $L(\Lambda_0)$ , where, by taking into account (3.2), the weight and the degree of a sequence is given by

(3.3) 
$$|p| = -\sum_{j=1}^{\infty} j H(p(j+1) \otimes p(j)),$$

(3.4) 
$$\operatorname{wt}(p) = \sum_{j=1}^{\infty} \operatorname{wt}(p(j)).$$

We want to interpret this result in terms of colored partitions, and, in order to do that, let us think of colored partitions in a different way: Let  $\leq$  be an order on  $\Gamma$  defined as

$$1 \succ 2 \succ 3 \succ 4 \succ 5 \succ 6 \succ 7 \succ 8 \succ 9,$$

and define an order on A by

$$i_{\beta} \preccurlyeq j_{\gamma}$$
 if either  $i < j$  or  $i = j, \beta \preccurlyeq \gamma$ .

This is a total order on A. For a colored partition

$$\nu = (j_1)_{\beta_1} \dots (j_s)_{\beta_s} \in \mathcal{P}(A)$$

we may assume that  $j_1 \leqslant \cdots \leqslant j_s < 0$  and that  $j_r = j_{r+1}$  implies  $\beta_r \preccurlyeq \beta_{r+1}$ , i.e.,  $(j_r)_{\beta_r} \preccurlyeq (j_{r+1})_{\beta_{r+1}}$ . Sometimes we shall denote a colored partition  $\nu$  as

$$\nu = ((j_1)_{\beta_1} \preccurlyeq (j_2)_{\beta_2} \preccurlyeq \cdots \preccurlyeq (j_s)_{\beta_s}).$$

We may visualize  $\nu$  by its Young diagram:

color 
$$\beta_1$$
  $\beta_2$   $\beta_2$   $\beta_3$   $\beta_4$   $\beta_5$   $\beta_6$   $\beta_7$   $\beta_8$   $\beta_8$ 

Then the total number of boxes  $||\nu||$  in the Young diagram of  $\nu$  is

$$||\nu|| = \sum_{m=1}^{s} -j_m$$
.

Since to each color  $\beta \in \Gamma$  we can associate its weight  $\operatorname{wt}(\beta) \in \mathfrak{h}^*$ , we define a weight  $\operatorname{wt}(\nu)$  of the colored partition  $\nu$  as

$$\operatorname{wt}(\nu) = \sum_{m=1}^{s} \operatorname{wt}(\beta_m) .$$

Now for a given path

$$p = \beta_1, \dots, \beta_{s-1}, \beta_s, \dots, 4, 4, 4, 4, \dots$$

we construct a colored partition  $\operatorname{part}_{\mathcal{D}}(p)$  in the following way: We start with a large enough s, i.e., an s such that  $\beta_j = 4$  for all  $j \geqslant s$ , and we set

$$-i_s = -i_{s+1} = \cdots = 0$$
,

and from there on

$$-i_{s-1} = E_{\beta_{s-1}\beta_s}, 
-i_{s-2} = E_{\beta_{s-2}\beta_{s-1}} + E_{\beta_{s-1}\beta_s}, 
\dots 
s-1$$

$$-i_1 = \sum_{r=1}^{s-1} E_{\beta_r \beta_{r+1}}.$$

Note that  $E_{44} = 0$ , so it does not matter with which  $s \gg 0$  we have started. Now we define a colored partition  $\operatorname{part}_{\mathcal{D}}(p)$  associated to p as

$$\operatorname{part}_{\mathcal{D}}(p) = (-i_1)_{\beta_1} \dots (-i_{s-1})_{\beta_{s-1}} (-i_s)_{\beta_s} \dots (0)_4 (0)_4 (0)_4 (0)_4 \dots,$$

where we identify the product of all  $(0)_4$  with  $1 \in \mathcal{P}(A)$ . It is easy to check that

(3.5) 
$$E_{\beta_r\beta_{r+1}} = 0$$
 implies  $\beta_r \preccurlyeq \beta_{r+1}$ ,

so by construction we have

$$\operatorname{part}_{\mathcal{D}}(p) = ((-i_1)_{\beta_1} \preccurlyeq (-i_2)_{\beta_2} \preccurlyeq \dots).$$

We may visualize the above construction of  $p \mapsto \operatorname{part}_{\mathcal{D}}(p)$  in terms of Young diagrams: We start from "the bottom" for large enough s and we add 0 boxes to a color  $\beta_s = 4$ . As we are finished with associating  $-i_{r+1}$  boxes to  $\beta_{r+1}$ , we associate to  $\beta_r$  an extra  $E_{\beta_r,\beta_{r+1}}$  boxes, so that  $\operatorname{part}_{\mathcal{D}}(p)$  satisfies difference conditions

$$-i_r = -i_{r+1} + E_{\beta_r, \beta_{r+1}}.$$

By counting the number of boxes we added at each stage, we see that the total number of boxes in  $\operatorname{part}_{\mathcal{D}}(p)$  equals -|p| given by (3.3), i.e.

(3.6) 
$$||\operatorname{part}_{\mathcal{D}}(p)|| = \sum_{r=1}^{s} r E_{\beta_r, \beta_{r+1}} = -|p|.$$

We also have (cf. (3.4) and (3.2))

(3.7) 
$$\operatorname{wt}(\operatorname{part}_{\mathcal{D}}(p)) = \sum_{r=1}^{s} \beta_r = \operatorname{wt}(p).$$

For a given colored partition  $\nu$  and a plain partition  $\Delta \in \mathcal{P}(\mathbb{Z}_{<0})$  (i.e. a partition "without colors"),

$$\nu = ((i_1)_{\beta_1} \preceq \cdots \preceq (i_s)_{\beta_s}),$$
  
$$\Delta = (j_1 \leqslant j_2 \leqslant \cdots \leqslant j_p),$$

 $i_s, j_p < 0$ , we define a colored partition  $\nu \oplus \Delta$  by

$$\nu \oplus \Delta = (k_1)_{\beta_1} \dots (k_r)_{\beta_r},$$

where  $r = \max\{s, p\}$ , with additional colors  $\beta_{s+1} = \cdots = \beta_p = 4$  in the case p > s, and

$$k_n = i_n + \sum_{-j_m \geqslant n} -1,$$

that is, to a Young diagram of  $\nu$  we add to each color  $\beta_1, \ldots, \beta_{-j_1}$  one box, then we add to each color  $\beta_1, \ldots, \beta_{-j_2}$  another one box, and so on. Of course, in the

case p > s we consider  $i_{s+1} = \cdots = i_p = 0$ . It is clear that in the case  $\nu = \operatorname{part}_{\mathcal{D}}(p)$  we have difference conditions

$$-i_r \geqslant -i_{r+1} + E_{\beta_r,\beta_{r+1}}$$
.

It is clear that the map

$$(p, \Delta) \mapsto \operatorname{part}_{\mathcal{D}}(p) \oplus \Delta, \qquad \mathcal{P}(\Lambda_0, \Gamma) \times \mathcal{P}(\mathbb{Z}_{<0}) \to \mathcal{P}(A)$$

is injective. Hence, by using (3.6) and (3.7), [KMN<sup>2</sup>, Proposition 4.6.4] implies

$$\sum_{\substack{\pi = \operatorname{part}_{\mathcal{D}}(p) \oplus \Delta}} e^{\operatorname{wt}(\pi) - ||\pi||\delta} = \prod_{r=1}^{\infty} (1 - e^{-r\delta})^{-1} \cdot e^{-\Lambda_0} \operatorname{ch} L(\Lambda_0),$$

where the sum runs over all pats  $p \in \mathcal{P}(\Lambda_0, \Gamma)$  and plain partitions  $\Delta \in \mathcal{P}(\mathbb{Z}_{<0})$ .

What we want to see is that every  $\pi = \operatorname{part}_{\mathcal{D}}(p) \oplus \Delta$  is in  $\mathcal{P}_{\mathcal{D}}$ . For that it is sufficient to show that

(3.8) 
$$(i_{\alpha} \preccurlyeq j_{\beta}) \in \mathcal{P}_{\mathcal{D}}$$
 if and only if  $|i - j| \geqslant E_{\alpha\beta}$ ,

$$(3.9) \quad i_{\alpha} \preccurlyeq j_{\beta} \preccurlyeq k_{\gamma}, \ |i - k| \leqslant 1, \ i_{\alpha}k_{\gamma} \in \mathcal{D} \quad \text{implies} \quad i_{\alpha}j_{\beta} \in \mathcal{D} \quad \text{or} \quad j_{\beta}k_{\gamma} \in \mathcal{D}.$$

It is easy to see, by using (3.5), that (3.8) holds. Now, in the presence of (3.8), we can easily check that (3.9) holds as well; in terms of E it reads as

$$(3.10) E_{\alpha\gamma} \leqslant E_{\alpha\beta} + E_{\beta\gamma},$$

or, equivalently, as

$$E_{\alpha\gamma} \leqslant E_{\alpha\beta}$$
 if  $E_{\beta\gamma} = 0$ ,  $E_{\alpha\gamma} \leqslant E_{\beta\gamma}$  if  $E_{\alpha\beta} = 0$ .

Hence every  $\pi = \operatorname{part}_{\mathcal{D}}(p) \oplus \Delta$  is in  $\mathcal{P}_{\mathcal{D}}$ . Moreover, every  $\pi \in \mathcal{P}_{\mathcal{D}}$  can be written in this way, and hence we have

$$\sum_{\pi \in \mathcal{P}_{\mathcal{D}}} e^{\operatorname{wt}(\pi) - ||\pi||\delta} = \prod_{r=1}^{\infty} (1 - e^{-r\delta})^{-1} \cdot e^{-\Lambda_0} \operatorname{ch} L(\Lambda_0).$$

Since the map  $(-i)_{\alpha} \mapsto |(-i)_{\alpha}|$  defined by (2.1) is the principal specialization, Theorem 2.1 holds.

# 4. Some remarks

Let  $\Gamma$  be a classical crystal with an energy function H with values in the set  $\{0,1,2\}$ . Then, as before, we can consider colored partitions  $\pi \in \mathcal{P}(A)$  with  $A = \Gamma \times \mathbb{Z}_{<0}$  and we can define the degree  $||\pi||$  and the  $\mathfrak{h}$ -weight of  $\pi$  as above, with  $\mathfrak{h} = \mathbb{C}$ -span $\{h_1,\ldots,h_\ell\}$  (see [KMN<sup>2</sup>]). If we set  $E_{\alpha\beta} = H(\beta \otimes \alpha) \in \{0,1,2\}$ , we can define

$$\mathcal{D} = \{ i_{\alpha} i_{\beta} \mid E_{\alpha\beta} E_{\beta\alpha} \geqslant 1 \} \bigcup \{ (i-1)_{\alpha} i_{\beta} \mid E_{\alpha\beta} = 2 \},$$

and we can consider colored partitions which satisfy difference  $\mathcal{D}$  conditions, that is,  $\pi \in \mathcal{P}_{\mathcal{D}}$ . Let us define a "character" of the partition ideal  $\mathcal{P}_{\mathcal{D}}$  as

$$\operatorname{ch}(\mathcal{P}_{\mathcal{D}}) = \sum_{\pi \in \mathcal{P}_{\mathcal{D}}} e^{\operatorname{wt}(\pi) - ||\pi||\delta}.$$

The map  $e^{-\alpha_i} \mapsto q$ ,  $i = 0, 1, \dots, \ell$ , defines  $e^{-\delta} \mapsto q^m$  and the principally specialized character

$$\operatorname{ch}_{q}(\mathcal{P}_{\mathcal{D}}) = \sum_{n=0}^{\infty} \sharp \{ |\pi| = n \mid \pi \in \mathcal{P}_{\mathcal{D}} \} q^{n},$$

where  $|(-j)_{\beta}| = mj - \langle \rho, \beta \rangle$ , with  $\langle \rho, \alpha_i \rangle = 1$  for  $i = 1, \dots, \ell$  (cf. [K]).

With this notation at hand we can write Theorem 2.1, for our particular choice of the  $A_2^{(1)}$ -crystal  $\Gamma$ , as

$$\operatorname{ch}_q(\mathcal{P}_{\mathcal{D}}) = \prod_{r=1}^{\infty} (1 - q^r)^{-1}.$$

Let  $\Gamma$  be a perfect crystal for  $\tilde{\mathfrak{g}} = \mathfrak{sl}(n,\mathbb{C})^{\sim}$ ,  $n \geq 4$ , coming from the tensor product of the vector representation and its dual. Then there is an energy function H taking values in  $\{0,1,2\}$  and the ground state path for the basic  $\tilde{\mathfrak{g}}$ -module is a constant sequence p(j) = a, with  $E_{aa} = H(a \otimes a) = 0$  and  $\operatorname{wt}(a) = 0$ . Moreover, there is a total order  $\leq$  on  $\Gamma$  such that (3.5) holds. So the same proof would go through, and the above identity would hold, if we could show (3.10). For example, in the case of  $A_3^{(1)}$ -crystal

$$41 \xrightarrow{0} 11 \xrightarrow{0} 14$$
,  $21 \xrightarrow{0} 24$ ,  $31 \xrightarrow{0} 34$ ,  $42 \xrightarrow{0} 12$ ,  $43 \xrightarrow{0} 13$ 

all these properties hold, including (3.10), and we have an identity of the above form. On the other side, let  $\Gamma = \{1, 2, 3, 4\}$  be the  $A_1^{(1)}$ -crystal

$$\begin{array}{ccc}
1 & \xrightarrow{1} & 3 \\
\uparrow 0 & & \downarrow \\
2 & \longleftarrow & 4
\end{array}$$

with the energy matrix

$$E = \begin{pmatrix} 2 & 1 & 2 & 2 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 2 \\ 0 & 1 & 0 & 2 \end{pmatrix}.$$

The KMN<sup>2</sup> crystal base character formula is proved under the assumption that the rank of  $\mathfrak{g}$  is at least two, but still many results also hold for  $\tilde{\mathfrak{g}} = \mathfrak{sl}(2,\mathbb{C})^{\sim}$ . So it is reasonable to ask whether an analogue of Theorem 2.1 holds as well, i.e., whether

$$\operatorname{ch}_q(\mathcal{P}_{\mathcal{D}}) = \prod_{r=1}^{\infty} (1 - q^r)^{-1} \quad ?$$

Note that here the principal specialization reads

$$(-i)_1 \mapsto 2i - 1$$
,  $(-i)_2 \mapsto 2i$ ,  $(-i)_3 \mapsto 2i$ ,  $(-i)_4 \mapsto 2i + 1$ .

What is surprising is that there are other identities of a similar form, but which are not related to the crystal base theory, at least not in any obvious way: consider an "almost perfect"  $A_1^{(1)}$ -crystal

$$1 \stackrel{\stackrel{1}{\longleftrightarrow}}{\longleftrightarrow} 2 \stackrel{\stackrel{1}{\longleftrightarrow}}{\longleftrightarrow} 3$$

with the energy matrix

$$E = \begin{pmatrix} 2 & 2 & 2 \\ 1 & 1 & 2 \\ 0 & 1 & 2 \end{pmatrix} .$$

Then we have

(4.1) 
$$\operatorname{ch}_{q}(\mathcal{P}_{\mathcal{D}}) = \prod_{r \text{ odd}} (1 - q^{r})^{-1}.$$

Note that here the principal specialization reads

$$(-i)_1 \mapsto 2i - 1, \quad (-i)_2 \mapsto 2i, \quad (-i)_3 \mapsto 2i + 1.$$

Moreover, if we define a map  $(-i)_{\alpha} \mapsto |(-i)_{\alpha}|$  for i > 0 by

$$(-i)_1 \mapsto 3i - 2, \quad (-i)_2 \mapsto 3i, \quad (-i)_3 \mapsto 3i + 2,$$

(i.e. if we take the (1,2)-specialization of  $ch(\mathcal{P}_{\mathcal{D}})$ ), then we have a Capparelli identity (see [C1]–[C3], [A2])

(4.2) 
$$\sum_{n=0}^{\infty} \sharp \{ |\pi| = n \mid \pi \in \mathcal{P}_{\mathcal{D}} \} q^n = \prod_{r \equiv 1,3,5,6 \mod 6} (1+q^r).$$

Both (4.1) and (4.2) are proved in [MP2] as specializations of the character formula for the basic  $\mathfrak{sl}(2,\mathbb{C})^{\sim}$ -module written in the form

$$e^{-\Lambda_0} \operatorname{ch} L(\Lambda_0) = \operatorname{ch}(\mathcal{P}_{\mathcal{D}}).$$

The character formula itself is proved by using the Lepowsky-Wilson approach, the proof being quite parallel to [MP1]. The set  $\mathcal{D}$  is originally defined as the set of leading terms for the vertex operator algebra defining relations for the basic  $\mathfrak{sl}(2,\mathbb{C})^{\sim}$ -module (cf. [MP2, Section 6]).

So it seems that interesting combinatorial properties of  $\operatorname{ch}(\mathcal{P}_{\mathcal{D}})$  go beyond the crystal base character formula [KMN<sup>2</sup>, Proposition 4.6.4], at least when the relation  $E_{\alpha\gamma} \leq E_{\alpha\beta} + E_{\beta\gamma}$  holds and when there is a total order  $\leq$  on  $\Gamma$  such that  $E_{\alpha\beta} = 0$  implies  $\alpha \leq \beta$ . Further indications for this provide the results in [P2] and the examples below.

Although the one by one matrices have nothing to do with crystals, the notion of difference  $\mathcal{D}$  condition still makes sense for  $\Gamma = \{1\}$  and  $E = (E_{11}) = (2)$ ; it is simply the difference 2 condition and  $\operatorname{ch}_q(\mathcal{P}_{\mathcal{D}})$ , defined via  $|(-i)_1| = i$ , is the sum side of a Rogers-Ramanujan identity. On the other hand, for  $E = (E_{11}) = (1)$  the difference  $\mathcal{D}$  condition defines partitions in distinct parts. Of course, the second case is much simpler, and it has an equally simple analogue for  $\Gamma = \{1,2\}$  and  $E = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ : consider an "almost perfect"  $A_1^{(1)}$ -crystal

$$1 \stackrel{\stackrel{1}{\longleftrightarrow}}{\longleftrightarrow} 2$$

with the energy matrix  $E_{\alpha\beta} = H(\beta \otimes \alpha)$  chosen to be E. Then

$$(-j)_1 \mapsto 2j - \frac{1}{2}, \qquad (-j)_2 \mapsto 2j + \frac{1}{2}$$

is the principal specialization and we have an identity for partitions in half-integers

$$\operatorname{ch}_q(\mathcal{P}_{\mathcal{D}}) = \prod_{r \geqslant 1} \left( 1 + q^{r + \frac{1}{2}} \right).$$

In the case  $E = \begin{pmatrix} 2 & 2 \\ 1 & 2 \end{pmatrix}$  the principal specialization of  $\mathcal{P}_{\mathcal{D}}$  gives partitions in half-integers satisfying difference 3 condition, but I am not aware of any formula that would express  $\operatorname{ch}_q(\mathcal{P}_{\mathcal{D}})$  as an infinite product.

Finally, the formulation of Rogers-Ramanujan type identities for colored partitions in terms of energy functions for crystals was also motivated by a desire to understand better an identity for the basic  $A_2^{(1)}$ -module obtained in [MP3]. The set  $\mathcal{D}$  of difference conditions is defined as the set of leading terms of relations for the basic module, but can be defined as well in the following way: consider the weighted  $A_2$ -crystal  $\Gamma' = \Gamma \setminus \{4\}$ 

It is not possible do define 0-arrows which would turn  $\Gamma'$  into an  $A_2^{(1)}$ -crystal with an energy function. But still, we can define a "difference condition matrix"  $(E_{\alpha\beta})_{\alpha,\beta\in\Gamma'}$ 

$$E = \begin{pmatrix} 2 & 2 & 2 & 2 & 2 & 2 & 2 & 2 & 2 \\ 1 & 2 & 1 & 2 & 1 & 2 & 2 & 2 & 2 \\ 1 & 1 & 2 & 1 & 2 & 2 & 2 & 2 & 2 \\ 0 & 1 & 1 & 1 & 1 & 1 & 2 & 2 & 2 \\ 1 & 1 & 1 & 1 & 1 & 2 & 1 & 2 & 2 \\ 0 & 1 & 0 & 1 & 1 & 2 & 1 & 2 & 2 \\ 0 & 0 & 1 & 1 & 1 & 1 & 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 2 & 2 \end{pmatrix}$$

and we can define a subset  $\mathcal{D}$  in  $\mathcal{P}(A)$ ,  $A = \Gamma' \times \mathbb{Z}_{<0}$ , by

$$\mathcal{D} = \{i_{\alpha}i_{\beta} \mid E_{\alpha\beta}E_{\beta\alpha} \geqslant 1\} \bigcup \{(i-1)_{\alpha}i_{\beta} \mid E_{\alpha\beta} = 2\}$$
$$\bigcup \{(i-1)_{3}i_{5}i_{1}\} \bigcup \{(i-1)_{9}(i-1)_{5}i_{7}\}.$$

Then, as proved in [MP3], we have

$$e^{-\Lambda_0} \operatorname{ch} L(\Lambda_0) = \operatorname{ch}(\mathcal{P}_{\mathcal{D}}).$$

By taking the principal specialization (2.1) without the color 4, we obtain a combinatorial identity

$$\operatorname{ch}_q(\mathcal{P}_{\mathcal{D}}) = \prod_{r \not\equiv 0 \mod 3} (1 - q^r)^{-1}.$$

As it happens, the above "difference 2 conditions"  $\{(i-1)_{\alpha}i_{\beta} \mid E_{\alpha\beta} = 2\}$  are the same as in the case of partitions discussed in Section 2. One is tempted to think that this is more than a mere coincidence and that the difference conditions  $\{(i-1)_3 i_5 i_1\} \bigcup \{(i-1)_9 (i-1)_5 i_7\}$  are some sort of "corrections" of the fact that  $\Gamma'$  is not a perfect  $A_2^{(1)}$ -crystal. With this regard it may be interesting to note that the energy matrix E in Section 2 is invariant under the change

$$1 \leftrightarrow 1$$
,  $2 \leftrightarrow 3$ ,  $5 \leftrightarrow 6$ ,  $7 \leftrightarrow 8$ ,  $9 \leftrightarrow 9$ 

and  $4 \leftrightarrow 4$ . Here we have  $E_{51} = 0$ ,  $E_{61} = 1$  and  $E_{95} = 0$ ,  $E_{96} = 1$ . So the "interaction" between parts  $i_6 i_1 \in \mathcal{D}$  and  $i_5 i_1 \notin \mathcal{D}$  is not symmetrical, the later is "compensated" with a weaker requirement  $(i-1)_3 i_5 i_1 \in \mathcal{D}$ . Likewise  $i_9 i_6 \in \mathcal{D}$  "corresponds" to a weaker  $i_9 i_5 (i+1)_7 \in \mathcal{D}$ . It should be said that the present proof of the above identity has no connections with the crystal base theory.

# REFERENCES

- [A1] G. E. Andrews, *The theory of partitions*, Encyclopedia of math. and appl., Addison-Wesley, Amsterdam, 1976.
- [A2] G. E. Andrews, Schur's theorem, Capparelli's conjecture and the q-trinomial coefficients, in: Proc. Rademacher Centenary Conf. (1992), Contemporary Math. 167, 1994, pp. 141–154.
- [C1] S. Capparelli, On some representations of twisted affine Lie algebras and combinatorial identities, J. Algebra 154 (1993), 335–355.
- [C2] S. Capparelli, A combinatorial proof of a partition identity related to the level 3 representations of a twisted affine Lie algebra, Comm. Algebra 23 (1995), 2959–2969.

- S. Capparelli, A construction of the level 3 modules for the affine Lie algebra  $A_2^{(2)}$  and [C3]a new combinatorial identity of the Rogers-Ramanujan type, Trans. Amer. Math. Soc. **348** (1996), 481–501.
- [DJKMO] E. Date, M. Jimbo, A. Kuniba, T. Miwa and M. Okado, Path space realization of the basic representation of  $A_n^{(1)}$ , in: Infinite dimensional Lie algebras and groups, Advanced series in Math. Phys. Vol. 7 (V. G. Kac, ed.), World Scientific, Singapore, 1989, pp. 108–123.
- [K]V. G. Kac, Infinite-dimensional Lie algebras 3rd ed., Cambridge Univ. Press, Cambridge, 1990.
- [KMN<sup>2</sup>] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki, Affine crystals and vertex models, Internat. J. Modern Phys. A, Vol. 7, Suppl. 1A, Proceedings of the RIMS Research Project 1991, "Infinite Analysis", World Scientific, Singapore, 1992, pp. 449–484.
- [L]J. Lepowsky, Lectures on Kac-Moody Lie algebras, Université de Paris VI, 1978.
- J. Lepowsky and M. Primc, Structure of the standard modules for the affine Lie algebra [LP]  $A_1^{(1)}$ , Contemporary Math. **46** (1985).
- J. Lepowsky and R. L. Wilson, The structure of standard modules, I: Universal algebras [LW] and the Rogers-Ramanujan identities, Invent. Math. 77 (1984), 199-290; II: The case  $A_1^{(1)}$ , principal gradation, Invent. Math. **79** (1985), 417–442.
- M. Mandia, Structure of the level one standard modules for the affine Lie algebras [Ma]  $B_l^{(1)}$ ,  $F_4^{(1)}$  and  $G_2^{(1)}$ , Memoirs American Math. Soc. **362** (1987). A. Meurman and M. Primc, Annihilating ideals of standard modules of  $\mathfrak{sl}(2,\mathbb{C})^{\sim}$  and
- [MP1] combinatorial identities, Advances in Math. 64 (1987), 177-240.
- [MP2]A. Meurman and M. Primc, Annihilating fields of standard modules of  $\mathfrak{sl}(2,\mathbb{C})^{\sim}$  and combinatorial identities (to appear in Memoirs American Math. Soc.).
- [MP3]A. Meurman and M. Prime, A basis of the basic  $\mathfrak{sl}(3,\mathbb{C})^{\sim}$ -module (preprint).
- K. C. Misra, Level one standard modules for affine symplectic Lie algebras, Math. Ann. [Mi] **287** (1990), 287–302.
- M. Prime, Vertex operator construction of standard modules for  $A_n^{(1)}$ , Pacific J. Math. [P1] **162** (1994), 143–187.
- [P2]M. Prime, Basic representations for classical affine Lie algebras (preprint).

Univ. of Zagreb, Dept. of Mathematics, Bijenička 30, Zagreb, Croatia E-mail address: primc@cromath.math.hr