Weak Hopf Algebras I. Integral Theory and C^* -structure

GABRIELLA BÖHM¹, FLORIAN NILL², KORNÉL SZLACHÁNYI³

Abstract

We give an introduction to the theory of weak Hopf algebras proposed recently as a coassociative alternative of weak quasi-Hopf algebras. We follow an axiomatic approach keeping as close as possible to the "classical" theory of Hopf algebras. The emphasis is put on the new structure related to the presence of canonical subalgebras A^L and A^R in any weak Hopf algebra A that play the role of non-commutative numbers in many respects. A theory of integrals is developed in which we show how the algebraic properties of A, such as the Frobenius property, or semisimplicity, or innerness of the square of the antipode, are related to the existence of non-degenerate, normalized, or Haar integrals. In case of C^* -weak Hopf algebras we prove the existence of a unique Haar measure $h \in A$ and of a canonical grouplike element $g \in A$ implementing the square of the antipode and factorizing into left and right elements $g = g_L g_R^{-1}$, $g_L \in A^L$, $g_R \in A^R$. Further discussion of the C^* -case will be presented in Part II.

To appear in J. Algebra

¹ Research Institute for Particle and Nuclear Physics, Budapest, H-1525 Budapest 114, P.O.B. 49, Hungary E-mail: BGABR@rmki.kfki.hu

Supported by the Hungarian Scientific Research Fund, OTKA – T 016 233

² Institut für Theoretische Physik, FU–Berlin Arnimallee 14, D-14195 Berlin, Germany E-mail: NILL@mail.physik.fu-berlin.de

Supported by the DFG, SFB 288 "Differentialgeometrie und Quantenphysik"

³ Research Institute for Particle and Nuclear Physics, Budapest, H-1525 Budapest 114, P.O.B. 49, Hungary E-mail: SZLACH@rmki.kfki.hu

Supported by the Hungarian Scientific Research Fund, OTKA – T 020 285.

Contents

1	Introduction	2
2	The Weak Hopf Calculus2.1The axioms2.2Weak bialgebras2.3Weak Hopf algebras2.4The "trivial" representation	$ \begin{array}{c} 4 \\ 4 \\ 6 \\ 9 \\ 12 \end{array} $
3	Weak Hopf Modules and Integral Theory3.1Integrals in weak Hopf algebras3.2Weak Hopf modules3.3Restrictions on the algebraic structure3.4Non-degenerate integrals3.52-sided non-degenerate integrals3.6Haar integrals	14 14 16 21 22 26 29
4 A	4.1First consequences of the C^* -structure4.2The Haar measure and selfduality4.3The canonical grouplike elementAppendix:The Weak Hopf Algebra $B \otimes B^{op}$	 31 31 33 35 38 30
	References	- 39

1 Introduction

Weak Hopf algebras have been proposed recently [2, 20, 14] as a new generalization of ordinary Hopf algebras that replaces Ocneanu's paragroup [16], in the depth 2 case, with a concrete "Hopf algebraic" object. The earlier proposals of *face algebras* [8] or *quantum groupoids* [17] are actually weak Hopf algebras even if not the most general ones. Also, the (finite dimensional) generalized Kac algebras of T. Yamanouchi [25] are weak Hopf algebras in our sense [14], albeit with an involutive antipode.

In contrast to other Hopf algebraic constructions such as the quasi Hopf algebras [6] or the weak quasi Hopf algebras and rational Hopf algebras [11, 22, 7] weak Hopf algebras are *coassociative*. This allows one to define actions, coactions, and crossed products as easily as in the Hopf algebra case. On the other hand weak Hopf algebras have "weaker" axioms related to the unit and counit: The comultiplication is non-unital, $\Delta(1) \neq 1 \otimes 1$ (like in weak quasi Hopf algebras) and the counit is only "weakly" multiplicative, $\varepsilon(xy) = \varepsilon(x1_{(1)})\varepsilon(1_{(2)}y)$. This kind of "weakness" is the "strength" of weak Hopf algebras because it allows (even in the finite dimensional and semisimple case) the weak Hopf algebra to possess non-integral (quantum) dimensions.

Thus weak Hopf algebras are not special cases of weak quasi Hopf algebras and also not more general than them. Nevertheless, in situations where only the representation category of the quantum group matters, these two concepts are equivalent. This is, of course, not surprising in view of MacLane's Theorem on the equivalence of relaxed and strict monoidal categories [12]. In fact not all of the potential of this theorem is utilized by weak Hopf algebras because their representation category is not quite strict: Only the associator is trivial but not the left and right isomorphisms of the monoidal unit. Although a general analysis clarifying the role of representation categories of weak Hopf algebras within the set of monoidal categories is still missing the examples constructed in [2] using Ocneanu's cocycle suggest that they play a rather fundamental role, as long as they can accomodate to arbitrary 6j-symbols.

So far weak Hopf algebras have been considered only under the additional assumption of finite dimensionality. Although a good deal of the results can be generalized to the infinite dimensional case, finite dimension is particularly attractive because it implies selfduality. Just like finite Abelian groups or finite dimensional Hopf algebras, the finite dimensional weak Hopf algebras (WHA) are *selfdual* in the following sense. If A is a WHA then its dual space \hat{A} is canonically equipped with a weak Hopf algebra structure. Furthermore this duality is reflexive, $(\hat{A})^{\uparrow} \cong A$. This is a feature which makes WHAs more natural objects of study than either finite (non-Abelian) groups or finite dimensional (weak) quasi Hopf algebras.

The main motivation for studying WHAs comes from quantum field theory and operator algebras and consists roughly of the following two symmetry problems.

- **I.** If $N \subset M$ is an inclusion of algebras satisfying certain conditions then find a (unique) "quantum group" G and an action of G on M such that $N = M^G$, the invariant subalgebra.
- **II.** The dual problem is to find a "quantum group" \hat{G} acting on N such that M is isomorphic to the crossed product $N \rtimes \hat{G}$.

Of course, determining the appropriate notion of "quantum group", as well as its action, is part of the problem. If $N \subset M$ is a finite index irreducible depth 2 inclusion of von Neumann factors then the answer is known by [10] to be a finite dimensional C^* -Hopf algebra. In [15] we will show that if we allow the inclusion to be reducible and N and M to have arbitrary finite dimensional centers then the appropriate "quantum group" is a C^* -weak Hopf algebra. Even in case of inclusions of certain associative (non-*) algebras the notion of a WHA over an arbitrary field K, introduced in this paper, may provide a useful invariant.

In Section 2 we introduce the axioms of weak bialgebras and weak Hopf algebras over a field K and discuss their consequences. If K = C, the complex field, then these axioms are equivalent to those of [20]. The present axioms have the advantage of being manifestly selfdual and almost each of them having an ancestor among the Hopf algebra axioms which it generalizes. In discussing the consequences particular attention is paid to the canonical subalgebras A^L and A^R present in any WHA both of which reducing to the scalars K1 if A is a Hopf algebra. From many point of views these subalgebras behave like non-commutative generalizations of numbers. Just to mention some: 1. A^L and A^R are separable K-algebras. 2. The trivial left A-module is a representation on the K-space A^L (or on A^R). 3. The dual weak Hopf algebra \hat{A} have left and right subalgebras \hat{A}^L and \hat{A}^R that are isomorphic to A^R and A^L , respectively. Of course, in order to realize the idea of A^L and A^R being "non-commutative numbers" one should completely get rid of the field K from the outset. As yet we have no concrete proposal for this scenario.

Section 3 is devoted to the study of integrals in weak Hopf algebras. Using the notion of *weak Hopf modules* which is a generalization of the Hopf modules [1, 19] we show that non-zero integrals exist. A weak Hopf version of Maschke's Theorem characterizes semisimple WHAs as those possessing normalized integrals. An other important class of WHAs are those which are Frobenius algebras. They are characterized by possessing non-degenerate left integrals. This class is a selfdual class by the Duality Theorem of non-degenerate integrals. We conclude with giving necessary and sufficient criteria for the existence of Haar integrals, i.e. normalized non-degenerate 2-sided integrals in a WHA.

Section 5 contains the basic properties of weak C^* -Hopf algebras such as the existence of a Haar integral h and a canonical grouplike element $g \ge 0$ implementing S^2 and the modular automorphism of the Haar measure. As a consequence of the existence of Haar measures the dual of a C^* -weak Hopf algebra is a C^* -weak Hopf algebra again. Further analysis of C^* -WHAs will be given in Part II where we discuss the representation category and a notion of dimension which turns out to be non-commutative in case of solitonic representations [3].

2 The Weak Hopf Calculus

2.1 The axioms

Definition 2.1 A weak bialgebra (WBA) is a quintuple $(A, \mu, u, \Delta, \varepsilon)$ satisfying Axioms 1, 2, and 3 below. If $(A, \mu, u, \Delta, \varepsilon, S)$ satisfies Axioms 1, 2, 3, and 4 below it is called a weak Hopf algebra (WHA).

Axiom 1. A is a finite dimensional associative algebra over a field K with multiplication $\mu: A \otimes A \to A$ and unit $u: K \to A$. I.e. μ and u are K-linear and satisfy

Associativity:
$$\mu \circ (\mu \otimes \mathrm{id}) = \mu \circ (\mathrm{id} \otimes \mu)$$
 (A.1)

Unit property: $\mu \circ (u \otimes id) = id = \mu \circ (id \otimes u)$ (A.2)

(Later on we will suppress μ and u, just write xy for $\mu(x, y)$ and use the *unit element* 1 := u(1) instead of u.)

Axiom 2. A is a coalgebra over K with comultiplication $\Delta: A \to A \otimes A$ and counit $\varepsilon: A \to K$. I.e. Δ and ε are K-linear and satisfy

Coassociativity:
$$(\Delta \otimes id) \circ \Delta = (id \otimes \Delta) \circ \Delta$$
 (A.3)

Counit property:
$$(\varepsilon \otimes id) \circ \Delta = id = (id \otimes \varepsilon) \circ \Delta$$
 (A.4)

Axiom 3. For compatibility of the algebra and coalgebra structures we assume

Multiplicativity of the coproduct: For all $x, y \in A$

$$\Delta(xy) = \Delta(x)\Delta(y) \tag{A.5}$$

Weak multiplicativity of the counit: For all $x, y, z \in A$

$$\varepsilon(xyz) = \varepsilon(xy_{(1)})\varepsilon(y_{(2)}z) \tag{A.6a}$$

$$\varepsilon(xyz) = \varepsilon(xy_{(2)})\varepsilon(y_{(1)}z) \tag{A.6b}$$

Weak comultiplicativity of the unit:

$$\Delta^2(1) = (\Delta(1) \otimes 1)(1 \otimes \Delta(1)) \tag{A.7a}$$

$$\Delta^2(1) = (1 \otimes \Delta(1))(\Delta(1) \otimes 1) \tag{A.7b}$$

Axiom 4. There exists a K-linear map $S: A \to A$, called the *antipode*, satisfying the following

Antipode axioms: For all $x \in A$

$$x_{(1)}S(x_{(2)}) = \varepsilon(1_{(1)}x)1_{(2)} \tag{A.8a}$$

$$S(x_{(1)})x_{(2)} = 1_{(1)}\varepsilon(x1_{(2)}) \tag{A.8b}$$

$$S(x_{(1)})x_{(2)}S(x_{(3)}) = S(x)$$
(A.9)

In eqs. (A.6-9) we used a standard suffix notation for (iterated) coproducts, omitting as usual summation indices and a summation symbol.

In the terminology of [14] $(A, \mu, u, \Delta, \varepsilon)$ is called a *weak bialgebra* if it satisfies the Axioms (A.1-5). There a weak bialgebra is called *monoidal* if it satisfies (A.6) and it is called *comonoidal* if it satisfies (A.7). As has been explored in detail in [14], these (co)monoidality axioms are precisely designed to render the category of A-modules (the category of A-comodules, respectively) monoidal.

The dual of a weak bialgebra (weak Hopf algebra) A is the dual space $\hat{A} := \text{Hom}_{K}(A, K)$ equipped with structure maps $\hat{\mu}, \hat{u}, \hat{\Delta}, \hat{\varepsilon}$ ($, \hat{S}$) defined by transposing the structure maps of Aby means of the canonical pairing $\langle , \rangle : \hat{A} \times A \to K$:

$$\begin{array}{lll} \langle \varphi \psi, x \rangle & := & \langle \varphi \otimes \psi, \Delta(x) \rangle \\ \langle \hat{1}, x \rangle & := & \varepsilon(x) \\ \langle \hat{\Delta}(\varphi), x \otimes y \rangle & := & \langle \varphi, xy \rangle \\ & \hat{\varepsilon}(\varphi) & := & \langle \varphi, 1 \rangle \\ \langle \hat{S}(\varphi), x \rangle & := & \langle \varphi, S(x) \rangle \end{array}$$

where $\varphi, \psi \in \hat{A}$ and $x, y \in A$.

Let f and g be maps from the m-fold tensor product $A^{\otimes m}$ to the n-fold tensor product $A^{\otimes n}$ such that they are composites of tensor products of the structure maps $\mu, u, \Delta, \varepsilon, S$ and of the twist maps τ_{ij} interchanging the *i*-th and the *j*-th A factors. Then the equality f = gis called an A-statement. Similarly one defines the \hat{A} -statements. Now every A-statement Q :: f = q determines an equivalent \hat{A} -statement Q^T :: $f^T = q^T$ obtained by reversing the order of composition and replacing μ with $\hat{\Delta}$, u with $\hat{\varepsilon}$, Δ with $\hat{\mu}$, ε with \hat{u} , and S with \hat{S} . The statement Q^T is called the *transpose* of Q.If we now substitute $\mu, u, \Delta, \varepsilon, S$, respectively in place of $\hat{\mu}, \hat{u}, \hat{\Delta}, \hat{\varepsilon}, \hat{S}$ in the statement Q^T we obtain a new A-statement Q^{\sim} :: $f^{\sim} = q^{\sim}$ which is not equivalent to Q in general. This Q^{\sim} will be called the *dual* of Q. For example one can easily verify that the WBA axioms satisfy $(A.1)^{\sim} = (A.3), (A.2)^{\sim} = (A.4), (A.5)^{\sim} = (A.5),$ $(A.6a)^{\sim} = (A.7a)$, and $(A.6b)^{\sim} = (A.7b)$. Thus the weak bialgebra axioms form a selfdual set of statements. This implies that the dual of a WBA is a WBA, too. The same holds for weak Hopf algebras, since each one of the antipode axioms is a selfdual statement. As a consequence of selfduality if Q is a true statement in a WBA or in a WHA then Q^{\sim} is also true there. This principle extends also to statements involving both A and A structure maps and canonical pairing(s).

As has been proven in [14], the above selfdual set of WHA axioms are equivalent to the non-selfdual set of axioms given in [20]. In this work we will gradually reproduce all axioms of [20] as a consequence of the present ones.

For a weak Hopf algebra $(A, 1, \Delta, \varepsilon, S)$ the following conditions are equivalent

- A is a Hopf algebra;
- $\Delta(1) = 1 \otimes 1$;
- $\varepsilon(xy) = \varepsilon(x)\varepsilon(y)$;
- $S(x_{(1)})x_{(2)} = 1\varepsilon(x)$;
- $x_{(1)}S(x_{(2)}) = 1\varepsilon(x)$.

The proof of these assertions are either trivial or will become trivial after acquainting the weak Hopf calculus developed in the next subsections, see also [14].

2.2Weak bialgebras

In a WBA define the maps $\sqcap^L, \sqcap^R: A \to A$ by the formulae

$$\sqcap^{L}(x) := \varepsilon(1_{(1)}x)1_{(2)} , \qquad \sqcap^{R}(x) := 1_{(1)}\varepsilon(x1_{(2)})$$
(2.1)

and introduce the notation $A^L := \sqcap^L(A), A^R := \sqcap^R(A)$ for their images. The analogue objects in the dual bialgebra \hat{A} will be denoted by $\hat{\sqcap}^L, \hat{\sqcap}^R, \hat{A}^L$, and \hat{A}^R , respectively.

Substituting y = 1 in Axiom (A.6b) one obtains immediately the identities

$$\varepsilon(x \sqcap^{L} (y)) = \varepsilon(xy) \qquad (2.2a) \qquad \qquad \varepsilon(\sqcap^{R} (x)y) = \varepsilon(xy) \qquad (2.2b) \sqcap^{L} \circ \sqcap^{L} = \sqcap^{L} \qquad (2.3a) \qquad \qquad \sqcap^{R} \circ \sqcap^{R} = \sqcap^{R} \qquad (2.3b)$$

As a first application of the duality principle take¹ the duals of Eqns(2.2a-b),

$$1_{(1)} \otimes \sqcap^{L}(1_{(2)}) = 1_{(1)} \otimes 1_{(2)} = \sqcap^{R}(1_{(1)}) \otimes 1_{(2)}$$
.

Then these are identities in any WBA. It follows that

$$\Delta(1) \in A^R \otimes A^L . \tag{2.4}$$

Q.e.d.

Lemma 2.2 The counit defines a non-degenerate bilinear form

$$x^L \in A^L, y^R \in A^R \mapsto \varepsilon(y^R x^L) \in K$$

Hence $A^L \cong A^R$ as K-spaces.

Proof:

$$\begin{split} \varepsilon(y^R x^L) &= 0 \ \forall y^R \in A^R \quad \Longrightarrow \quad x^L = \varepsilon(\mathbf{1}_{(1)} x^L) \mathbf{1}_{(2)} = 0 \\ \varepsilon(y^R x^L) &= 0 \ \forall x^L \in A^L \quad \Longrightarrow \quad y^R = \mathbf{1}_{(1)} \varepsilon(y^R \mathbf{1}_{(2)}) = 0 \end{split}$$

where we used (2.4).

Returning to Eqns(2.2a-b) and substituting them into the definitions (2.1) one obtains

$$\Box^{L}(x \Box^{L}(y)) = \Box^{L}(xy) \qquad (2.5a) \qquad \qquad \Box^{R}(\Box^{R}(x)y) = \Box^{R}(xy) \qquad (2.5b)$$

The duals of (2.5a-b),

$$\Delta(A^L) \subset A \otimes A^L \qquad (2.6a) \qquad \qquad \Delta(A^R) \subset A^R \otimes A \qquad (2.6b)$$

tell us that A^L and A^R are left, respectively right coideals in the coalgebra A. Using Axiom (A.7b) we can obtain explicit expressions for these coproducts

$$\Delta(x^L) = \varepsilon(1_{(1)}x^L)1_{(2)} \otimes 1_{(3)} = \varepsilon(1_{(1')}x^L)1_{(1)}1_{(2')} \otimes 1_{(2)} = 1_{(1)}x^L \otimes 1_{(2)}$$
(2.7a)

$$\Delta(x^R) = \mathbf{1}_{(1)} \otimes \mathbf{1}_{(2)} \varepsilon(x^R \mathbf{1}_{(3)}) = \mathbf{1}_{(1)} \otimes \mathbf{1}_{(1')} \mathbf{1}_{(2)} \varepsilon(x^R \mathbf{1}_{(2')}) = \mathbf{1}_{(1)} \otimes x^R \mathbf{1}_{(2)}$$
(2.7b)

where x^L and x^R are meant to denote arbitrary elements of A^L , resp. A^R .

¹In taking the transpose of a statement with $\sqcap^{L/R}$ use the fact that in a WBA $\langle \varphi, \sqcap^L(x) \rangle = \langle \hat{1}_{(1)} \otimes \hat{1}_{(2)}, 1_{(1)} \otimes x \rangle \langle \varphi, 1_{(2)} \rangle = \langle \hat{\sqcap}^L(\varphi), x \rangle$ and similarly $\langle \varphi, \sqcap^R(x) \rangle = \langle \hat{\sqcap}^R(\varphi), x \rangle$.

Lemma 2.3 For all $x \in A$ we have the identities

$$x_{(1)} \otimes \sqcap^{L}(x_{(2)}) = 1_{(1)} x \otimes 1_{(2)}$$
 (2.8a)

$$\Box^{R}(x_{(1)}) \otimes x_{(2)} = 1_{(1)} \otimes x_{(2)}$$
(2.8b)

Proof: Using Axiom (A.7b) one obtains

$$\begin{array}{rcl} x_{(1)} \otimes \varepsilon(1_{(1)}x_{(2)})1_{(2)} &=& 1_{(1')}x_{(1)}\varepsilon(1_{(1)}1_{(2')}x_{(2)}) \otimes 1_{(2)} = 1_{(1)}x_{(1)}\varepsilon(1_{(2)}x_{(2)}) \otimes 1_{(3)} = \\ &=& 1_{(1)}x \otimes 1_{(2)} \\ 1_{(1)}\varepsilon(x_{(1)}1_{(2)}) \otimes x_{(2)} &=& 1_{(1)} \otimes \varepsilon(x_{(1)}1_{(1')}1_{(2)})x_{(2)}1_{(2')} = 1_{(1)} \otimes \varepsilon(x_{(1)}1_{(2)})x_{(2)}1_{(3)} = \\ &=& 1_{(1)} \otimes x1_{(2)} \end{array}$$

Q.e.d.

As a consequence we obtain the dual statements

$$x \sqcap^{L} (y) = \varepsilon(x_{(1)}y)x_{(2)} \qquad (2.9a) \qquad \qquad \sqcap^{R}(x)y = y_{(1)}\varepsilon(xy_{(2)}) . \qquad (2.9b)$$

Proposition 2.4 Let A be a WBA. Then A^L and A^R are subalgebras of A containing 1 and

$$x^L y^R = y^R x^L$$
 for all $x^L \in A^L$ and $y^R \in A^R$. (2.10)

Proof: Eqns(2.8a-b) imply the relations

$$1_{(1)}1_{(1')} \otimes 1_{(2)} \otimes 1_{(2')} = 1_{(1)} \otimes \sqcap^L(1_{(2)}) \otimes 1_{(3)}$$
(2.11a)

$$1_{(1)} \otimes 1_{(1')} \otimes 1_{(2)} 1_{(2')} = 1_{(1)} \otimes \sqcap^R (1_{(2)}) \otimes 1_{(3)}$$
(2.11b)

Now either Axiom (A.7a) or (A.7b) show that on the RHS of (2.11a) the first tensor factor belongs to A^R and on the RHS of (2.11b) the last factor belongs to A^L . This is sufficient for A^R , respectively A^L to be closed under multiplication. Hence they are algebras. Obviously $1 \in A^L \cap A^R$ since $\sqcap^L(1) = 1 = \sqcap^R(1)$. In order to see commutativity of left and right elements just compare Axioms (A.7a) and (A.7b). Q.e.d.

As the duals of the statements that A^L and A^R are subalgebras we obtain that $\text{Ker} \sqcap^L$ and $\text{Ker} \sqcap^R$ are coideals of the coalgebra A, i.e.

$$\Delta(\operatorname{Ker} \sqcap^{C}) \subset A \otimes \operatorname{Ker} \sqcap^{C} + \operatorname{Ker} \sqcap^{C} \otimes A, \qquad (2.12)$$

$$\varepsilon(\operatorname{Ker} \sqcap^{C}) = 0, \qquad C = L, R.$$

On the other hand, being the annihilator of the left coideal \hat{A}^L , $\operatorname{Ker} \sqcap^L$ is a left ideal of the algebra A and similarly, $\operatorname{Ker} \sqcap^R$ is a right ideal.

Lemma 2.5 Consider A^L and A as left A^L -modules by left multiplication. Then $\sqcap^L : A \to A^L$ is a left A^L -module map. Analogously, $\sqcap^R : A \to A^R$ is a right A^R -module map. That is to say

$$\Box^{L} (\Box^{L}(x)y) = \Box^{L}(x) \Box^{L}(y)$$
(2.13a)

$$\Box^{R}(x \Box^{R}(y)) = \Box^{R}(x) \Box^{R}(y)$$
(2.13b)

hold true for all $x, y \in A$.

Proof: At first use the definition of $\sqcap^{L/R}$, then Eqn(2.2a-b), and finally Eqn(2.7a-b):

Q.e.d.

Our next assertion about WBA-s establishes a canonical isomorphism between the left (right) subalgebra of A and the right (left) subalgebra of \hat{A} . Since the existence of a common non-trivial subalgebra of A and \hat{A} for Hopf algebras is by far not typical, this result is the first hint towards the fundamental role A^L and A^R will play in the theory of WHAs.

In order to formulate the statement we introduce the Sweedler arrow notation

$$x \rightharpoonup \varphi := \varphi_{(1)} \langle \varphi_{(2)}, x \rangle , \qquad \varphi \leftharpoonup x := \langle \varphi_{(1)}, x \rangle \varphi_{(2)} . \tag{2.14}$$

Since A is the dual WBA of \hat{A} , the Sweedler arrows $\varphi \rightharpoonup x$ and $x \leftarrow \varphi$ are also defined.

Lemma 2.6 The map $\kappa_A^L: x^L \mapsto (x^L \to \hat{1})$ is an algebra isomorphism from A^L onto \hat{A}^R . The map $\kappa_A^R: x^R \mapsto (\hat{1} \leftarrow x^R)$ is an algebra isomorphism from A^R onto \hat{A}^L . Furthermore, the restriction of the canonical pairing to $\hat{A}^L \times A^L$, $\hat{A}^R \times A^R$, $\hat{A}^L \times A^R$, or to $\hat{A}^R \times A^L$ is non-degenerate.

Proof: Using Eqns(2.11a-b) and the defining properties $\langle \varphi \leftarrow x, y \rangle = \langle \varphi, xy \rangle, \dots$ etc. of the Sweedler arrows one can easily verify that

$$\begin{aligned} (x^{L} \rightarrow \hat{1})(y^{L} \rightarrow \hat{1}) &= \hat{1}_{(1)} \hat{1}_{(1')} \langle \hat{1}_{(2)}, x^{L} \rangle \langle \hat{1}_{(2')}, y^{L} \rangle = \hat{1}_{(1)} \langle \hat{1}_{(2)}, x^{L} \rangle \langle \hat{1}_{(3)}, y^{L} \rangle = \\ &= x^{L} y^{L} \rightarrow \hat{1} \end{aligned}$$

$$(2.15)$$

$$\begin{aligned} (\hat{1} \leftarrow x^R)(\hat{1} \leftarrow y^R) &= \langle \hat{1}_{(1)}, x^R \rangle \langle \hat{1}_{(1')}, y^L \rangle \hat{1}_{(2)} \hat{1}_{(2')} &= \langle \hat{1}_{(1)}, x^R \rangle \langle \hat{1}_{(2)}, y^R \rangle \hat{1}_{(3)} \\ &= \hat{1} \leftarrow x^R y^R \end{aligned}$$
 (2.16)

$$(\hat{1} - x^R) \rightarrow 1 = 1_{(1)} \langle \hat{1} - x^R, 1_{(2)} \rangle = 1_{(1)} \varepsilon(x^R 1_{(2)}) = x^R$$
(2.17)

$$1 - (x^{L} \rightarrow \hat{1}) = \langle 1_{(1)}, x^{L} \rightarrow \hat{1} \rangle 1_{(2)} = \varepsilon (1_{(1)} x^{L}) 1_{(2)} = x^{L}$$
(2.18)

Thus κ_A^L (κ_A^R) is an algebra map with inverse $\kappa_{\hat{A}}^R$ $(\kappa_{\hat{A}}^L)$. As for the non-degeneracy

$$\begin{split} \langle \varphi^R, x^L \rangle &= 0 \; \forall \varphi^R \implies x^L = \langle \hat{1}, x^L \mathbf{1}_{(1)} \rangle \mathbf{1}_{(2)} = \langle \mathbf{1}_{(1)} \rightharpoonup \hat{1}, x^L \rangle \mathbf{1}_{(2)} = 0 \\ \langle \varphi^L, x^L \rangle &= 0 \; \forall \varphi^L \implies x^L = \langle \hat{1}, \mathbf{1}_{(1)} x^L \rangle \mathbf{1}_{(2)} = \langle \hat{1} \leftarrow \mathbf{1}_{(1)}, x^L \rangle \mathbf{1}_{(2)} = 0 \end{split}$$

and the transpose of these prove the claim.

If $\{b_i\}$ is a K-basis of A and $\{\beta^i\} \subset \hat{A}$ is its dual basis, $\langle \beta^i, b_j \rangle = \delta_{ij}$, then

$$\sum_{i} \sqcap^{L}(b_{i}) \otimes \beta^{i} = \sum_{i} b_{i} \otimes \widehat{\sqcap}^{L}(\beta^{i}) = 1 - \widehat{1}_{(1)} \otimes \widehat{1}_{(2)}$$

$$(2.19a)$$

$$\sum_{i} \sqcap^{R}(b_{i}) \otimes \beta^{i} = \sum_{i} b_{i} \otimes \widehat{\sqcap}^{R}(\beta^{i}) = 1_{(1)} \otimes 1_{(2)} \rightharpoonup \widehat{1}$$
(2.19b)

This can be easily seen by pairing both hand sides of any of these equations with $\varphi \otimes x$ and apply the definitions (2.1).

The four arrow identities of the next Scholium will be frequently used in later computations.

Q.e.d.

Scholium 2.7 Let A be a WBA. Then for all $\varphi \in \hat{A}$, $x^L \in A^L$, and $x^R \in A^R$

$$x^{L} \rightharpoonup \varphi = (x^{L} \rightharpoonup \hat{1})\varphi \tag{2.20a}$$

$$\varphi - x^R = \varphi(\hat{1} - x^R) . \qquad (2.20b)$$

$$\varphi - x^L = (\hat{1} - x^L)\varphi \tag{2.21a}$$

$$x^R \rightarrow \varphi = \varphi(x^R \rightarrow \hat{1})$$
. (2.21b)

2.3 Weak Hopf algebras

In this Subsection we will show how the existence of an antipode relates \sqcap^L , A^L with \sqcap^R , A^R and derive the expected properties of S that have been axioms in earlier formulations. The two most important results will be invertibility of the antipode and separability of the algebras A^L and A^R . Let us start with the question of uniqueness of the antipode.

Lemma 2.8 The unit, the counit, and the antipode, if exist, are unique. I.e. if $(A, \mu, u, \Delta, \varepsilon, S)$ and $(A, \mu, u', \Delta, \varepsilon', S')$ are both weak Hopf algebras then u' = u, $\varepsilon' = \varepsilon$, and S' = S.

Proof: The uniqueness of the unit and the counit are obvious. Therefore \sqcap^L and \sqcap^R are common in these two WHA-s. In order to prove S' = S introduce the convolution product

$$(f \diamond g)(x) := f(x_{(1)})g(x_{(2)}), \qquad x \in A$$
 (2.22)

on functions $f, g \in \operatorname{Hom}_{K}(A, A)$. This is an associative operation in terms of which the antipode axioms take the form

$$\mathrm{id} \diamond S = \sqcap^L, \qquad S \diamond \mathrm{id} \diamond S = S, \qquad S \diamond \mathrm{id} = \sqcap^R.$$

Now S' satisfies the same equations with the same \sqcap^L, \sqcap^R , therefore

$$S' = S' \diamond \mathrm{id} \diamond S' = S' \diamond \sqcap^L = S' \diamond \mathrm{id} \diamond S = \sqcap^R \diamond S = S \diamond \mathrm{id} \diamond S = S \ .$$

Q.e.d.

As a preparation for the Theorem below notice that the definitions (2.1) have counterparts involving the antipode:

$$\Box^{L}(x) = \varepsilon(S(x)1_{(1)})1_{(2)} \qquad (2.23a) \qquad \qquad \Box^{R}(x) = 1_{(1)}\varepsilon(1_{(2)}S(x)) \qquad (2.23b)$$

As a matter of fact

$$\Box^{L}(x) = \varepsilon(1_{(1)} \Box^{L}(x))1_{(2)} = \varepsilon(\Box^{L}(x)1_{(1)})1_{(2)} = \varepsilon(x_{(1)}S(x_{(2)})1_{(1)})1_{(2)} =$$

= $\varepsilon(\Box^{R}(x_{(1)})S(x_{(2)})1_{(1)})1_{(2)} = \varepsilon(S(x_{(1)})x_{(2)}S(x_{(3)})1_{(1)})1_{(2)} =$
= $\varepsilon(S(x)1_{(1)})1_{(2)}$

where in the subsequent equations (2.1), (2.10), (A.8a), (2.2b), (A.8b), and finally (A.9) have been used. Eqn(2.23b) can be proven analogously. As the duals of (2.23a-b) we have automatically the identities

$$\Box^{L}(x) = S(1_{(1)})\varepsilon(1_{(2)}x) \qquad (2.24a) \qquad \qquad \Box^{R}(x) = \varepsilon(x1_{(1)})S(1_{(2)}). \qquad (2.24b)$$

Lemma 2.9 In a WHA A the following identities hold:

$$\Box^L \circ S = \Box^L \circ \Box^R = S \circ \Box^R \tag{2.25a}$$

$$\square^R \circ S = \square^R \circ \square^L = S \circ \square^L \tag{2.25b}$$

Q.e.d.

Proof: It is sufficient to prove the first equalities in (2.25a) and (2.25b) because the second ones then follow by duality.

$$\Box^{L} \circ S(x) = \varepsilon(1_{(1)}S(x))1_{(2)} = \varepsilon(1_{(1)}S(x_{(1)})x_{(2)}S(x_{(3)}))1_{(2)} =$$

= $\varepsilon(1_{(1)}S(x_{(1)}) \Box^{L}(x_{(2)}))1_{(2)} = \varepsilon(1_{(1)}S(x_{(1)})x_{(2)})1_{(2)}$
= $\Box^{L} \circ \Box^{R}(x).$

In a similar way one can verify $\Box^R \circ S = \Box^R \circ \Box^L$.

The above Lemma implies that $S(A^R) \subset A^L$ and $S(A^L) \subset A^R$. On the other hand Eqns(2.24a-b) say that $A^L \subset S(A^R)$ and $A^R \subset S(A^L)$. Therefore the antipode maps A^L onto A^R bijectively and maps A^R onto A^L bijectively.

Theorem 2.10 Let A be a WHA. Then the antipode is antimultiplicative and anti comultiplicative,

$$S(xy) = S(y)S(x) \qquad x, y \in A , \qquad (2.26)$$

$$S(x)_{(1)} \otimes S(x)_{(2)} = S(x_{(2)}) \otimes S(x_{(1)}) \qquad x \in A , \qquad (2.27)$$

and the restrictions $S|_{A^L}$ and $S|_{A^R}$ are bijections such that

$$S(A^L) = A^R$$
, $S(A^R) = A^L$. (2.28)

The unit and the counit are S-invariant,

$$S(1) = 1$$
, $(2.29a)$ $\varepsilon \circ S = \varepsilon$. $(2.29b)$

Furthermore $S: A \to A$ is invertible.

Proof: We have already shown (2.28). (2.27) is the dual of (2.26) and (2.29a) is the dual of (2.29b).

$$\begin{split} S(xy) &= S(x_{(1)}y_{(1)})x_{(2)}y_{(2)}S(x_{(3)}y_{(3)}) = S(x_{(1)}y_{(1)}) \sqcap^{L} (x_{(2)} \sqcap^{L} (y_{(2)})) = \\ &= S(x_{(1)}y_{(1)})x_{(2)} \sqcap^{L} (y_{(2)})S(x_{(3)}) = \sqcap^{R} (\sqcap^{R}(x_{(1)})y_{(1)})S(y_{(2)})S(x_{(2)}) = \\ &= S(y_{(1)}) \sqcap^{R} (x_{(1)})y_{(2)}S(y_{(3)})S(x_{(3)}) = S(y_{(1)})y_{(2)}S(y_{(3)})S(x_{(1)})x_{(2)}S(x_{(3)}) = \\ &= S(y)S(x) \;. \end{split}$$

Next we prove (2.29b).

$$\begin{split} \varepsilon(S(x)) &= \varepsilon(S(x_{(1)})x_{(2)}S(x_{(3)})) = \varepsilon(S(x_{(1)})\sqcap^L(x_{(2)})) = \varepsilon(S(x_{(1)})x_{(2)}) = \varepsilon(\sqcap^R(x)) = \\ &= \varepsilon(x) \; . \end{split}$$

In order to prove invertibility of S notice that the descending chain $A \supset S(A) \supset S^2(A) \supset \ldots$ of WHAs all contain 1 by (2.29a). This implies the existence of $n \in \mathbb{N}$ such that

$$1 \in S^{n+1}(A) = S^n(A) \subset S^{n-1}(A)$$
.

We want to show that this implies $S^n(A) = S^{n-1}(A)$. Replacing A by $S^{n-1}(A)$ it is therefore enough to prove invertibility of S under the additional assumption $S^2(A) = S(A)$, implying

$$\operatorname{Ker} S \cap S(A) = 0$$

In this case let $\overline{S} := S|_{S(A)}$, then $\overline{S}: S(A) \to S(A)$ is bijective and

$$P_S := \bar{S}^{-1} \circ S : A \to S(A)$$

is a multiplicative idempotent satisfying

$$P_S(xS(y)) = P_S(x)S(y), \quad x, y \in A.$$

By (2.28) $A^{L,R} \subset S(A)$. Now taking into account the identity $x = x_{(1)}S(x_{(2)})x_{(3)}$, which follows directly from axioms (A.8a) and (A.4), then using also $P_S(1) = 1$ we have

$$P_{S}(x) = P_{S}(x_{(1)}S(x_{(2)})x_{(3)}) = P_{S}(x_{(1)})S(x_{(2)})x_{(3)} = P_{S}(x_{(1)}S(x_{(2)}))x_{(3)} = = P_{S}(1)x_{(1)}S(x_{(2)})x_{(3)} = x,$$

so Ker $P_S = \text{Ker } S = 0.$

We are now able to derive (versions of) the original antipode axioms of [2, 20]:

$$\begin{aligned} x_{(1)} \otimes x_{(2)} S(x_{(3)}) &= 1_{(1)} x \otimes 1_{(2)} \\ S(x_{(1)}) x_{(2)} \otimes x_{(3)} &= 1_{(1)} \otimes x_{1_{(2)}} \end{aligned}$$
(2.30a)
(2.30b)

Q.e.d.

$$S(x_{(1)})x_{(2)} \otimes x_{(3)} = 1_{(1)} \otimes x_{(2)}$$
 (2.30b)

$$x_{(1)} \otimes S(x_{(2)}) x_{(3)} = x \mathbf{1}_{(1)} \otimes S(\mathbf{1}_{(2)})$$
(2.30c)

$$x_{(1)}S(x_{(2)}) \otimes x_{(3)} = S(1_{(1)}) \otimes 1_{(2)}x$$
(2.30d)

The first two are just rewritings of the bialgebra identities (2.8a-b). The second two are more delicate.

$$\begin{array}{rcl} x_{(1)} \otimes S(x_{(2)})x_{(3)} &=& x_{(1)} \otimes \varepsilon(x_{(2)}1_{(1)})S(1_{(2)}) = x_{(1)}1_{(1')} \otimes \varepsilon(x_{(2)}1_{(2')}1_{(1)})S(1_{(2)}) = \\ &=& x_{(1)}1_{(1)}\varepsilon(x_{(2)}1_{(2)}) \otimes S(1_{(3)}) = x1_{(1)} \otimes S(1_{(2)}) \\ x_{(1)}S(x_{(2)}) \otimes x_{(3)} &=& S(1_{(1)})\varepsilon(1_{(2)}x_{(1)}) \otimes x_{(2)} = S(1_{(1)})\varepsilon(1_{(2)}1_{(1')}x_{(1)}) \otimes 1_{(2')}x_{(2)} = \\ &=& S(1_{(1)}) \otimes \varepsilon(1_{(2)}x_{(1)})1_{(3)}x_{(2)} = S(1_{(1)}) \otimes 1_{(2)}x \end{array}$$

The following Proposition also holds, if A is just a WBA, see [14].

Proposition 2.11 Let A be a WHA over K. Then A^L and A^R are separable K-algebras, in particular, they are semisimple.

Proof: Recall that an algebra A is separable if and only if there exists a $q \in A \otimes A$ such that $(x \otimes 1)q = q(1 \otimes x)$ holds for all $x \in A$ and furthermore $\mu(q) = 1$, where μ denotes the multiplication map of A [18]. Such a q will be called a separable idempotent². So, our proof will consist of showing that $q^L = S(1_{(1)}) \otimes 1_{(2)} \in A^L \otimes A^L$ and $q^R = 1_{(1)} \otimes S(1_{(2)}) \in A^R \otimes A^R$ are separable idempotents of A^L and A^R , respectively. In fact we will prove the somewhat more general identities

$$x_{(1)}y^R \otimes x_{(2)} = x_{(1)} \otimes x_{(2)}S(y^R)$$
(2.31a)

$$x_{(1)} \otimes y^L x_{(2)} = S(y^L) x_{(1)} \otimes x_{(2)}$$
 (2.31b)

²In fact q is an idempotent only if considered as an element of $A \otimes A^{op}$.

valid for all $x \in A$ and $y^L \in A^L$, $y^R \in A^R$. Pairing the LHS of (2.31a) with $\varphi \otimes \psi$, we obtain

$$\begin{split} \langle \varphi \otimes \psi, LHS \rangle &= \langle \varphi(y^R \rightharpoonup \hat{1}), x_{(1)} \rangle \langle \psi, x_{(2)} \rangle = \langle \varphi, x_{(1)} \rangle \langle (S(y^R) \rightharpoonup \hat{1}) \psi, x_{(2)} \rangle = \\ &= \langle \varphi \otimes \psi, RHS \rangle \;. \end{split}$$

The proof of (2.31b) is simply the mirror image of the above argument. Q.e.d.

2.4 The "trivial" representation

Since the counit of a WHA is in general not an algebra map, weak Hopf algebras may be lacking of any 1-dimensional representation. Nevertheless the axioms ensure that any WHA A has a distinguished representation providing a unit object for the (relaxed) monoidal category of left A-modules. We shall discuss this category in detail in [3]. Now we concentrate only on the properties of this representation. We remark that the trivial representation exists already in WBA's [14] and therefore the use of the antipode in this subsection is not obligatory.

Since the algebras $A^{L/R}$ occur on the right hand side of Axioms (A.8a-b) where in ordinary Hopf algebras the trivial representation stands, one expects that the "trivial representation" of WHAs must be a non-trivial representation acting on either one of the algebras $A^{L/R}$ or $\hat{A}^{L/R}$.

Lemma 2.12 The following left A-modules are isomorphic.

$$\begin{array}{rcl} {}_{A}\hat{A}^{R} & :: & the \ vector \ space \ \hat{A}^{R} & with \ action & x \cdot \varphi^{R} := x \rightharpoonup \varphi^{R} \\ {}_{A}\hat{A}^{L} & :: & the \ vector \ space \ \hat{A}^{L} & with \ action & x \cdot \varphi^{L} := \varphi^{L} \leftarrow S(x) \\ {}_{A}A^{L} & :: & the \ vector \ space \ A^{L} & with \ action & x \cdot y^{L} := \sqcap^{L}(xy^{L}) \\ {}_{A}A^{R} & :: & the \ vector \ space \ A^{R} & with \ action & x \cdot y^{R} := \sqcap^{R}(y^{R}S(x)) \end{array}$$

Proof: $\hat{S}: \hat{A}^L \to \hat{A}^R$ is an isomorphism of vector spaces and $\hat{S}(\varphi \leftarrow S(x)) = x \to \hat{S}(\varphi)$ is a general WHA identity. This proves the isomorphism of the first two A-modules. Similarly, $S: A^L \to A^R$ is an isomorphism of vector spaces and $S(\sqcap^L(xy)) = \sqcap^R(S(y)S(x))$ is a WHA identity. This proves the isomorphism of the last two A-modules.

In order to show the isomorphism of ${}_A\hat{A}^R$ with ${}_AA^L$ consider the bijection $B: \hat{A}^R \to A^L$, $B(\varphi^R) := 1 \leftarrow \varphi^R$. Then

$$\begin{split} B(x \rightharpoonup \varphi^R) &= 1 \leftharpoonup (x \rightharpoonup \varphi^R) = \langle 1_{(1)} x, \varphi^R \rangle 1_{(2)} = \\ &= \langle 1_{(1)} (x \leftharpoonup \varphi^R), \hat{1} \rangle 1_{(2)} = \sqcap^L (x \leftharpoondown \varphi^R) = \sqcap^L (x (1 \leftharpoondown \varphi^R)) = \\ &= \sqcap^L (x B(\varphi^R)) \end{split}$$

hence B is a left A-module map. Here, in the last-but-one equality we have used one of the four arrow identities of Scholium 2.7. Q.e.d.

Definition 2.13 By the trivial representation of the WHA A we mean the cyclic left A-module $V_{\varepsilon} := {}_{A}\hat{A}^{R}$ with A-action $D_{\varepsilon}: A \to End_{K}\hat{A}^{R}, D_{\varepsilon}(x)\varphi := x \rightharpoonup \varphi$.

The third and fourth A-modules of the above Lemma demonstrate that the restriction of the trivial representation to A^L (A^R) is equivalent to its left regular representation, hence faithful. This is one of the instances where $A^{L/R}$ appears in the role of a ground "field".

Later we will need the following strengthening of Lemma 2.6.

Lemma 2.14 Let A be a WHA and introduce the notation $Z^L := A^L \cap Center A$, $Z^R := A^R \cap Center A$, and $Z := A^L \cap A^R$. Then the isomorphism (of algebras) $\kappa_A^L : A^L \to \hat{A}^R$ restricts to an isomorphism $Z^L \to \hat{Z}$ and the isomorphism $\kappa^R_A: A^R \to \hat{A}^L$ restricts to the isomorphism $Z^R \to \hat{Z}$. Therefore

$$Z^{L} \rightarrow \hat{1} = \hat{Z} = \hat{1} \leftarrow Z^{R}$$
$$Z \rightarrow \hat{1} = \hat{Z}^{R}, \qquad \hat{Z}^{L} = \hat{1} \leftarrow Z$$

The two isomorphisms have a common restriction to the hypercenter $Hypercenter A := Z^L \cap Z^R$ and yields an isomorphism Hypercenter $A \rightarrow$ Hypercenter \hat{A} .

 $Proof: \text{ Notice that for } c \in \text{Center } A \ \hat{1} \leftarrow c = c \rightharpoonup \hat{1}. \text{ Therefore } x^L \in Z^L \Rightarrow x^L \rightharpoonup \hat{1} = \hat{1} \leftarrow x^L \in \hat{Z}.$ This proves $\kappa_A^L(Z^L) \subset Z$.

If $z \in Z$ then $(z \rightarrow \hat{1})\varphi = z \rightarrow \varphi$ by (2.20a) and $z \rightarrow \varphi = \varphi(z \rightarrow \hat{1})$ by (2.21b). Hence $z \rightarrow \hat{1}$ is central. This proves $\kappa_A^L(Z) \subset \hat{Z}^R$. Since $(\kappa_A^L)^{-1} = \kappa_{\hat{A}}^R$, the analogue inclusions $\kappa_A^R(Z^R) \subset \hat{Z}$ and $\kappa_A^R(Z) \subset \hat{Z}^L$ complete the

proof. Q.e.d.

The unusual feature of the trivial representation of WHA-s is that it can be decomposable. But this can occur only if the left and right subalgebras of the dual have non-trivial intersection as the next Proposition claims.

Proposition 2.15 Let A be a WHA, let $(V_{\varepsilon}, D_{\varepsilon})$ be its trivial representation as in Definition 2.13. Then

$$EndV_{\varepsilon} = D_{\varepsilon}(Z^{L}) = D_{\varepsilon}(Z^{R}) , \qquad (2.32)$$

where $EndV_{\varepsilon}$ denotes the algebra of A-module endomorphisms of V_{ε} .

Proof: Let $T \in \text{End} V_{\varepsilon}$ then $T(x \rightarrow \hat{1}) = x \rightarrow T(\hat{1})$, for $x \in A$, in particular

$$\begin{array}{rcl} T(x^L \rightarrow \hat{1}) &=& x^L \rightarrow T(\hat{1}) &=& (x^L \rightarrow \hat{1})T(\hat{1}) \\ T(x^L \rightarrow \hat{1}) &=& T(S^{-1}(x^L) \rightarrow \hat{1}) = S^{-1}(x^L) \rightarrow T(\hat{1}) &=& T(\hat{1})(x^L \rightarrow \hat{1}) \end{array}$$

where we have made use of Eqns (2.20a) and (2.21b). Since by Lemma 2.6 $A^L \rightarrow \hat{1} = \hat{A}^R$, $\zeta := T(\hat{1}) \in \text{Center } \hat{A}^R \text{ and } T(\varphi^R) = \varphi \zeta.$ Thus $x \to \zeta = T(x \to \hat{1}) = (x \to \hat{1}) \zeta$ holds for all $x \in A$. It follows that

$$\langle \hat{\sqcap}^L(\zeta), x \rangle = \langle \zeta, 1 \leftarrow (x \rightharpoonup \hat{1}) \rangle = \langle (x \rightharpoonup \hat{1}) \zeta, 1 \rangle = \langle x \rightharpoonup \zeta, 1 \rangle = \langle \zeta, x \rangle ,$$

i.e. $\zeta \in \hat{A}^L \cap \hat{A}^R \equiv \hat{Z}$. Now by Lemma 2.14 there exists a $z^L \in Z^L$ such that $\zeta = z^L \rightharpoonup \hat{1}$. We can conclude that

$$T(\varphi^R) = \zeta \varphi^R = (z^L \rightharpoonup \hat{1}) \varphi^R = z^L \rightharpoonup \varphi^R = D_{\varepsilon}(z^L) \varphi^R$$

i.e. $T = D_{\varepsilon}(z^L)$. This proves End $V_{\varepsilon} \subset D_{\varepsilon}(Z^L)$. The opposite inclusion is trivial since $D_{\varepsilon}(Z^L) \subset \operatorname{Center}(D_{\varepsilon}(A)).$ This finishes the proof of $\operatorname{End} V_{\varepsilon} = D_{\varepsilon}(Z^L).$

Showing the other statement $\operatorname{End} V_{\varepsilon} = D_{\varepsilon}(Z^R)$ one proceeds as above but chooses a $z^R \in$ Z^R such that $\zeta = \hat{1} - z^R$. Then

$$\begin{array}{rcl} T(\varphi^R) &=& \varphi^R(\hat{1} \leftharpoonup z^R) = \varphi^R(z^R \rightharpoonup \hat{1}) = z^R \rightharpoonup \varphi^R = \\ &=& D_{\varepsilon}(z^R)\varphi^R \end{array}$$

completes the proof.

Q.e.d.

Notice that the above Proposition does not imply that the trivial A-module is semisimple. It does imply, however, that V_{ε} has a decomposition $V_{\varepsilon} \cong \bigoplus_{\nu} V_{\nu}$ into indecomposable A-modules in which the indecomposables are disjoint, i.e. Hom $(V_{\mu}, V_{\nu}) = 0$ for all $\mu \neq \nu$.

Definition 2.16 If $Z^L = K1$, or equivalently, if the trivial representation is indecomposable then the WHA is called pure.

The name "pure" comes from the C^* -setting when the trivial representation arises from the positive linear functional ε by the GNS construction. Thus A is pure iff ε is pure.

Nota bene pureness is not a selfdual notion, duals of pure WHA-s may not be pure. Clearly, A is pure iff $Z^L \cong Z^R$ is trivial but \hat{A} is pure iff Z is trivial.

3 Weak Hopf Modules and Integral Theory

As in Hopf algebras so in weak Hopf algebras the integrals play a decisive role in the structure analysis of these algebras. Using integrals we can formulate conditions for the algebra to be Frobenius, symmetric, or semisimple, and study questions related to innerness of S^2 or S^4 . Furthermore we will be able to characterize those WHAs that have Haar measures. In deriving the basic properties of integrals the weak generalization of the the Fundamental Theorem of Hopf modules is very useful. Unfortunately, it seems to be less powerful than in Hopf algebra theory (cf. [13]) where it implies the existence of non-degenerate integrals. It is an open problem yet whether all WHAs are Frobenius algebras. We can prove, however, that all of them are quasi-Frobenius algebras.

3.1 Integrals in weak Hopf algebras

The following definition provides the weak Hopf generalization of the well known notion of integrals in a Hopf algebra [19].

Definition 3.1 A left (right) integral in a weak Hopf algebra A is an element $l \in A$ ($r \in A$) satisfying

$$xl = \sqcap^{L}(x)l \qquad \left(rx = r \sqcap^{R}(x) \right)$$
(3.1)

for all $x \in A$. The space of left (right) integrals in A is denoted by $\mathcal{I}^{L}(A)$ ($\mathcal{I}^{R}(A)$). Elements of $\mathcal{I} := \mathcal{I}^{L}(A) \cap \mathcal{I}^{R}(A)$ are called two-sided integrals. A left or right integral in A is called non-degenerate if it defines a non-degenerate functional on \hat{A} . $l \in \mathcal{I}^{L}(A)$ is called normalized if $\sqcap^{L}(l) = 1$, $r \in \mathcal{I}^{R}(A)$ is called normalized if $\sqcap^{R}(r) = 1$.

Some equivalent formulations of left (right) integrals are gathered in the next

Lemma 3.2 Let A be a weak Hopf algebra. Then the following statements for an element $l \in A$ are equivalent:

- a) $l \in \mathcal{I}^L(A)$
- **b)** $l_{(1)} \otimes x l_{(2)} = S(x) l_{(1)} \otimes l_{(2)}$ for all $x \in A$
- c) $l \rightarrow \hat{A} \subset \hat{A}^L$
- **d)** $(\varphi \leftarrow x) \rightharpoonup l = S(x)(\varphi \rightharpoonup l)$ for all $\varphi \in \hat{A}$ and $x \in A$
- e) (Ker \sqcap^L)l = 0
- f) $S(l) \in \mathcal{I}^R(A)$

Proof: **a**) \Rightarrow **b**): Using (2.30b) and (2.7a) we have $l_{(1)} \otimes xl_{(2)} = [S(x_{(1)}) \otimes 1]\Delta(x_{(2)}l) = [S(x_{(1)})x_{(2)}S(x_{(3)}) \otimes 1]\Delta(l) = S(x)l_{(1)} \otimes l_{(2)}$. **b**) \Rightarrow **a**): $xl = x_{(1)}l_{(1)}\varepsilon(x_{(2)}l_{(2)}) = x_{(1)}S(x_{(2)})l_{(1)}\varepsilon(l_{(2)}) = \sqcap^{L}(x)l$. **a**) \Leftrightarrow **c**): For an $l \in A$ the equation $\langle l \rightharpoonup \varphi, x \rangle = \langle \cap^{L}(l \rightharpoonup \varphi), x \rangle$ is clearly equivalent to the equation $\langle \varphi, xl \rangle = \langle \varphi, \sqcap^{L}(x)l \rangle$. **b**) \Leftrightarrow **d**): By pairing the 2nd tensor factor of **b**) with an arbitrary $\varphi \in \hat{A}$. **a**) \Rightarrow **e**): is obvious. **e**) \Rightarrow **a**): $xl = [x - \sqcap^{L}(x)]l + \sqcap^{L}(x)l = \sqcap^{L}(x)l$. **f**) \Leftrightarrow **a**): This follows by applying S to (3.1). Q.e.d.

Definition 3.1 as well as Lemma 3.2 provide rather technical characterizations of integrals. The next argument sheds some light on their real nature. Consider the left A-module map ε_R from the left regular A-module to the trivial A-module given by acting with the trivial representation on the cyclic vector $\hat{1}$:

$$\varepsilon_R : {}_A A \to {}_A \hat{A}^R
 x \mapsto (x \rightarrow \hat{1}) .$$
(3.2)

The existence of this (non-zero) map shows that Hom $({}_{A}A, {}_{A}\hat{A}^{R})$ is non-zero. However, there is in general no guarantee that Hom $({}_{A}\hat{A}^{R}, {}_{A}A)$ is non-zero. Left integrals are precisely the objects that label the possible homomorphisms of the latter type.

Lemma 3.3 Left integrals l in A are in one-to-one correspondence with left A-module homomorphisms $f: {}_{A}\hat{A}^{R} \to {}_{A}A$. The correspondence is given by $f \mapsto f(\hat{1}) \in \mathcal{I}^{L}$. What is more the above map provides an isomorphism $\mathcal{I}_{A}^{L} \cong Hom({}_{A}\hat{A}^{R}, {}_{A}A)$ of right A-modules. In other words \mathcal{I}_{A}^{L} is isomorphic to the A-dual of the trivial left A-module.

Proof: If $f \in \text{Hom}\left({}_{A}\hat{A}^{R}, {}_{A}A\right)$ then $xf(\hat{1}) = f(x \rightarrow \hat{1}) = f(\sqcap^{L}(x) \rightarrow \hat{1}) = \sqcap^{L}(x)f(\hat{1})$, hence $f(\hat{1}) \in \mathcal{I}^{L}$. This is obviously a right A-module map. It is invertible since for $l \in \mathcal{I}^{L}$ the map $f_{l}: \hat{A}^{R} \rightarrow A, f_{l}(\varphi^{R}) := (1 \leftarrow \varphi^{R})l$ is a left A-module map and satisfies $f_{l}(\hat{1}) = l$. Q.e.d.

The identification of \mathcal{I}^L with Hom $({}_A\hat{A}^R, {}_AA)$ yields an A-valued bilinear form ${}_A\hat{A}^R \times \mathcal{I}^L_A \to A$ given by evaluation, $(\varphi^R, l) \mapsto f_l(\varphi^R)$. Replacing ${}_A\hat{A}^R$ with ${}_AA^L$ using the isomorphism of Lemma 2.12 we obtain that this bilinear form is nothing but multiplication in A

$${}_{A}A^{L} \times \mathcal{I}^{L}_{A} \to {}_{A}A_{A} , \qquad (x^{L}, l) \mapsto x^{L}l$$

$$(3.3)$$

and it is an A-A bimodule map. We claim that (3.3) is a non-degenerate bilinear form. From one side, $x^L l = 0 \ \forall x^L \in A^L \Rightarrow l = 0$, this is trivial. From the other side we will be able to prove this after having established that WHAs are quasi-Frobenius algebras in Thm. 3.11. As a matter of fact by Theorem 61.2 of [4] the left annihilator of the right annihilator of the left ideal Ker \sqcap^L is Ker \sqcap^L itself. Now by Lemma 3.2.e) the right annihilator of Ker \sqcap^L is just \mathcal{I}^L . Thus $x^L l = 0 \ \forall l \in \mathcal{I}^L \Rightarrow x^L = 0$ follows.

Now we turn to an other characterization of left integrals that is related to conditional expectations. Notice at first that if $\lambda \in \mathcal{I}^L(\hat{A})$ then the map $E_{\lambda}: x \mapsto \lambda \rightharpoonup x$ is an $A^L \cdot A^L$ -bimodule map from A into A^L commuting with the right \hat{A} -action on A. In fact, all such maps arise from a left integral, as the following Lemma shows.

Lemma 3.4 The left integrals $\lambda \in \mathcal{I}^{L}(\hat{A})$ are in one-to-one correspondence with right \hat{A} module maps $E \in Hom(A_{\hat{A}}, A_{\hat{A}}^{L})$ via

$$\begin{array}{rccc} \lambda & \mapsto & E_{\lambda} \\ E & \mapsto & \varepsilon \circ E \end{array}$$

Proof: If λ is a left integral then E_{λ} is a right \hat{A} -module map and maps into A^{L} by Lemma 3.2.c).

Now let $E \in \text{Hom}(A_{\hat{A}}, A_{\hat{A}}^L)$. Then

$$E(x) = \varepsilon(1_{(1)}E(x))1_{(2)} = \varepsilon \circ E(S^{-1}(1_{(1)})x)1_{(2)}$$

where we used the fact that a right \hat{A} -module map is an A^L - A^L -bimodule map by (2.20b) and (2.21a). Hence

$$E(x) = \varepsilon \circ E(x_{(3)}) x_{(1)} S(x_{(2)}) = \sqcap^{L} (\lambda \rightharpoonup x)$$

where $\lambda := \varepsilon \circ E$. It remained to show that λ is a left integral.

$$\langle \varphi \lambda, x \rangle = \varepsilon (E(x \leftarrow \varphi)) = \varepsilon (E(x) \leftarrow \varphi) = \langle \widehat{\sqcap}^L(\varphi) \lambda, x \rangle$$

which proves the claim.

Q.e.d.

The characterization of left integrals λ as "conditional expectations" E_{λ} provides a link to the theory of inclusions and "Jones extensions" [15].

The properties of the normalized and the non-degenerate left integrals will be discussed in later Subsections. Here we only remark that λ is non-degenerate iff E_{λ} is non-degenerate and λ is normalized iff E_{λ} is unital.

There are two twisting operations $A \mapsto A^{op}$ and $A \mapsto A_{cop}$ that produce WHAs from WHAs. In the first one the multiplication μ is replaced with opposite multiplication $\mu^{op}(x, y) = \mu(y, x)$ while in the second the coproduct is replaced by $\Delta^{op}(x) = x_{(2)} \otimes x_{(1)}$. In both cases the antipode is replaced by S^{-1} . The left and right subalgebras/integrals and the dual WHAs of the resulting four twisted versions of a WHA A are related to those of A as follows.

	\sqcap^L	\sqcap^R	A^L	A^R	\mathcal{I}^L	\mathcal{I}^R	Â
$A = A(\mu, \Delta, S)$	\sqcap^L	\sqcap^R	A^L	A^R	\mathcal{I}^L	\mathcal{I}^R	Â
$A^{op} = A(\mu^{op}, \Delta, S^{-1})$	$S^{-1} \circ \sqcap^R$	$S^{-1} \circ \sqcap^L$	A^L	A^R	\mathcal{I}^R	\mathcal{I}^L	\hat{A}_{cop}
$A_{cop} = A(\mu, \Delta^{op}, S^{-1})$	$S^{-1} \circ \sqcap^L$	$S^{-1} \circ \sqcap^R$	A^R	A^L	\mathcal{I}^L	\mathcal{I}^R	\hat{A}^{op}
$A^{op}_{cop} = A(\mu^{op}, \Delta^{op}, S)$	\sqcap^R	\sqcap^L	A^R	A^L	\mathcal{I}^R	\mathcal{I}^L	\hat{A}_{cop}^{op}

As an application of the table we give here the twisted versions of the identity of Lemma 3.2.d):

$$(\varphi - x) \rightarrow l = S(x)(\varphi \rightarrow l) \tag{3.4a}$$

$$(x \rightarrow \varphi) \rightarrow r = (\varphi \rightarrow r)S^{-1}(x)$$
 (3.4b)

$$l \leftarrow (\varphi \leftarrow x) = S^{-1}(x)(l \leftarrow \varphi) \tag{3.4c}$$

$$r \leftarrow (x \rightharpoonup \varphi) = (r \leftarrow \varphi)S(x) \tag{3.4d}$$

for all $x \in A$, $\varphi \in \hat{A}$, $l \in \mathcal{I}^L$, and $r \in \mathcal{I}^R$.

3.2 Weak Hopf modules

Let A be a WHA. Recall that a left A-module is a K-linear space M carrying a left action of the algebra A, denoted by $x \in A, m \in M \mapsto x \cdot m$. A right A-module is a left module M of the opposite algebra A^{op} with action denoted by $x \in A, m \in M \mapsto m \cdot x$. Since A is unital, all modules are assumed to be non-degenerate, i.e. 1 acts as the identity. The left A-module M is called faithful if $x \cdot m = 0, \forall m \in M$ implies x = 0. The A-modules know nothing about the coalgebra structure of A. The left A-comodules M in turn are the comodules of the coalgebra A and carry no information about the algebra structure of A. The left coaction is denoted by $m \mapsto m_{-1} \otimes m_0 \in A \otimes M$. One defines the right A-comodules analogously and denotes the coaction as $m \mapsto m_0 \otimes m_1 \in M \otimes A$.

Because of the finite dimensionality of A there is a one-to-one correspondence between left (right) A-coactions on M and right (left) \hat{A} -actions on M given by

$$m \cdot \varphi = \langle \varphi, m_{-1} \rangle m_0 , \qquad m_{-1} \otimes m_0 = \sum_i b_i \otimes m \cdot \beta^i$$

$$(3.5)$$

$$\varphi \cdot m = m_0 \langle \varphi, m_1 \rangle$$
, $m_0 \otimes m_1 = \sum_i \beta^i \cdot m \otimes b_i$. (3.6)

Here $\{b_i\}$ denotes an arbitrary basis of A and $\{\beta^i\}$ is its dual basis: $\langle\beta^i, b_j\rangle = \delta_{ij}$. There are 8 basic examples of A modules with the target space M being either A itself or its dual \hat{A} . These are the following.

where the Sweedler arrow notation (2.14) has been used. They all are faithful and nondegenerate due to the existence of a unit and a counit. To each of the A-modules in the above list there is a corresponding \hat{A} -comodule denoted by the same symbol. This identification is justified also by the fact that $N \subset M$ is an A-submodule if and only if it is an \hat{A} -subcomodule.

By analogy with our definition of left integrals, the space of *invariants* of a left A-module M is defined to be the subspace

Inv
$$M := \{ m \in M \mid x \cdot m = \sqcap^{L}(x) \cdot m, \forall x \in A \}$$
 (3.7)

By the same methods as in Lemma 3.3, Inv M is linearly isomorphic to Hom $({}_A\hat{A}^R, {}_AM)$ via

$$\operatorname{Inv} M = \{ f(\hat{1}) \, | \, f \in \operatorname{Hom} (_A \hat{A}^R, _A M) \}.$$
(3.8)

By duality, we define the *coinvariants* of a right A-comodule M as

$$\operatorname{Coinv} M := \{ m \in M \,|\, m_0 \otimes m_1 = m_0 \otimes \sqcap^L(m_1) \}$$

$$(3.9)$$

Thus, $m \in \operatorname{Coinv} M \Leftrightarrow m_0 \otimes m_1 \in M \otimes A^L$ and for a left A-module M, the invariants Inv $M \subset M$ coincide with $\operatorname{Coinv} M \subset M$ considered as an \hat{A} -comodule. Similarly, for a right A-module (left A-comodule) M the invariants (coinvariants) are

Inv
$$M = \{ m \in M \mid m \cdot x = m \cdot \sqcap^R(x), \forall x \in A \}$$
 (3.10)

Coinv
$$M = \{ m \in M | m_{-1} \otimes m_0 = \sqcap^R (m_{-1}) \otimes m_0 \}.$$
 (3.11)

Notice that the (co)invariants do not form a sub(co)module, even not an $A^{L/R}$ -submodule.

Scholium 3.5 The invariants of the left (right) regular A-module are precisely the left (right) integrals of A:

 $\operatorname{Inv}_A A = \mathcal{I}^L(A) , \qquad \operatorname{Inv} A_A = \mathcal{I}^R(A) .$

The invariants of $_A\hat{A}$ and \hat{A}_A , on the other hand, yield the left and right subalgebras, respectively:

$$\operatorname{Inv}_A \hat{A} = \hat{A}^L$$
, $\operatorname{Inv} \hat{A}_A = \hat{A}^R$.

Investigating the structure of the mixed modules ${}_{\hat{A}}\hat{A}^A$ and ${}^{\hat{A}}\hat{A}_{\hat{A}}$, that incorporate the whole bialgebra structure of A, one arrives to a weak generalization of the notion of Hopf modules [1, 19].

Definition 3.6 A right weak Hopf module (right WHM) over A is a right A module M which is also a right A-comodule such that the compatibility relation

$$(m \cdot x)_0 \otimes (m \cdot x)_1 = m_0 \cdot x_{(1)} \otimes m_1 x_{(2)}$$
(3.12)

holds for $x \in A$, $m \in M$.

Lemma 3.7 Let M be a right WHM over A. Then for all $m \in M$

i) $m_0 \cdot \sqcap^R(m_1) = m$

ii) Coinv $M = \{m \in M \mid m_0 \otimes m_1 = m \cdot 1_{(1)} \otimes 1_{(2)}\}$ and Coinv M is a right A^L submodule. iii) $E(m) := m_0 \cdot S(m_1)$ provides a projection $E : M \to \text{Coinv } M$.

Proof: i) Let M be a right WHM over A. Since

$$m_0 \varepsilon(m_1 x) = m_0 \cdot \mathbf{1}_{(1)} \varepsilon(m_1 \mathbf{1}_{(2)} x) = m_0 \cdot \mathbf{1}_{(1)} \varepsilon(m_1 \mathbf{1}_{(2)}) \varepsilon(\mathbf{1}_{(3)} x) = m \cdot \mathbf{1}_{(1)} \varepsilon(\mathbf{1}_{(2)} x)$$
(3.13)

for all $x \in A$, we have $m_0 \otimes \hat{1} \leftarrow m_1 = m \cdot 1_{(1)} \otimes \hat{1} \leftarrow 1_{(2)}$, so that

$$m_0 \cdot \sqcap^R(m_1) = m_0 \cdot [(\hat{1} \leftarrow m_1) \rightharpoonup 1] = m \cdot 1_{(1)} [(\hat{1} \leftarrow 1_{(2)}) \rightharpoonup 1] = m \cdot 1_{(1)} \sqcap^R (1_{(2)}) = m.$$
(3.14)

ii) The inclusion \supset follows from (2.4). Conversely, if $m \in \text{Coinv } M$ then $m_0 \otimes m_1 \in M \otimes A^L$, implying by (2.3a), (2.10) and (3.13)

$$m_0 \otimes m_1 = m_0 \otimes \varepsilon(m_1 \mathbf{1}_{(1)}) \mathbf{1}_{(2)} = m \cdot \mathbf{1}_{(1)} \otimes \varepsilon(\mathbf{1}_{(2)} \mathbf{1}_{(1')}) \mathbf{1}_{(2')} = m \cdot \mathbf{1}_{(1)} \otimes \mathbf{1}_{(2)}.$$

Also $\Delta(A^L) \subset A \otimes A^L$ and therefore Coinv M is a right A^L -submodule.

iii) To check that $n := m_0 \cdot S(m_1)$ is a coinvariant for all $m \in M$ we compute

$$n_0 \otimes n_1 = (m_0 \otimes m_1)(\cdot (S(m_2)_{(1)} \otimes S(m_2)_{(2)}) = = m_0 \cdot S(m_3) \otimes m_1 S(m_2) = m_0 \cdot S(1_{(2)}m_1) \otimes S(1_{(1)}) = = n \cdot 1_{(1)} \otimes 1_{(2)}.$$

Since for $n \in \text{Coinv } M$ we have $n_0 \cdot S(n_1) = n \cdot 1_{(1)}S(1_{(2)}) = n$, E is a projection onto Coinv M. Q.e.d.

Example 3.8 The right weak Hopf module $_{\hat{A}}\hat{A}^A$

As a linear space the module is the dual WHA \hat{A} . The right action and coaction are

$$\varphi \cdot x := S(x) \rightharpoonup \varphi, \qquad \varphi_0 \otimes \varphi_1 := \sum_i \beta^i \varphi \otimes b_i.$$
 (3.15)

Clearly, the right A-coaction is dual to the left \hat{A} -multiplication and therefore counital and right-coassociative. The compatibility condition (3.12) can be seen as follows.

$$(\varphi \cdot x)_0 \otimes (\varphi \cdot x)_1 = \sum_i \beta^i (S(x) \rightharpoonup \varphi) \otimes b_i =$$

$$= \sum_{i} \beta^{i} \left[S(x_{(1)}) \rightarrow x_{(2)} S(x_{(3)}) \rightarrow \varphi \right] \otimes b_{i} =$$

$$= \sum_{i} \beta^{i} \left[S(x_{(1)}) \rightarrow (x_{(2)} \rightarrow \hat{1}) \varphi \right] \otimes b_{i} =$$

$$= \sum_{i} \beta^{i} \left[S(x_{(2)}) x_{(3)} \rightarrow \hat{1} \right] \left[S(x_{(1)}) \rightarrow \varphi \right] \otimes b_{i} =$$

$$= \sum_{i} \left[S(x_{(2)}) x_{(3)} \rightarrow \beta^{i} \right] \left[S(x_{(1)}) \rightarrow \varphi \right] \otimes b_{i} =$$

$$= \sum_{i} S(x_{(1)}) \rightarrow \left[(x_{(2)} \rightarrow \beta^{i}) \varphi \right] \otimes b_{i} =$$

$$= \sum_{i} S(x_{(1)}) \rightarrow (\beta^{i} \varphi) \otimes b_{i} x_{(2)} =$$

$$= \varphi_{0} \cdot x_{(1)} \otimes \varphi_{1} x_{(2)} .$$

The A-coinvariants of this WHM coincide with the \hat{A} -invariants of the dual left regular \hat{A} -module $_{\hat{A}}\hat{A}$ and therefore with the space of left integrals in \hat{A} by Scholium 3.5,

$$\operatorname{Coinv}\left(_{\hat{A}}\hat{A}^{A}\right) = \mathcal{I}^{L}(\hat{A}).$$
(3.16)

The Fundamental Theorem of Hopf modules generalizes to the weak case as follows.

Theorem 3.9 Let A be a WHA, M be a right WHM over A and let N = Coinv M denote the set of coinvariants of M. Since N is a right A^L submodule, one can form the A^L -module tensor product $N \underset{A^L}{\otimes} A$ and make it into a right WHM by the definitions

$$(n \otimes a) \cdot x := n \otimes ax \tag{3.17a}$$

$$(n \otimes a)_0 \otimes (n \otimes a)_1 := (n \otimes a_{(1)}) \otimes a_{(2)}$$

$$(3.17b)$$

where $a, x \in A, n \in N$. Then the map

$$\alpha: N \underset{A^{L}}{\otimes} A \to M, \quad n \otimes x \mapsto n \cdot x \tag{3.18}$$

is an isomorphism of right WHM-s.

Recall that an *isomorphism of WHM-s* is just a module isomorphism which is a comodule isomorphism at the same time.

Proof: That α is a module map and comodule map is easy to verify. To construct the inverse define

$$\beta : M \to N \underset{A^L}{\otimes} A$$

$$\beta(m) = m_0 \cdot S(m_1) \otimes m_2 \equiv E(m_0) \otimes m_1 . \qquad (3.19)$$

Then β is obviously a comodule map. We show that it is also a module map.

We are left with showing that on the one hand

$$\begin{array}{lll} \beta \circ \alpha(n \otimes x) &=& \beta(n \cdot x) = \beta(n) \cdot x = n \cdot 1_{(1)} S(1_{(2)}) \otimes 1_{(3)} x = \\ &=& n \otimes 1_{(1)} S(1_{(2)}) 1_{(3)} x = \\ &=& n \otimes x \end{array}$$

and on the other hand

$$\alpha \circ \beta(m) = m_0 \cdot S(m_1)m_2 = m$$

where in the last equality Lemma 3.7i) has been used.

Q.e.d.

Applying this Theorem to the WHM of Example 3.8 we obtain the right WHM isomorphism

$${}_{\hat{A}}\hat{A}^A \cong \mathcal{I}^L(\hat{A}) \underset{A^L}{\otimes} {}_{\hat{A}}A_A .$$

$$(3.20)$$

Corollary 3.10 In any WHA A the space of left integrals $\mathcal{I}^{L}(A) = S(\mathcal{I}^{R}(A))$ is non-zero and $\mathcal{I}^{L}(\hat{A})$ is the dual of $\mathcal{I}^{R}(A)$ with respect to the restriction of the canonical pairing. Moreover, choosing a basis $\{\lambda^{a}\}$ in $\mathcal{I}^{L}(\hat{A})$ and taking its dual basis $\{r_{a}\}$ in $\mathcal{I}^{R}(A)$, we have

$$\hat{1} = \sum_{a} S(r_a) \rightarrow \lambda_a \tag{3.21}$$

$$1 = \sum_{a} r_a - \hat{S}(\lambda_a) \tag{3.22}$$

Proof: $\mathcal{I}^{L}(\hat{A}) \neq 0$ follows from (3.20). By inspecting the form of the projection $E: M \to N$ in Example 3.8 we get a projection $L: A \to \mathcal{I}^{L}(A)$ onto the left integrals,

$$L(x) = \sum_{i} \hat{S}^{2}(\beta^{i}) \rightharpoonup (b_{i}x) . \qquad (3.23)$$

Therefore the projection to the right integrals is

$$R(x) = S \circ L \circ S^{-1}(x) = \sum_{i} (xb_i) - \hat{S}^2(\beta^i) .$$
(3.24)

Similar expressions define the projections \hat{L} and \hat{R} to the dual integrals. Now it is easy to check that

$$\langle \hat{L}(\varphi), x \rangle = \langle \varphi, R(x) \rangle$$
 (3.25)

proving the non-degeneracy of the restriction of the canonical pairing to $\mathcal{I}^{L}(\hat{A}) \times \mathcal{I}^{R}(A)$.

The dual bases satisfy $\langle \lambda^a, r_b \rangle = \delta_{ab}$ therefore

$$\langle \lambda^a, xS(r_a) \rangle = \langle \hat{L}(\beta^i), xS(b_i) \rangle = \langle S^2(b_j) \rightharpoonup \beta^j \beta^i, xS(b_i) \rangle = \langle \beta^k, xS \circ \sqcap^R(b_k) \rangle = (3.26)$$

= $\langle 1_{(2)} \rightharpoonup \hat{1}, xS(1_{(1)}) \rangle$ (3.27)

where in the last step we used (2.19b). This proves (3.21). (3.22) is the twisted version in A_{cop}^{op} . Q.e.d.

3.3 Restrictions on the algebraic structure

The existence of a weak Hopf structure on the K-algebra A involves certain restrictions on the algebra A, just like in case of Hopf algebras. In this subsection we show that any WHA A is quasi-Frobenius, i.e. self-injective. The notions of semisimple and separable algebras coincide within the class of WHAs. Moreover, we prove an analogue of Maschke's Theorem which claims that A is semisimple if and only if it has normalized left integrals.

Theorem 3.11 Every weak Hopf algebra over a field K is a quasi-Frobenius algebra.

Proof: By Theorem 61.2 of [4] it is sufficient to prove that the left regular A-module ${}_{A}A$ is injective. By the Nagao–Nakayama theorem injectivity of a left A-module is equivalent to that it is a direct sum of K-duals of principal indecomposable right A-modules. Since \hat{A}^A is the K-dual of ${}_{A}A$, we need to show that \hat{A}^A is a direct sum of principal indecomposable right A-modules, i.e. that \hat{A}^A is projective. This in turn is a consequence of the Fundamental Theorem of WHM-s.

As a matter of fact we have the right A-module isomorphisms

$$\hat{A}^A \cong \mathcal{I}^L(\hat{A}) \underset{A^L}{\otimes} A_A \cong P(\mathcal{I}^L(\hat{A}) \underset{\kappa}{\otimes} A_A)$$
(3.28)

the first of which is the consequence of the Fundamental Theorem of the right WHM $_{\hat{A}}\hat{A}^{A}$, the second of which is a rather simple property of the amalgamated tensor product with respect to the separable algebra A^{L} . In order to explain the projection P here we make a digression.

Lemma 3.12 Define the map $P: \mathcal{I}^{L}(\hat{A}) \underset{K}{\otimes} A \to \mathcal{I}^{L}(\hat{A}) \underset{K}{\otimes} A$ by

$$P(\lambda \otimes x) := S^2(1_{(1)}) \rightharpoonup \lambda \otimes 1_{(2)}x .$$
(3.29)

Then $P \circ P = P$ and Ker P coincides with $Ker \pi$ of the canonical projection π from the free right A module $\mathcal{I}^{L}(\hat{A}) \underset{K}{\otimes} A_{A}$ onto $\mathcal{I}^{L}(\hat{A}) \underset{A}{\otimes} A_{A}$. Therefore

$$P(\mathcal{I}^{L}(\hat{A}) \underset{K}{\otimes} A_{A}) \xrightarrow{\pi \mid_{ImP}} \mathcal{I}^{L}(\hat{A}) \underset{A^{L}}{\otimes} A_{A}$$
(3.30)

is an isomorphism of right A-modules.

Proof: The kernel of the canonical projection is

$$\operatorname{Ker} \pi = \operatorname{Span}_{K} \{ (\lambda \otimes x^{L} y) - (S(x^{L}) \rightharpoonup \lambda \otimes x) \mid \lambda \otimes y \in \mathcal{I}^{L}(\hat{A}) \otimes A, \ x^{L} \in A^{L} \}$$
(3.31)

If $\sum_{i} (\lambda_i \otimes x_i) \in \text{Ker } \pi$ then obviously $\sum_{i} S^2(1_{(1)}) \rightharpoonup \lambda_i \otimes 1_{(2)} x_i = 0$, therefore $\text{Ker } \pi \subset \text{Ker } P$. Now assume $\sum \lambda_i \otimes x_i \in \text{Ker } P$. Then

$$\sum_{i} \lambda_i \otimes x_i = \sum_{i} \lambda_i \otimes S(1_{(1)}) 1_{(2)} x_i =$$
$$\sum_{i} [\lambda_i \otimes S(1_{(1)}) (1_{(2)} x_i) - S^2(1_{(1)}) \rightharpoonup \lambda_i \otimes 1_{(2)} x_i] \in \operatorname{Ker} \pi$$

This proves $\operatorname{Ker} \pi = \operatorname{Ker} P$. That P is a projection and a right A-module map is trivial to verify. Therefore $\pi|_{\operatorname{Im} P}$ is an A-module isomorphism. Q.e.d.

Back to the Proof of Theorem 3.11: In virtue of the above Lemma the amalgamated tensor product $\mathcal{I}^{L}(\hat{A}) \underset{A^{L}}{\otimes} A_{A}$ is the direct summand of a free A-module, hence projective. By Equation 3.28 this is isomorphic to \hat{A}^{A} . This proves projectivity of \hat{A}^{A} , hence injectivity of $_{A}A$. Q.e.d.

The equivalence of (c) and (d) of the next Theorem provides a weak Hopf version of Maschke's Theorem known for Hopf algebras as well [9]. Below we denote $\varepsilon_R(x) := x \rightarrow \hat{1}$.

Theorem 3.13 The following conditions on a WHA A over K are equivalent:

a) A is semisimple.

b) In the category of left A-modules the following exact sequence is split

 $0 \longrightarrow Ker \varepsilon_R \longrightarrow {}_{A}A \xrightarrow{\varepsilon_R} {}_{A}\hat{A}^R \longrightarrow 0$

c) There exists a normalized left integral $l \in A$.

d) A is a separable K-algebra.

Proof: (a \Rightarrow c): If A is semisimple, then Ker $\sqcap^L \equiv$ Ker ε_R being a left ideal there exists $p = p^2 \in A$ such that Ker $\sqcap^L = Ap$, whence l = 1 - p is a normalized left integral by Lemma 3.2.e.

(b \Leftrightarrow c): Let $F \in \text{Hom}(_A \hat{A}^R, _A A)$ be such that $\varepsilon_R \circ F = \text{id}$. Then $xF(\hat{1}) = F(x \rightharpoonup \hat{1}) = \Box^L(x)F(\hat{1})$, for $x \in A$, therefore $F(\hat{1}) \in \mathcal{I}^L(A)$. Moreover, $\hat{1} = \varepsilon_R(F(\hat{1})) = F(\hat{1}) \rightharpoonup \hat{1}$ implying $\Box^L(F(\hat{1})) = 1$. Conversely, if $l \in \mathcal{I}^L$ is a normalized left integral then $F \in \text{Hom}(_A \hat{A}^R, _A A)$ given by $F(\varepsilon_R(x)) := xl$ satisfies $\varepsilon_R \circ F = \text{id}$.

 $(c\Rightarrow d)$: Let l be a normalized left integral. Then $q = l_{(1)} \otimes S(l_{(2)})$ is a separating idempotent for A. As a matter of fact $\mu(q) = 1$ follows from the normalization $\sqcap^{L}(l) = 1$ while $(x \otimes 1)q = q(1 \otimes x)$ is precisely the left integral property of Lemma 3.2.b.

 $(d\Rightarrow a)$: This is a standard result [18]. Q.e.d.

3.4 Non-degenerate integrals

Until now we have not been able to decide whether the WHM Theorem of Subsection 3.2 implies the existence of non-degenerate integrals, as it does in case of Hopf algebras. In the present subsection we will show that the existence of non-degenerate integrals in the WHA \hat{A} is equivalent to the existence of non-degenerate functionals on A, i.e. that A is a Frobenius algebra. As a byproduct we obtain that the class of Frobenius WHAs is selfdual.

The space \mathcal{I}^{R} of right integrals can be viewed as a K-module, as a left A^{L} -module ${}_{A^{L}}\mathcal{I}^{R}$ by left multiplication, and as a left A-module ${}_{A}\mathcal{I}^{R}$ since it is a left ideal of A. From the latter point of view ${}_{A}\mathcal{I}^{R}$ is the dual of the trivial right A-module, ${}_{A}\mathcal{I}^{R} \cong \text{Hom}(\hat{A}^{L}_{A}, A_{A})$, by a twisted version of Lemma 3.3. As a K-module \mathcal{I}^{R} has $\hat{\mathcal{I}}^{L}$ as its K-dual, $\hat{\mathcal{I}}^{L} \cong \text{Hom}(_{K}\mathcal{I}^{R}, _{K}K)$ the isomorphism being given by the restriction of the canonical pairing (see Corollary 3.10). The next Lemma shows that $\hat{\mathcal{I}}^{L}$ is also the A^{L} -dual of \mathcal{I}^{R} with right A^{L} -module structure precisely the one needed in Example 3.8, i.e. $\lambda \cdot x^{L} = S(x^{L}) \rightarrow \lambda$.

Lemma 3.14 The A^L -valued bilinear form

$$(,)_{A^L} : \mathcal{I}^R \times \hat{\mathcal{I}}^L \to A^L, \qquad (r,\lambda)_{A^L} = \lambda \rightharpoonup r$$

$$(3.32)$$

provides an isomorphism of right A^L -modules

$$(\hat{\mathcal{I}}^L)^{A^L} \to Hom(_{A^L}\mathcal{I}^R, {}_{A^L}A^L) , \qquad \lambda \mapsto (r \mapsto (r, \lambda)_{A^L}) , \qquad (3.33)$$

i.e. $(\hat{\mathcal{I}}^L)^{A^L}$ is the A^L -dual of ${}_{A^L}\mathcal{I}^R$.

Proof: At first verify the following properties of the A^L -valued bilinear form.

$$(x^L \cdot r, \lambda)_{A^L} = x^L(r, \lambda)_{A^L}$$
(3.34a)

$$(r, \lambda \cdot x^L)_{A^L} = (r, \lambda)_{A^L} x^L$$
(3.34b)

$$(r,\lambda)_{A^L} = 0 \ \forall r \in \mathcal{I}^R \quad \Rightarrow \quad \lambda = 0$$

$$(3.34c)$$

The first two are simple WHA identities. The third one follows from the relation $\varepsilon((r, \lambda)_{AL}) =$ $\langle \lambda, r \rangle$ and from non-degeneracy of the canonical pairing \langle , \rangle on $\hat{\mathcal{I}}^L \times \mathcal{I}^R$ (Corollary 3.10). Now properties (a) and (b) tell us that $\lambda \mapsto (., \lambda)_{A^L}$ is indeed the required A^L -module map and (c) ensures that it is injective. In order to show that it is surjective it is sufficient to find finite sets of elements $\{r_a\}$ in \mathcal{I}^R and $\{\lambda^a\}$ in $\hat{\mathcal{I}}^L$ such that

$$\sum_{a} (r, \lambda^a)_{A^L} r_a = r \qquad \forall r \in \mathcal{I}^R .$$
(3.35)

For if such elements exist then any $f \in \text{Hom}\left(_{A^L}\mathcal{I}^R, _{A^L}A^L\right)$ can be written as $f = \sum_a \lambda^a \cdot f(r_a)$. As a matter of fact

$$f(r) = f(\sum_{a} (r, \lambda^a)_{A^L} \cdot r_a) = \sum_{a} (r, \lambda^a)_{A^L} f(r_a) = (r, \sum_{a} \lambda^a \cdot f(r_a))_{A^L}$$

for all $r \in \mathcal{I}^R$. Now we claim that a pair of dual bases $\{r_a\}$ of \mathcal{I}^R and $\{\lambda^a\}$ of $\hat{\mathcal{I}}^L$, in the sense of K-duality, i.e. $\langle \lambda^a, r_b \rangle = \delta_{ab}$, also satisfies (3.35). As a matter of fact for $\lambda \in \hat{\mathcal{I}}^L$ we have

$$\langle \lambda, \sum_{a} (\lambda^{a} \rightarrow r) r_{a} \rangle = \varepsilon (\sum_{a} (\lambda^{a} \rightarrow r) (\lambda \rightarrow r_{a})) = \sum_{a} \langle \lambda^{a}, r(\lambda \rightarrow r_{a}) \rangle =$$

$$= \sum_{a} \langle \lambda^{a}, r(S(r_{a}) \leftarrow \hat{S}^{-1}(\lambda)) \rangle = \sum_{a} \langle \lambda^{a}_{(1)}, r \rangle \langle \hat{S}^{-1}(\lambda) \lambda^{a}_{(2)}, S(r_{a}) \rangle =$$

$$= \sum_{a} \langle \lambda \lambda^{a}_{(1)}, r \rangle \langle \lambda^{a}_{(2)}, S(r_{a}) \rangle = \langle \lambda \left[\sum_{a} S(r_{a}) \rightarrow \lambda^{a} \right], r \rangle = \langle \lambda, r \rangle$$

where in the last equality (3.21) has been used.

Q.e.d.

Notice that Eqn(3.35) means that ${}_{A^L}\mathcal{I}^R$ is finitely generated projective³. Therefore by a general result (see e.g. [5]) Î

$$\tilde{\mathcal{L}}^L \bigotimes_{A^L} \mathcal{I}^R \cong \operatorname{End}_{A^L} \mathcal{I}^R .$$
 (3.36)

On the other hand the isomorphism α of the WHM Theorem, if restricted to $\hat{\mathcal{I}}^L \bigotimes_{A^L} \mathcal{I}^R$, yields an isomorphism onto \hat{A}^L . Thus we have the composition

$$\mathcal{E}\colon A^L \longrightarrow \hat{A}^L \xrightarrow{\alpha^{-1}} \hat{\mathcal{I}}^L \underset{A^L}{\otimes} \mathcal{I}^R \longrightarrow \text{End}_{A^L} \mathcal{I}^R \tag{3.37}$$

³Although this is clear from the fact that A^{L} is semisimple, constructing the concrete bases λ^{a} , r_{a} was not in vain since it will help to compute the commutant in (3.38).

of isomorphisms. Evaluating it explicitly we obtain

$$r \cdot \mathcal{E}(x^{L}) = r \cdot \left(\sum_{ij} S^{2}(b_{i}) \rightharpoonup (\beta^{i} \beta^{j} (\hat{1} \leftarrow x^{L})) \otimes b_{j} \right) = r \cdot \left(\sum_{j} \hat{L}(\beta^{j} (\hat{1} \leftarrow x^{L})) \otimes b_{j} \right) =$$
$$= r \cdot (\sum_{a} \lambda^{a} \otimes S(x^{L})r_{a}) = \sum_{a} (r, \lambda^{a})_{A^{L}} S(x^{L})r_{a} =$$
$$= S(x^{L})r \quad \forall r \in \mathcal{I}^{R}, \ x^{L} \in A^{L}.$$
(3.38)

This proves our next

Proposition 3.15 The left modules ${}_{A^L}\mathcal{I}^R$ and ${}_{A^R}\mathcal{I}^R$ are faithful and the endomorphism algebra of ${}_{A^L}\mathcal{I}^R$ consists of left multiplications with elements of A^R . Therefore

$$End_{A^L}\mathcal{I}^R \cong A^L$$
, as algebras. (3.39)

The set $Sec A^L$ of equivalence classes of simple left A^L -modules will be called the *sectors* of A^L . For $a \in Sec A^L$ let V_a be a simple module from the class a and let $\mathcal{D}_a = \operatorname{End} V_a$ be the corresponding division algebra. Then by the Wedderburn structure theorem $A^L \cong \bigoplus_a M_{n_a}(\mathcal{D}_a)$. Let m_a denote the multiplicity of V_a in the semisimple module ${}_{AL}\mathcal{I}^R$. Then End ${}_{AL}\mathcal{I}^R \cong \bigoplus_a M_{m_a}(\mathcal{D}_a)$ which is, by the Proposition, isomorphic to A^L . This is possible only if there is a permutation

$$\mathcal{L}: \operatorname{Sec} A^L \to \operatorname{Sec} A^L, \quad \text{such that } n_{\tilde{a}} = m_a \text{ and } \mathcal{D}_{\tilde{a}} = \mathcal{D}_a.$$
(3.40)

This means that \mathcal{I}^R , as an A^L -End ${}_{A^L}\mathcal{I}^R$ bimodule, can be identified with a direct sum of matrices,

$${}_{A^L} \mathcal{I}^R \cong \oplus_a \operatorname{Mat} \left(n_a \times m_a, \mathcal{D}_a \right) \,. \tag{3.41}$$

This allows us to compute its K-dimension and apply the Cauchy-Schwarz inequality to obtain the bound

$$\dim_K \mathcal{I}^R = \sum_a (\dim_K \mathcal{D}_a) n_a m_a \leq \sum_a (\dim_K \mathcal{D}_a) n_a^2 = \dim_K A^L .$$
(3.42)

Equality holds here iff $m_a = n_a$, $a \in Sec A^L$, i.e. iff ${}_{A^L}\mathcal{I}^R \cong {}_{A^L}A^L$. Now we are ready to prove the following

Theorem 3.16 Let A be a WHA over the field K. Then the following conditions are equivalent.

- i) A is a Frobenius algebra;
- ii) $\dim_K \mathcal{I}^R = \dim_K A^L;$
- **iii)** Non-degenerate integrals exist in A;
- iv) \hat{A} is a Frobenius algebra.

Proof: (i) \Rightarrow (ii) If $_{A}A \cong _{A}\hat{A}$ then their invariants $\mathcal{I}(_{A}A) = \mathcal{I}^{L}$ and $\mathcal{I}(_{A}\hat{A}) = \hat{A}^{L}$, respectively (see Scholium 3.5), are isomorphic as K-spaces. (ii) \Rightarrow (iii) As we have seen above the K-space isomorphism of \mathcal{I}^{R} and A^{L} implies that $_{A^{L}}\mathcal{I}^{R}$ is isomorphic to the left regular module $_{A^{L}}A^{L}$. Since the latter is cyclic, there exists a cyclic vector $r \in _{A^{L}}\mathcal{I}^{R}$. Thus l := S(r) is cyclic in

 $(\mathcal{I}^L)^{\hat{A}^L}$. As a matter of fact $\mathcal{I}^L = S(\mathcal{I}^R) = S(A^L r) = lA^R = \hat{S}(\hat{A}^L) \rightarrow l$. Now interchanging the roles of A and \hat{A} in the WHM Theorem

$$A = \alpha(\mathcal{I}^L \underset{\hat{A}^L}{\otimes} \hat{A}) = \hat{S}(\hat{A}) \rightharpoonup \mathcal{I}^L = \hat{S}(\hat{A}) \rightharpoonup (\hat{S}(\hat{A}^L) \rightharpoonup l) = \hat{A} \rightharpoonup l \ ,$$

hence l is a non-degenerate left integral in A. (iii) \Rightarrow (iv) is obvious since l is a non-degenerate functional on \hat{A} . (iv) \Rightarrow (i) Repeat the arguments above from (i) to (iv) with A replaced by \hat{A} . Q.e.d.

Weak Hopf algebras satisfying any one of the conditions of the above Theorem will be called *Frobenius WHAs*. Note that since semisimple algebras are Frobenius, in a semisimple WHA there exist both normalized and non-degenerate integrals, although there may be no integral sharing both properties. 4

As an immediate consequence of the above considerations we have

Scholium 3.17 The following properties for $l \in \mathcal{I}^L$ $(r \in \mathcal{I}^R)$ are equivalent:

- i) l (r) is non-degenerate;
- **ii)** *l* is separating for $\mathcal{I}_{AL,R}^L$ (*r* is separating for $_{AL,R}\mathcal{I}^R$);
- iii) *l* is cyclic for $\mathcal{I}_{AL,R}^L$ (*r* is cyclic for $_{A^{L,R}}\mathcal{I}^R$).

In a Frobenius WHA A the group of invertible elements A^R_{\times} of A^R acts on the set $\mathcal{I}^L_*(A)$ of non-degenerate left integrals transitively and freely. Similar statement holds for the non-degenerate right integrals \mathcal{I}^R_* ,

$$\mathcal{I}^L_* = l A^R_\times , \qquad \mathcal{I}^R_* = A^L_\times r \tag{3.43}$$

for any $l \in \mathcal{I}^L_*$ and $r \in \mathcal{I}^R_*$. Similar relation for the dual integrals shows that there are one-to-one correspondences between non-degenerate integrals of A and of \hat{A} . The Theorem below selects a distinguished "natural" one-to-one correspondence.

Theorem 3.18 Let A be a WHA and $l \in \mathcal{I}^{L}(A)$ be a left integral. If there exists a $\lambda \in \hat{A}$ such that $\lambda \rightharpoonup l = 1$ then it is unique, it is a left integral in \hat{A} , and both l and λ are non-degenerate. Moreover $l \rightharpoonup \lambda = \hat{1}$. Such a pair (l, λ) will be called a dual pair of left integrals.

Similarly, elements $r \in \mathcal{I}^R_*$ and $\lambda \in \hat{\mathcal{I}}^L_*$ are in one-to-one correspondence by either one of the equivalent relations $\lambda \rightharpoonup r = 1$ or $\lambda \leftarrow r = \hat{1}$.

Proof: By Lemma 3.2.d) if l is a left integral such that $\lambda \rightarrow l = 1$ then⁵ $l_R \circ \lambda_L = S$. Since S is invertible, both l_R and λ_L are invertible, i.e. l and λ are non-degenerate and λ is unique. To show that $\lambda \in \mathcal{I}^L(\hat{A})$

$$\varphi \lambda \rightarrow l = \varphi \rightarrow 1 = \hat{\sqcap}^L(\varphi) \lambda \rightarrow l , \quad \varphi \in \hat{A}$$

⁴As an example consider $M_2(\mathbb{Z}_2)$, the semisimple algebra of two by two matrices over the field of mod2 residue classes. Fix a set of matrix units $\{e_{ij}\}$ and introduce the coproduct $\Delta(e_{ij}) := e_{ij} \otimes e_{ij}$. Then we have two normalized left integrals $l_j = \sum_i e_{ij}$ for j = 1, 2 neither of which is non-degenerate. The only non-degenerate left integral is $l = l_1 + l_2$ for which however $\sqcap^L(l) = 0$.

⁵Here we use the standard notations $f_L, f_R: A \to \hat{A}$ defined by $f_L(x) := f \leftarrow x$ and $f_R(x) := x \rightharpoonup f$ for any $f \in \hat{A}$.

suffices since l_R is a bijection. It remains to show that $l \rightarrow \lambda = \hat{1}$ which eventually justifies the term "dual" left integral. For $l \in \mathcal{I}^L(A)$ and $\lambda \in \hat{A}$ we have

$$\begin{split} \lambda &\rightharpoonup l = 1 \quad \Leftrightarrow \quad x l_{(1)} \langle \lambda, l_{(2)} \rangle = x \quad x \in A \\ &\Leftrightarrow \quad \langle \lambda, x l_{(2)} \rangle S^{-1}(l_{(1)}) = x \quad x \in A \\ &\Rightarrow \quad \langle l \rightharpoonup \lambda, x \rangle = \varepsilon(x) \quad x \in A \\ &\Leftrightarrow \quad l \rightharpoonup \lambda = \hat{1} \end{split}$$

The duality between $\hat{\mathcal{I}}^L$ and \mathcal{I}^R follows from the above duality between $\hat{\mathcal{I}}^L$ and \mathcal{I}^L by passing from A to A^{op} . The other two twisted versions of the Theorem are not spelled out explicitly. They can also be obtained by applying the antipode to the above relations. Q.e.d.

Recall that the quasibasis of a non-degenerate functional f on A is an element $\sum_i a_i \otimes b_i \in A \otimes A$ such that (cf. [23])

$$\sum_{i} f(xa_{i})b_{i} = x = \sum_{i} a_{i}f(b_{i}x), \quad x \in A.$$
(3.44)

(If K is a field then this just means that $\{b_i\}$ is a K-basis of A and $\{a_i\}$ is its dual basis w.r.t. f.) In other words $\sum_i a_i \otimes b_i$ is simply the expression $\sum_i f_R^{-1}(\beta^i) \otimes b_i$ of the inverse of $f_R: A \to \hat{A}$ as an element of $A \otimes A$. The index of f is then defined by Index $f := \sum_i a_i b_i$ which belongs to Center A. Now let (l, λ) be a dual pair of left integrals. Then the quasibasis of λ is $l_{(2)} \otimes S^{-1}(l_{(1)})$ and

Index
$$\lambda = S^{-1} \circ \sqcap^L(l) \in Z^R$$
. (3.45)

In particular a non-degenerate left integral l is normalized if and only if its dual has index 1.

3.5 2-sided non-degenerate integrals

The space of 2-sided integrals $\mathcal{I}(A) := \mathcal{I}^L(A) \cap \mathcal{I}^R(A)$ in a weak Hopf algebra A is a possibly zero subalgebra of A. The assumption $\mathcal{I}(A) \neq 0$ is independent of the assumption $\mathcal{I}_*^L(A) \neq \emptyset$ since already Hopf algebras provide examples [21] for $\mathcal{I}_*^L(A) \neq 0$ and $\mathcal{I}(A) = 0$. In this Subsection we will make the stronger assumption $\mathcal{I}_*(A) := \mathcal{I}_*^L(A) \cap \mathcal{I}(A) \neq \emptyset$ and study some of the consequences. The main result will be finding a criterion for a WHA to be a symmetric algebra.

At first we observe that if a non-degenerate 2-sided integral j exists then the subspace of 2-sided integrals is obtained from j by the action of the central subalgebra $Z^R = A^R \cap \text{Center } A$,

$$\mathcal{I} = j Z^R, \quad \mathcal{I}_* = j Z^R_{\times} \quad \text{for any} \quad j \in \mathcal{I}_* .$$
(3.46)

As a matter of fact if $i \in \mathcal{I}$ then *i* is a left integral therefore there exists an $x^R \in \mathcal{A}^R$ such that $i = jx^R$. Thus for all $y \in A$ we have $jx^R \sqcap^R (y) = jx^R y = j \sqcap^R (x^R y)$. Since *j* is separating for the right A^R -action, $x^R \sqcap^R (y) = \sqcap^R (x^R y)$. Therefore

$$x^{R}S(y) = x^{R}S(y_{(1)})y_{(2)}S(y_{(3)}) = S(y_{(1)})x^{R}y_{(2)}S(y_{(3)}) = S(1_{(1)}y)x^{R}1_{(2)} = S(y)x^{R}y_{(2)}S(y_{(3)}) = S(1_{(1)}y)x^{R}y_{(2)}S(y_{(3)}) = S(1_{(1)}y)x^{R}y_{(3)}S(y_{(3)}) =$$

hence x^R is central.

Next we recall some facts about "modular automorphisms". Let A be a finite dimensional Frobenius algebra over a field K and let $f: A \to K$ be a non-degenerate functional. Then the modular automorphism of f is defined to be the unique $\theta_f \in \text{Aut } A$ such that

$$f(xy) = f(y\theta_f(x)), \qquad x, y \in A.$$
(3.47)

It is worth to give two other equivalent definitions of θ :

$$f - x = \theta_f(x) \rightarrow f , \qquad x \in A \tag{3.48}$$

or simply

$$\theta_f = f_R^{-1} \circ f_L . aga{3.49}$$

Since any two non-degenerate functionals f and g are related by $g = x \rightarrow f$, with $x \in A_{\times}$, the equivalence class $\theta_A := [\theta_f]$ of θ_f modulo inner automorphisms is independent of the choice of f. If A is a WHA which is Frobenius then one may ask the question whether $\theta_A = [S^2]$.

Definition 3.19 A non-degenerate functional $f: A \to K$ over a WHA A is called a q-trace if $\theta_f = S^2$.

In the term "q-trace" the letter "q" has no individual meaning. One may as well read it as "skew trace" although we do not denie that our motivation came from the theory of q-deformed Hopf algebras.

Lemma 3.20 In a WHA A let l be a non-degenerate left integral. Then S(l) = l if and only if its dual left integral λ is a q-trace.

Proof: $\theta_{\lambda} = S^2$ is equivalent to that the quasibasis of λ satisfies

$$l_{(2)} \otimes S^{-1}(l_{(1)}) = S(l_{(1)}) \otimes l_{(2)} .$$
(3.50)

Applying S to the second tensor factor we obtain $\Delta(l) = \Delta(S(l))$ which yields l = S(l) by the existence of a counit. Q.e.d.

Lemma 3.21 If non-degenerate 2-sided integrals exist then all 2-sided integrals $i \in \mathcal{I}(A)$ are S-invariant, S(i) = i.

Proof: If we can show only that the non-degenerate 2-sided integrals are S-invariant then we are ready since $j = S(j) \in \mathcal{I}_*$ implies $S(jz^R) = S(z^R)j = z^Rj = jz^R$ for all $z^R \in Z^R$.

So let $j \in \mathcal{I}_*$. Then $S(j) \in \mathcal{I}_*$ thus there exists an invertible $z \in Z^R$ such that S(j) = jz. Let λ be the dual of j as a left integral. Then for arbitrary $x \in A$ and for $z^L = S^{-1}(z^{-1})$

$$z^{L}S(x) = z^{L}(\lambda - x) \rightarrow j = (\lambda - x) \rightarrow z^{L}j = (\lambda - x) \rightarrow S^{-1}(j)$$

$$S^{2}(x)z^{-1} = j - \hat{S}^{-1}(\lambda - x) = j_{L} \circ \hat{S}^{-1} \circ \lambda_{L}(x) = \lambda_{R}^{-1} \circ \lambda_{L}(x) =$$

$$= \theta_{\lambda}(x)$$

Therefore $z^{-1} = \theta_{\lambda}(1) = 1$ and j is S-invariant.

Theorem 3.22 The WHA A over K is a symmetric algebra if and only if it has non-degenerate 2-sided integrals and the square of the antipode is an inner automorphism.

Proof: Let A be a symmetric WHA, and $\tau \in \hat{A}$ be a non-degenerate trace. Then there exists a unique $i \in A$ such that $i \rightharpoonup \tau = \hat{1} = \tau \leftarrow i$. We claim that i is a 2-sided integral. As a matter of fact

$$\begin{array}{rcl} xi \rightharpoonup \tau &=& x \rightharpoonup \hat{1} &=& \sqcap^{L}(x) \rightharpoonup \hat{1} &=& \sqcap^{L}(x)i \rightharpoonup \tau \\ \tau \leftharpoonup ix &=& \hat{1} \leftharpoonup x &=& \hat{1} \leftharpoonup \sqcap^{R}(x) &=& \tau \leftharpoonup i \sqcap^{R}(x) \end{array}$$

Q.e.d.

so by non-degeneracy of τ , $i \in \mathcal{I}$. This integral i is also non-degenerate: For any $x^R \in A^R$ one has $ix^R \rightharpoonup \tau = i \rightharpoonup \tau \leftarrow x^R = \hat{1} \leftarrow x^R$, hence i is separating for $\mathcal{I}_{A^R}^L$ so non-degenerate by Scholium 3.17.

The innerness of S^2 in a symmetric algebra follows if we can construct a non-degenerate functional on A the modular automorphism of which is S^2 . By Lemma 3.21 *i* is S-invariant so by Lemma 3.20 χ , the dual left integral to *i*, is such a non-degenerate q-trace.

Conversely, let $S^2 = \operatorname{Ad}_g$ with some $g \in A_{\times}$ and $i \in \mathcal{I}_*$. Denoting the dual left integral of i by χ again, $g^{-1} \rightharpoonup \chi$ is a non-degenerate trace. Q.e.d.

We close this Subsection with a result arising from assuming the existence of non-degenerate 2-sided integrals in both A and \hat{A} . Although the arising structure is reminiscent to that of the "distinguished grouplike element" in Hopf algebra theory it is not a generalization of that.

Proposition 3.23 Let A be a WHA and assume that both $\mathcal{I}_*(A)$ and $\mathcal{I}_*(\hat{A})$ are non-empty. Then S^4 is inner and the square of θ_A is the identity in Out A. Moreover and more explicitly, for $\hat{h} \in \mathcal{I}_*(\hat{A})$ there exist invertible elements $a_L \in A^L$ and $\alpha_L \in \hat{A}^L$ such that, with the notations $a_R = S(a_L)$ and $\alpha_R = \hat{S}(\alpha_L)$, we have

$$Ad_{a_L a_R^{-1}} = S^4 (3.51)$$

$$Ad_{a_L a_R} = \theta_{\hat{h}}^2 \tag{3.52}$$

$$a_L a_R^{-1} \rightharpoonup \psi \leftharpoonup a_L a_R^{-1} = \alpha_L \alpha_R \psi \alpha_R^{-1} \alpha_L^{-1}, \quad \psi \in \hat{A} .$$

$$(3.53)$$

Proof: Choose $h \in \mathcal{I}_*(A)$ and $\hat{h} \in \mathcal{I}_*(\hat{A})$ and let λ be the dual of h and l be that of \hat{h} , as left integrals. Define

$$a_L = \hat{h} \rightharpoonup h$$
, $\alpha_L = h \rightharpoonup \hat{h}$. (3.54)

Then

$$\hat{1} - a_L = \langle \hat{1}_{(1)} \hat{h}, h \rangle \hat{1}_{(2)} = \langle \hat{h}_{(1)}, h \rangle \hat{h}_{(2)} \hat{S}(\hat{h}_{(3)}) = \hat{\sqcap}^L (\hat{h} - h) = \hat{S}(\hat{h} - h) = \alpha_L$$
(3.55)

and introducing a_R and α_R as above

$$la_L = l \leftarrow \alpha_R = l \leftarrow (\hat{h} \leftarrow h) = h(l \leftarrow \hat{h}) = h(\hat{S}^2(\hat{h}) \rightharpoonup l) = h$$
(3.56)

where q-trace property of l and \hat{S} -invariance of \hat{h} have been used. Similarly,

$$\alpha_R \rightharpoonup l = h = la_R \tag{3.57}$$

$$a_R \rightharpoonup \lambda = \hat{h} = \lambda \alpha_R \tag{3.58}$$

$$\lambda \leftarrow a_R = \hat{h} = \lambda \alpha_L. \tag{3.59}$$

Non-degeneracy of \hat{h} and h now imply invertibility of a_L , a_R , α_L , and α_R . Hence Eq(3.53) readily follows.

We can now compute the modular automorphism of \hat{h} using the information $\theta_{\lambda} = S^2$.

$$\hat{h} \leftarrow x = a_R \rightharpoonup \lambda \leftarrow x = a_R \theta_\lambda(x) \rightharpoonup \lambda \quad \Rightarrow \quad \theta_{\hat{h}} = \operatorname{Ad}_{a_R} \circ S^2 \tag{3.60}$$

Computing $\hat{S}(\hat{h} \leftarrow x)$ in two different ways

$$\begin{split} \hat{S}(\hat{h} \leftarrow x) &= S^{-1}(x) \rightharpoonup \hat{h} = \hat{h} \leftarrow \theta_{\hat{h}}^{-1}(S^{-1}(x)) \\ &= \hat{S}(\theta_{\hat{h}}(x) \rightharpoonup \hat{h}) = \hat{h} \leftarrow S^{-1}(\theta_{\hat{h}}(x)) \end{split}$$

yields

$$S^{-1} \circ \theta_{\hat{h}} = \theta_{\hat{h}}^{-1} \circ S^{-1} \tag{3.61}$$

Q.e.d.

and finally

$$\operatorname{Ad}_{a_L} \circ S^{-2} = \theta_{\hat{h}} = \operatorname{Ad}_{a_R} \circ S^2$$
(3.62)

from which (3.51) and (3.52) follow immediately.

3.6 Haar integrals

Since finite dimensional weak Hopf algebras do not go beyond the "compact" and "discrete" case, the following very conservative definition of Haar measure will suffice.

Definition 3.24 An element h of a WHA A is called a Haar integral in A or Haar measure on \hat{A} if h is a normalized 2-sided integral, i.e. $h \in \mathcal{I}(A)$ and $\Box^{L}(h) = \Box^{R}(h) = 1$.

Obviously, if Haar integral exists then it is a unique S-invariant idempotent. As a matter of fact let h and h' be Haar integrals. Then $h' = \sqcap^L(h)h' = hh' = h \sqcap^R(h') = h$. In particular $h^2 = h$. S-invariance follows from uniqueness since S(h) is a always a Haar integral if h is.

In finding criteria for the existence of Haar measure in A an important role will be played by a special element $\chi \in \hat{A}$ the definition of which was inspired by similar computations in Hopf algebra theory [21]:

$$\chi := \sum_{i} \beta^{i} - S^{-2}(b_{i}) \equiv \hat{L}'(\hat{1}), \qquad (3.63)$$

where $\{b_i\}$ and $\{\beta^i\}$ are dual bases of A and \hat{A} , respectively, and $\hat{L}': \hat{A} \to \hat{A}$ is given by $\hat{L}'(\psi) := \sum_i \beta^i \psi \leftarrow S^{-2}(b_i)$. Note that \hat{L}' is the "cop" version of the dual analogue \hat{L} of the projection (3.23) onto the space of left integrals. Hence χ is a left integral in $(\hat{A})_{cop}$ and therefore in \hat{A} .

As we will see below if χ is non-degenerate and a q-trace then its dual left integral will automatically be the Haar measure. In order to see that it is a q-trace let Tr_A be the standard trace on $\operatorname{End}_K A$ and introduce the notation $Q_-(x)y := yx$. Then for $x \in A$ we have

$$\begin{split} \chi(x) &= \sum_{i} \langle \beta^{i}, S^{-2}(b_{i})x \rangle = \operatorname{Tr}_{A} Q_{-}(x) \circ S^{-2} \\ \chi(xy) &= \operatorname{Tr}_{A} Q_{-}(y) \circ Q_{-}(x) \circ S^{-2} = \operatorname{Tr}_{A} Q_{-}(y) \circ S^{-2} \circ Q_{-}(S^{2}(x)) = \\ &= \chi(yS^{2}(x)). \end{split}$$

The next Lemma will be crucial in deciding whether χ is non-degenerate.

Lemma 3.25 Let l be a left integral in a WHA A and let $\chi \in \hat{A}$ be the q-trace left integral defined in Eq(3.63). Then

$$l \rightharpoonup \chi = \hat{S}^2(\hat{1} \leftarrow l) . \tag{3.64}$$

Proof: Using the q-trace property of χ and then (2.19a)

$$\begin{split} l &\rightharpoonup \chi &= \sum_{i} \beta^{i} - S^{-2}(b_{i}l) = (\hat{1} - 1_{(1)}) - S^{-2}(1_{(2)}l) = \\ &= \langle \hat{1}_{(1)}, S^{-1}(1_{(2)}) 1_{(1)} \rangle \hat{1}_{(2)} - S^{-2}(l) = \hat{1} - S^{-2}(l) = \hat{S}^{2}(\hat{1} - l) \end{split}$$

$$Q.e.d.$$

Proposition 3.26 Let A be a weak Hopf algebra over a field K and let χ be given by (3.63).

- i) The Haar integral $h \in A$ exists if and only if χ is non-degenerate, in which case (h, χ) is a dual pair of left integrals. In particular Haar integrals are non-degenerate.
- ii) A left integral $l \in \mathcal{I}^L(A)$ is a Haar integral if and only if $\sqcap^R(l) = 1$.

The characterization of Haar measures under (ii) is so simple that it could be well used as a definition of Haar measure. Notice that in that case the formal difference between the notions of normalized left integral and Haar measure were so tiny (change \sqcap^L for \sqcap^R) that it would smear out the big conceptual difference: The existence of normalized left integrals is equivalent to semisimplicity while the existence of Haar measures is much stronger.

Proof: ii) Assume $l \in \mathcal{I}^{L}(A)$ satisfies $\sqcap^{R}(l) = 1$. Then by Lemma 3.25 $l \rightarrow \chi = \hat{1}$. Therefore the duality Theorem (Thm 3.18) implies that (l, χ) is a dual pair of non-degenerate left integrals. Since χ is a q-trace, Lemma 3.20 shows that l is an S invariant non-degenerate left integral. Furthermore $\sqcap^{L}(l) = \sqcap^{L}(S(l)) = S \circ \sqcap^{R}(l) = 1$. Thus l is a Haar integral. Now assume h is a Haar integral. Then obviously h is a left integral satisfying $\sqcap^{R}(h) = 1$.

i) The "only if" part follows from the proof of (ii). Assume χ is non-degenerate and let h be its dual left integral. Then by Lemma 3.20 h is 2-sided and by Lemma 3.25 it is normalized. *Q.e.d.*

However simple, the criteria of the above Proposition are very difficult to verify in concrete situations. So it is worth looking for other criteria even if they are not applicable in full generality.

Theorem 3.27 Let A be a WHA over an algebraically closed field K. Then a necessary and sufficient condition for the existence of Haar measure $h \in A$ is that A is semisimple and there exists a $g \in A_{\times}$ such that $gxg^{-1} = S^2(x)$ for $x \in A$ and $tr D_r(g^{-1}) \neq 0$ for all irreducible representation D_r of A.

The assumption on K is used only to ensure that A is split semisimple, $A = \bigoplus_r M_{n_r}(K)$, once knowing that it is semisimple. In particular there will be a K-basis $\{e_r^{\alpha\beta}\}$ for A obeying matrix unit relations.

Proof: Sufficiency: Let $\tau: A \to K$ be the trace with trace vector $\tau_r = \operatorname{tr} D_r(g^{-1})$. Then τ is non-degenerate and has as quasibasis the element

$$\sum_{i} x_i \otimes y_i := \sum_{r} \frac{1}{\tau_r} \sum_{\alpha,\beta=1}^{n_r} e_r^{\alpha\beta} \otimes e_r^{\beta\alpha}$$

Notice that $\sum_i x_i g^{-1} y_i = 1$. Now we define $\chi' := g \rightharpoonup \tau$ and claim that χ' coincides with the χ of Eq(3.63). As a matter of fact

$$\begin{split} \chi'(x) &= \tau(gx) = \sum_{i} \tau(x_i g^{-1} y_i gx) = \sum \tau(x_i S^{-2}(y_i) x) = \\ &= \sum_{i} \langle \beta_i, S^{-2}(b_i) x \rangle = \chi(x) \end{split}$$

where we used the fact that the dual of the basis $b_i = y_i$ is $\beta_i = \tau \leftarrow x_i$. Since χ' was nondegenerate by construction, we conclude that the χ of Eq(3.63) is non-degenerate and therefore its dual left integral l has $\sqcap^R(l) = 1$ by Eq(3.64). Therefore l is a Haar measure.

Necessity: If $h \in A$ is a Haar measure then A is semisimple by Theorem 3.13. Therefore A is a symmetric algebra and $\theta_A = \text{id}$. This means that θ_{ψ} is inner for all non-degenerate

functional ψ . In particular $\theta_{\chi} = S^2$ is inner where χ is the dual left integral of h. Choose a $g \in A_{\times}$ implementing S^2 and construct the non-degenerate trace $\tau := g^{-1} \rightharpoonup \chi$.

$$\tau(x) = \chi(xg^{-1}) = \operatorname{Tr}_A Q_-(xg^{-1}) \circ S^{-2} = \operatorname{Tr}_A Q_+(g^{-1}) \circ Q_-(x)$$

where $Q_+(x)y := xy$ is left multiplication on A. Choosing a matrix unit basis to evaluate the trace we obtain

$$\tau(x) = \sum_r \operatorname{tr} D_r(g^{-1}) \operatorname{tr} D_r(x)$$

and by non-degeneracy of τ all components tr $D_r(g^{-1})$ of the trace vector are non-vanishing. *Q.e.d.*

4 C*-Weak Hopf Algebras

In this Section we introduce the C^* -structure in WHAs which is inevitable if WHAs are to be used as symmetries of inclusions of von Neumann algebras, in particular in quantum field theory. Utilizing the results of Sections 2 and 3 we establish the existence of two canonical elements in any C^* -WHA, the Haar measure h and the canonical grouplike element g. While the Haar measure is well known for C^* -Hopf algebras, the canonical grouplike element cannot be recognized in finite dimensional Hopf algebras because it is always equal to 1. This is related to involutivity of the antipode in finite dimensional C^* -Hopf algebras [24]. The very fact that C^* -WHAs can have non-involutive antipodes provides the sufficient flexibility for the emergence of non-integer dimensions.

4.1 First consequences of the C*-structure

Definition 4.1 We define a *-WHA as a WHA $(A, 1, \Delta, \varepsilon, S)$ over the complex numbers \mathbb{C} together with an antilinear involution * such that

- i) (A, *) is a *-algebra,
- ii) Δ is a *-algebra map, i.e. $(x^*)_{(1)} \otimes (x^*)_{(2)} = (x_{(1)})^* \otimes (x_{(2)})^*$ for all $x \in A$.

By uniqueness of the unit, counit, and the antipode (see Lemma 2.8) we have the following additional relations in a *-WHA.

$$1^* = 1, \quad \varepsilon(x^*) = \overline{\varepsilon(x)}, \quad S(x^*)^* = S^{-1}(x).$$
 (4.1)

Now it is easy to check that the projections \sqcap^L and \sqcap^R satisfy

$$\Box^{L}(x)^{*} = \Box^{L}(S(x)^{*}), \qquad \Box^{R}(x)^{*} = \Box^{R}(S(x)^{*})$$
(4.2)

therefore A^L and A^R are *-subalgebras of A. As an elementary exercise we obtain selfduality of the *-WHA:

Scholium 4.2 Let A be a *-WHA and define a star operation on its dual as follows

$$\langle \varphi^*, x \rangle =: \overline{\langle \varphi, S(x)^* \rangle} .$$
 (4.3)

Then \hat{A} with this star operation becomes a *-WHA.

For a *-WHA A the canonical isomorphisms $\kappa_A^L: A^L \to \hat{A}^R$ and $\kappa_A^R: A^R \to \hat{A}^L$ of Lemma 2.6 become *-algebra isomorphisms.

We omit the discussion of further properties of *-WHAs and turn to the most important case of C^* -WHAs.

Definition 4.3 A^* -WHA A possessing a faithful *-representation is called a C*-weak Hopf algebra, or C*-WHA for short.

Being a finite dimensional C^* -algebra any C^* -WHA can be uniquely characterized, as an algebra, by the dimensions $n_r \in \mathbb{N}$ of its blocks where r is running over the finite set Sec A of equivalence classes of irreducible representations (i.e. the sectors) of A.

$$A \cong \bigoplus_{r \in \mathcal{S}ec \, A} M_{n_r} , \qquad M_{n_r} = \operatorname{Mat}(n_r, \mathbb{C}) .$$

$$(4.4)$$

 A^L and A^R are unital *-subalgebras therefore they are C*-algebras as well and we have natural numbers n_a , $a \in Sec A^L$ and n_b , $b \in Sec A^R$ characterizing the type of A^L and A^R , respectively.

$$A^c \cong \bigoplus_{a \in Sec A^c} M_{n_a} , \qquad c = L, R .$$

$$(4.5)$$

The antiisomorphism $S: A^L \to A^R$ establishes a bijection $a \mapsto \bar{a}$ of the blocks of A^L to the blocks of A^R such that $n_{\bar{a}} = n_a$. (We consider Sec A^L , Sec A^R , and Sec A as disjoint sets which allows to use one function n.)

The following elementary but important Proposition will be the basic ingredient in proving both the existence of Haar measures and rigidity of the representation category of C^* -WHAs.

Proposition 4.4 Let A be a finite dimensional C^* -algebra and $S: A \to A^{op}$ an algebra isomorphism such that $(* \circ S)^2 = id_A$. Then there exists $g \in A_{\times}$ such that

- i) $g \ge 0$
- ii) $gxg^{-1} = S^2(x), x \in A$
- iii) $tr_r(g) = tr_r(g^{-1}), r \in Sec A$
- iv) $S(g) = g^{-1}$

where tr_r denotes trace in the irreducible representation D_r . An element $g \in A$ satisfying only the first three properties is already unique.

Proof: The restriction $S|_{\text{Center }A}$ is an algebra automorphism therefore acts on the minimal central idempotents e_r as $S(e_r) = e_{\bar{r}}$ where $r \mapsto \bar{r}$ is a permutation of Sec A. Since $e_r^* = e_r$ and $* \circ S$ is an involution, $r \mapsto \bar{r}$ is an involution.

Choose matrix units $\{e_r^{\alpha\beta}\}$ for the C^* -algebra A and define the antiautomorphism $S_0: A \to A$ by $S_0(e_r^{\alpha\beta}) := e_{\overline{r}}^{\beta\alpha}$. Then $S_0^2 = \operatorname{id}_A$ and $* \circ S_0 = S_0 \circ *$. Since $S \circ S_0$ is an automorphism of Athat acts as the identity on the centre, there exists $C \in A$ invertible such that $S = \operatorname{Ad}_C \circ S_0$. It follows that

$${}^{*} \circ S(x) = C^{-1*} S_0(x^*) C^*, \quad ({}^{*} \circ S)^2(x) = C^{-1*} S_0(C^{-1}) x S_0(C) C^* = x, \tag{4.6}$$

therefore $S_0(C)C^*$ is central and so is its adjoint $K := CS_0(C^*) = S(C^*)C$.

$$S^{2}(x) = CS_{0}(CS_{0}(x)C^{-1})C^{-1} = CS_{0}(C^{-1})xS_{0}(C)C^{-1}, \quad x \in A$$
(4.7)

hence $T := CS_0(C^{-1}) = CC^*[S_0(C)C^*]^{-1} = CC^*K^{-1*}$ implements S^2 and its polar decomposition takes the form

$$T = ug', \qquad u = K^{-1*} (K^* K)^{1/2}, \quad g' = C (K^* K)^{-1/2} C^*.$$
 (4.8)

Using centrality of the unitary part and the computations $S(T) = S(C)S^2(C^{-1}) = S(C)TC^{-1}T^{-1} = T^{-1}$ and $S(K) = S_0(K) = C^*S_0(C) = S_0(C)C^* = K^*$ we obtain that g' is positive invertible, implements S^2 , and satisfies $S(g') = g'^{-1}$. These latter three properties, however, do not fix g completely. If c is positive, central, and satisfies $S(c) = c^{-1}$ then g = g'c will also satisfy the above three properties. Now defining

$$g := g'c$$
 where $c = \sum_{r} e_r \left(\frac{\operatorname{tr}_r(g'^{-1})}{\operatorname{tr}_r(g')}\right)^{1/2}$ (4.9)

it is easy to verify that g obeys (i–iv) of the Proposition. If $f \in A$ satisfies only (i), (ii), and (iii) then f = gc where c is positive invertible, central, and satisfies $D_r(c) = D_r(c)^{-1}$ for all irrep D_r . Hence c = 1, proving uniqueness of g. Q.e.d.

4.2 The Haar measure and selfduality

Recall that the Haar measure in a WHA A has been defined in Definition 3.24 as the unique element $h \in A$ making the integral $\int \varphi := \langle \varphi, h \rangle$ of a function $\varphi : A \to \mathbb{C}$ to be a non-degenerate functional invariant under left and right translations and normalized according to $\int \varphi^L = \hat{\varepsilon}(\varphi^L)$ for $\varphi^L \in \hat{A}^L$. The sufficient conditions for its existence given by Theorem 3.27 will be used here to prove the next Theorem.

Theorem 4.5 In a C^{*}-WHA A Haar measure $h \in A$ exists. It is selfadjoint, $h^* = h$, and such that

$$(\varphi,\psi) := \langle \varphi^*\psi, h \rangle, \qquad \varphi,\psi \in \hat{A} , \qquad (4.10)$$

is a scalar product on \hat{A} making \hat{A} a Hilbert space and making the left regular module ${}_{\hat{A}}\hat{A}$ a faithful *-representation of the *-WHA \hat{A} . Thus \hat{A} is a C*-WHA, too.

Proof: A being a finite dimensional C^* -algebra is semisimple. By Proposition 4.4 there exists a g implementing S^2 . This g was shown to be positive and invertible, hence tr $D_r(g^{-1}) > 0$ for all $r \in Sec A$. Therefore all the conditions of Theorem 3.27 are satisfied and Haar measure h exists.

Since h is non-degenerate, (,) is a non-degenerate sesquilinear form on \hat{A} . So it remains to show positivity. By the equality

$$(\psi,\psi) = \langle \psi^*\psi, h \rangle = \overline{\langle \psi, S(h_{(1)})^* \rangle} \langle \psi, h_{(2)} \rangle$$
(4.11)

positivity of (,) follows if we can show that $(S \otimes id) \circ \Delta(h)$ belongs to the positive cone

$$\mathcal{P} = \left\{ \sum_{k} a_{k}^{*} \otimes a_{k} \mid a_{k} \in A \right\} \subset A \otimes A .$$

$$(4.12)$$

Q.e.d.

Therefore the next Lemma will complete the proof.

Lemma 4.6 Choose matrix units $\{e_q^{\alpha\beta}\}$ for A and let g denote the element determined in Proposition 4.4. If furthermore $\sum_i x_i \otimes y_i$ is the quasibasis of the trace $\tau: A \to \mathbb{C}$ with trace vector $\tau_q = tr_q(g^{-1})$ then

$$S(h_{(1)}) \otimes h_{(2)} = \sum_{i} x_{i} \otimes g^{-1} y_{i} = \sum_{q \in Sec \ A} \frac{1}{\tau_{q}} \sum_{\alpha\beta} e_{q}^{\alpha\beta} g^{-1/2} \otimes g^{-1/2} e_{q}^{\beta\alpha}$$
(4.13)

$$h_{(1)} \otimes S(h_{(2)}) = \sum_{i} x_{i}g \otimes y_{i} = \sum_{q \in \mathcal{S}ec \ A} \frac{1}{\tau_{q}} \sum_{\alpha\beta} e_{q}^{\alpha\beta} g^{1/2} \otimes g^{1/2} e_{q}^{\beta\alpha}$$
(4.14)

Proof: The quasibasis of $\chi = g \rightharpoonup \tau$ is $\sum x_i g^{-1} \otimes y_i$ and since χ is the dual left integral of h, this quasibasis is equal to $h_{(2)} \otimes S^{-1}(h_{(1)})$. This implies the first row. By property **iii**) of Proposition 4.4 τ is an S-invariant trace, therefore its quasibasis can also be written as $\sum_i y_i \otimes x_i = \sum_i S^{-1}(x_i) \otimes S^{-1}(y_i)$. Thus the second row follows from the first. Q.e.d. From now on $h \in A$ will always denote the Haar measure of A and $\hat{h} \in \hat{A}$ that of \hat{A} .

Lemma 4.7 In a C^{*}-WHA A the counit is a positive linear functional, $\varepsilon(x^*x) \ge 0$, $x \in A$.

Proof:

$$\varepsilon(x^*x) = \varepsilon(x^*1_{(1)})\varepsilon(1_{(2)}1_{(2')})\varepsilon(1_{(1')}x) = \varepsilon(\sqcap^L(x)^*\sqcap^L(x)) = \langle \hat{h}, \sqcap^L(x)^*\sqcap^L(x) \rangle \ge 0,$$

where we have used $\hat{h}|_{A^L} = \varepsilon|_{A^L}$, which follows from $\langle \hat{h}, x^L \rangle = \langle \hat{\sqcap}^L(\hat{h}), x^L \rangle = \langle \hat{1}, x^L \rangle$ for all $x^L \in A^L$.

A being semisimple the trivial representation V_{ε} decomposes into irreducibles V_q each of them with multiplicity 1 by Proposition 2.15. The sectors $q \in Sec A$ occuring in V_{ε} with non-zero multiplicity will be called *vacuum sectors*.

$$V_{\varepsilon} \cong \bigoplus_{q \in \mathcal{V}ac \ A} V_q \ . \tag{4.15}$$

By Proposition 2.15 there is a bijection $q \mapsto z_q^L$ from the set $\mathcal{V}ac A$ of vacuum sectors to the set of minimal projections in Z^L such that, with $z_q^R := S(z_q^L)$, we have

$$D_{\varepsilon}(z_q^L) = D_{\varepsilon}(e_q) = D_{\varepsilon}(z_q^R)$$
(4.16)

$$z_q^L = \sqcap^L(e_q) \qquad \qquad \sqcap^R(e_q) = z_q^R \tag{4.17}$$

where e_q denotes the minimal central projection in A supporting the irreducible vacuum representation D_q .

Lemma 4.8 $D_r(h)$ is a 1-dimensional projection for $r \in Vac A$ and $D_r(h) = 0$ if r is not a vacuum sector. The algebra of 2-sided integrals is generated by minimal projections h_q

$$\mathcal{I}(A) = hAh = \operatorname{Span}\{h_q \mid q \in \operatorname{Vac} A\}, \qquad h_q = he_q .$$
(4.18)

The non-degenerate 2-sided integrals are presisely the invertible elements: $\mathcal{I}_*(A) = \mathcal{I}(A)_{\times}$.

Proof: If $D_r(h) \neq 0$ then pick up a non-zero vector v_r from the subspace $D_r(h)V_r$ of the irreducible A-module V_r and define

$$T: A^L \to V_r$$
, $Tx^L := D_r(x^L)v_r$. (4.19)

This map is a non-zero left A-module map if we equip A^L with the structure of the trivial A-module ${}_AA^L$ introduced in Lemma 2.12. Indeed,

$$D_r(x)Tx^L = D_r(xx^Lh)v_r = D_r(\sqcap^L(xx^L)h)v_r = T \sqcap^L(xx^L) .$$
(4.20)

Therefore $r \in \operatorname{Vac} A$. This proves that $D_r(h) = 0$ for $r \notin \operatorname{Vac} A$.

Now let the Haar integral act on the trivial left A-module ${}_A \hat{A}^R$.

$$D_{\varepsilon}(h)\varphi^{R} = h \rightharpoonup \varphi^{R} \in \hat{A}^{L} \cap \hat{A}^{R} \equiv \hat{Z} .$$

$$(4.21)$$

Thus $D_{\varepsilon}(h): \hat{A}^R \to \hat{Z}$ is a projection, onto. If z^L is a minimal projection in Z^L then $z^L \to \hat{1}$ is a minimal projection in \hat{Z} by Lemma 2.14. Hence $D_{\varepsilon}(z^Lh)$ maps \hat{A}^R onto $(z^L \to \hat{1})\hat{Z} \cong C$. This proves that $D_{\varepsilon}(z^Lh)$, the restriction of which is precisely $D_q(h)$ for some $q \in \mathcal{V}ac A$, is a 1-dimensional projection. If $i \in \mathcal{I}$ then by the 2-sided normalization of h one can write i = hih. Conversely, hxh is a 2-sided integral for all $x \in A$. This proves the remaining assertions. Q.e.d.

The Haar measure provides conditional expectations

$$E^L: A \to A^L, \qquad E^L(x) = \hat{h} \rightharpoonup x$$

$$(4.22)$$

$$E^R: A \to A^R, \qquad E^R(x) = x - \hat{h}$$

$$(4.23)$$

As a matter of fact by Lemma 3.2.c) the image of E^L is in A^L since \hat{h} is a left integral. E^L is unit preserving since \hat{h} is normalized. Finally, E^L is positive since \hat{h} is positive and Δ is a *-algebra map.

4.3 The canonical grouplike element

In this Subsection we investigate further properties of the element g of Proposition 4.4. We show that it is always a product of left and right elements, implying its grouplikeness immediately, and obtain expressions for the modular automorphisms of the Haar measures of A and \hat{A} .

Proposition 4.9 In a C^* -WHA A there exists a unique $g \in A$ such that

- i) $g \ge 0$ and invertible,
- ii) $gxg^{-1} = S^2(x)$ for all $x \in A$,
- iii) $h_{(2)} \otimes h_{(1)} = h_{(1)} \otimes gh_{(2)}g$.

Proof: Existence: Let g be the (unique) element defined by the conditions of Proposition 4.4. As in the proof of Lemma 4.6 let τ be the S-invariant trace with trace vector $\tau_q = \operatorname{tr}_q(g)$ and $\sum x_i \otimes y_i$ be its quasibasis. Then

$$h_{(2)} \otimes h_{(1)} = \sum_{i} x_i \otimes S(g^{-1}y_i) = \sum_{i} S^{-1}(y_i) \otimes x_i g =$$
(4.24)

$$= \sum_{i} S^{-1}(y_i g^{-1}) \otimes g x_i g = \sum_{i} g S^{-1}(y_i) \otimes g x_i g =$$
(4.25)

$$= \sum_{i} S(g^{-1}y_{i}) \otimes gx_{i}g = h_{(1)} \otimes gh_{(2)}g$$
(4.26)

Uniqueness: Let g and g' satisfy i), ii), and iii). Then g' = gc with c central, positive, and invertible. Furthermore, since iii) is equivalent to

$$\langle \varphi \psi, h \rangle = \langle \psi(g \rightharpoonup \varphi \leftharpoonup g), h \rangle, \qquad (4.27)$$

non-degeneracy of h implies

$$g' \rightharpoonup \varphi \leftarrow g' = g \rightharpoonup \varphi \leftarrow g , \qquad \varphi \in \hat{A} .$$
 (4.28)

Therefore $c^2 \rightharpoonup \varphi \equiv c \rightharpoonup \varphi \leftarrow c = \varphi$ for all $\varphi \in \hat{A}$. Thus $c^2 = 1$ and, by positivity, c = 1. Q.e.d.

Notice that property **iii**) of Proposition 4.9 is equivalent to that the modular automorphism of the Haar functional $\varphi \mapsto \varphi(h)$ is expressible in the form

$$\theta_h(\psi) = g \rightharpoonup \psi \leftharpoonup g , \qquad \psi \in \hat{A} .$$
(4.29)

Definition 4.10 Let A be a C^* -weak Hopf algebra. Then the unique element $g \in A$ determined either by the conditions of Proposition 4.4 or by the conditions of Proposition 4.9 is called the canonical grouplike element of A.

As one may suspect the canonical grouplike element is grouplike in the sense of

Definition 4.11 An element x of a WHA A is called grouplike if

$$\Delta(x) = x \mathbf{1}_{(1)} \otimes x \mathbf{1}_{(2)} = \mathbf{1}_{(1)} x \otimes \mathbf{1}_{(2)} x \tag{4.30}$$

$$S(x)x = 1$$
. (4.31)

We remark that if (4.30) holds then condition (4.31) is equivalent to the assumption that x is invertible. One should emphasize that grouplike elements are not always like group elements if a *-operation is present. Namely we allow for x not to be unitary. Thus there can be positive grouplike elements, for example, in a C^* -WHA.

If x is an invertible element factorizable as $x_L x_R^{-1}$ with $x_L \in A^L$ and $x_R = S(x_L) = S^{-1}(x_L)$ then x is automatically grouplike. As a matter of fact $\Delta(x) = x_L \mathbf{1}_{(1)} \otimes x_R^{-1} \mathbf{1}_{(2)} = x x_R \mathbf{1}_{(1)} \otimes x_R^{-1} \mathbf{1}_{(2)} = x \mathbf{1}_{(1)} \otimes x \mathbf{1}_{(2)}$. Now it follows from the next Lemma that the canonical grouplike element g is grouplike.

Lemma 4.12 In a weak C^* -Hopf algebra A the elements $h \leftarrow \hat{h}$ and $\hat{h} \rightarrow h$ are positive and invertible. The canonical grouplike element of A can be factorized as

$$g = g_L g_R^{-1} , \qquad where \qquad (4.32)$$

$$g_L := (\hat{h} \rightarrow h)^{1/2} , \qquad g_R = (h - \hat{h})^{1/2}$$

$$(4.33)$$

Proof: $\hat{h} \rightarrow h = E^L(h) = E^L(h^*h) \ge 0$ and similarly $h \leftarrow \hat{h} \ge 0$ by positivity of the conditional expectations (4.22). Invertibility follows from the existence of the dual left integral χ since $(h \leftarrow \hat{h}) \rightarrow \chi = \hat{S}(\hat{h})(h \rightarrow \chi) = \hat{h}$ can hold for the non-degenerate χ and \hat{h} only if $h \leftarrow \hat{h}$ is invertible. Thus $\hat{h} \rightarrow h = S(h \leftarrow \hat{h})$ is invertible, too.

The next point is to observe that the three elements $\hat{h} \rightarrow h$, $h \leftarrow \hat{h}$, and g commute with each other. For g and any one of the others this follows from the fact that $\hat{h} \rightarrow h$ and $h \leftarrow \hat{h}$ are invariant under S^2 . For the commutativity of the remaining two notice that one of them belongs to A^L the other to A^R . Now compare the following expressions:

$$\hat{h} = (h \leftarrow \hat{h}) \rightharpoonup \chi = (h \leftarrow \hat{h})g \rightharpoonup \tau ,$$

$$(4.34)$$

$$\hat{h} = \hat{S}^{-1}(\hat{h}) = \tau \leftharpoonup g^{-1}(\hat{h} \rightharpoonup h) = g^{-1}(\hat{h} \rightharpoonup h) \rightharpoonup \tau .$$

$$(4.35)$$

By non-degeneracy of τ we obtain

$$\begin{array}{rcl} (\hat{h} \rightharpoonup h)g^{-1} &=& (h \leftharpoonup \hat{h})g\\ (\hat{h} \rightharpoonup h)(h \leftharpoonup \hat{h})^{-1} &=& g^2 \end{array}$$

and taking the (positive) square root the Lemma is proven. Q.e.d.

Lemma 4.13 The left-right components of the canonical grouplike element g of A and \hat{g} of \hat{A} obey the following identities.

$$\hat{g}_L = \hat{1} - g_L = \hat{1} - g_R \qquad g_L = 1 - \hat{g}_L = 1 - \hat{g}_R \qquad (4.36)$$

$$\hat{g}_R = g_R \rightharpoonup 1 = g_L \rightharpoonup 1$$
 $g_R = \hat{g}_R \rightharpoonup 1 = \hat{g}_L \rightharpoonup 1$ (4.37)

$$S(g_L) = g_R = S^{-1}(g_L)$$
 $S(\hat{g}_L) = \hat{g}_R = S^{-1}(\hat{g}_L)$ (4.38)

Proof: Since $g_L \in A^L$ and $g_R \in A^R$, they commute and both of them are invariant under $S^2 = \operatorname{Ad}_q$. So are the C^{*}-algebras generated by each of them, pointwise. Hence $S(g_L^{1/2})^* =$ $S^{-1}(g_L^{1/2}) = S(g_L^{1/2})$ therefore $S(g_L) = S(g_L^{1/2})^2 \ge 0$. On the other hand $S(g_L)^2 = S(g_L^2) = g_R^2$, therefore $S(g_L)$ is the positive square root of g_R^2 , i.e. $S(g_L) = g_R$.

Next we want to show that $\hat{1} \leftarrow (\hat{h} \rightarrow h) = \hat{h} \rightarrow \hat{h}$. Since both hand sides belong to \hat{A}^L , the identity

$$\begin{split} \langle \hat{1} \leftarrow (\hat{h} \rightarrow h), \, x^R \rangle &= \varepsilon((\hat{h} \rightarrow h)S(x^R)) = \varepsilon(\hat{h} \rightarrow hS(x^R)) = \\ &= \langle \hat{h} \leftarrow h, \, S(x^R) \rangle = \langle h \rightarrow \hat{h}, \, x^R \rangle \;, \end{split}$$

valid for $x^R \in A^R$, suffices. Therefore $\hat{1} \leftarrow g_L^2 = \hat{g}_L^2$, or $\hat{1} \leftarrow g_R^2 = \hat{g}_L^2$. Now use the fact that $A^R \ni x^R \mapsto (\hat{1} \leftarrow x^R) \in \hat{A}^L$ is a *-algebra isomorphism. Hence passing to the square roots we obtain $1 \leftarrow g_R = \hat{g}_L$. All the remaining identities are simple consequences of this. Q.e.d.

Proposition 4.14 Let A be a C^{*}-WHA with dual \hat{A} and let $h \in A$, $\hat{h} \in \hat{A}$ be the corresponding Haar measures. Then

- i) the modular automorphism of the Haar functional \hat{h} is implemented by $g_L g_R$, i.e. for all $x \in A$ we have $\theta_{\hat{h}}(x) = g_L g_R x g_R^{-1} g_L^{-1}$;
- ii) the dual left integral of h can be expressed as $\chi = \hat{h}\hat{g}_{B}^{-2}$;
- iii) the S-invariant trace functional $\tau = g^{-1} \rightharpoonup \chi$ and the Haar functional \hat{h} are related by

$$\tau = \hat{g}_L^{-1} \hat{h} \hat{g}_R^{-1} \tag{4.39}$$

$$\hat{h} = g_L g_R \rightharpoonup \tau \tag{4.40}$$

Proof: i): Using identities like $\hat{g}_L \rightarrow x = g_R x$,...etc, which follow from Scholium 2.7, one can easily verify $\hat{g} \rightharpoonup x \leftarrow \hat{g} = g_L g_R x g_R^{-1} g_L^{-1}$, for $x \in A$.

ii): The identity $\hat{h} \rightarrow h = g_L^2 = 1 \leftarrow \hat{g}_R^2$ implies $1 = \hat{h} \rightarrow h \leftarrow \hat{g}_R^{-2} = \hat{h} \rightarrow hg_L^{-2}$, hence $hg_L^{-2} = hg_R^{-2}$ is the dual left integral of \hat{h} . By duality, $\hat{h}\hat{g}_R^{-2}$ is the dual left integral χ of h. iii): $\tau = g^{-1} \rightarrow \hat{h}\hat{g}_R^{-2} = \hat{g}_R^{-1}(\hat{h}\hat{g}_R^{-2})\hat{g}_R = \hat{g}_L^{-1}\hat{h}\hat{g}_R^{-1}$ and $\tau = g^{-1} \rightarrow (g_R^{-2} \rightarrow \hat{h}) = g_L^{-1}g_R^{-1} \rightarrow \hat{h}$

completes the proof. Q.e.d.

Cyclicity and separability of the vector h in the right $A^{L,R}$ -module \mathcal{I}^{L} (cf. Scholium 3.17) allows us to introduce \hat{A}^{R} -valued "Radon-Nikodym derivatives" of left integrals l with respect to the Haar measure. At first note that $l = \sqcap^L(h)l = hl = h \sqcap^R (l) = hS^{-1}(\sqcap^R(l))$ therefore using Scholium 2.7 we have

$$\langle \varphi, l \rangle = \langle \varphi \rho_R, h \rangle = \langle \rho_L \varphi, h \rangle$$

$$(4.41)$$

$$G^{-1}(\Box^R(l)) \rightarrow \hat{1} = \hat{G}^2(z_1)$$

where $\rho_R = \Box^R(l) \rightharpoonup \hat{1}$ and $\rho_L = S^{-1}(\Box^R(l)) \rightharpoonup \hat{1} = \hat{S}^2(\rho_R)$.

Proposition 4.15 The bijections $\mathcal{I}^L(A) \to \hat{A}^R$ provided by the left and right Radon-Nikodym derivatives $l \mapsto \rho_L$ and $l \mapsto \rho_R$, respectively, obey the following properties.

- i) l is non-degenerate iff $\rho_{R,L}$ is invertible.
- ii) If l is non-degenerate then l is normalized iff $l^2 = l$.
- **iii)** l is of positive type, i.e. $\langle \varphi^* \varphi, l \rangle \geq 0$ for all $\varphi \in \hat{A}$, iff $\sqcap^R(l) \geq 0$ iff $\rho_R \in \hat{g}_R^{1/2} \hat{A}_+^R \hat{g}_R^{-1/2}$ where \hat{A}_+^R is the cone of positive elements in \hat{A}^R . In this case $\rho_L = \rho_R^*$ and there exists a $\xi \in \hat{A}$ such that $\langle \varphi, l \rangle = \langle \xi^* \varphi \xi, h \rangle$ for $\varphi \in \hat{A}$.
- iv) Let λ be the dual left integral of l. Then the Radon-Nikodym derivatives of λ and l are related by $\sqcap^R(l)(\widehat{\sqcap}^R(\lambda) \rightarrow 1) = g_R^{-2}$.

Proof: i) follows from cyclicity of h in $\mathcal{I}_{A^R}^L$. ii): $l^2 = l$ implies $(\sqcap^L(l) - 1)l = 0$ and acting with $\lambda \rightarrow$, where λ is the dual left integral of l, one obtains $\sqcap^L(l) = 1$. The converse implication is trivial. iii): As in the proof of Theorem 4.5 l is of positive type iff $S(l_{(1)}) \otimes l_{(2)}$ belongs to the positive cone (4.12). If it does then $\sqcap^R(l) = S(l_{(1)})l_{(2)} \ge 0$. Now assume $\sqcap^R(l) \ge 0$. Then introducing $\xi = \sqcap^R(l)^{1/2} \rightarrow \hat{1}$ we have $\sqcap^R(l)^{1/2} = \xi \rightarrow 1$, $S^{-1}(\sqcap^R(l)^{1/2}) = S(\xi \rightarrow 1)^* = (1 \leftarrow \hat{S}^{-1}(\xi))^* = 1 \leftarrow \xi^*$ therefore $l = hS^{-1}(\sqcap^R(l)^{1/2}) \sqcap^R(l)^{1/2} = \xi \rightarrow h \leftarrow \xi^*$ proving that l is of positive type. It remained to reformulate positivity of $\sqcap^R(l)$ in terms of ρ_R . Use the fact that the antimultiplicative map $x^R \mapsto (x^R \rightarrow \hat{1})$ from A^R to \hat{A}^R sends the *-operation into a new involution, $x^{R*} \rightarrow \hat{1} = (S^{-1}(x^R) \rightarrow \hat{1})^* = (S^{-2}(x^R) \rightarrow \hat{1})^* = (g_R x^R g_R^{-1} \rightarrow \hat{1})^* = \hat{g}_R(x^R \rightarrow \hat{1})^* \hat{g}_R^{-1}$. Therefore the equality $\sqcap^R(l) = x^{R*}x^R$ for some $x^R \in A^R$ is equivalent to the equality $\rho_R = (x^R \rightarrow \hat{1})(x^{R*} \rightarrow \hat{1}) = \hat{g}_R^{1/2} \eta \eta^* \hat{g}_R^{-1/2}$ with $\eta = \hat{g}_R^{-1/2}(x^R \rightarrow \hat{1})\hat{g}_R^{1/2} \in \hat{A}^R$. iv) follows by an elementary calculus starting from the identity $1 = \lambda \rightarrow l = \hat{h} \cap^R(\lambda) \rightarrow hS^{-1}(\sqcap^R(l))$.

A Appendix: The Weak Hopf Algebra $B \otimes B^{op}$

Let B be a separable algebra over the field K and let $E: B \to K$ be a non-degenerate functional with index 1. These are the data needed for constructing a WHA structure on the algebra $B \otimes B^{op}$. For a similar construction of a WBA see [14].

At first choose a basis $\{e_i\}$ of B over K and let $\{f_i\}$ be its dual basis w.r.t. E, i.e. $E(e_i f_j) = \delta_{ij}$. Then

- a) $\sum_i f_i \otimes e_i \in B \otimes B$ is independent of the choice of $\{e_i\}$;
- **b)** $\sum_i E(xf_i)e_i = x = \sum_i f_i E(e_i x), x \in B;$
- c) $\sum f_i e_i = 1;$
- d) $\sum_i x f_i \otimes e_i = \sum_i f_i \otimes e_i x, x \in B;$
- e) if θ denotes the modular automorphism of E, i.e. $E(xy) = E(y\theta(x)), x, y \in B$, then

$$\sum_{i} f_i \otimes x e_i = \sum_{i} f_i \theta(x) \otimes e_i, \quad x \in B$$

f) $\sum_i f_i \otimes e_i = \sum_i e_i \otimes \theta^{-1}(f_i) = \sum_i \theta(e_i) \otimes f_i.$

The algebra $B \otimes B^{op}$ is the K-space $B \otimes B$ with multiplication $(a \otimes b)(x \otimes y) := (ax \otimes yb)$. Its WHA structure is defined by

$$\Delta(x \otimes y) = \sum_{i} (x \otimes f_i) \otimes (e_i \otimes y)$$
(A.1)

$$\varepsilon(x \otimes y) = E(xy) \tag{A.2}$$

$$S(x \otimes y) = y \otimes \theta(x) \tag{A.3}$$

The verification of the WHA axioms is left to the reader. The left and right subalgebras of $B \otimes B^{op}$ are $B \otimes 1$ and $1 \otimes B$, respectively, because we have

$$\sqcap^{L} (x \otimes y) = xy \otimes 1, \qquad \sqcap^{R} (x \otimes y) = 1 \otimes y\theta(x) . \tag{A.4}$$

Let A be an arbitrary WHA over K. Then $A^L A^R$ is a sub-WHA with hypercenter $A^L \cap A^R$. Thus $A^L A^R$ decomposes into a direct sum of WHA-s each summand being isomorphic to a WHA of the type $B \otimes B^{op}$.

Since $B \otimes B^{op}$ is separable, by Theorem 3.13, it must contain a normalized left integral. Indeed,

$$l := \sum_{i} f_i \otimes e_i \equiv S^2(1_{(2)}) 1_{(1)}$$
(A.5)

is such a left integral. What is more, it is non-degenerate.

Before looking for Haar integrals some remarks about innerness of θ are in order. The quantity $q = \sum_i e_i f_i$ always implements θ^{-1} , i.e. $xq = q\theta(x)$ for $x \in B$, but it is not necessarily invertible. (For example for $B = M_2(\mathbb{Z}_2)$ and for any non-degenerate functional E the q is identically zero.) In fact q is invertible iff the left regular trace on B is non-degenerate (especially if K is of characteristic zero). Fortunately one can circumvent this nuisance by using the existence of a non-degenerate trace tr on any separable algebra B (see [5]). Then the Radon-Nykodim derivative γ of E w.r.t. tr provides an invertible element implementing θ ,

$$E(x) = \operatorname{tr}(x\gamma) , \qquad \theta(x) = \gamma x \gamma^{-1} , \quad x \in B .$$
 (A.6)

This proves that θ is inner and therefore so is the square of the antipode, $S^2 = \theta \otimes \theta$.

Omitting the details we can now formulate the condition for the existence of the Haar measures h and \hat{h} as follows. Haar measure in $B \otimes B^{op}$ exists iff $\sum_i f_i \gamma^2 e_i$ is invertible and Haar measure in $B \otimes B^{op}$ exists iff $E(1_B) \neq 0$.

References

- [1] E. Abe, *Hopf Algebras*, Cambridge University Press, Cambridge, 1980
- [2] G. Böhm, K. Szlachányi, A Coassociative C*-Quantum Group with Nonintegral Dimensions, Lett. Math. Phys. 35, 437 (1996)
- [3] G. Böhm, F. Nill, K. Szlachányi, Weak Hopf Algebras II: Representation Theory, Dimensions, and Markov Traces, in preparation
- C. W. Curtis, I. Reiner, Representation Theory of Finite Groups and Associative Algebras, John Wiley & Sons, Inc., 1962
- [5] C. W. Curtis, I. Reiner, Methods of Representation Theory, John Wiley & Sons
- [6] V.G. Drinfeld, Quasi-Hopf Algebras, Leningrad Math. J. 1, 1419-1457 (1990)
- [7] J. Fuchs, A. Ganchev, P. Vecsernyés, Rational Hopf algebras: Polynomial equations, gauge fixing, and low dimensional examples, *Int. J. Modern Physics* A 10, 3431 (1995)
- [8] T. Hayashi, Quantum group symmetry of partition functions of IRF models and its application to Jones' index theory, Commun. Math. Phys. 157, 331-345 (1993)
- [9] R. G. Larson, M. Sweedler, An associative orthogonal bilinear form for Hopf algebras, Amer. J. Math. 91, 75-93 (1969)

- [10] R. Longo, A duality for Hopf algebras and for subfactors I, Commun. Math. Phys. 159, 133-150 (1994)
- [11] G. Mack and V. Schomerus, Quasi Hopf quantum symmetry in quantum theory, Nucl. Phys. B370, 185 (1992)
- [12] S. MacLane, Categories for the Working Mathematician, Springer 1971
- [13] S. Montgomery, Hopf Algebras and their Actions on Rings, CBMS Series of the Amer. Math. Soc., No. 82, 1993
- [14] F. Nill, Axioms for Weak Bialgebras, q-alg/9805104
- [15] F. Nill, K. Szlachányi, H.-W. Wiesbrock, Weak Hopf Algebras and Reducible Jones Inclusions of Depth 2. I. From crossed products to Jones towers, q-alg/9805nnn
- [16] A.Ocneanu, Quantized groups, string algebras, and Galois theory for algebras, in Operator Algebras and Applications, Vol. 2, eds.: D.E. Evans et al., London Math. Soc. Lect. Notes 135, Cambridge 1988
- [17] A. Ocneanu, Quantum Cohomology, Quantum Groupoids, and Subfactors, unpublished talk presented at the First Caribic School of Mathematics and Theoretical Physics, Guadeloupe 1993
- [18] R. S. Pierce, Associative Algebras, (Graduate Texts in Matematics 88) Springer-Verlag 1982
- [19] M.E. Sweedler, *Hopf algebras*, Benjamin 1969
- [20] K. Szlachányi, Weak Hopf Algebras, in Operator Algebras and Quantum Field Theory, eds.: S. Doplicher, R. Longo, J. E. Roberts, and L. Zsidó, International Press (1996)
- [21] A. Van Daele, The Haar measure on finite quantum groups, to appear in *Proc. Amer. Math. Soc.*
- [22] P. Vecsernyés, On the Quantum Symmetry of the Chiral Ising Model, Nucl. Phys. B 415 [FS], 557-588 (1994)
- [23] Y. Watatani, Index for C^* -subalgebras, Memoirs of the AMS No. 424 (1990)
- [24] S.L. Woronovicz, Compact Matrix Pseudogroups, Commun. Math. Phys. 111, 613-665 (1987)
- [25] T. Yamanouchi, Duality for Generalized Kac Algebras and a Characterization of Finite Groupoid Algebras, Journal of Algebra 163, 9-50 (1994)