Comment on "Shadowability of Statistical Averages in Chaotic Systems"

Lai *et al.* [1] investigate systems with a stable periodic orbit and a coexisting chaotic saddle. They claim that, by adding white Gaussian noise (GN), averages change with an algebraic scaling law above a certain noise threshold and argue that this leads to a breakdown of shadowability of averages. Here, we show that (i) shadowability is not well defined and even if it were, no breakdown occurs. We clarify misconceptions on (ii) thresholds for GN. We further point out (iii) misconceptions on the effect of noise on averages. Finally, we show that (iv) the lack of a proper threshold can lead to any (meaningless) scaling exponent.

(i) Shadowing deals with macroscopic error propagation of noise bounded by a small value (e.g. computer accuracy of 10^{-16}), comparing a noisy and a 'true' trajectory [2]. Since the authors use GN, one cannot properly speak of shadowing. However, even for bounded noise, their claims are still doubtful. In a recent work, the error of an *average* was found to be amplified by a factor of up to 10^{12} , although the trajectory never exits the attractor [3]. This renders a reliable computation truly unfeasible. In [1], though, the effect on the averages is only of the same order as the variation of the noise level *D*. Furthermore, the averages in Figs. 1a and 2a in [1] do change even less below $D = 10^{-2.5}$. Therefore, shadowability is not compromised at all.

(ii) The authors of [1] mention a threshold for GN above which the periodic orbit and the chaotic saddle would become connected. Yet, such a threshold does not exist, since the mean first exit time $\langle \tau \rangle$ is given by Kramers' law $\langle \tau \rangle \sim \exp(\frac{\Delta U}{D})$, where *D* is the noise variance and ΔU is either the potential [4] or, for nonequilibrium and chaotic systems, the quasipotential difference [5]. Thus, $\langle \tau \rangle$ varies with *D*, yielding a different average for every noise level.

(iii) Generally, averages depend on the noise for all noise levels, implying that no threshold can exist, even with only one metastable state. For the linear map $x_{n+1} = ax_n + b + \xi_n$ with a fixed point $x_* = \frac{b}{1-a}$ and white GN one gets $\langle x^2 \rangle = x_*^2 + \frac{1}{1-a^2} D$. This is pictured in Fig. 1a, fitting perfectly the data. Although the average appears to be constant for low noise, it depends, in fact, on the noise for all D. The same applies also to nonlinear systems (cf. fit in Fig. 1b) and bounded noise.

(iv) Because the value of the threshold is arbitrary, any scaling can be achieved, just by tuning D_c , as fittings of the form of Eq. (1) of [1] are very sensitive to the value of D_c . To verify this, we show in Fig. 1b the logistic map $x_{n+1} = ax_n(1-x_n) + D\xi_n$ with $\langle \xi_n, \xi_m \rangle = \delta_{nm}$ as in [1].

The putative threshold of $D_c = 10^{-5}$ is marked by an arrow. The graph is evidently not constant there. With $D_c \approx 7 \cdot 10^{-6}$ (first arrow), a scaling $\Delta G \sim (D - D_c)^{\alpha}$ results, with ΔG of [1]. This yields $\alpha \approx 2.1$ (Fig. 1c), in clear contrast to $\alpha \approx 1$ reported in [1]. Therefore, no reliable (i. e. any) exponent can be obtained, since D_c is not well defined from the outset.




FIG. 1: (a) $\langle x^2 \rangle$ for the linear map vs. D with a = 0.7and b = 0.75 (circles) and analytical fit $\langle x^2 \rangle = x_\star^2 + \frac{1}{1-a^2}D$ (full line). (b) $\langle \sin x \rangle$ for the logistic map with a = 3.8008and a polynomial fit for $D \leq 1.4 \cdot 10^{-5}$ (full line) (c) Scaling $\Delta G \sim (D-D_c)^{\alpha}$ with $D_c \approx 7 \cdot 10^{-6}$ (circles) and least squares fit (full line). All averages are for 10^8 iterations.

As to the reply, the authors claim to have "a theoretical justification for the existence of a threshold" through $D_c = \sqrt{\Delta \Phi / \ln \chi^{-1}}$, with $\Delta \Phi$ the quasipotential (see (ii)) and χ the probability resolution. Contrary to Ref. [5] of the reply, where χ drops out since only proportionalities under parameter variation are considered, D_c in [1] depends for fixed parameters on χ and hence α is not uniquely defined. Applied to (iv) with $D_c = 10^{-5}$ of [1] and $\Delta \Phi \approx 7.8 \cdot 10^{-10}$ (not shown) yields $\chi = 0.0004$. But a finer resolution $\chi = 10^{-7}$ gives $D_c = 7 \cdot 10^{-6}$, with $\alpha \approx 2.1$ (cf. (iv)), whereas $\chi = 10^{-2}$ results in $D_c = 1.3 \cdot 10^{-5}$ and $\alpha \approx 0.21$ (not shown). Again, this invalidates any proper scaling.

Suso Kraut

Instituto de Física, Universidade de São Paulo Caixa Postal 66318, 05315-970 São Paulo, Brazil

- [1] Y. C. Lai et al., Phys. Rev. Lett. 89, 184101 (2002).
- [2] C. Grebogi *et al.*, Phys. Rev. Lett. **65**, 1527 (1990).
- [3] T. D. Sauer, Phys. Rev. E 65, 036220 (2002).
- [4] H. A. Kramers, Physica 7, 284 (1940).
- [5] R. Graham and T. Tél, Phys. Rev. Lett. 52, 9 (1984);
 R. Graham, A. Hamm, and T. Tél, *ibid* 66, 3089 (1991).