

From collective rhythm to adaptive synchronization

Debin Huang

Department of Mathematics, Shanghai University, Shanghai 200436, P.R. China

(E-mail: dbhuang@staff.shu.edu.cn)

A novel viewpoint, i.e., adaptive synchronization, is proposed to explore collective rhythm observed in many complex, self-organizing systems. We show that a simple adaptive coupling is able to tip arrays of oscillators towards collective synchronization. Two arrays of simply coupled Hindmarsh-Rose chaotic neurons are used to illustrate cooperative dynamics of neural activity like the central pattern generators, which supplies a new idea for biological experiments and numerical simulations. The results enhance the viewpoint that chaos is a necessary ingredient in life, and indicate that such small-world adaptive coupling may be a universal essence of the collective rhythm observed in nature.

TODAY one of the main unsolved problems in science is how to apperceive and study complex, self-organizing systems. A puzzling characteristic in these systems is the spontaneous collective rhythm, i.e., so-called collective synchronization. This remarkable phenomenon has been observed extensively in nature, ranging from inorganic systems to organic systems, e.g., Christiaan Huygens' two synchronization clocks, wobbly bridges, the oscillating uniformly Josephson junctions, the Bose-Einstein condensate (BEC), emerging coherence in chemical oscillators, fireflies synchronizing spontaneously their flashes, self-synchronization of the cardiac pacemaker cells, the cooperative pattern in network of neurons, animals' gaits, groups of women with the mutually synchronized menstrual cycles, an audience clapping in sync, and coherent moving states recently observed in highway traffic, etc^[1-7]. In the collective rhythm, each individual itself executes two different behaviors, i.e., periodic oscillating and aperiodic (i.e., chaotic) oscillating. The collective rhythm of periodic oscillators emerges in many fields, while the collective chaotic synchronization is observed in the network of neurons and two weakly coupled BECs, etc. As is well known, there exist infinitely many unstable periodic orbits embedded in a chaotic attractor, meanwhile due to the extreme sensitivity of initial values chaotic systems intrinsically defy synchronization. Therefore from this sense the collective synchronization of chaotic oscillators seems to be more difficult and significant than that of periodic oscillators.

Although collective synchronization has been observed in various concrete experiments, some key theoretical problems remain open for scientists, especially for mathematicians. Polymath Norbert Wiener ever tried to develop a mathematical model of collection of oscillators in the late 1950s but little fruit was obtained^[8]. The theoretical breakthrough came from A.T. Winfree’s pioneering work in 1966^[9], where he developed a mathematical model to study large populations of periodic oscillators, and obtained some important results although stymied by the difficulty of solving the mathematical model. A crucial breakthrough came in 1975 when Y. Curamoto refined and simplified the mathematical framework developed by Winfree^[10], where an exactly solvable mean-field model of coupled oscillators was proposed. Since these pioneering studies lots of theoretical works have been accomplished^[11–19]. Note that only the case of coupled periodic oscillators was investigated in the literature. Through the large amount of effort in idealized mathematical models there has been a common acknowledgement for the complex problem on collective rhythm: individual oscillators are coupled together, and there exists a critical value for the intensity of mutual coupling, below which anarchy prevails and above which coherence with the collective rhythm reigns. However, in this viewpoint two points deserve to investigate further. One is how the individual oscillators are coupled together, and the other is how complex, self-organizing systems produce the threshold of coupling to reach collective synchronization. In the previous theoretical work, the weak couplings were almost global, e.g., mean-field coupling, where each oscillator interacts with the rest of oscillators. However from the perspective of biology (especially neurobiology) such all-to-all coupling is too complicated to benefit to realize promptly the corresponding function. As for the problem on the critical intensity of coupling, it is still far away from scientists.

In my opinion, the architecture of interaction of weakly coupling in the collection of oscillators should be as simple as possible, and the critical coupling strength should be reached self-adaptively due to the spontaneousness of collective rhythm. This speculation is inspired by the provocative words “self-adaption creates complexity” which is a basic idea of complex adaptive systems (CAS) theory, meanwhile this idea is motivated directly by considering the collective chaos synchronization. In network of chaotic oscillators the mathematical model must be exactly unsolvable, but as what introduced above almost all the previously obtained results on collective synchronization of coupled oscillators are based on the solvability of the mathematical model. In this paper, a simple mathematical framework is proposed to confirm this speculation, where a simple adaptive coupling is used to produce the collective synchronization of arrays of oscillators.

Mathematical model and results

Our analysis will be limited to an array of oscillators that are all strictly identical, which ignores the diversity. Suppose that the dynamical behavior of each oscillator is governed by an n -dimensional differential equation

$$\dot{x} = f(x), x \in \mathbb{R}^n, \tag{1}$$

where $f(x)$ is a differentiable nonlinear vector function. We firstly consider cycle-

type coupling in a ring of N oscillators,

$$\dot{x}_i = f(x_i) - \epsilon_{i,i-1}(x_i - x_{i-1}), \quad i = 1, 2, \dots, N, \quad (2)$$

where $x_i = (x_{i,1}, x_{i,2}, \dots, x_{i,n}) \in \mathbf{R}^n$, $x_0 \equiv x_N$, $\epsilon_{i,i-1} = (\epsilon_{i,i-1,1}, \epsilon_{i,i-1,2}, \dots, \epsilon_{i,i-1,n})$, and $\epsilon_{i,i-1}(x_i - x_{i-1}) \equiv (\epsilon_{i,i-1,1}(x_{i,1} - x_{i-1,1}), \epsilon_{i,i-1,2}(x_{i,2} - x_{i-1,2}), \dots, \epsilon_{i,i-1,n}(x_{i,n} - x_{i-1,n}))$. This model indicates that the connection topology is cycle-type, and each individual oscillator is coupled only to its nearest neighbor. Here the coupling intensity $\epsilon_{i,i-1}$ varies adaptively according to the following update law (this is very crucial!)

$$\dot{\epsilon}_{i,i-1,j} = \gamma_{i,i-1,j}(x_{i,j} - x_{i-1,j})^2, \quad i = 1, 2, \dots, N, \quad j = 1, 2, \dots, n, \quad (3)$$

where $\gamma_{i,i-1,j}$ are arbitrary positive constants. Next we consider another simple configuration of network, i.e., star-type coupling,

$$\dot{x}_1 = f(x_1), \quad \dot{x}_i = f(x_i) - \epsilon_{i,1}(x_i - x_1), \quad i = 2, 3, \dots, N, \quad (4)$$

where all notations are as those above, and the coupling strength $\epsilon_{i,1}$ varies according to the update law

$$\dot{\epsilon}_{i,1,j} = \gamma_{i,1,j}(x_{i,j} - x_{1,j})^2, \quad i = 2, \dots, N, \quad j = 1, 2, \dots, n. \quad (5)$$

Such star-type coupling admits the characteristics of small-world networks, where the first oscillator, x_1 , represents a hub.

Using the well-known Lasalle's invariance principle in mathematics, one may prove that for arbitrary bounded solutions of system (2)-(3) (or (4)-(5)) as $t \rightarrow \infty$ $x_i(t) \rightarrow x_1(t)$, $i = 2, \dots, N$, and the coupling strength will converges to a constant dependent on the initial values. The idea of proof is similar to that in [20,21]. The results indicate that the above two coupling schemes can produce the collective synchronization of N oscillators, and such collective synchronization is nonlinear stable and robust against the effect of noise. Moreover the variable coupling strength reaches self-adaptively a value which just corresponds to the threshold for arising collective rhythm, meanwhile the small weak coupling may be obtained by adjusting the dissipation parameter $\gamma_{i,i-1,j}$. In addition, note that in the present coupling scheme the mutual interaction will vanish once the collective synchronization is achieved, which implies that the collective rhythm doesn't change dynamical behavior of each individual. In particular, when the oscillator (1) is chaotic, thanks to the global attraction and nonhyperbolicity of chaotic attractor these coupling schemes will be more simple and the global collective synchronization will be realized more easily, see the examples below.

Collective synchronization of arrays of chaotic neurons

It has been observed in neurobiological experiments and in numerical simulations that individual neurons show chaotic spiking-bursting, while ensemble of such irregularly bursting neurons can produce cooperatively coherent, rhythmical bursting, i.e., synchronized pattern of neural activity^[22-26]. Such cooperative property

plays a crucial role in neural activity, for example the central pattern generators (CPG) controlling the rhythmic motor behavior of animals. An important question is how neural assemblies produce and control regular rhythm. In the literature, some global coupling schemes like mean-field coupling have been used to explore the interesting sync pattern^[27–30]. Here we will attempt to apply the proposed adaptive synchronization to reveal this complex, self-organizing phenomenon. To do it, we choose the famous Hindmarsh-Rose neuron model^[31], a third-order ordinary differential equation with one slow variable modelling chaotic spiking-bursting of neuron, as dynamical equations of individual neurons. Firstly we consider a network of neurons as the following cycle-type synaptic coupling scheme,

$$\begin{aligned}\dot{X}_i &= Y_i + 3X_i^2 - X_i^3 - Z_i + I - \epsilon_{i,i-1}(X_i - X_{i-1}), \\ \dot{Y}_i &= 1 - 5X_i^2 - Y_i, \quad \dot{Z}_i = -rZ_i + 4r(X_i + 1.6), \quad i = 1, 2, \dots, N,\end{aligned}\tag{6}$$

with

$$\dot{\epsilon}_{i,i-1} = \gamma_{i,i-1}(X_i - X_{i-1})^2,\tag{7}$$

where $X_0 \equiv X_N$, $r = 0.0012$, and the external input $I = 3.281$. Each neuron is characterized by three time-dependent variables: the membrane potential X_i , the recovery variable Y_i and the slow adaptation current Z_i . $\epsilon_{i,i-1}$ represents the self-adaptive strength of the synaptic coupling between the i th neuron and the $(i-1)$ th neuron. Similarly, the star-type coupling network of neurons with the first neuron being the hub is given as

$$\begin{aligned}\dot{X}_i &= Y_i + 3X_i^2 - X_i^3 - Z_i + I - \epsilon_{i,i-1}(X_i - X_1), \\ \dot{Y}_i &= 1 - 5X_i^2 - Y_i, \quad \dot{Z}_i = -rZ_i + 4r(X_i + 1.6), \quad i = 1, 2, \dots, N,\end{aligned}\tag{8}$$

with $\epsilon_{11} \equiv 0$, and otherwise

$$\dot{\epsilon}_{i,1} = \gamma_{i,1}(X_i - X_1)^2, \quad i = 2, 3, \dots, N.\tag{9}$$

For the sake of simplicity, we set the parameters $\gamma_{ij} \equiv 0.1$ and $N = 5$. The adaptive collective synchronization of the two networks of chaotic neurons is shown numerically in Figs. 1-4. Comparing the above two coupling schemes (see Fig. 4 legend), we find that the adaptive synchronization in the star-type network with the characteristics of small-world is achieved more easily, which confirms again the strong synchronizability of small-world networks^[32,33]. In conclusion, such adaptive collective synchronization is more simple than those based on the global coupling schemes in literature, which supplies a new idea for biological experiments and numerical simulations. Moreover the result supports the well-known neural theory of high-level brain function proposed by D. Hebb in 1949: Learning takes place by the adaptive adjustment of the conductances of the connections between neurons^[34].

As the above statements, besides the self-adaption of individual neurons a key reason why such simple synaptic coupling schemes (i.e., only coupling one variable representing the membrane potential) can produce so interesting collective synchronization of neural activity is the chaotic characteristic of neural dynamics. This result throws highly light to the well-known viewpoint that chaos is a necessary

ingredient in life. Obviously the present adaptive collective synchronization can be generalized to the other networks, such as the multi-star-type networks and the higher-dimensional lattice networks. In particular, such adaptive coupling can be extended to the complex networks like small-world networks and scale-free networks [32,33,35,36]. Combining the characteristics of small-world networks, such as quick signal-propagation and strong synchronizability, etc., we speculate that such small-world adaptive coupling may be a universal essence of collective rhythm observed in nature. Therefore we hope that this work will inspire further studies of complex, self-organizing systems.

References

- [1] Strogatz, S.H. & Stewart, I. Coupled oscillators and biological synchronization. *Sic. Am.* **269**, 102-108 (1993).
- [2] Helbing, D. & Huberman, B.A. Coherent moving states in highway traffic. *Nature* **396**, 738-740 (1998).
- [3] Winfree, A.T. *The Geometry of Biological Time* (2nd edn, Springer, New York, 2001).
- [4] Kiss, I.Z., Zhai, Y. & Hudson, J.L. Emerging Coherence in a Population of Chemical Oscillators. *Science* **296**, 1676-1678 (2002).
- [5] Winfree, A.T. On emerging coherence. *Science* **298**, 2336-2337 (2002).
- [6] Nadis, S. All together now. *Nature* **421**, 780-782 (2003).
- [7] Strogatz, S.H. *Sync: the emerging science of spontaneous order* (Hyperion, New York, 2003).
- [8] Wiener, N. *Cybernetics* (2nd edn, MIT Press, Cambridge, Massachusetts, 1961).
- [9] Winfree, A.T. Biological rhythms and the behavior of populations of coupled oscillators. *J. Theor. Biol.* **16**, 15-42 (1967).
- [10] Kuramoto, Y. *Chemical Oscillations, Waves, and Turbulence* (Springer, Berlin, 1984).
- [11] Lumer, E.D. & Huberman, B.A. Hierarchical dynamics in large assemblies of interacting oscillators. *Phys. Lett. A* **160**, 227-232 (1991).
- [12] Strogatz, S.H., Mirollo, R.E. & Matthews, P.C. Coupled nonlinear oscillators below the synchronization threshold: Relaxation by generalized Landau damping. *Phys. Rev. Lett.* **68**, 2730-2733 (1992).
- [13] Daido, H. Generic scaling at the onset of macroscopic mutual entrainment in limit-cycle oscillators with uniform all-to-all coupling. *Phys. Rev. Lett.* **73**, 760-763 (1994).
- [14] Crawford, J.D. Scaling and singularities in the entrainment of globally coupled oscillators. *Phys. Rev. Lett.* **74**, 4341-4344 (1995).

- [15] Wu, C.H. & Chua, L.O. Synchronization in an array of linearly coupled dynamical systems. *IEEE Trans. Circuits Syst.-I* **42**, 430-447 (1995).
- [16] Wiesenfeld, K., Colet, P. & Strogatz, S.H. Synchronization transition in a disordered Josephson series array. *Phys. Rev. Lett.* **76**, 404-407 (1996).
- [17] Daido, H. Onset of cooperative entrainment in limit-cycle oscillators with uniform all-to-all interactions: bifurcation of the order function. *Physica D* **91**, 24-66 (1996).
- [18] Crawford, J.D. & Davies, K.T.R. Synchronization of globally coupled phase oscillators: singularities and scaling for general couplings. *Physica D* **125**, 1-46 (1999).
- [19] Strogatz, S.H. From Kuramoto to Crawford: exploring the onset of synchronization in populations of coupled oscillators. *Physica D* **143**, 1-20 (2000).
- [20] Huang Debin. Stabilizing near-nonhyperbolic chaos and its potential applications in neuroscience. *Phys. Rev. Lett.* (To appear, 2004).
- [21] Huang Debin. A simple adaptive controller for high-quality chaos synchronization (Preprint, 2004).
- [22] Gray, C.M., Koning, P., Engel, A.K. & Singer, W. Oscillatory responses in cat visual cortex exhibit inter-columnar synchronization with reflects global stimulus properties. *Nature* **338**, 334-337 (1989).
- [23] Elson, R.C. & Selverston, A.I. Slow and fast synaptic inhibition worked by pattern-generating neurons of the gastric mill network in spiny lobsters. *J. Neurophysiol.* **74**, 1996-2001 (1995).
- [24] Gray, C.M. & McCormick, D.A. Chattering cells: superficial pyramidal neurons contributing to the generation of synchronous oscillations in the visual cortex. *Scienc* **274**, 109-113 (1996).
- [25] Izhikevich, E.M. Synchronization of elliptic bursters. *SIAM Rev.* **43**, 315-344 (2001).
- [26] Samonds, J.M., Allison, J.D., Brown, H.A. & Bonds, A.B. Cooperative synchronized assemblies enhance orientation discrimination. *Proc. Natl. Acad. Sci. USA* **101**, 6722-6727 (2004).
- [27] Hansel, D. & Sompolinsky, H. Synchronization and computation in a chaotic neural network. *Phys. Rev. Lett.* **68**, 718-721 (1992). Bazhenov, M.;
- [28] Huerta, R., Rabinovich, M.I. & Sejnowski, T. Cooperative behavior of a chain of synaptically coupled chaotic neurons. *Physica D* **116**, 392-400 (1998).
- [29] Rulkov, N.F. Regularization of synchronized chaotic bursts. *Phys. Rev. Lett.* **86**, 183-186 (2001).
- [30] Rosenblum, M.G. & Pikovsky, A.S. Controlling synchronization in an ensemble of globally coupled oscillators. *Phys. Rev. Lett.* **92**, 114102 (2004).

- [31] Hindmarsh, J.L. & Rose, R.M. A model of neuronal bursting using three coupled first order differential equations. *Proc. R. Soc. London B* **221**, 87-102 (1984).
- [32] Watts, D.J. & Strogatz, S.H. Collective dynamics of “small-world” networks. *Nature* **393**, 440-442 (1998).
- [33] Strogatz, S.H. Exploring complex networks. *Nature* **410**, 268-276 (2001).
- [34] Hebb, D. *The organization of behavior* (New York: Wiley, 1949).
- [35] Barabasi, A.L. & Albert, L. Emergence of scaling in random networks. *Science* **286**, 509-512 (1999).
- [36] Albert, L. & Barabasi, A.L. Statistical mechanics of complex networks. *Rev. Mod. Phys.* **74**, 47-97 (2002).

Acknowledgments

This work is supported by the National Natural Science Foundation of China (Division of Mathematical Sciences).




Figure.1. The collective synchronization emerging in the cycle-type neural network (6). The collective synchronization in the simply model with 5-coupled chaotic neurons is shown by calculating synchronization error in system (6), where $e_{X,i} \equiv X_i - X_1$, $e_{Y,i} \equiv Y_i - Y_1$ and $e_{Z,i} \equiv Z_i - Z_1, i = 2, 3, 4, 5$ respectively denoting the synchronization errors in each variable converge to zero. Here the initial values of variables are set as $(0.2, 3, 0.7, 0.1, 4, 0.8, 0.3, 2, 0.6, 0.1, 2, 0.7, 0.3, 4, 0.6)$.




Figure 2. The temporal evolution of the mutual synaptic coupling strength $\epsilon_{i,i-1}$, $i = 1, 2, 3, 4, 5$ in system (6)-(7). The mutual synaptic coupling strength between neurons tends eventually to a small constant in the course of arising the collective synchronization in Fig. 1. The evolution is governed by the adaptive law (7), where the dissipation parameter is uniformly chosen as $\gamma_{i,i-1} = 0.1$, $i = 1, 2, 3, 4, 5$, and the initial coupling strength is uniformly set as 0.




Figure 3. The collective synchronization emerging in the star-type network (8) with 5-coupled neurons. The collective synchronization in the star-type network of coupled Hindmarsh-Rose chaotic neurons is shown by numerically calculating synchronization error in system (8), where $e_{X,i} \equiv X_i - X_1$, $e_{Y,i} \equiv Y_i - Y_1$ and $e_{Z,i} \equiv Z_i - Z_1, i = 2, 3, 4, 5$ respectively denoting the synchronization errors in each variable tend to zero. Here the initial values of variables are set as (0.2, 3, 0.7, 0.1, 4, 0.8, 0.3, 2, 0.6, 0.1, 2, 0.7, 0.3, 4, 0.6).




Figure 4. The temporal evolution of the mutual synaptic coupling strength $\epsilon_{i,1}$, $i = 2, 3, 4, 5$ in system (8)-(9). According to the update law (9) the mutual synaptic coupling intensity between individual neurons and the hub neuron self-adaptively approaches the threshold for producing the collective synchronization in Fig. 3, where the dissipation parameter is uniformly chosen as $\gamma_{i,1} = 0.1$, $i = 2, 3, 4, 5$, and the initial coupling strength is uniformly set as 0. Obviously here mean coupling strength is smaller than that in Fig. 2.