

Poincaré Invariance, Cluster Properties, and Particle Production

W. N. Polyzou^{*†‡}
The University of Iowa

October 29, 2018

Abstract

A method is presented for constructing a class of Poincaré invariant quantum mechanical models of systems of a finite number of degrees of freedom that satisfy cluster separability, the spectral condition, but do not conserve particle number. The class of models includes the relativistic Lee model [1] and relativistic isobar models.

Any reasonable quantum theory of strongly interacting particles should be Poincaré invariant, and should satisfy cluster properties and the spectral condition. These conditions are physical requirements. In addition, the theory should be able to model reactions that change particle number. These conditions require the existence of a unitary representation of the Poincaré group $\hat{U}(\Lambda, a)$ [2] on the model Hilbert space \mathcal{H} that satisfies a cluster separability condition and the spectral condition.

While these requirements are dictated by physics, it is difficult to find well-defined mathematical models for realistic systems that are consistent with all of these properties. Relativistic quantum theory of N -particles[1][3] is a mathematically well-defined theory where these conditions can be satisfied. A mathematically well-defined theory is required to formulate high-precision ab-initio calculations, which are important for strongly interacting systems.

This talk explains how to extend the relativistic quantum theory of N -particles [1][3] to treat a class of models with particle production. This class of models has a bounded number of degrees of freedom and includes the relativistic Lee model [1] and relativistic isobar models.

The key observation is that mass is not superselected in relativistic quantum theory. To contrast this with non-relativistic quantum theory, consider a reaction involving a fixed number of constituent particles, where a bound target is

^{*}*E-mail address:* polyzou@uiowa.edu

[†]This work Supported by the U.S. Department of Energy, Nuclear Physics Division, contract DE-FG02-86ER40286.

[‡]Contributed talk to the XVIII European Conference on Few-Body Problems

broken up into a number of subsystems. In this reaction, the mass and number of particles in the initial and final states is different. This reaction cannot be consistently modeled with a non-relativistic quantum theory because it violates Galilean invariance if the Galilean and inertial masses are identified.

The difference between the above reaction and a general production reaction is the presence of a conserved number of constituent particles, which limits the number of degrees of freedom. Motivated by this observation, this talk considers production reactions that have a set of conserved quantum numbers which are associated with particles called “atomic partons”. Additional bare particles with *composite* atomic-parton quantum numbers are introduced. The model Hilbert space is a direct sum of tensor products of bare and atomic particle subspaces with the same atomic-parton numbers. The dynamics is restricted to conserve atomic-parton number.

For example, a model where the atomic partons are a nucleon and two pions could have a bare Δ , with the quantum numbers of a nucleon and a pion, and a bare ρ , with the quantum numbers of two pions. Introducing bare particles does not mean that these particles exist as stable particles. Stable particles correspond to point eigenstates of the mass operator. This talk investigates the modifications of the fixed N -construction [1][3] which are needed to treat this type of particle production.

In the above example the conserved atomic-parton numbers are $\mathcal{N} := \{N_N, N_\pi\}$. The Hilbert space, $\mathcal{H}_{\mathcal{N}}$, is a direct sum of tensor products of irreducible representation spaces of the Poincaré group, where the particles in each tensor product have atomic parton number \mathcal{N}

$$\mathcal{H}_{\mathcal{N}} = \oplus_i (\otimes \mathcal{H}_{a_i}); \text{ i.e. } \mathcal{H}_{\{1,1\}} = (\mathcal{H}_N \otimes \mathcal{H}_\pi) \oplus \mathcal{H}_\Delta.$$

There is a natural partial ordering on the quantum numbers \mathcal{N} given by $\mathcal{N} \leq \mathcal{N}'$ if $N_i \leq N'_i$ for each type, i , of atomic parton. The space $\mathcal{H}_{\mathcal{N}}$ has a natural non-interacting representation of the Poincaré group, $\hat{U}_0(\Lambda, a)$

For each partition, b , of the atomic partons into disjoint equivalence classes, b_i , there is a factorization of the model Hilbert space, $\mathcal{H}_{\mathcal{N}}$, into a direct sum of a tensor product, \mathcal{H}_b , of n_b proper subsystem Hilbert spaces, \mathcal{H}_{b_i} , with $\mathcal{N}_i < \mathcal{N}$, and a residual space, \mathcal{H}^b , corresponding to bare particles with quantum numbers in more than one equivalence class:

$$\mathcal{H}_{\mathcal{N}} = \mathcal{H}_b \oplus \mathcal{H}^b \quad \mathcal{H}_b := \otimes_{b_i \in b} \mathcal{H}_{b_i}.$$

The difference between the construction outlined in this talk and the construction for fixed N in [1][3] involves the modifications that are necessary to treat the residual space \mathcal{H}^b . The asymptotic dynamics and translation operators associated with partition b are needed to formulate cluster properties. On $\mathcal{H} = \mathcal{H}_b \oplus \mathcal{H}^b$ these operators can be expressed in terms of subsystem representations $\hat{U}_{b_i}(\Lambda, a)$ as:

$$\hat{U}_b(\Lambda, a) := \begin{pmatrix} \otimes \hat{U}_{b_i}(I, y_i) & 0 \\ 0 & \hat{I}^b \end{pmatrix} \quad \hat{T}_b(y_1, \dots, y_k) := \begin{pmatrix} \otimes_i \hat{U}_{b_i}(I, y_i) & 0 \\ 0 & \hat{I}^b \end{pmatrix}. \quad (1)$$

The projection on \mathcal{H}_b ,

$$\hat{\Pi}_b := \begin{pmatrix} \hat{I} & 0 \\ 0 & 0 \end{pmatrix},$$

is used to define a modified formulation of the cluster property

$$\lim_{\min(y_i - y_j)^2 \rightarrow \infty} \|[\hat{U}(\Lambda, x) - \hat{U}_b(\Lambda, x)]\hat{T}_b(y_1, \dots, y_k)\hat{\Pi}_b|\psi\rangle\| = 0. \quad (2)$$

This condition requires more than a range condition on the interaction; it also requires that part of the operator that survives asymptotically is a tensor product of proper-subsystem representations [3].

In this formulation \mathcal{N} -body interactions, $\hat{V}_{\mathcal{N}}$, satisfy

$$\lim_{\min(y_i - y_j)^2 \rightarrow \infty} \|\hat{V}_{\mathcal{N}}\hat{T}_b(y_1, \dots, y_k)\hat{\Pi}_b|\psi\rangle\| = 0 \quad \forall b.$$

The dynamics of all proper subsystems determine the dynamics of the full system up to an \mathcal{N} -body interaction.

The construction of a two atomic parton dynamics uses *Wigner's forms of the dynamics* [4][3]; Clebsch-Gordan coefficients of the Poincaré group are used to construct an irreducible basis for the non-interacting system. Interactions are added to the non-interacting mass that are block-diagonal in the non-interacting spin, \hat{j}_0 , and commute with and are independent of the commuting functions, \hat{F}_0 , of the Poincaré generators that label vectors in irreducible subspaces. It follows that simultaneous eigenstates of the mass, \hat{M} , \hat{j}_0 , and \hat{F}_0 define an irreducible basis for an interacting representation of the Poincaré group. In this representation the mass eigenvalue and non-interacting spin are the new Casimir operators.

The many-body construction is recursive. The representations $\hat{U}_b(\Lambda, a)$ are constructed from tensor products of proper subsystem, ($\mathcal{N}_i < \mathcal{N}$), representations. If $\hat{U}_b(\Lambda, a)$ is decomposed into irreducible representations, the spin \hat{j} , and the vector operators, \hat{F} , depend on the interaction. If the order of constructing irreducible representations and adding interactions is reversed, the result is a mass operator, \hat{M}_b , that commutes with \hat{j}_0 , and \hat{F}_0 , and is independent of \hat{F}_0 . Simultaneous eigenstates of \hat{M}_b , \hat{j}_0 , and \hat{F}_0 transform irreducibly and define a representation, $\bar{U}_b(\Lambda, a)$, of the Poincaré group. The representations $\bar{U}_b(\Lambda, a)$ and $\hat{U}_b(\Lambda, a)$ are related by a unitary scattering equivalence \hat{B}_b [1][4] of the form:

$$\hat{B}_b := \begin{pmatrix} \hat{A}_b & 0 \\ 0 & \hat{I}^b \end{pmatrix} \quad \bar{U}_b(\Lambda, a) = \hat{B}_b \hat{U}_b(\Lambda, a) \hat{B}_b^\dagger \quad (3)$$

which satisfies

$$\lim_{t \rightarrow \pm\infty} \|[(\hat{B}_b - \hat{I})\hat{\Pi}_0 \hat{U}_0(I, t)|\psi\rangle\| = 0,$$

where $\hat{\Pi}_0$ is the projector on the \mathcal{N} -atomic parton subspace of \mathcal{H} . For systems of more than three atomic partons, the freedom to choose the operators \hat{B}_b must be utilized to ensure that the \hat{B}_b are also consistent with cluster properties.

The techniques used the N -particle case[1][4] can be applied to the modified operators (1) and (3).

All interactions appear in the $\bar{U}_b(\Lambda, a)$ as b ranges over all partitions with at least two equivalence classes, except the \mathcal{N} -body interactions. The mass operators \bar{M}_b for the different $\bar{U}_b(\Lambda, a)$ can be combined, using an Ursell cluster expansion, to define

$$\bar{M} := \sum_b (-)^{n_b} (n_b - 1)! \hat{B}_b \hat{M}_b \hat{B}_b^\dagger,$$

which commutes with the non-interacting \mathcal{N} -body spin \hat{j}_0 and the non-interacting \hat{F}_0 . Simultaneous eigenstates of \bar{M} , \hat{j}_0 , \hat{F}_0 define a basis that transforms irreducibly and thus a representation $\bar{U}(\Lambda, a)$ of the \mathcal{N} -body dynamics.

The representation $\bar{U}(\Lambda, a)$ fails to satisfy the cluster property (2) because of the transformations \hat{B}_b appearing in Eq. (3). This can be remedied by defining

$$\hat{B} = \frac{1 - i\hat{\beta}}{1 + i\hat{\beta}} \quad \hat{\beta} = \sum_b (-1)^{n_b} (n_b - 1)! \hat{\beta}_b$$

$$\hat{\beta}_b := i \frac{\hat{B}_b - \hat{I}}{\hat{B}_b + \hat{I}}.$$

When the Cayley transforms, $\hat{\beta}_b$, are bounded the operator \hat{B} is a scattering equivalence and the representation

$$\hat{U}(\Lambda, a) := \hat{B}^\dagger \bar{U}(\Lambda, a) \hat{B}$$

satisfies cluster properties (2) for suitable short-ranged [3] interactions.

This construction shows how the method for constructing relativistic quantum theory of a fixed number of particles can be extended to treat a class reactions with particle production. Conservation of atomic parton number allowed the separation of the difficulties associated with particle production from the difficulties associated with having an infinite number of degrees of freedom.

References

- [1] F. Coester and W.N. Polyzou: Phys. Rev. D **26** 1348 (1982)
- [2] E. P. Wigner: Ann. Math. **40**, 141 (1939)
- [3] W.N. Polyzou: J. Math. Phys. In press; nucl-th/0201013
- [4] W. Polyzou: Ann. Phys. 193, 367(1989)