

# Method of multiple internal reflections in description of $\alpha$ -decay of nucleus \*

Sergei P. Maydanyuk<sup>†</sup>, Vladislav S. Olkhovsky<sup>‡</sup> and Sergei V. Belchikov<sup>§</sup>

*Institute for Nuclear Research, National Academy of Sciences of Ukraine*

*prosp. Nauki, 47, Kiev-28, 03680, Ukraine*

18 Dec 2001

A spatial-time method of description of  $\alpha$ -decay of a nucleus is presented. The method is based on the quantum mechanical model of the  $\alpha$ -decay of a compound nuclear system with the barrier.  $\alpha$ -decay dynamics is described on the basis of multiple internal reflections of wave packets from boundaries of the decay barrier. The method allows to study this process in details at any time moment or point of space. According to the method, some time before the time moment of the nuclear decay the nucleus makes oscillations. In the calculations of the time duration of the  $\alpha$ -decay of the compound nucleus we propose to take into account the time durations of these oscillations. The method of the multiple internal reflections allows to calculate the time of such oscillations before the  $\alpha$ -decay and the total  $\alpha$ -decay time.

**PACS numbers:** 03.40.Kf, 03.50.De, 03.65.Nk, 41.20.Jb

**Keywords:** alpha-decay, multiple internal reflections, wave packet, tunneling, times, penetrability and reflection coefficients

An approach for description of a motion of a non-relativistic particle above a potential barrier in the one-dimensional problem with taking into account of multiple internal reflections of plane waves, which describe the propagation of this particle, between boundaries of this barrier was studied in number of papers [1, 2, 3] and has been known for the long time. Here, each such wave was considered as stationary, on their basis only the stationary solutions for wave functions (WF) were obtained and there was a difficulty for description of tunneling of the particle under the barrier.

We have generalised this approach, developing the nonstationary method of solution of the tunneling problem of the non-relativistic particle through the one-dimensional barrier, which is constructed with use of the multiple internal reflections of nonstationary wave packets (WP)

---

\*The paper is published (in Russian) in Journ. "Problems of atomic science and technology" (RFNC-VNIIEF, Sarov, Russia), Vol. 1 (2), p.16–19 (2002)

<sup>†</sup>E-mail: maidan@kinr.kiev.ua

<sup>‡</sup>E-mail: olkhovsk@kinr.kiev.ua

<sup>§</sup>E-mail: belchik@kinr.kiev.ua

between the barrier boundaries [4]<sup>1</sup>. The non-stationary use of WPs in the processes of the multiple internal reflections, fulfilled on the basis of papers [6, 7, 8], allows from the physical point of view to motivate correctly the approach for tunneling description on the basis of the multiple internal reflections. The method allows to study in details the tunneling of the particle at any point of space and to calculate different times during tunneling (or propagation). In calculation of the time durations of tunneling this method has shown itself as simple and effective enough.

After developing the method formalism for the spherically symmetrical problems, with its help a problem of elastic scattering of the non-relativistic particle upon nucleus was solved and some new characteristics, describing such process, were obtained [4].

As a further development of the time analysis of nucleus processes [4, 6, 7, 8], for the first time we present a non-stationary method of the description of the nuclear  $\alpha$ -decay on the basis of the multiple internal reflections of the wave packets relatively on the barrier boundaries. The method is constructed on the basis of the quantum mechanical one-particle model of the decay of a compound nucleus with use of the barrier [9] and allows to fulfill the space-time analysis of the decay process. With its help, some new stationary and time parameters for the description of the  $\alpha$ -decay of the nucleus are found.

Now we describe briefly the method in solution of the one-dimensional problem of the particle tunneling through the square barrier of the following form:

$$V(x) = \begin{cases} 0, & \text{at } x < 0 & \text{(region I);} \\ V_1, & \text{at } 0 < x < a & \text{(region II);} \\ 0, & \text{at } x > a & \text{(region III).} \end{cases} \quad (1)$$

Let's assume, that the particle with the mass  $m$ , falling on the barrier from the left side and tunneling through it at the energy level  $E$ , which locates below than the barrier heights  $V_1$ . A general stationary solution for WF for this process has the form:

$$\varphi(x) = \begin{cases} e^{ikx} + A_R e^{-ikx}, & \text{at } x < 0 & \text{(region I);} \\ \alpha e^{\xi x} + \beta e^{-\xi x}, & \text{at } 0 < x < a & \text{(region II);} \\ A_T e^{ikx}, & \text{at } x > a & \text{(region III),} \end{cases} \quad (2)$$

where  $k = \frac{1}{\hbar} \sqrt{2mE}$ ,  $\xi = \frac{1}{\hbar} \sqrt{2m(V_1 - E)}$  and  $\alpha$ ,  $\beta$ ,  $A_T$ ,  $A_R$  are the normalization factors.

According to the method, we describe tunneling of the particle under the barrier with use of the WP sequentially by steps of its transitions through each boundary. In the beginning, on the first step we consider the particle, which is incident upon the first boundary of the barrier at point  $x = 0$  from the left side. This particle is reflected from the boundary with certain probability and transits through this boundary with another certain probability, tunneling inside the barrier further. One can describe the incident, transitted and reflected propagations (with taking into account of tunneling) of the particle relatively of the first boundary of the

---

<sup>1</sup>in [5] the extended version of this method is presented, where its proof, main definitions and peculiarities are given in study of the simplest problem of tunneling, its application to the analysis of the particle tunneling through the one-dimensional barriers of more complicated forms and to the analysis of the particle in its spherically symmetric scattering upon nucleus are studied, tunneling of photons is studied by this method (for the first time)

barrier with use of the WP of the following form:

$$\Psi_i(x, t) = \int_0^{+\infty} g(E - \bar{E}) \varphi_i(k, x) e^{-iEt/\hbar} dE, \quad (3)$$

where  $\bar{E}$  is the average energy of the particle. The incident, transmitted and reflected stationary components  $\varphi_i(x)$  can be found from

$$\varphi(x) = \begin{cases} e^{ikx} + A_R^0 e^{-ikx}, & \text{at } x < 0; \\ \beta^0 e^{-\xi x}, & \text{at } 0 < x < a. \end{cases} \quad (4)$$

The constant factors  $A_R^0$  and  $\beta^0$  are determined from the continuity conditions of the non-stationary WF (constructed on the basis of the WP in regions I and II) and its derivative at point  $x = 0$ .

On the second step, we consider the particle, tunneling along the region II and incidenting upon the second boundary of the barrier at point  $x = a$ . It transits through this boundary with the certain probability and is reflected from it with the certain probability also. One can describe this process with use of the WP of the form (3), where one need to use such stationary WF:

$$\varphi(x) = \begin{cases} \alpha^0 e^{\xi x} + \beta^0 e^{-\xi x}, & \text{at } 0 < x < a; \\ A_T^0 e^{ikx}, & \text{at } x > a. \end{cases} \quad (5)$$

The constant factors  $\alpha^0$  and  $A_T^0$  are obtained from the continuity conditions of the WF and its derivative at point  $x = a$ .

On the third step, the particle, reflected from the second boundary of the barrier, tunnels along the region II back and is incident upon the first boundary at point  $x = 0$ . It transits through this boundary with the certain probability and is reflected from it with another probability. One can describe this process with use of the WP (3), where one can use the stationary WF of the form:

$$\varphi(x) = \begin{cases} \alpha^0 e^{\xi x} + \beta^1 e^{-\xi x}, & \text{at } 0 < x < a; \\ A_R^1 e^{-ikx}, & \text{at } x < 0. \end{cases} \quad (6)$$

The constant factors  $\beta^1$  and  $A_R^1$  can be calculated from the continuity conditions of the WF and its derivatives at point  $x = 0$ .

Considering further the process of tunneling (and propagating) of the particle by such a way, we conclude, that any following step of the particle transition or reflection relatively the boundary can be reduced to one of two steps, considered above. For the constant factors one can construct the recurrence relations.

We describe the propagation and reflection of the particle relatively the barrier on the basis of total transmitted and reflected WPs of the form:

$$\begin{aligned} \Psi_{tr}(x, t) &= \int_0^{+\infty} g(E - \bar{E}) \sum A_T^{(i)} e^{ikx - iEt/\hbar} dE, & \text{at } x > a; \\ \Psi_{ref}(x, t) &= \int_0^{+\infty} g(E - \bar{E}) \sum A_R^{(i)} e^{ikx - iEt/\hbar} dE, & \text{at } x < 0. \end{aligned} \quad (7)$$

Here, one can calculate the sums of the factors  $A_T^{(i)}$  and  $A_R^{(i)}$  on the basis of the recurrence relations found earlier:

$$\begin{aligned} \sum_{i=0}^{+\infty} A_T^{(i)} &= \frac{i4k\xi e^{-\xi a - ika}}{F}; & F &= (k^2 - \xi^2)D_- + 2ik\xi D_+; \\ \sum_{i=0}^{+\infty} A_R^{(i)} &= \frac{i4k\xi e^{-\xi a - ika}}{F}; & D_{\pm} &= 1 \pm e^{-2\xi a}. \end{aligned} \quad (8)$$

All expressions for the constant factors sums, obtained by the method of multiple internal reflections, coincide with the corresponding coefficients, calculated with use of standard stationary approaches. Using the transformation

$$i\xi \rightarrow k_2, \quad (9)$$

where  $k_2 = \frac{1}{\hbar} \sqrt{2m(E - V_1)}$ , the expressions for the constant factors at each step, the expressions for the stationary WFs at each step, the total expressions (8) transform into the corresponding expressions for the problem of the particle propagating above the barrier, coinciding with the solutions of the problems, studied in [1, 2, 3].

We fulfill time analysis of the particle tunneling (or propagating) relatively the barrier on the basis of the method of stationary phase [4, 6, 7, 8]. For obtaining of the tunneling and reflecting times it is needed to know the expressions of the incident, transmitted and reflected WPs relatively the barrier. In their calculation the method of multiple internal reflections has shown itself as the most effective in a comparison with other known methods. This perspective is not visible in solution of the one-dimensional problem of the particle tunneling through the rectangular barrier, but is shown brightly in the solution of the problems of tunneling through the barriers of the more complicated forms. For the one-dimensional problem with the rectangular barrier (1) one can find by the method of multiple internal reflections both the components of the incident, transmitted and reflected WPs (see (7) and (8)), calculate many other  $n$ -multiple WPs, and determine on their basis many other detailed time characteristics.

Now we formulate briefly an approach for a space-time description of  $\alpha$ -decay of a nucleus, based on the method of multiple internal reflections. We consider the decay process on the basis of the quantum mechanical one-particle model of the decay of a compound system with use of the barrier of the following form:

$$V(r) = \begin{cases} -V_0, & \text{at } r < R_1 & \text{(region I);} \\ V_1, & \text{at } R_1 < r < R_2 & \text{(region II);} \\ 0, & \text{at } r > R_2 & \text{(region III).} \end{cases} \quad (10)$$

Let's study a case, when the moment is  $l = 0$  and the decay occurs at the energy levels, located below the barrier heights (i. e. the  $\alpha$ -particle tunnels through the barrier from the region I into the region III). A general solution of the stationary WF has the following form:

$$\varphi(r, \theta, \phi) = \frac{\chi(r)}{r} Y_{00}(\theta, \phi); \quad \chi(r) = \begin{cases} A(e^{-ik_1 r} + e^{ik_1 r}), & \text{at } r < R_1; \\ \alpha e^{\xi r} + \beta e^{-\xi r}, & \text{at } R_1 < r < R_2; \\ S e^{ikr}, & \text{at } r > R_2, \end{cases} \quad (11)$$

where  $Y_{lm}(\theta, \phi)$  is the spherical function,  $k_1 = \frac{1}{\hbar} \sqrt{2m(E + V_0)}$ ,  $\xi = \frac{1}{\hbar} \sqrt{2m(V_1 - E)}$ .

According to the method, we describe the tunneling of the  $\alpha$ -particle through the barrier with use of WP consecutively by steps of its propagation relatively each boundary, like the one-dimensional problem. At the first step, we study the beginning of the decay of the compound nucleus. We consider the  $\alpha$ -particle, which propagates inside the region I and is incident upon the internal boundary of the barrier at point  $r = R_1$ . Here, we determine the expression for the incident WP, which we use in definition of the time moment of the  $\alpha$ -particle fall upon the barrier on the inside (*the time moment of the beginning of the decay of the compound nucleus*, according to the method). At the first step, the transmitted and reflected WPs start the processes of the  $\alpha$ -particle leaving from the barrier region and its delay inside the internal region I (the region of the compound nucleus).

Further, we study the tunneling of the particle through the barrier and its oscillations inside the region I at the following steps. Here, one can reduce any step of the WP propagation relatively the barrier boundaries to one of 4 independent steps. With use of their analysis the recurrence relations for the calculation of the constant factors for the arbitrary step  $n$  are constructed. The total non-stationary WF in the each region, which takes into account the multiple reflections (and transitions) of WP relatively the barrier boundaries, can be present in the form of series of the convergent and divergent WPs, which are coincided and can be calculated by use of the recurrence relations obtained earlier. In result, we find the incident, total transmitted and reflected WPs relatively the barrier:

$$\begin{aligned}\chi_{inc}(r, t) &= \int_0^{+\infty} g(E - \bar{E})\Theta(V_1 - E)A_{inc}e^{ikr - \frac{iEt}{\hbar}} dE, \\ \chi_{tr}(r, t) &= \int_0^{+\infty} g(E - \bar{E})\Theta(V_1 - E)A_{inc}C_{tr}e^{ikr - \frac{iEt}{\hbar}} dE, \\ \chi_{ref}(r, t) &= \int_0^{+\infty} g(E - \bar{E})\Theta(V_1 - E)A_{inc}C_{ref}e^{-ikr - \frac{iEt}{\hbar}} dE,\end{aligned}\quad (12)$$

where

$$\begin{aligned}C_{tr} &= \frac{i4k_1\xi}{(k + i\xi)(k_1 + i\xi + (k_1 - i\xi)\exp(2ik_1R_1))} \times \\ &\times \left(1 - \frac{(i\xi - k)i\xi - k_1 - (i\xi + k_1)\exp(2ik_1R_1)}{(i\xi + k)i\xi + k_1 - (i\xi - k_1)\exp(2ik_1R_1)} \exp(2\xi(R_1 - R_2))\right)^{-1}; \\ C_{ref} &= \frac{C_1}{C_2}; \\ C_1 &= \frac{i\xi - k_1}{i\xi + k_1} \exp(2ikR_1) \left(1 - \frac{(i\xi - k)(i\xi - k_1)}{(i\xi + k)(i\xi + k_1)} \exp(2\xi(R_1 - R_2))\right) + \\ &+ \frac{4ik\xi(k - i\xi)}{(k + i\xi)^2(k_1 + i\xi)} \exp(2\xi(R_1 - R_2) + 2ik_1R_1); \\ C_2 &= \left(1 + \frac{k_1 - i\xi}{k_1 + i\xi} \exp(2ikR_1)\right) \left(1 - \frac{(i\xi - k)(i\xi - k_1)}{(i\xi + k)(i\xi + k_1)} \exp(2\xi(R_1 - R_2))\right) + \\ &+ \frac{4ik\xi(k - i\xi)}{(k + i\xi)^2(k_1 + i\xi)} \exp(2\xi(R_1 - R_2) + 2ik_1R_1),\end{aligned}\quad (14)$$

and  $\Theta$ -function satisfies to the condition:

$$\Theta(\eta) = \begin{cases} 0, & \text{at } \eta < 0, \\ 1, & \text{at } \eta > 0. \end{cases}\quad (15)$$

On the basis of these expressions one can calculate the coefficients of penetrability and reflection of the  $\alpha$ -particle relatively the barrier.

We fulfill the time analysis of the  $\alpha$ -decay of the compound nucleus on the basis of papers [4, 6, 7, 8]. Here, we calculate the *time of the  $\alpha$ -decay of the compound nucleus*, which we define as a difference between the time moment of the  $\alpha$ -particle leaving from the barrier region at point  $r = R_2$  (*the time moment of finishing of the decay of the compound nucleus*, according to the method) and the time moment of the beginning of the  $\alpha$ -decay of the compound nucleus. The time moment of the  $\alpha$ -particle leaving from the barrier we define on the basis of the expression for the total transmitted WP of the form (12) with taking into account of (13). According to the calculations, the time of the  $\alpha$ -decay of the compound nucleus has the form:

$$\tau_{tr} = \hbar \frac{\partial}{\partial E} \arg C_{tr}. \quad (16)$$

In accordance with the analysis of the consequent transmissions and reflections of the multiple WP relatively the barrier boundaries, we note that the total WP leaves the region I not at the time moment of the  $\alpha$ -decay of the compound nucleus, but after the some period of time. One can suppose, that at the beginning of the decay the particle makes oscillations inside the region I (this follows from the analysis of WF in this region). A probability of its existence inside the region I decreases evenly, but at the beginning of the decay process it is larger than inside the regions II and III. We define the delay time as a difference between the time moment of the falling of the total WP inside the region I upon the internal boundary of the barrier at point  $r = R_1$  and the time moment of the beginning of the  $\alpha$ -decay of the compound nucleus.

One can calculate the factor  $A_{inc}$ , used in (12), from the normalization condition of WF, and it does not influence on the values of the penetrability and reflection coefficients, and on the values of the times of the  $\alpha$ -decay and delay.

In conclusion we note, that the approach for description of the particle tunneling (and propagating) through the barrier on the basis of the processes of the multiple internal reflections is non-stationary and it reflects *wave properties* of the particle in its tunneling (or propagation). Applying of the method of multiple internal reflections, which uses such approach, to the description of the  $\alpha$ -decay of the spherical nuclei is the further development of the method of multiple internal reflections, which was applied earlier for solution of the problems of elastic scattering of particles upon spherical nuclei and the one-dimensional problems of tunneling of the particles through barriers of the different forms. In the problem of the nuclear  $\alpha$ -decay with its use the effect of the oscillations of the compound nucleus is opened, which takes place at the beginning of the decay process. We propose to take into account such effect in calculations of the decay time of the compound nucleus.

- [1] K. W. McVoy, L. Heller and M. Bolsterli, *Reviews of Modern Physics* **39**, 245 (1967).
- [2] A. Anderson, *Multiple scattering approach to one-dimensional potential problems*, *American Journal of Physics* **57** (3), 230–235 (1989).
- [3] J. H. Fermor, *Quantum-mechanical tunneling*, *American Journal of Physics* **34**, 1168–1170 (1966).

- [4] V. S. Olkhovsky and S. P. Maydanyuk, *Method of multiple internal reflections in description of tunneling evolution through barriers*, Ukrainian Physical Journal **45** (10), 1262–1269 (2000), [arXiv:nucl-th/0406035].
- [5] S. P. Maydanyuk, V. S. Olkhovsky and A. K. Zaichenko, *The method of multiple internal reflections in description of tunneling evolution of nonrelativistic particles and photons*, Journal of Physics Study **6** (1), 1–16 (2002), [arXiv:nucl-th/0407108].
- [6] V. S. Olkhovsky and E. Recami, *Recent developments in the time analysis of tunneling processes*, Physics Reports **214** (6), 339–356 (1992).
- [7] F. Cardone, R. Mignani and V. S. Olkhovsky, *About superluminal propagation of an electromagnetic wavepacket inside a rectangular waveguide*, Journal de Physique. I (France) **7** (10), 1211–1219 (1997).
- [8] V. S. Olkhovsky and S. P. Maydanyuk, *On the evolution of particle transitions from one well to another in a double-well potential* Journal of Physics Study **3** (1), 12–24 (1999); [arxiv:quant-ph/0311128].
- [9] A. I. Baz, Y. B. Zeldovich and A. M. Perelomov, *Rasseyanie, reaktsii i raspadi v nerelyativistskoi kvantovoi mehanike* (Nauka, Moskva, 1971) — [in Russian].