# Backbending phenomena in light nuclei at $A \sim 60$ mass region

S. U. El-Kameesy,<sup>1,\*</sup> H. H. Alharbi,<sup>2,†</sup> and H. A. Alhendi<sup>3,‡</sup>

<sup>1</sup>Department of Physics, Faculty of Science, Ain-Shams University, Cairo, Egypt.

<sup>2</sup>National Center for Mathematics and Physics, KACST, P.O. Box 6086, Riyadh 11442, Saudi Arabia,

<sup>3</sup>Department of Physics and Astronomy, College of Science,

King Saud University, P.O. Box 2455, Riyadh 11454, Saudi Arabia

(Dated: December 15, 2018)

Recent studies of the backbending phenomenon in medium light weight nuclei near  $A \sim 60$  expanded greatly our interest about how the single particle orbits are nonlinearly affected by the collective motion. As a consequence we have applied a modified version of the exponential model with the inclusion of paring correlation to describe the energy spectra of the ground state bands and/or the backbending phenomenon in mass region at  $A \sim 60$ . A firm conclusion is obtained concerning the successful validity of the proposed modified model in describing the backbending phenomenon in this region. Comparison with different theoretical descriptions is discussed.

PACS numbers: 21.10.-k, 21.60.+v, 21.90.+f

### I. INTRODUCTION

Investigations of the ground state bands of nuclei at  $A \sim 60$  mass region have recently become a particularly interesting subject in nuclear structure studies [1–5]. These nuclei exhibit a range of interesting features, including oblate and prolate deformations as well as rapid variations in shape as a function of both spin and particle number. The sudden disappearance of E2 strength at certain spins indicates a shape change and requires the inclusion of upper pf configuration [6]. The shell model predictions have allowed calculations in the full fp model space. These calculations have shown that the collective properties of rotor like energies, backbending and large B(E2) values can be reproduced.

Hara *et al* [5]. studied the backbending mechanism of <sup>48</sup>Cr within the projected shell model (PSM) [7], which has been successful in describing well deformed heavy nuclei and those of transitional region [8], it was concluded that the backbending in <sup>48</sup>Cr is due to a band crossing involving an excited band built on simultaneously broken pairs of neutrons and protons in the intruder subshell  $f_{7/2}$ . This result differs from that of Tanaka *et al* [9] based on the Cranked Hartee–Fock–Bogoliubov (CHFB), which claims that the backbending in <sup>48</sup>Cr is not due to level crossing.

The application of the generator coordinate method (GCM) has showed that the backbending in  $^{48}$ Cr can be interpreted as due to crossing between the deformed and spherical bands [7]. Accordingly, while the backbending phenomenon in medium heavy nuclei are well described and commonly understood as a band crossing phenomenon involving strong pairing correlation [10], the origin of backbending in medium light nuclei has been debated. Additionally, the role of the paring force in the backbending phenomenon is not clearly outlined.

The interest of the  $1f_{7/2}$  nuclei has been extended to levels above the  $1f_{7/2}$  band termination, in particular in connection with a possible building up of superdeformation [11], where the quality of SM calculations is probably not as good because of the possible contributions from the SD shell and the  $1g_{9/2}$  orbital [2]. Cranking model analysis of <sup>80</sup>Br energy levels reveals a signature inversion at a spin of  $12\hbar$  and a probably neutron alignment at  $\hbar\omega \approx 0.7$  MeV. The results are discussed within the framework of the systematics of similar bands in the lighter Br isotopes and the cranked–shell model [12].

The present study has been initiated due to the above contradictions of different models in describing the backbending phenomenon in medium light nuclei. Our study is based on applying a modified version of the exponential model with pairing attenuation [13]. It is hoped by such work to have a strong evidence that the pairing force contribution still plays an effective role in the backbending mechanism in this mass region at  $A \sim 60$ . In the following section we briefly present the model and in the next section the model is applied to some even-even medium light nuclei. Finally the last section contains our conclusion.

<sup>\*</sup>Electronic address: saimrkhamisy@yahoo.com

<sup>&</sup>lt;sup>†</sup>Electronic address: alharbi@kacst.edu.sa

<sup>&</sup>lt;sup>‡</sup>Electronic address: alhendi@ksu.edu.sa

 $\mathbf{2}$ 

### II. MODEL DESCRIPTION

Sood and Jain [14] have previously developed an exponential model based on the exponential dependence of the nuclear moment of inertia on pairing correlation [15]. They gave the following relation:

$$E(I) = \frac{\hbar^2}{2\varphi_0} I(I+1) Exp\left[\Delta_0 \left(1 - \frac{I}{I_c}\right)^{\frac{1}{2}}\right]$$
(1)

Excellent results have been obtained by means of this approach in describing the ground state bands in deformed nuclei up to the point where backbending occurs. They selected the value  $18\hbar$  for  $I_C$  as an input cutoff corresponds to the point where the rotational frequency  $\omega$  in the  $\varphi - \omega^2$  plots reaches a minimum value and the pairing correlations disappear completely. Zhou and Zheng [17] have demonstrated that  $I_C = 85\hbar$  is a suitable choice in their calculations concerning superdeformed bands near  $A \sim 190$ , since the pairing correlation in that region is still strong event at very high rotational frequency. For medium light nuclei,  $I_C$  can take values smaller than  $18\hbar$  because the backbending phenomenon in this region ( $A \approx 60$ ) lies at spin  $I \approx 10\hbar$  [4]. These works led us to use a suitable  $I_C$  values to represent both the variation of the moment of inertia and the paring correlation and to give the model the ability to describe well the  $\varphi - \omega^2$  plot regions, in particular the forward and down-bending regions, which lie after the backbending region. The modified version of the exponential model with pairing attenuation has the following form [13, 16]:

$$E(I) = \frac{\hbar^2}{2\varphi_0} I(I+1) Exp\left[\Delta_0 \left(1 - \frac{I}{I_c}\right)^{\frac{1}{\nu}}\right]$$
(2)

Where  $\varphi_0$ ,  $\Delta_0$  and  $\nu$  are the free parameters of the model, which are adjusted to give a least-square fit to the experimental data. This approach is supported by Ma and Rasmussen suggestion that there is an exponential dependence of the moment of inertia on the parameter  $\nu$  for a wide range  $\nu$  values [18].

## III. APPLICATION TO SOME EVEN-EVEN MEDIUM LIGHT NUCLEI

The anomalous behavior, i.e. backbending of several medium light even-even nuclei (Zn, Ge, Se, Kr, Sr), has been studied using our improved modified version of the exponential model with pairing attenuation. The parameters of the model (Table 1) were determined by means of a least-square fitting procedure involving the experimental known energy levels [19].

The plots of the calculated data of  $2\varphi_I/\hbar^2$  versus  $(\hbar\omega)^2$  for these isotopes are given in figure 1, where the experimental data are also presented. From the excitation energies E(I) of the yrast bands we deduce the moment of inertia and the squared rotational frequency  $\omega^2$  by using the well-known relations

$$\frac{2\varphi}{\hbar^2} = \frac{4I - 2}{E(I) - E(I - 2)} , \qquad (3)$$

and

$$(\hbar\omega)^{2} = (I^{2} - I + 1) \left[\frac{E(I) - E(I - 2)}{2I - 1}\right]^{2}$$
(4)

In figure 1 the experimental data show a clear evidence of backbending phenomenon in all the presented nuclei at  $I = 8 - 12\hbar$ . It is clear from the same figure that the predictions of the applied improved exponential model reproduce very well the backbending phenomenon in those nuclei and its application improves as A increases. This result may give an indication that the pairing force contribution to the backbending phenomenon increases as A increases in the mass region under investigation. Another noticeable success of the model is shown in Figure 1 "concerning <sup>68</sup>Ge, <sup>72</sup>Se, <sup>78</sup>Kr and <sup>80</sup>Sr" where the forward and down-bending regions are very well described by its calculations.

 $\mathbf{I}_{c}$ Nucleus  $2\varphi_0/\hbar^2$  $\mathbf{\Delta}_0$ ν  $^{50}\mathrm{Cr}$ 39.8033 1.66542 0.678977 26 $^{62}$ Zn 16.1483 1.30409 0.313542 18  $^{64}\mathrm{Ge}$  $15.5511 \ 1.38726 \ 0.234353 \ 18$  $^{68}\mathrm{Ge}$ 25.516 1.90454 0.401219 20  $^{74}\mathrm{Kr}$ 47.5087 1.03889 0.428975 40  $^{78}{
m Kr}$  $42.5699 \ 1.31551 \ 0.124932 \ 80$  $^{80}$ Kr 35.0581 1.53869 0.452602 20  $^{72}\mathrm{Se}$  $38.7645 \ 1.45564 \ 0.375074 \ 30$  $^{74}\mathrm{Se}$ 40.435 1.26539 0.446761 28  $^{76}\mathrm{Se}$ 40.7577 1.60714 0.180245 50 $^{78}\mathrm{Se}$  $33.3453 \ 1.50423 \ 0.466925 \ 20$  $^{78}\mathrm{Sr}$ 40.4515 0.68491 0.125873 80 $^{80}\mathrm{Sr}$  $43.9731 \ 1.03228 \ 0.146307 \ 80$  $^{82}\mathrm{Sr}$  $48.9001 \ 1.25575 \ 0.173081 \ 80$ 

TABLE I: The fitting parameters of the present model.

FIG. 1: Calculated and observed moment of inertia  $2\varphi/\hbar^2$  vs.  $(\hbar\omega)^2$  for yrast levels of some light nuclei. The dots represent experimental values.

# IV. CONCLUSION

The present results of the improved exponential model with pairing attenuation give a firm confirmation that the backbending in medium light nuclei at low spins  $(I = 8 - 12\hbar)$  can be interpreted due to paring force which supports the band crossing mechanism in analogy with the earlier calculations [5] based on the projected shell model (PSM) and the generator coordinator method (GCM). Furthermore, our simple modified formula is able to describe well the forward and down-bending regions of the  $\varphi - \omega^2$  plots.

- [1] J.A. Cameron et al., Phys. Lett. B 235, 239 (1990).
- [2] F. Brandolini *et al.*, Nucl. Phys. A **642**, 387 (1998).
- [3] E. Caurier, A.P. Zuker, A. Poves, and G. Martinez-Pinedo, Phys. Rev. C 50, 225 (1994); E. Caurier, F. Nowacki and A. Poves, Phys.Rev.Lett. 95, 042502 (2005).
- [4] V. Velázquez, J.G. Hirsch, Y. Sun, Nucl. Phys. A 686, 129 (2001).
- [5] K. Hara, Y. Sun, and T. Mizusaki, Phys. Rev. Lett. 83, 1922 (1999).
- [6] J.A. Cameron *et al.*, Phys. Rev C 58, 808 (1998).
- [7] K. Hara, Y. Sun, Int. J. Mod. Phys. E 4, 637 (1995); G.-L. Long and Y. Sun, Phys. Rev. C 63, 021305 (2001).
- [8] J. A. Sheikh, K. Hara, Phys. Rev. Lett. 82, 3968 (1999).
- [9] T. Tanaka, K. Iwasawa, and F. Sakata, Phys. Rev. C 58, 2765 (1998).
- [10] R.A. Sorenson, Nucl. Phys. A 269, 301 (1976).
- [11] S.M. Lenzi et al., Phys. Rev. C 56, 1313 (1997).
- [12] I. Ray et al., Nucl. Phys. A 678, 258 (2000).
- [13] H. A. Alhendi, H. H. Alharbi, S. U. El-Kameesy, "Improved Exponential model with pairing attenuation and the backbending phenomenon", nucl-th/0409065. (Submitted)
- [14] P. C. Sood and A. K. Jain, Phys Rev. C 18, 1906 (1978).
- [15] A. Bohr and R. Mottelson, Nuclear structure. Voll II, Benjamin, W.A. Inc., New York, Amesterdam (1975).
- [16] H. H. Alharbi, H. A. Alhendi, and S. U. El-Kameesy, J. Physics G, (to appear)
- [17] Shan-Gui Zhou and Chunkai Zheng, Phys. Rev. C 55, 2324 (1997).
- [18] C.W. Ma and J.O. Rasmussen, Phys. Rev. C 9, 1083 (1974).
- [19] J.K. Tuli, Evaluated nuclear structure data file, Nucl. Instr. Meth. in Phys. Res. A 369, 506 (1996).

