

Non-linear mean field dynamics in the nuclear spinodal zone

G.F. Burgio, M. Baldo, and A. Rapisarda

Università di Catania and INFN Sezione di Catania

Corso Italia 57, I-95129 Catania, Italy

(June 4th 1993 - Catania University preprint no.93/20)

Abstract

We demonstrate, by numerical simulations, that the dynamics of nuclear matter mean field inside the spinodal region is chaotic. Spontaneous symmetry-breaking - no explicit fluctuating term is considered - occurs leading to wild unpredictable density fluctuations. A proper recipe to calculate an average Lyapunov exponent in this multidimensional phase space is introduced. The latter is calculated for different values of the density in order to characterize in a quantitative way the chaotic and regular regions. It is argued that the mean field chaoticity can be the main mechanism of the nuclear multifragmentation occurring in the intermediate energy reactions.

25.70.Pq, 24.60.Lz, 21.65.+f

arXiv:nucl-th/9306008v1 7 Jun 1993

Heavy ion collisions in the intermediate energy regime between 20 and 100 MeV/nucleon can result in multifragment emissions, where a number of intermediate mass fragments (IMF : $Z \geq 2$) is observed. This disassembly of the nuclear system, which can include a substantial fraction of the whole colliding system, falls under the generic name of "nuclear multifragmentation", and is presently under intense study in different laboratories. The main mechanism which is responsible of this phenomenon has not yet been determined. Both statistical models and dynamical simulations [12] have been used to describe the cluster formation in the multifragmentation regime. The role of mechanical instabilities, either of the Raleigh type [2], or associated with ring formation [3], has been also stressed recently. These calculations provide suggestions on the possible dominant path followed by the nuclear system along the multifragmentation process. The onset of instability in the spinodal zone of the nuclear matter Equation of State (EOS) has been studied in ref. [4], by analyzing the corresponding linear response for imaginary frequency as a function of the wave number and of the density.

In this letter we establish, by numerical simulations, that the dynamical evolution of nuclear matter mean field in the spinodal region is chaotic. We estimate the corresponding Lyapunov exponent which characterizes the rate of divergence between the trajectories in density space. This result can have far reaching consequences for the whole picture of the multifragmentation process. In particular, a) the relevance of the mean trajectory, which is an essential ingredient in most of the simulations with kinetic equations, appears questionable if the spinodal region is reached; b) the onset of chaoticity implies, as we explicitly verify numerically, strong non-linearity of the evolution, and therefore strong coupling between different modes; c) the occurrence of chaotic motion suggests that thermodynamical quantities, like entropy, are not necessarily appropriate for characterizing the system dynamics, but they should be substituted by others, based on chaotic evolution concepts, like the Kolmogorov entropy [5]. Other more quantitative consequences will be discussed in the following. The numerical analysis is performed by solving the Vlasov equation

$$\frac{\partial f}{\partial t} + \{H[f], f\} = 0 \quad (1)$$

on a two-dimensional lattice, according to the method of ref. [6]. In eq. (1) $H[f]$ is the self-consistent effective one-body Hamiltonian. In this approach the one-body density distribution is represented on a grid at variance with the more commonly used test particle methods [7,8]. The former has been preferred since it allows a better control of the numerical accuracy and conservation laws. The single particle phase space is divided into cells and nuclear matter is confined in a large torus with periodic boundary conditions. The side lengths of the torus are equal to $L_x = 51 \text{ fm}$ and $L_y = 15 \text{ fm}$. In order to achieve sufficient accuracy, very small cells having $\delta x = \frac{1}{3} \text{ fm}$ and $\delta p = 40 \text{ MeV}/c$ are employed. The time integration step was chosen equal to $0.5 \text{ fm}/c$. We adopt a simplified Skyrme interaction for the effective one-body field which is averaged over the y-direction [6]. We employ a Fermi momentum at saturation of $P_F = 260 \text{ MeV}/c$, which gives a density in two dimensions equal to $\rho_0 = 0.55 \text{ fm}^{-2}$. The mean field evolution is treated by means of a standard matrix technique [6]. The main indication of chaotic behaviour of a dynamical system is the extreme sensitivity of its evolution to the initial conditions. This is illustrated in figs. 1a) and 1b). At the initial time nuclear matter presents a density wave, with an amplitude $\Delta\rho_1 = 10^{-2}\rho_0$, and with a wave number $k = 2\pi(5/L_x)$. The local momentum distribution is assumed to be the one of a Fermi gas at a temperature of 3 MeV . This corresponds to a very small zero sound oscillation. The initial conditions are varied in a very simple way, namely by changing the density ρ by an amount $\Delta\rho = 10^{-4}\rho_0$. No appreciable change of the results are obtained with different prescriptions of modifying the initial conditions, as discussed later. We checked that in all the calculations here presented the total energy was conserved up to 1% and that the particle number remains constant during the whole evolution. For a density equal to the saturation density ρ_0 , fig.1 (a), the density profiles of nuclear matter corresponding to different initial conditions keep close to each other for the whole considered time interval. This is typical of the "regular" region of a dynamical system, and indicates the stability of the dynamics with respect to small perturbations. On

the contrary, if the density ρ is chosen well inside the spinodal region, in this case $\rho = \frac{1}{2}\rho_0$, the density profiles, after a short time, differ completely between each other, even if initially they are close within a part over 10^4 . This is the typical behaviour of a dynamical system in a chaotic region, where very small initial perturbations are rapidly amplified. The relevant feature of fig.1 (b) is that the two density profiles do not differ simply in their amplitudes, as it would occur if the linear response regime would be valid, but mainly in their shape, which can occur only in a regime of strong non-linearity. We have checked that the results are stable with respect to the increase of the precision in the numerical integration. In fact, a decrease of the time integration step does not affect the results, and therefore the divergence in shape of the trajectories is an intrinsic dynamical property of the nuclear mean field in this region. The chaotic dynamics is intimately related with the strong non-linearity of the evolution, which implies a strong coupling among the different modes. This can be checked by Fourier transforming the density profiles at different times. We found that the different wave numbers increase their mixing more and more as the evolution proceeds. At the latest stage a wide range of wave numbers result to be excited, despite the fact that only one wave number k is initially populated. This point is illustrated in fig.2 for a single trajectory corresponding to the full curve of fig.1 (b). The final wave number spectral density has therefore little resemblance with the initial one, and this feature was found to be independent from the particular initial k . In a linear regime, on the contrary, the system would be equivalent to a set of independent harmonic oscillators, one for each wave number k , with an imaginary frequency, corresponding to the unstable region. Such a system is integrable, and the corresponding flow in phase space is regular. It is important to note that the mixing takes place in an interval of time of about $20 fm/c$, which is short with respect to a typical collision time or expansion time, characteristic of heavy ion collisions at intermediate energy. The calculations here presented are in two dimensions, but the characteristic times of chaotic dynamics are expected to be little dependent on dimensionality. A behaviour similar to that shown in fig.1 (b) was found for a single trajectory in the case of ref. [6] where a dissipative term of Langevin type has been added to the Vlasov equation.

In principle a sensitive dependence on the initial conditions should be present also in that case. Calculations in that direction are in progress. To perform a more quantitative analysis, we follow the standard procedure in the theory of dynamical systems of characterizing the divergence among the phase space trajectories by a set of Lyapunov exponents [5]. Let $d(t)$ be the distance between two trajectories, along an unstable direction in phase space, at a given time t , and define

$$\lambda(t) = \frac{\log(d(t)/d_0)}{t}, \quad (2)$$

where $d_0 = d(0)$. Then the corresponding Lyapunov exponent is obtained by the limit

$$\lambda_\infty = \lim_{d_0 \rightarrow 0} \lim_{t \rightarrow \infty} \lambda(t) \quad (3)$$

provided the limit exists. The very definition of eq. (2) implies that asymptotically for large time and small enough perturbation the divergence is essentially exponential, namely that the equation of motion for the distance between the two trajectories is a linear one. This by no means implies that each trajectory follows linear equations of motion. In numerical applications one has to select a series of small values of d_0 , a set of increasing times t and check that the corresponding values of $\lambda(t)$ stabilize around a definite value λ_∞ . In the present case the actual number of degrees of freedom and therefore the number of Lyapunov exponents is just equal to the number of cells in which the phase space is divided. In order to calculate an average value of the Lyapunov exponent, it is convenient to choose a distance which is connected with a global description of the system. We have therefore chosen as distance between two trajectories the difference in norm between their density profiles

$$d(t) = \sum_i |\rho_i^{(1)}(t) - \rho_i^{(2)}(t)|/N_c, \quad (4)$$

where the index i runs over the N_c cells in ordinary space, and $\rho_i^{(1)}, \rho_i^{(2)}$ are the densities in the cell i for the trajectories 1 and 2 respectively. This definition neglects possible differences which can occur in momentum space between the two trajectories. The definition of eq. (4) is sufficient for the present analysis. It should include the contribution of all the unstable

modes, in particular the ones which dominate for large t . In fig.3 (a) it is displayed the time evolution of $\lambda(t)$ for different values of ρ/ρ_0 , taking an initial $d_0 = 10^{-4}\rho$. It was checked that for d_0 values of this size, $\lambda(t)$ stabilizes and is independent of d_0 . One can distinguish two well defined regimes. For densities ρ well inside the spinodal region, the time dependence of $\lambda(t)$ is quite weak, resulting in the almost straight lines appearing at the top of fig.3 (a). This behaviour unambiguously indicates that the Lyapunov exponent in this case is well defined and that the system evolves in a genuine chaotic way. As the density increases and approaches the upper limit of the spinodal region, the general trend starts to change. In this case $\lambda(t)$ becomes a decreasing function of t which approaches rapidly very small values, indicating that the trajectories actually do not diverge between each other. The general behaviour appearing in fig.3 (a) is typical of dynamical systems possessing a regular and a chaotic region [5]. The actual value of $\lambda(t)$ can be considered a measure of the non-linearity and chaoticity of the system [9]. It has to be also stressed that, when the dynamics is non-linear, the Lyapunov exponent λ_∞ has not to be confused with the rate of increase of the density fluctuations. In the present case we found that, for a given trajectory, the density fluctuations increase with a characteristic time [6] substantially longer than the divergence characteristic time $1/\lambda_\infty$ [6], by a factor ranging from 2 to 3, according to the density and the initial conditions. In fig.3 (b) we show the Lyapunov exponent λ_∞ calculated at different densities. For density values close to the spinodal boundary $\lambda(t)$ shows some oscillations even for the largest times considered. However in this region the values are very small and the general trend of fig.3 (b) does not depend on this uncertainty. The value of λ_∞ has a flat maximum around the density $\rho = 0.35\rho_0$. The curve reported in fig.3 (b) is essentially independent from k , which is in sharp contrast with the linear regime behaviour, for which the imaginary frequency is mainly a linear function of k . It can be instructive to compare the characteristic times $\tau_{MF} = \hbar/\lambda_\infty$, which defines the time scale of the divergence between mean field trajectories, with the single particle characteristic time $\tau_{sp} = \hbar/E_F$, being E_F the Fermi energy at the given density. This is done in table I for a set of densities. One can see that for densities $\rho \leq 0.4\rho_0$ the divergence time is smaller than the

single particle time, and therefore in this region the notion itself of mean field ceases to have any validity. In this region it appears more appropriate to speak of a fully many particle chaotic behaviour [10]. It has to be stressed that, despite the density is small, the mean distance between particles is only slightly larger than at normal density, and therefore the particles can still strongly interact. In central collisions between heavy ions at intermediate energy nuclear matter is expected to be compressed at the initial stage of the reaction, until a highly excited composite nuclear system is formed. Numerous calculations [8], based on kinetic equations as well as on molecular dynamic models [11], show that the momentum distribution is close to spherical symmetry and not too far from thermodynamic equilibrium. In this first stage of the reaction the dynamics is insensitive to small modifications of the initial conditions. Once the maximum compression is reached, nuclear matter starts to expand and can merge into the spinodal region. At this point the dynamics should change dramatically. According to the results presented in this paper, chaotic dynamics sets in, and the corresponding large density fluctuations dominate the cluster formations. The degree of chaoticity and fluctuations depends of course on the details of the dynamics which brings the reaction inside the spinodal region. The ratio between the expansion characteristic time and τ_{MF} could indicate whether chaos has the time to develop. The multifragmentation regime should appear in the energy interval where the chaoticity and the corresponding fluctuations are strong enough to produce several IMF. A confirmation of this scenario could be obtained by means of numerical simulations of heavy ion reactions. Unfortunately, at the moment, the extension of the calculations presented in this paper to the case of collisions between heavy ions appears problematic even in two dimensions. Thus one should find the signature of chaotic dynamics using more phenomenological approaches. The study of fragment size fluctuations and their statistical properties [12], in particular along the lines of the intermittency analysis [13], can be one of the viable possibilities. However, the problem is still completely open and demands a careful analysis, both theoretically and experimentally.

REFERENCES

- [1] D.H.E. Gross and X.Z. Zhang , Phys Lett. **161 B** , 47 (1982); J.P.Bondorf, R.H.Donangelo, I.N.Mishustin, C.J.Pethick, H.Schultz and K.Sneppen, Nucl. Phys. **A443**, 321 (1985); G.Peilert, H.Stöcker, J.Randrup and W.Greiner, Phys. Lett. **B260**, 271 (1991); C.Dorso and J.Randrup, preprint LBL-33120; M.Colonna, M.Di Toro, V.Latora, Nucl. Phys. **545**, 111c (1992).
- [2] L.Moretto, K.Tso, N.Colonna, G.Wozniak, Phys. Rev. Lett. **69**, 1884 (1992).
- [3] W.Bauer, G.Bertsch and H.Schultz, Phys. Rev. Lett. **69**, 1888 (1992).
- [4] H. Heiselberg C.J. Pethick and D.G. Ravenhall Phys. Rev. Lett. **61**, 818 (1988), and Ann. Phys. **223**, 37 (1993); M.Colonna, Ph.Chomaz and J.Randrup, GANIL preprint 1992.
- [5] G.Benettin, L.Galgani and J-M.Strelcyn, Phys. Rev. **A14**, 2338 (1976); A.J.Lichtenberg and M.A.Lieberman, *Regular and Stochastic motion*, Springer-Verlag (1983).
- [6] G.F. Burgio, Ph. Chomaz and J. Randrup, Phys. Rev. Lett. **69**, 885 (1992); G.F. Burgio, Ph. Chomaz and M. Colonna, Catania preprint 1993.
- [7] C.Grégoire, B.Remaud, F.Séville, L.Vinet and Y.Raffray, Nucl. Phys. **A465**, 317 (1987).
- [8] G.Bertsch and S.Gupta, Phys. Rep. **160**, 189 (1988); A. Bonasera, F. Gulminelli and J. Molitoris, Phys. Rep. in preparation.
- [9] It can be instructive to analyze what should result for $\lambda(t)$ if the linear regime would be valid, adopting the present procedure of modifying the initial conditions. In this case the variation $\Delta\rho$ of the density is equivalent to the addition of a mode at $k = 0$. In the linear regime this mode has zero frequency and propagates independently from the mode k , therefore at time t , one has approximately $d(t) \approx \Delta\rho + |\Delta\rho_1 (\exp(\omega + \Delta\omega)t - \exp\omega t)|$, where ω is the growth rate of the mode k at the density ρ and $\omega + \Delta\omega$ the corresponding one at $\rho + \Delta\rho$. For small enough t , and not

too close to the spinodal boundary, $\lambda(t) = \log(d(t)/\Delta\rho)/t \approx \Delta\rho_1\Delta\omega/\Delta\rho \approx \omega(\Delta\rho_1/\rho)$, which is negligibly small, since $\Delta\rho_1 \ll \rho$. The value of $\lambda(t)$ keeps close to zero, for $t \leq \omega^{-1}$, and then after it increases towards the value ω , which can be considered the Lyapunov exponent of the mode k in the linear regime. The actual behaviour of $\lambda(t)$ in the spinodal zone is quite different and this demonstrates that the mean field dynamics cannot be described within the linear approximation in this region.

[10] V.G.Zelevinsky, Nucl. Phys. **A553**, 125c (1993).

[11] Aichelin and H.Stöcker, Phys. Lett. **B163**, 59 (1985)

[12] D.H.E. Gross, Nucl. Phys. **A553**, 175c (1993);

[13] M.Ploszajczak and A.Tucholski, Phys. Rev. Lett. **65**, 1539 (1990).

TABLES

TABLE I. Comparison between the characteristic times $\tau_{MF} = 1/\lambda_\infty$ for the mean field and $\tau_{sp} = \hbar/E_F$ for the single particle at different values of the nuclear density. See text.

ρ/ρ_0	τ_{sp} fm/c	τ_{MF} fm/c
0.7	7.76	1538.
0.6	9.06	28.57
0.5	10.87	13.34
0.4	13.59	9.92
0.3	18.12	9.06
0.2	27.17	10.00

FIGURES

FIG. 1. The evolution for two close trajectories (dotted and full curve) for the nuclear matter density profile is illustrated as a function of time. Two different cases are shown for values of the initial density, (a) $\rho/\rho_0=1$ and (b) $\rho/\rho_0=0.5$. While in case (a) the infinitesimal initial difference keeps constant in time and it is hardly visible, in case (b) it grows exponentially. Note the different scale used in fig.1(a) and 1(b).

FIG. 2. Power spectrum in momentum space of the density profile shown in fig.1(b) as full curve.

FIG. 3. (a) The behaviour of $\lambda(t)$ is plotted vs. time for different values of the initial density. The constant value obtained for $\rho/\rho_0=0.4, 0.5, 0.6$ allows to define the Lyapunov exponent λ_∞ and to classify the chaoticity of the mean field at $T=3$ MeV. At variance for $\rho/\rho_0=0.7$ $\lambda(t)$ is a decreasing function of t and the mean field has a regular linear evolution. (b) The Lyapunov exponent λ_∞ is displayed for different values of the initial density. See text.